

“Trimethylguanosine nucleoside inhibits cross-linking between Snurportin 1 and m3G-capped U1 snRNA” Bahia, D., Aviñó, A., Eritja, R., Darzynkiewicz, E., Bach-Elias, M. Nucleosides, Nucleotides & Nucleic Acids, 25(8), 909-923 (2006). doi: 10.1080/15257770600793901.

---

## Trimethylguanosine nucleoside inhibits cross-linking between Snurportin 1 and m3G-capped U1 snRNA

Diana Bahia<sup>1,§</sup>, Anna Aviñó<sup>2</sup>, Ramon Eritja<sup>2</sup>, Edward Darzynkiewicz<sup>3</sup>, and Montserrat Bach-Elias<sup>1\*</sup>

<sup>1</sup>IIBB-CSIC (Instituto de Investigaciones Biomédicas de Barcelona-Consejo Superior de Investigaciones Científicas), c/Jorge Girona Salgado 18-26, 08034 Barcelona, Spain

<sup>2</sup> Institut de Biología Molecular de Barcelona, CSIC (IBMB-CSIC), Jordi Girona 18-26, 08034 Barcelona

<sup>3</sup> Institute of Experimental Physics, Department of Biophysics, Warsaw University, Warsaw, Poland).

Correspondence to: Dr. Montse Bach-Elias, IIBB-CSIC, c/ Jorge Girona Salgado 18-26, 08034 Barcelona, Spain.

Tel: +34-93-4006134;

Fax: +34-93-2045904;

E-mail: [mbebmc@cid.csic.es](mailto:mbebmc@cid.csic.es)

<sup>§</sup> Present address: Departamento de Genética, ESALQ, USP, Av. Pádua Dias, 11, 13400-970, Piracicaba, SP, Brazil.

### Abstract

Macromolecular nuclear import is an energy and signal-dependent process. The best characterised type of nuclear import consists of proteins carrying the classical NLS that is mediated by the heterodimeric receptor importin  $\alpha/\beta$ . Spliceosomal snRNPs U1, U2, U4 and U5 nuclear import depend both on the 5' terminal m3G (trimethylguanosine) cap structure of the U snRNA and the Sm core domain. Snurportin 1 recognises the m3G-cap structure of m3G-capped U snRNPs. In this report, we show how a synthesised trimethylguanosine nucleoside affects the binding of Snurportin 1 to m3G-capped U1 snRNA in a UV-cross-linking assay. The data indicated that TMG nucleoside is an essential component required in the recognition by Snurportin 1; thus, suggesting that interaction of Snurportin 1 with U1 snRNA is not strictly dependent on the presence of whole cap structure, but rather on the presence of TMG nucleoside structure. These results indicate that the free nucleoside TMG could be a candidate to be an inhibitor of the interaction between Snurportin and U snRNAs. We also show the behaviour of free TMG nucleoside in *in vitro* U snRNPs nuclear import

**KEYWORDS:** Trimethylguanosine nucleoside, TMG, m<sub>3</sub>GpppG cap, Snurportin 1, nuclear import, UsnRNPs, UV-cross-linking assay.  
Abbreviations used: TMG, trimethylguanosine

## **INTRODUCTION**

The selective transport of proteins and ribonucleoproteins between the nucleus and the cytoplasm of eukaryotic cells is an important activity that integrates the functions of protein synthesis, RNA genesis and DNA replication. Both protein and nucleoprotein complexes are actively transported through the nuclear pore complex (NPC), embedded in the nuclear envelope, a process characterised by energy and signal dependence, thus indicating that it is mediated by carriers (reviewed in Gorlich and Mattaj (1)).

A major breakthrough in the study of nuclear import was the development of effective *in vitro* systems (2). Macromolecular import has been well established for proteins that carry a short basic nuclear localisation signal (NLS). In a first step, the NLS is recognised by a soluble heterodimeric receptor known as the importin  $\alpha/\beta$  heterodimer. The importin  $\alpha$  subunit (60 kDa) functions as an anchor, termed as adapter between NLS peptide and importin  $\beta$ : importin  $\alpha$  binds to the NLS protein; at the same time importin  $\alpha$  binds to the IBB domain of importin  $\beta$  (97 kDa). The NLS-receptor complex is then targeted to the NPC via importin  $\beta$ . This complex is subsequently translocated through the NPC, by an energy-dependent mechanism, with the help of Ran-GTPase and NTF2.

A second class of NLS is the M9 domain, which is present in a heterogeneous-nuclear hnRNP A1. The M9 domain is enriched in glycine and aromatic residues and can bind directly to its receptor transportin, thus, in this protein import pathway there is no requirement of an adapter (importin  $\alpha$ -like) for the binding (3). Other import signals, like the classical NLS, have been identified. They are composed mainly of basic amino acids and were found in ribosome proteins and histone H1 (4; 5). In addition, the U1A and U2B'' import does not follow the NLS model and requires a new mediator of nuclear protein import (6).

A third class of import signal was identified in U snRNPs (U small nuclear ribonucleoproteins). The U snRNPs are essential components of the splicing machinery assembled in a complex sequence of events in both the nuclear and the cytoplasmic compartments (see reviews in Will and Luhrmann (7; 8)). The trimethylguanosine (m<sub>3</sub>G) cap is the hallmark of the U snRNAs (U small nuclear RNAs family) (9). Major U snRNPs U1, U2, U4/U6.U5 are essential spliceosomal factors and they are complexes of U snRNAs and a very characteristic set of common and specific polypeptides (10). The common polypeptides (B', B, D1, D2, D3, E, F and G) form a common structure present in all the spliceosomal U snRNPs, the so-called Sm core. This event requires the survival of motor neuron (SMN) protein complex's activity (8;11-13). With the sole exception of U6 snRNA, which is capped with MeGTP (14), all the major U snRNAs contain the m<sub>3</sub>G cap in human cells. Shortly after polymerase II transcription, UsnRNAs are capped with a 7-methylguanosine (m<sup>7</sup>G) structure (15). The newly transcribed UsnRNAs are exported to the cytoplasm by binding to an adaptor protein called PHAX (16). The stable

formation of Sm core in the cytoplasm promotes the hypermethylation of the m<sup>7</sup>G cap to a 2,2,7-trimethylguanosine-cap (m<sub>3</sub>G) (17). Both the Sm binding site domain and the m<sub>3</sub>G cap constitute the bipartite nuclear localization signal that targets the U snRNPs to the nucleus (18;19).

Snurportin 1, a 45 kDa protein, mediates nuclear import of m<sub>3</sub>G-capped uridine-rich small nuclear RNPs (U snRNPs) (20). Snurportin is thus an adaptor that recognises the m<sub>3</sub>G-cap structure and links UsnRNPs to importin β for subsequent transport to the nucleus (20;21). Once in the nucleus, Snurportin 1 returns to the cytoplasm after each round of import. The re-export of Snurportin 1 is mediated by CRM1 (22). The import of U snRNPs does not need importin α (23). Furthermore, the depletion of importin α stimulates the nuclear import of U snRNPs, indicating competition between U snRNP nuclear import factors and importin α (23)

Previously, our group has synthesised high amounts of non-commercial TMG (trimethylguanosine) nucleoside (24) by using a modified synthetic procedure described for nucleosides conjugates (25). Moreover, we showed that the synthesised TMG product is immunologically identical to the m<sub>3</sub>G cap of the UsnRNPs. These experiments used antibodies raised against TMG-KLH conjugates. Anti-TMG antibodies precipitated the U snRNAs and this pull-down could be competed with free TMG (24). Here, we show how TMG nucleoside affects the binding of Snurportin to m<sub>3</sub>G-capped U1 snRNA in a new UV-cross-linking assay created based on previous procedures (20;26). We next attempted to test the effect caused by the addition of TMG nucleoside in an in vitro UsnRNPs import using Hela digitonin permeabilised cells. Surprisingly, TMG not only did not block U snRNPs import, but in fact stimulated it, when added in concentrations lower than 0.5 mM. When 1 and 2 mM of free TMG were added, no U snRNPs import stimulation was observed. We discuss the aspects involved in the behaviour of free TMG in nuclear import and cross-linking assays.

## RESULTS AND DISCUSSION

### **Snurportin1 binds only to m<sub>3</sub>G cap in the new cross-linking assay**

The purified his-tagged Snurportin 1 has an apparent molecular weight of 47 kDa (Fig. 1, lane 1) in 15% SDS-PAGE gel. In this gel, we could observe that Snurportin 1 has been well expressed and purified through nickel-NTA agarose and was suitable for using in further experiments. We have created a new UV-cross-linking assay based on previous publications (20;26) and tested its efficiency in the binding of Snurportin 1 to m<sup>7</sup>G and m<sub>3</sub>G caps. U1 snRNA was transcribed using radioactive label and capped with m<sup>7</sup>G and m<sub>3</sub>G nucleotides and was then mixed with recombinant Snurportin 1.

After UV irradiation and RNase 1 treatment, the samples were subjected to SDS-PAGE and the protein was visualised by autoradiography. Snurportin 1 clearly bound to the m<sub>3</sub>G cap as indicated by the signal observed (Fig. 2, lane 2); however, Snurportin 1 did not bind to m<sup>7</sup>G cap (Fig. 2, lane 1). Thereby, indicating the efficiency of our optimised assay. Furthermore, we tested whether the nucleoside synthesised here, i.e. the TMG

nucleoside, is able to compete with the binding of Snurportin 1 to the m<sub>3</sub>GpppG-capped U1 snRNA, as free m<sub>3</sub>GpppG cap does.

### **Trimethylguanosine nucleoside (TMG) inhibits cross-linking between Snurportin 1 and m<sub>3</sub>G-capped U1 snRNA**

Free m<sub>3</sub>GpppG cap can compete with the binding of Snurportin 1 to m<sub>3</sub>GpppG-capped U1 snRNA. In Fig 3A,B, we observe the competition with 1.0 (Fig 3A, lane 2), 0.1 (Fig 3B, lane 2) and 0.01 mM (Fig 3B, lane 3) of m<sub>3</sub>GpppG cap. By comparing the disappearance of Snurportin 1 cross-linked bands (3A, lane 2; 3B lane 2) with the cross-linked snurportin band (3A and B, lane 1) we could conclude that 1 and 0.1 mM m<sub>3</sub>GpppG cap produced a total inhibition of binding. These data verify that the binding of Snurportin 1 to m<sub>3</sub>GpppG cap, in the cross-linking assay as created here, has the same specificity of m<sub>3</sub>G cap oligo (m<sub>3</sub>GpppAmpUmpA) observed previously (20). TMG was synthesised from m<sup>2,2</sup>G (27) by methylation of position 7 of the ring with dimethyl sulphate (24). Surprisingly, free TMG nucleoside was also able to inhibit the binding between Snurportin 1 and m<sub>3</sub>G cap in UV-cross-linking assays when used in 1 and 0.1 mM as competitor (Fig 3, lanes 4 and 5), although it could be anticipated from the crystal structure of 2,2,7-trimethyl-GpppG bound to snurportin 1 (cap-binding fragment) published very recently (28). In the present assay, only in the presence of 1 mM TMG a total inhibition of binding was obtained. In fact, TMG nucleoside has a decreased strength of interaction with snurportin 1 of approximately one order of magnitude.

This effect may be due to the presence of whole nucleotide structure of m<sub>3</sub>GpppG. We also observed that m<sup>2,2</sup>G (2,2-dimethylguanosine) nucleoside was unable to inhibit the UV-cross-linking between Snurportin 1 and m<sub>3</sub>GpppG-capped U1 snRNA (not shown), confirming that methylation in position 7 of the guanosine ring is essential in this respect (20;24). Therefore, these data strongly indicate that TMG nucleoside is an essential component required in the recognition by Snurportin 1; thus, suggesting that interaction of Snurportin 1 with U1 snRNA is not strictly dependent on the presence of whole cap structure, but rather on the presence of TMG nucleoside structure. However, whether artificial constructs made with TMG nucleoside can exhibit the same binding avidity with native Snurportin 1 is still an open question. Moreover, the next step will be to characterise the binding properties of TMG nucleoside in in vitro nuclear import assays.

### **U snRNP import is stimulated by trimethylguanosine nucleoside (TMG) in concentrations under 0.5 mM.**

It is known that U snRNP import is effectively blocked by trimethylguanosine (m<sub>3</sub>G) cap analog m<sup>2,2,7</sup>GTP but it is not affected by the monomethylguanosine cap analog m<sup>7</sup>GTP (23). Both effects may be seen in Figure 4 B and C-D, respectively. The nuclear fluorescence seen in Fig 4A, due to nuclear import of fluorescent U snRNPs, has decreased in B, by adding cap m<sub>3</sub>GpppG. On the other hand, the panels C and D shows that nuclear fluorescence is the same of control A, thus demonstrating that the adding of m<sup>7</sup>GpppG cap has not affected the import rate of fluorescent U snRNPs into the nucleus.

In this direction we tested if our compound, the free TMG, could alter an in vitro U snRNPs import. So, we verified that at concentrations higher than 0.5 mM there is U snRNP import inhibition, mimicking the effect observed in the presence of m<sub>3</sub>GpppG cap analog, but surprisingly, under this concentration we observed a stimulation of U snRNPs import.

Figure 4 also shows an U snRNPs in vitro import assay in the presence of free TMG. Surprisingly, TMG not only did not block U snRNPs import, but in fact stimulated it (Figure 4 compare F with control E) when used with the molarity of 0.5 mM. Nevertheless, when 1 and 2 mM of free TMG were added (Figure 4, G and H, respectively) no U snRNPs import stimulation was observed. In conclusion, when TMG is added in low concentrations (under 0.5 mM, not shown), it may stimulates U snRNPs nuclear import. These experiments have been repeated several times, including various TMG concentrations till 0.5 mM, and both the stimulation and inhibitory effects observed were the same.

### **U snRNP import is blocked by trimethylguanosine nucleoside (TMG) in concentrations of 1 mM.**

A fluorescent BSA conjugated with a peptide containing the reverse NLS sequence (“SLN”) was also used to check the veracity of the effect caused by free TMG in a U snRNPs nuclear import assay. As the SLN signal is not recognised by the protein import pathway, it was used to discard the possibility of a passive diffusion of the conjugated. When TMG is used at a final concentration of 2 mM it seems to be toxic to the cells as shown when it was assayed with BSA-SLN, leading to a passive diffusion of fluorescent conjugate into the nucleus (Fig 5, D). This effect was monitored by fluorescence in the nucleus (Fig 5, D compared with the control A). BSA-SLN in in vitro import assays either alone (Fig 5 A) or with the addition of 1 mM (Fig 5, panel C) or 0.5 mM (Fig 5, panel B) of TMG nucleoside did not lead to increased nuclear fluorescence. A slight fluorescence can be seen around the nuclear membrane. These experiments confirm that both the stimulation of import of U snRNPs in the presence of 0.5 mM TMG free nucleoside, and the blocking of U snRNPs import in the presence of 1 mM TMG is due to an active import of the karyophile.

TMG has an unstable imidazole ring, which is a property of this product, and the imidazole degradation does not depend on the product's synthesis procedure. The degradation of the imidazole ring is due to two different reactions: depurination, a reaction catalyzed by acids; and the opening of the imidazole ring, by susceptibility to nucleophilic attack in basic pH. Both reactions are favored by the methylation at position 7. Some amount of the products obtained from the imidazole degradation may be present in our preparations; they could be toxic to the nuclear membrane and the stimulation effect could be simply due to a passive diffusion through broken nuclear membranes. The results observed in the presence of 2 mM TMG indicate that the imidazole degradation is affecting the nuclear membrane, thus leading to a passive nuclear diffusion of BSA-SLN. Nevertheless, the results observed in the presence of 0.5 and 1 mM TMG compared with the negative control BSA-SLN indicated that the minor imidazole degradation products

was not affecting the membrane at the concentration range, thus BSA-SLN is not obtained in the nucleus by passive diffusion in presence of several concentrations of TMG nucleoside. The inhibition effect indicates that TMG at concentrations higher than 0.5 mM really mimics the U snRNPs cap and inhibits the active transport. The stimulation effect under 0.5 mM is so far not easily explained; we can suggest that TMG would have some effect on the molecular attraction and concentration of other factors, including those involved in import, a role already suggested for other protein-protein interaction systems (29-33).

Future experiments will undoubtedly help to elucidate the roles of TMG nucleoside in the molecular mechanisms behind these processes.

## **Materials and Methods**

The  $m^7G$ -cap analogue was purchased from Biolabs.  $m_3GpppG$  was synthesised as described elsewhere (34;35), Radio-labelled nucleotide triphosphates were obtained from Amersham. The RNase inhibitor was obtained from Gibco and the SP6 polymerase used for the pHU1 transcription was from Boehringer. Recombinant Snurportin 1 was a kind gift from Dr Martin Hetzer. All nuclear import procedures can also be found in Dingwall and Palacios (36).

### **Recombinat Snurportin expression and purification**

Snurportin 1 was cloned into the BamHI-XmaI sites of pQE30 (Qiagen), expressed with an  $NH_2$ -terminal His tag and purified on nickel-NTA agarose followed by dialysis in 20 mM Hepes-KOH, pH 7.5, 200 mM NaCl, 2 mM magnesium acetate, 250 mM sucrose.

### **TMG nucleoside synthesis**

TMG nucleoside synthesis was performed as follows: the TMG nucleoside precursor was  $m^{2,2}G$ , which was synthesised as described previously (27). The synthesis of TMG nucleoside used in the cross-linking assays was performed essentially as described previously (24).

### **UV-cross-linking assay**

The following technique was created based on previous citations (1;22). pHU1A (37) was transcribed with  $m_3G$  and  $m^7G$  caps in a transcription assay where  $\alpha^{32}UTP$  (and no cold UTP) was used. The transcripts were purified through centri spin 10 (Princeton) columns equilibrated with Roeder 200 mM KCl (10 mM HEPES pH 7.9; 1.5 mM  $MgCl_2$ ; 200 mM KCl; 500 mM DTT; 500 mM PMSF; 5% glycerol) buffer. To visualise the binding between Snurportin 1 and  $m_3G$  by UV-cross-linking, 1 pmol of newly transcribed  $m_3GpppG$ -capped U1 snRNA ( $250-350 \times 10^4$  counts per min) was incubated for 15 min on ice with 1.5  $\mu g$  of recombinant Snurportin 1 in a total volume of 10  $\mu l$ . Reaction mixtures were irradiated at 254 nm with a Sylvania G8T5 germicidal UV lamp for 5 min at a distance of 2 cm, and then placed on ice for 2 min. To each reaction, 1  $\mu l$  RNase A (3 mg/ml) was added and the samples were incubated for 30 min at 37°C. In competition assays, recombinant Snurportin 1 was mixed with 0.1 and 1 mM  $m_3G$  cap or TMG nucleoside on ice for 30 min prior to UV-irradiation. Cross-linked proteins were

separated on 12.5 or 15% SDS-PAGE gels and visualised by autoradiography (Film KODAK BioMax MS in an intensify Kodak Biomax MS screen).

### **Preparation of fluorescent karyophiles**

U snRNPs were purified from HeLa nuclear extracts by using an immunoaffinity column anti-trimethylguanosine (anti-m<sub>3</sub>G) as described by (38). U snRNPs were concentrated to 1 mg/ml with Filtron 30K (Pall Filtron). In the last step of concentration U snRNPs were resuspended in 100 mM borate buffer, pH 7.5. FLUOS (Boehringer) was prepared fresh in DMSO (dimethylsulfoxid) and added to U snRNPs in 100-fold molar excess. The reaction proceeded for 45 min at room temperature with rotation in the darkness, and was stopped by adding Tris to 10 mM, pH 7.5. The labeled U snRNPs were separated from the free dye by loading the sample in a Bioget p60 column previously equilibrated with 1 X transport buffer-TB-(20 mM Hepes-KOH, pH 7.3; 110 mM KOAc; 5 mM NaAc; 2 mM MgAc; 1.0 mM EGTA; 1 mM DTT; 0.1 mM PMSF; 1 µg/ml aprotinin; 1 µg/ml pepstatin A, 1 µg/ml leupeptin, 5% v/v sucrose). Fluorescent U snRNPs were repurified again in an anti-m<sub>3</sub>G column and eluted using 15 mM 7-methylguanosine in TB.

BSA was cross-linked to a non-functional peptide with the reverse NLS sequence (SLN) (15-20 peptides per molecule), as described for BSA-NLS (39). Conjugation to the carrier protein was obtained through the *N*-terminal cysteine residue and the glycine residues serve as a flexible hinge or spacer to separate the peptide from the protein itself. BSA-NLS conjugate was also labelled with FLUOS following the manufacturer's instructions.

### **Preparation of permeabilised cells**

The cells were cultivated in a Dulbecco's modified Eagles's medium (Gibco), supplemented with 10% fetal calf serum and penicillin/streptomycin (Gibco-BRL) at 37°C, 5% CO<sub>2</sub>, and transferred to 10 plastic flasks (125 m<sup>2</sup>). They were grown to approximately 70% confluence and harvested using 5 ml of a non-enzymatic cell dissociation solution (Sigma) for 10 min at 37°C. The cells were detached from the walls and then transferred to a 50 ml Falcon tube. The subsequent steps were carried out on ice. The cells were washed 3 times with ice-cold permeabilisation buffer-PB (50 mM HEPES-KOH, pH 7.3; 50 mM KOAc; 8 mM MgCl<sub>2</sub>; 2 mM EGTA) and resuspended to a cell density of 1-5 X 10<sup>6</sup> cell/ml in ice-cold PB. Digitonin (Fluka) was then added to 50 µg/ml and the cells were left on ice for exactly 5 min. Afterwards, the cells were washed three times with PB and finally resuspended at a density of 5-10 X 10<sup>6</sup> cell/ml in TB that contained 5% DMSO. The cells were then aliquoted into cold eppendorf tubes and the tubes were frozen slowly in a Styrofoam box for 1 hr at -20°C and then stored at -80°C.

### **Nuclear import in vitro assay**

Both U snRNPs and BSA-SLN in vitro import reactions have always contained 1 µl of a 5 mg/ml nucleoplasmic core (NPC), obtained as described elsewhere (40), 1 µl of a 10 X energy-regenerating system (the final concentration in the assay must be 0.5 mM ATP, 0.5 mM GTP, 10 mM creatine phosphate, 50 µg/ml creatine phosphokinase); 1 µl of digitonin-permeabilised cells (1 X 10<sup>4</sup> cells); TB 1 X (from 5 ml transport buffer plus 5 µl DTT 1 M; 50 µl Trasylol 500000 KIE Bayer; 5 µl pepstatin 1 mg/ml; 5 µl PMSF 1 M;

1  $\mu$ l leupeptin 5 mg/ml; 250 mM sucrose) with or without the factors to be tested. All the purified TMG-binding proteins fractions were previously desalted through micro bio-spin chromatography columns (BioRad bio-gel P-6) in order to change the buffer to TB and remove free TMG nucleoside.

BSA-SLN import was performed in a final volume of 10  $\mu$ l always containing 2  $\mu$ l of *Xenopus* egg extract (41) previously spun at 13.000 rpm for 10 min- and 1  $\mu$ l of a 1/10 dilution of BSA-SLN in TB (at a final concentration of 70  $\mu$ g/ml). First, the BSA-SLN was premixed with the *Xenopus* extract and the factors and incubated on ice for 5 min. Then, a mixture was prepared with the cells and the NPc and then added to the first premix. To start the import reaction the energy mix was added and the reaction proceeded for 10 min, at room temperature. For U snRNPs import, each reaction was performed in 15-17  $\mu$ l and also contained 6  $\mu$ l of importin  $\alpha$ -depleted *Xenopus* extract (described below), 0.5  $\mu$ l of tRNA (to a final concentration of 0.2 mg/ml), and 5  $\mu$ l of fluorescent U snRNPs (to a final concentration of 20  $\mu$ g/ml). Initially, U snRNPs were premixed with *Xenopus* extract and then the factors, and the mixture were left on ice for 15 min. Then a mixture was prepared with tRNA, NPc and digitonin-permeabilised cells which was added to the eppendorfs containing the first premix. In order to start the reaction, the energy mix was added and the reaction was proceeded for 60-90 min at room temperature, in the dark. When free nucleotides and TMG nucleoside were added to the assay they were mixed with U snRNPs and *Xenopus* extract before leaving the mixture on ice.

Both reactions were stopped by fixing with 200  $\mu$ l of 8 % (w/v) paraformaldehyde on ice together with 200  $\mu$ l DAPI dye (330 ng/ $\mu$ l) in TB. The fixed reaction was layered over 700  $\mu$ l of 30 % sucrose in a tube containing a poly(lysine)-coated cover slip (MW 30,000-70,000) at 1 mg/ml in water. After centrifugation (3.000 rpm, 15 min.) the cover slip was recovered, washed with PBS and mounted on top of a drop of Vectashield mounting medium (Vector Laboratories, Burlingame CA) and viewed directly. Images acquisition and the quantification of nuclear fluorescence were performed as described (36;39).

### **Xenopus extracts**

The *Xenopus* extract (41) used in U snRNPs import was importin  $\alpha$  depleted by passing through an immunoaffinity column with an antibody against an *N*-terminal peptide from *Xenopus* importin  $\alpha$  as described (23).

### **Acknowledgments**

We thank Drs. Iain Mattaj and Martin Hetzer for technical help and discussion. We are indebted to Dr. Renato A. Mortara for his help with confocal images. We are very grateful to Dr. Martin Hetzer for providing recombinant Snurportin 1. This work was supported by Plan Nacional BFU2005-00701 and the Polish Committee for Scientific Research (KBN) # 6 P04A 055 17. D.B. was a recipient of a CNPq Brazilian fellowship, and EMBO and FEBS short-term fellowships.


### **REFERENCES**

1. Gorlich,D. and Mattaj,I.W. (1996) Nucleocytoplasmic transport. *Science*, 271, 1513-1518.
2. Adam,S.A., Marr,R.S. and Gerace,L. (1990) Nuclear protein import in permeabilized mammalian cells requires soluble cytoplasmic factors. *J.Cell Biol.*, 111, 807-816.
3. Pollard,V.W., Michael,W.M., Nakielny,S., Siomi,M.C., Wang,F. and Dreyfuss,G. (1996) A novel receptor-mediated nuclear protein import pathway. *Cell*, 86, 985-994.
4. Jakel,S. and Gorlich,D. (1998) Importin beta, transportin, RanBP5 and RanBP7 mediate nuclear import of ribosomal proteins in mammalian cells. *EMBO J.*, 17, 4491-4502.
5. Jakel,S., Albig,W., Kutay,U., Bischoff,F.R., Schwamborn,K., Doenecke,D. and Gorlich,D. (1999) The importin beta/importin 7 heterodimer is a functional nuclear import receptor for histone H1. *EMBO J.*, 18, 2411-2423.
6. Hetzer,M. and Mattaj,I.W. (2000) An ATP-dependent, Ran-independent mechanism for nuclear import of the U1A and U2B" spliceosome proteins. *J.Cell Biol.*, 148, 293-303.
7. Will,C.L. and Luhrmann,R. (1997) Protein functions in pre-mRNA splicing. *Curr.Opin.Cell Biol.*, 9, 320-328.
8. Will,C.L. and Luhrmann,R. (2001) Spliceosomal UsnRNP biogenesis, structure and function. *Curr.Opin.Cell Biol.*, 13, 290-301.
9. Birnstiel,M.L.ed. Structure and Function of major and minor small ribonucleoprotein particles. 1988. Berlin, Springer-Verlag Press.
10. Luhrmann,R., Kastner,B. and Bach,M. (1990) Structure of spliceosomal snRNPs and their role in pre-mRNA splicing. *Biochim.Biophys.Acta*, 1087, 265-292.
11. Paushkin,S., Gubitz,A.K., Massenet,S. and Dreyfuss,G. (2002) The SMN complex, an assemblyosome of ribonucleoproteins. *Curr.Opin.Cell Biol.*, 14, 305-312.
12. Narayanan,U., Ospina,J.K., Frey,M.R., Hebert,M.D. and Matera,A.G. (2002) SMN, the spinal muscular atrophy protein, forms a pre-import snRNP complex with snurportin1 and importin beta. *Hum.Mol.Genet.*, 11, 1785-1795.
13. Massenet,S., Pellizzoni,L., Paushkin,S., Mattaj,I.W. and Dreyfuss,G. (2002) The SMN complex is associated with snRNPs throughout their cytoplasmic assembly pathway. *Mol.Cell Biol.*, 22, 6533-6541.
14. Vankan,P., McGuigan,C. and Mattaj,I.W. (1990) Domains of U4 and U6 snRNAs required for snRNP assembly and splicing complementation in *Xenopus* oocytes. *EMBO J.*, 9, 3397-3404.
15. Hamm,J. and Mattaj,I.W. (1990) Monomethylated cap structures facilitate RNA export from the nucleus. *Cell*, 63, 109-118.
16. Ohno,M., Segref,A., Bachi,A., Wilm,M. and Mattaj,I.W. (2000) PHAX, a mediator of U snRNA nuclear export whose activity is regulated by phosphorylation. *Cell*, 101, 187-198.
17. Mattaj,I.W. (1986) Cap trimethylation of U snRNA is cytoplasmic and dependent on U snRNP protein binding. *Cell*, 46, 905-911.
18. Fischer,U. and Luhrmann,R. (1990) An essential signaling role for the m3G cap in the transport of U1 snRNP to the nucleus. *Science*, 249, 786-790.

19. Hamm,J., Darzynkiewicz,E., Tahara,S.M. and Mattaj,I.W. (1990) The trimethylguanosine cap structure of U1 snRNA is a component of a bipartite nuclear targeting signal. *Cell*, 62, 569-577.
20. Huber,J., Cronshagen,U., Kadokura,M., Marshallsay,C., Wada,T., Sekine,M. and Luhrmann,R. (1998) Snurportin1, an m3G-cap-specific nuclear import receptor with a novel domain structure. *EMBO J.*, 17, 4114-4126.
21. Huber,J., Dickmanns,A. and Luhrmann,R. (2002) The importin-beta binding domain of snurportin1 is responsible for the Ran- and energy-independent nuclear import of spliceosomal U snRNPs in vitro. *J.Cell Biol.*, 156, 467-479.
22. Paraskeva,E., Izaurrealde,E., Bischoff,F.R., Huber,J., Kutay,U., Hartmann,E., Luhrmann,R. and Gorlich,D. (1999) CRM1-mediated recycling of snurportin 1 to the cytoplasm. *J.Cell Biol.*, 145, 255-264.
23. Palacios,I., Hetzer,M., Adam,S.A. and Mattaj,I.W. (1997) Nuclear import of U snRNPs requires importin beta. *EMBO J.*, 16, 6783-6792.
24. Espuny,R., Bahia,D., Barretto Cicarelli,R.M., Codony,C., Khaouja,A., Avino,A.M., Eritja,R. and Bach-Elias,M. (1999) Preparation of N<sup>2</sup>, N<sup>2</sup>,7-trimethylguanosine affinity columns. *Nucleosides Nucleotides*, 18, 125-136.
25. Erlanger,B.F. and Beiser,S.M. (1964) Antibodies specific for ribonucleosides and ribonucleotides and their reaction with dna. *Proc.Natl.Acad.Sci.U.S.A.*, 52, 68-74.
26. Woppmann,A., Rinke,J. and Luhrmann,R. (1988) Direct cross-linking of snRNP proteins F and 70K to snRNAs by ultra-violet radiation in situ. *Nucleic Acids Res.*, 16, 10985-11004.
27. Avino,A.M., Mayordomo,A., Espuny,R., Bach,M. and Eritja,R. (1995) A Convenient Method for the Preparation of N-2,N-2-Dimethylguanosine. *Nucleosides & Nucleotides*, 14, 1613-1617.
28. Strasser,A., Dickmanns,A., Luhrmann,R. and Ficner,R. (2005) Structural basis for m3G-cap-mediated nuclear import of spliceosomal UsnRNPs by snurportin1. *EMBO J.*, 24, 2235-2243.
29. Jones,S., Daley,D.T., Luscombe,N.M., Berman,H.M. and Thornton,J.M. (2001) Protein-RNA interactions: a structural analysis. *Nucleic Acids Res.*, 29, 943-954.
30. Jones,S., Marin,A. and Thornton,J.M. (2000) Protein domain interfaces: characterization and comparison with oligomeric protein interfaces. *Protein Eng*, 13, 77-82.
31. Jones,S. and Thornton,J.M. (1997) Prediction of protein-protein interaction sites using patch analysis. *J.Mol.Biol.*, 272, 133-143.
32. Jones,S. and Thornton,J.M. (1997) Analysis of protein-protein interaction sites using surface patches. *J.Mol.Biol.*, 272, 121-132.
33. Jones,S. and Thornton,J.M. (1996) Principles of protein-protein interactions. *Proc.Natl.Acad.Sci.U.S.A.*, 93, 13-20.
34. Izaurrealde,E., Lewis,J., McGuigan,C., Jankowska,M., Darzynkiewicz,E. and Mattaj,I.W. (1994) A nuclear cap binding protein complex involved in pre-mRNA splicing. *Cell*, 78, 657-668.
35. Iwase,R., Sekine,M., Tokumoto,Y., Ohshima,Y. and Hata,T. (1989) Synthesis of N<sup>2</sup>, N<sup>2</sup>, 7-trimethylguanosine cap derivatives. *Nucleic Acids Res.*, 17, 8979-8989.

36. Dingwall,C. and Palacios,I. (1998) In vitro systems for the reconstitution of snRNP and protein nuclear import. *Methods Cell Biol.*, 53, 517-543.
37. Patton,J.R., Patterson,R.J. and Pederson,T. (1987) Reconstitution of the U1 small nuclear ribonucleoprotein particle. *Mol.Cell Biol.*, 7, 4030-4037.
38. Bach,M., Bringmann,P. and Luhrmann,R. (1990) Purification of small nuclear ribonucleoprotein particles with antibodies against modified nucleosides of small nuclear RNAs. *Methods Enzymol.*, 181, 232-257.
39. Palacios,I., Weis,K., Klebe,C., Mattaj,I.W. and Dingwall,C. (1996) RAN/TC4 mutants identify a common requirement for snRNP and protein import into the nucleus. *J.Cell Biol.*, 133, 485-494.
40. Gorlich,D., Prehn,S., Laskey,R.A. and Hartmann,E. (1994) Isolation of a protein that is essential for the first step of nuclear protein import. *Cell*, 79, 767-778.
41. Newmeyer,D.D., Lucocq,J.M., Burglin,T.R. and De Robertis,E.M. (1986) Assembly in vitro of nuclei active in nuclear protein transport: ATP is required for nucleoplasmin accumulation. *EMBO J.*, 5, 501-510.


FIGURES



**Fig. 1**

**Fig 1.** Expression and purification of recombinant Snurportin 1. Snurportin 1 was expressed in pQE30 (Qiagen) as his-tagged recombinant protein and purified through nickel-NTA agarose. Snurportin 1 was submitted to 15% SDS-PAGE gel in order to check its integrity and purity. 5  $\mu$ g of purified recombinant Snurportin 1 (Lane 1); protein molecular weight markers (Lane 2). The numbers on the right indicate the molecular weights in kDa. The gel was stained with Coomassie.


## Cross-linked to Snurportin 1



**Fig. 2**


**Fig 2.** UV cross-linking assay. U1 snRNA was transcribed with both m<sup>7</sup>GpppG and m<sub>3</sub>GpppG caps in an assay with only <sup>32</sup>UTP (no cold UTP). To the cross-linking assay, 1.5 µg of Snurportin 1 was incubated with radio-labelled U1 snRNAs on ice in a total volume of 10 µl for 15 min. Both mixtures were irradiated with 254 nm UV Sylvania G8T5 for 5 min at a distance of 2 cm. The cross-linked Snurportin 1 was analysed on a 15% SDS-PAGE gel. Snurportin 1 cross-linked to m<sup>7</sup>GpppG-U1 snRNA (Lane 1) and to m<sub>3</sub>GpppG-U1 snRNA (Lane 2).

## **m<sub>3</sub>GpppG-U1snRNA cross-linked to Snurportin 1 and competed with**




**Fig. 3**

**Fig.3.** Snurportin1 binding to m<sub>3</sub>GpppG cap is inhibited in the presence of free TMG nucleoside. U1 snRNA was transcribed with m<sub>3</sub>GpppG cap, in an assay without UTP cold. To the cross-linking assay, 1.5 µg of Snurportin 1 was incubated with radio-labelled U1 snRNAs on ice in a total volume of 10 µl for 15 min. Both mixtures were irradiated with 254 nm UV Sylvania G8T5 for 5 min at a distance of 2 cm. In competition assays, Snurportin1 was mixed for 30 min on ice with either m<sub>3</sub>GpppG cap or free TMG nucleoside prior to UV irradiation. The cross-linked Snurportin 1 was analysed on a 12% SDS-PAGE gel. Fig 3A. Snurportin 1 cross-linked to m<sub>3</sub>GpppG-U1snRNA (Lane 1, see arrow) competed with 1 mM m<sub>3</sub>GpppG cap (Lane 2). Fig 3B. Snurportin 1 cross-linked to m<sub>3</sub>GpppG-U1snRNA (Lane 1) competed with 0.1 (Lane 2) and 0.01 (Lane 3) mM m<sub>3</sub>GpppG cap, or with 1 (Lane 4) and 0.1 (Lane 5) mM TMG nucleoside.



**Figure 4.** U snRNPs import in vitro assay in the presence of TMG nucleoside. Digitonin-permeabilised HeLa cells were incubated with purified and fluorescently labelled U snRNPs for 90 min, at room temperature, in the presence of *Xenopus* egg extracts importin  $\alpha$  depleted, nucleoplasmin core (NPC), GTP, ATP and an ATP regeneration system (see Material and Methods). A and E) U snRNPs import reaction control in the absence of free nucleosides or nucleotides; B) U snRNPs import in the presence of 2 mM  $m_3$ GpppG cap; C-D) U snRNPs import in the presence of 2 mM  $m_7$ GpppG; F-H) U snRNPs import in the presence of 0.5, 1 and 2 mM of TMG nucleoside, respectively. Magnification bars represent 20 micrometers. ABCD and EFGH represent different assays.



**Figure 5.** BSA-SLN import in vitro assay in the presence of TMG nucleoside. Digitonin-permeabilised HeLa cells were incubated with fluorescently labelled BSA-SLN for 10 min, at room temperature, in the presence of *Xenopus* egg extracts, nucleoplasmin core (NPc), GTP, ATP and an ATP regeneration system (see Material and Methods). A) BSA-SLN import in the absence of TMG (control); B-D) BSA-SLN Import in the presence 0.5, 1 and 2 mM of free TMG, respectively. Magnification bars for A and B represent 10 micrometers, for C and D, 20 micrometers.