

Astronomical radio reception techniques to detect the emission of molecular and short lived species in a cold plasma/gas chamber

J. Cernicharo¹, I. Tanarro², B. Aleman¹, V. Herrero², P. de Vicente³,
J.D. Gallego³,

¹ *ICMM, CSIC, C/Sor Juana Ines de la Cruz 3, 28049 Cantoblanco, Madrid (Spain)*

² *IEM, CSIC, C/Serrano 123, 28006 Madrid (Spain)*

³ *Observatorio de Yebes, IGN, Guadalajara (Spain)*

In this work we prove the viability of the use of standard radioastronomy receivers to detect rotational spectral lines emitted at room temperatures by different molecular precursors and plasma products introduced or generated in a low pressure plasma chamber, placed in the beam path of the 40 m Yebes observatory radio telescope operating in the Q band (41-49 GHz). Right and left circular polarization signals were sampled over 500 MHz bandwidth. The data acquisition was performed with a Fast Fourier Transform Spectrometer with a spectral resolution of 30 kHz. Depending on the weather conditions, the background for emission measurements came from the antenna pointing towards the zenith (clear blue sky) or from a near blackbody load at liquid N₂ temperature (cloudy or rainy weather), implying 42 K or 70 K temperatures, respectively, at 45 GHz.

The plasma was produced in a cylindrical, 25 cm diameter, 42 cm length, stainless steel vacuum chamber by an inductively coupled (5 - 60 W, 13.56 MHz) RF discharge, established through a refrigerated copper coil (4 turns, 10 cm length, 17 cm diameter) inserted axially and located where the gaussian beam of the receivers narrows to its minimum width. Windows of 75 μm thickness × 22 cm diameter made of Uplex were placed at both ends of the chamber at slightly divergent planes to avoid standing waves. A differentially pumped quadrupole mass spectrometer (0-100 amu) allowed to identify the stable species introduced or generated during the on/off plasma intervals and follow their time evolution. Gas pressures of ≈ 0.7-90 Pa produced similar column densities to those of a typical hot core ≈ 10¹⁵⁻¹⁶ cm⁻². A pressure range of ≈ 10-30 Pa at gas flow conditions proved to be more appropriate during plasma operation.

OCS and CH₃OH, which have quite different line intensities, were selected for static gas detection in the observing emission band, displaying maximum equivalent radiation temperatures of ≈ 4 K for carbonyl sulfide and 0.3 K for methanol. In the lower pressure regime, the linewidths were mainly due to thermal broadening and at high pressure they were dominated by pressure broadening. OCS and CS₂ were selected as plasma precursors of the CS radical, which emits also in the 41-49 GHz region and was regularly detected at different plasma conditions, with equivalent temperatures up to 4 K. O₂ discharges applied after sulfur deposition in the reactor walls by the previous S rich containing plasmas allowed to generate SO₂ by heterogeneous plasma processes and detect its rotational transitions in different vibrational (bending) states $v_2 = 0,1,2$; the intensity of the transitions from upper levels increasing with discharge power.

With the plasma ON no spurious signals were induced in the resulting spectra from electromagnetic coupling of the RF discharge with the IF signal generated by the receivers. In conclusion, these experiments confirm the applicability of standard radioastronomy receivers to detect molecular and short lived neutrals in plasma reactors and gas simulation chambers.