

103 mg of (-)-isocomene (**3**) identical with a natural sample. Compound **5** (40 mg) in Et₂O (2 ml) and H₂ atmosphere (room temp. and pres.) with 10% Pd/C afforded 37 mg of **3**.

Acknowledgements—The authors are indebted to Dr. José M^a Hernández Hernández, of our Laboratory, for MS measurements.

REFERENCES

- Pascual Teresa, J. de, Barrero, A. F., Caballero, E. and Medarde, M. (1979) *An. Quim.* **75**, 323.
- Pascual Teresa, J. de, Barrero, A. F. and San Feliciano, A. (1977) *An. Quim.* **73**, 1525.
- Bohlmann, F., Zdero, C. and Suwita, A. (1974) *Chem. Ber.* **107**, 1038.
- Khafagy, S. M., Nazmi Sabri, N. Abd El-Salam, N. A. and Seif El-Din, A. A. (1979) *Planta Med.* **35**, 186.
- Zalkow, L. H., Harris III, R. N., Van Derveer, D. and Bertrand, J. A. (1977) *J. Chem. Soc. D*: 456.
- Zalkow, L. H., Harris III, R. N. and Van Derveer, D. (1978) *J. Chem. Soc. D*: 420.
- Zalkow, L. H., Harris III, R. N. and Burke, N. I. (1979) *J. Nat. Products* **42**, 96.
- Bohlmann, F., Le Van, N., Van Cuong Pham, T., Jakupovic, J., Schuster, A., Zabel, V. and Watson, W. H. (1979) *Phytochemistry* **18**, 1831.
- Bohlmann, F. and Jakupovic, J. (1980) *Phytochemistry* **19**, 259.

Phytochemistry, 1981, Vol. 20, pp. 167–169. © Pergamon Press Ltd. Printed in England.

0031-9422/81/0101-0167 \$02.00/0

CARBON-13 NMR SPECTRA OF SOME *ENT*-ROSANE DITERPENOIDS

MARÍA C. GARCÍA-ÁLVAREZ,* BENJAMÍN RODRÍGUEZ,* SERAFÍN VALVERDE,* BRAULIO M. FRAGA† and ANTONIO G. GONZÁLEZ†

* Instituto de Química Orgánica, CSIC, Juan de la Cierva 3, Madrid-6, Spain and † Instituto de Productos Naturales Orgánicos, CSIC, Tenerife, Spain

(Received 17 March 1980)

Key Word Index—Diterpenoids; *ent*-rosa-5,15-diene; *ent*-8,5-friedo-pimara-5,15-diene derivatives; ¹³C NMR analysis.

Abstract—The ¹³C NMR signals of the parent hydrocarbon *ent*-rosa-5,15-diene (*ent*-8,5-friedo-pimara-5,15-diene) and some of their oxygenated derivatives have been assigned.

INTRODUCTION

¹³C NMR spectroscopy is an excellent tool for the structural elucidation of natural substances. In the diterpene field there is already a considerable amount of information which confirms this view. Podocarpanes [1], labdanes [2, 3], primaranes [4, 5], abietanes [6], *ent*-kauranes [7], *ent*-beyeranes [8, 9], *ent*-atisanes [10–12], *ent*-trachylobanes [13, 14] and *ent*-clerodanes [15–17] have been studied among others.

The availability of a series of *ent*-rosane derivatives and our continued interest [8–11] in the applications of ¹³C NMR spectroscopy has prompted us to carry out this study. Until now, only ¹³C NMR data of rosenonolactone and some of its derivatives [18], jesromotetrol [19], and partial results on rimuene [20] (the epimer at C-13 of compound **1**) have been reported. This paper presents the

¹³C NMR data for the parent hydrocarbon *ent*-rosa-5,15-diene (**1**) and the C-11 and/or C-15 and C-16 oxygenated derivatives (**2–8**), many of which are natural diterpenoids [21, 22].

RESULTS AND DISCUSSION

The assignments for the ¹³C NMR chemical shifts of the diterpenoids (Table 1) were made with the aid of off-resonance decoupled spectra and selective decoupling experiments, and further based on comparison of pairs of compounds, consideration of β, γ and δ substituent effects, general chemical shift arguments and literature data on related structures [4, 5, 18–20].

In general we have detected no unusual effects in this type of hydrocarbon skeleton. The presence of an 11-keto group (**8**), the different stereochemistry of the C-11

hydroxyl group (**6** and **7**) and the absence of any substitution in this carbon atom (**1** and **3**) are clearly reflected in the chemical shifts of C-1, C-8, C-10 and C-20 carbon atoms. On the other hand, a γ -gauche effect caused by the C-15 hydroxyl group is evidenced by the C-14 carbon atom chemical shifts (Table 1).

EXPERIMENTAL

All ^{13}C NMR spectra were obtained on a Varian XL-100 (15) spectrometer equipped with a Nicolet TT-100PFT accessory. The spectra were recorded at the base frequency of 25.2 MHz over a spectral width of 6 kHz with the solvent (commercial NMR grade CDCl_3) providing an internal deuterium lock. A total of 8 K data points were collected in the FID. A pulse width of $10.0\ \mu\text{s}$ was used to aid in the acquisition of the slower relaxing carbons. The total time delay between pulses was 1.3 sec. The proton-decoupled spectra were obtained using a square-wave modulated noise band centred in the proton spectrum. All samples were prepared as 0.3 M CDCl_3 solns in 12 mm sample tubes. The probe temp. in each case was 37°. Chemical shifts are accurate to $\pm 0.05\ \text{ppm}$.

Compounds **3** to **8** have been previously described [21, 22], whilst diterpenes **1** and **2** were obtained from compounds **3** and **4**, respectively, by literature methods [23].




Table 1. ^{13}C NMR chemical shifts (in ppm from TMS) of compounds **1-8**

	1	2	3	4	5	6	7	8
C-1	26.5	30.8	26.5	30.4	30.4	30.8	25.7	28.9
C-2	21.9	22.4	21.9	22.3	22.3	22.4	21.8	21.8
C-3	40.8	40.9	40.7	40.7	40.7	40.9	40.7	40.9
C-4	36.3	36.3	36.5	36.3	36.3	36.3†	35.9	36.2
C-5	145.7	146.6	145.6	146.1	146.2	146.6	145.7	146.2
C-6	116.2	115.6	116.0	115.5	115.5	115.4	115.4	114.4
C-7	30.2*	29.4*	29.3	29.0	29.2	29.4	30.5	28.9
C-8	36.1	36.3	35.6	35.8	35.9	35.7	29.4	37.7†
C-9	34.7†	40.5	34.6	39.3	39.4	40.5	38.7	49.1
C-10	47.3*	47.1*	47.1	46.0	46.1	46.9	38.0	38.5†
C-11	34.3	77.9	33.6	78.3	80.2*	77.6	72.2	214.7
C-12	32.4	38.6	30.2	34.8	34.6	35.5	34.7	43.7*
C-13	35.8†	37.3	35.8	37.3	37.5	36.4†	34.7	40.9
C-14	39.1	42.5	36.2	35.8	35.8	38.7	36.5	35.7
C-15	151.3	149.8	79.2	79.2	80.2	84.0	84.8	82.8
C-16	108.4	108.9	63.0	62.7	62.3	64.4	64.3	64.3
C-17	22.4	23.7	18.5	19.6	19.8	20.0	22.5	20.2
C-18	29.3‡	28.6†	29.1†	28.4†	28.4†	28.5‡	29.6†	28.7‡
C-19	29.7‡	30.0†	29.7†	29.9†	29.9†	30.0‡	29.7†	29.6‡
C-20	12.4*	5.9*	12.3*	7.2*	7.2	5.9	12.8*	12.2
11-OAc	—	—	—	170.1	170.5	—	—	—
	—	—	—	21.6*	21.7*	—	—	—
15-OAc	—	—	170.6‡	170.3‡	—	—	—	—
	—	—	20.9§	20.8§	—	—	—	—
16-OAc	—	—	170.3‡	170.1‡	—	—	—	—
	—	—	20.8§	20.7§	—	—	—	—
—CMe ₂ —	—	—	—	—	—	108.5	108.5	108.8
	—	—	—	—	—	26.2	26.2*	26.2*
	—	—	—	—	—	25.0	25.1*	25.0*

* These assignments have been confirmed by selective decoupling experiments.

†, ‡, § Values in any vertical column may be interchanged, but those given here are considered to be most likely.

ent-Rosa-5,15-diene (1). Thick oil, n_D^{10} 1.5137, $[\alpha]_D^{21} -42.4^\circ$ (c 1.33, CHCl_3). IR $\nu_{\text{max}}^{\text{film}}$ cm^{-1} : 3090, 3050, 1640, 915, 870, 830, 820, 805. $^1\text{H NMR}$ (CDCl_3) δ H-15 and 2H-16 as an ABX system: $\delta_A = 4.93$, $\delta_B = 5.00$, $\delta_X = 5.95$ ($J_{AB} = 2.6$, $J_{AX} = 10.6$, $J_{BX} = 18$ Hz), 5.60 (1H, *m*, $W_{1/2} = 7$ Hz, H-6), C-Me singlets at 1.09 (3H), 1.05 (6H) and 0.69 (3H). (Found: C, 88.27; H, 11.56. $\text{C}_{20}\text{H}_{32}$ requires: C, 88.16; H, 11.84%).

ent-11 β -Hydroxy-rosa-5,15-diene (2). Mp 99–102° (*n*-hexane), $[\alpha]_D^{21} -45.4^\circ$ (c 0.68, CHCl_3). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3290, 3095, 3060, 1640, 1020, 915, 825, 805. $^1\text{H NMR}$ (CDCl_3) δ H-15 and 2H-16 as an ABX system: $\delta_A = 4.91$, $\delta_B = 4.96$, $\delta_X = 5.88$ ($J_{AB} = 2.6$, $J_{AX} = 10.6$, $J_{BX} = 18$ Hz), 5.60 (1H, *m*, $W_{1/2} = 7$ Hz, H-6), 3.77 (1H, *q*, $J_{aa'} = 9.3$ Hz, $J_{ac'} = 6.6$ Hz, H-11), C-Me singlets at 1.07 (6H), 1.05 (3H) and 0.70 (3H). (Found: C, 83.31; H, 11.27. $\text{C}_{20}\text{H}_{32}\text{O}$ requires: C, 83.27; H, 11.18%).

Acknowledgements—We thank Miss M. D. Casado and Miss M. Plaza for recording the ^{13}C and $^1\text{H NMR}$ spectra and Mr. J. Prieto for elemental analyses. This work was supported in part by the Comisión Asesora de Investigación Científica y Técnica.

REFERENCES

1. Wahlberg, I., Almqvist, S. O., Nishida, T. and Enzell, C. (1975) *Acta Chem. Scand. Ser. B* **29**, 1047.
2. Almqvist, S. O., Enzell, C. R. and Wehrli, F. W. (1975) *Acta Chem. Scand. Ser. B* **29**, 695.
3. Buckwalter, B. L., Burfitt, I. R., Nagel, A. A., Wenkert, E. and Näf, F. (1975) *Helv. Chim. Acta* **58**, 1567.
4. Wenkert, E. and Buckwalter, B. L. (1972) *J. Am. Chem. Soc.* **94**, 4367.
5. Beier, R. (1978) *Org. Mag. Reson.* **11**, 586.
6. Nishida, T., Wahlberg, I. and Enzell, C. R. (1977) *Org. Mag. Reson.* **9**, 203.
7. Hanson, J. R., Siverns, M., Piozzi, F. and Savona, G. (1976) *J. Chem. Soc. Perkin Trans. 1*, 114.
8. von Carstenn-Lichterfelde, C., Pascual, C., Pons, J., Rabanal, R. M., Rodríguez, B. and Valverde, S. (1975) *Tetrahedron Letters* 3569.
9. von Carstenn-Lichterfelde, C., Pascual, C., Rabanal, R. M., Rodríguez, B. and Valverde, S. (1977) *Tetrahedron* **33**, 1989.
10. López, A., Márquez, C., Rabanal, R. M. and Valverde, S. (1978) *An. Quím.* **74**, 1100.
11. Rodríguez, B., Alemany, A. and Pinar, M. (1978) *Tetrahedron Letters* 3069.
12. Chalmers, A. A., Gorst-Allman, C. P. and Piacenza, L. P. L. (1977) *Tetrahedron Letters* 1665.
13. Cory, R. M. and Stothers, J. B. (1978) *Org. Mag. Reson.* **11**, 252.
14. Arnone, A., Mondelli, R. and Pyrek, J. St. (1979) *Org. Mag. Reson.* **12**, 429.
15. Wagner, H., Seitz, R., Lotter, H. and Herz, W. (1978) *J. Org. Chem.* **43**, 3339.
16. Gács-Baitz, E., Radics, L., Oganessian, G. B. and Mnatsakanian, V. A. (1978) *Phytochemistry* **17**, 1967.
17. Savona, G., Paternostro, M. P., Piozzi, F., Hanson, J. R., Hitchcock, P. B. and Thomas, S. A. (1978) *J. Chem. Soc. Perkin Trans. 1*, 643.
18. Dockerill, B., Hanson, J. R. and Siverns, M. (1978) *Phytochemistry* **17**, 572.
19. Smith, W. B. (1979) *J. Org. Chem.* **44**, 1631.
20. Buckwalter, B. L., Burfitt, I. R., Felkin, H., Joly-Goudket, M., Naemura, K., Salomon, M. F., Wenkert, E. and Wovkulich, P. M. (1978) *J. Am. Chem. Soc.* **100**, 6445.
21. Panizo, F. M., Rodríguez, B. and Valverde, S. (1974) *An. Quím.* **70**, 164.
22. de Quesada, T. G., Rodríguez, B. and Valverde, S. (1975) *Phytochemistry* **14**, 517.
23. Hanessian, S., Bargiotti, A. and La Rue, M. (1978) *Tetrahedron Letters* 737.