Production of Plantaricin NC8 by *Lactobacillus*
***plantarum* NC8 is Induced in the Presence of**
Different Types of Gram-positive Bacteria

Antonio Maldonado • José Luis Ruiz-Barba • Rufino Jiménez-Díaz

Address to which proofs are to be sent: Departamento de Biotecnología de Alimentos,
Instituto de la Grasa (CSIC), Avda. Padre García Tejero, 4, Aptdo. 1078, 41012 Seville,
Spain

(✉)**Corresponding author.** Phone: +34 954 69 10 54. Fax: +34 954 69 12 62. e-mail:
rjimenez@cica.es

Antonio Maldonado · José Luis Ruiz-Barba · Rufino Jiménez-Díaz (✉)
Departamento de Biotecnología de Alimentos,
Instituto de la Grasa, Consejo Superior de Investigaciones Científicas,
Avda. Padre García Tejero, 4, Aptdo. 1078,
41012 Seville, Spain

Abstract

Lactobacillus plantarum NC8 was shown to produce plantaricin NC8 (PLNC8), a recently purified and genetically characterized inducible class IIb bacteriocin, only when it was co-cultured with other Gram-positive bacteria. Among 82 strains belonging to the genera *Bacillus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Listeria*, *Pediococcus*, *Staphylococcus* and *Streptococcus*, forty-one of them were shown to induce PLNC8 production in *L. plantarum* NC8. There was apparently no relationship between sensitivity of strains and their ability to induce the bacteriocin thus indicating that both the inducer and sensitive phenotypes may not be linked. In some instances, the induction was promoted by both living and heat-killed cells of the inducing bacteria. However, no PLNC8-inducing activity was found in their respective cell-free, pure culture supernatants. Inducer strains promoted also production of a PLNC8-autoinducing activity by *L. plantarum* NC8, which was found only in the cell-free co-culture supernatants showing inhibitory activity. This PLNC8-autoinducing activity was shown to be diffusible, heat resistant and of proteinaceous nature, being different from the bacteriocin itself. Our results suggest that the presence of specific Gram-positive bacteria act as an environmental stimulus activates PLNC8 production by *L. plantarum* NC8 as well as a PLNC8-autoinducing activity which triggers or maintains bacteriocin production in the absence of inducing cells.

Key words Autoinduction · Induction · *L. plantarum* · Plantaricin NC8

1 **Introduction**

2
3 Bacteriocins produced by lactic acid bacteria (LAB) are peptides or proteins that inhibit the
4 growth of strains and species usually related to or of species sharing similar nutritive
5 requirements with the bacteriocin-producing bacteria (de Vuyst 1994; de Vuyst and
6 Vandamme 1994; Jack et al. 1993; Klaenhammer 1993; Tagg et al. 1976). Therefore,
7 bacteriocin production seems to be a defence mechanism directed to compete for nutrients
8 against other bacteria in the same environment (Barefoot and Grinstead 1993; Dykes 1995;
9 Dykes and Hastings 1997; Riley 1998).

10 Several reports have shown that production of bacteriocins by some Gram-positive
11 bacteria is a regulated process. Production of nisin by *Lactococcus lactis* and subtilin by
12 *Bacillus subtilis*, respectively, two modified class I bacteriocins, is regulated by a cell-
13 density dependent quorum sensing mechanism (Gutowski-Ecked et al. 1994; Kleerebezem
14 et al. 1997; Kleerebezem and Quadri 2001; Klein et al. 1993; Kuipers et al. 1995; Kuipers
15 et al. 1998). In these cases, the bacteriocins are also autoinducers. Production of several
16 class II bacteriocins such as sakacin P by *Lactobacillus sake* LTH673 (Brurberg et al.
17 1997; Eisink et al. 1996), enterocins A and B by *Enterococcus faecium* CTC 492 (Nilsen et
18 al. 1997), plantaricin ABP-118 by *L. salivarius* subsp. *salivarius* UCC118 (Flynn et al.
19 2002), plantaricins E/F and J/K by *Lactobacillus plantarum* C11 (Diep et al. 1994, 1995,
20 and 1996; Moll et al. 1999), sakacin A by *L. sakei* Lb706 (Axelsson and Holck 1995; Diep
21 et al. 2000), and carnobacteriocins A and B2 by *Carnobacterium piscicola* LV17 (Franz et
22 al. 2000; Kleerebezem et al. 2001; Quadri et al. 1997; Saucier et al. 1995; Saucier et al.
23 1997) have been shown to be regulated via a quorum sensing mechanism mediated by
24 peptide pheromones or autoinducer peptides. However, in most of them, the quorum
25 sensing mechanism is not sufficient to maintain bacteriocin production in highly diluted

1 cultures (Kleerebezem et al. 1997; Kleerebezem and Quadri 2001; Nes et al. 1996; Nes and
2 Eijsink 1999). Since concentration of the peptide pheromones in those diluted cultures is
3 not sufficient to trigger bacteriocin production, the involvement of other unknown factors
4 can be assumed. Environmental factors such as pH, temperature, and other growth
5 conditions have been proposed to play an important role in regulation of bacteriocin
6 production (Diep et al. 1995; Diep et al. 2000; Kleerebezem et al. 1997; Kleerebezem and
7 Quadri 2001; Nes et al. 1996; Nes and Eijsink 1999; Nilsen et al. 1997; Saucier et al.
8 1995). Nevertheless, little or nothing is known about how these factors interact with the
9 regulatory systems that control bacteriocin production.

10 The presence of competing microorganisms has been reported as an environmental
11 factor affecting production of bacteriocins by some LAB (Barefoot et al. 1994a and 1994b;
12 Maldonado et al. 2003; Sip et al., 1998). Thus, production of lactacin B and divercin by
13 *Lactobacillus acidophilus* N2 and *Carnobacterium divergens* AS7, respectively, is
14 enhanced in the presence of the indicator strains *L. delbrueckii* ATCC 4797 (Barefoot et al.
15 1994a and 1994b) and *C. piscicola* NCDO 2765 (Sip et al. 1998), respectively. More
16 recently, the inducible bacteriocin plantaricin NC8 (PLNC8) produced by *L. plantarum*
17 NC8 has been purified and genetically characterized (Maldonado et al. 2003). This
18 bacteriocin consists of two distinct peptides, named PLNC8 α and PLNC8 β , whose
19 complementary action is necessary for full PLNC8 activity. In this work we show that, in
20 contrast to previously reported induced bacteriocins, PLNC8 is produced only after co-
21 culture with certain Gram-positive bacteria. As a response to the presence of the other
22 bacteria, a PLNC8-autoinducing activity which is most probably involved in the regulation
23 of PLNC8 production was also produced by *L. plantarum* NC8.

Materials and methods

Bacterial strains and media

The plantaricin NC8-producer *Lactobacillus plantarum* NC8, a strain originally isolated from grass silage (Aukrust and Blom 1992), has been described previously (Maldonado et al. 2003). It was maintained at -80°C as frozen stocks in de Man, Rogosa and Sharpe medium (MRS; Oxoid, Basingstoke, Hampshire, England) plus 20% (v/v) glycerol, and propagated in MRS broth at 30°C before use.

Strains tested as inducers of PLNC8 production by *L. plantarum* NC8 and as indicator microorganisms are listed in Table 1. All *Lactobacillus*, *Leuconostoc*, *Pediococcus* and *Streptococcus* strains were grown in MRS broth at 30°C , *Leuconostoc cremoris* DB1275 was grown at 25°C , *L. acidophilus* NCDO 1748, *L. casei* strains, *L. plantarum* NCDO 1193, *L. reuteri* DSM 20016 and *L. salivarius* NCFB 2747 were grown at 37°C , and *L. bulgaricus* ATCC 11842, *L. helveticus* ATCC 15009 and *Streptococcus thermophilus* strains were grown at 42°C . *Enterococcus* and *Lactococcus* strains were propagated in GM17 medium, consisting of M17 medium (Oxoid) plus 1% (w/v) of glucose at 30°C , *L. lactis* subsp. *cremoris* CNRZ 117 was grown at 25°C , and *Enterococcus faecalis* EF1 was grown at 37°C . *Listeria* strains, *Staphylococcus carnosus* MC1, and *Bacillus cereus* ATCC 9139 were grown in Brain Heart Infusion (BHI) medium (Oxoid) at 30, 42, and 37°C , respectively. All these strains were maintained as frozen stocks at -80°C in MRS, BHI, or GM17 plus 20% (v/v) glycerol and propagated twice in their corresponding broth medium before use.

Spontaneous rifampin-resistant (Rif^r, 30 $\mu\text{g/ml}$) derivatives of *L. lactis* MG1363 and *P. pentosaceus* FBB63, were isolated by sequential selection on MRS agar containing

1 increasing concentrations of the antibiotic in the range of 0.5 to 30 µg/ml (Sigma Chemical
2 Co., St. Louis, Mo.).

3 4 Bacteriocin assays

5
6 The agar drop diffusion test was used for detection of antimicrobial activity in MRS broth
7 cultures (Jiménez-Díaz et al. 1993). Lawns of each indicator strain (Table 1) were prepared
8 by inoculating ca. 10^5 cells in 4 ml of the respective soft-overlay (0.75% agar) medium
9 which was poured onto the surface of the respective agar plates.

10 The antimicrobial activity of PLNC8 was quantified in a microtiter plate assay system
11 (Geis et al. 1983), and *L. plantarum* 128/2 [PLNC8 sensitive (PLNC8^S)] was used as the
12 indicator strain (Jiménez-Díaz et al. 1995; Maldonado et al. 2003). One bacteriocin unit
13 (BU) was arbitrarily defined as the amount of bacteriocin that inhibited the growth of the
14 indicator strain by 50% (50% of the turbidity of the control culture without bacteriocin).
15 This was expressed as the reciprocal of the highest dilution exhibiting 50% inhibition of
16 the indicator strain per millilitre (BU/ml).

17 18 Testing *L. plantarum* NC8 for PLNC8 production

19
20 *Lactobacillus plantarum* NC8 MRS broth cultures were tested for PLNC8 production at
21 different phases of growth (lag-, exponential-, or stationary-phase), sizes of initial
22 inoculum (ranging from 10^2 to 10^8 CFU/ml), pH values (in the range of 4.0 to 8.0), and
23 growth temperatures (ranging from 25 to 35°C). In all cases, culture samples were
24 withdrawn at different phases of growth, centrifuged at $10,000 \times g$ for 10 min at 4°C,
25 adjusted to pH 7.0 with 5 M NaOH, and filter-sterilized through a 0.22-µm-pore-size

1 Millex-GV filter (Millipore Iberica, S.A., Madrid, Spain). Ten μ l of these cell-free
2 supernatants were spotted onto indicator lawns of *L. plantarum* 128/2. The plates were
3 incubated at the appropriate temperature for 16 h, and examined for inhibition zones.

4 Production of PLNC8 by *L. plantarum* NC8 in co-culture with the Gram-positive
5 bacteria listed in Table 1 was determined as described in Figure 1A: fresh MRS broth was
6 inoculated with 1-2% of an overnight culture of *L. plantarum* NC8 plus 0.5% of an
7 overnight culture of each strain to test for induction. The mixed cultures were held at 30°C
8 for 6-8 h, centrifuged, and the supernatants adjusted to pH 7.0 with 5 M NaOH, filter-
9 sterilized, and finally their inhibitory activity assayed by the agar drop diffusion test, using
10 *L. plantarum* 128/2 as the indicator strain. In addition, four of these cell-free supernatants
11 showing inhibitory activity against *L. plantarum* 128/2 were selected to further determine
12 their antimicrobial spectrum against all of the strains listed in Table 1 (see footnote *c* in
13 Table 1). In control assays, all of the strains used in the mixed cultures were propagated as
14 pure cultures in their respective media and then assayed for antimicrobial activity as
15 described above.

16
17 Determination of the number of inducer cells necessary to promote PLNC8 production by
18 *L. plantarum* NC8

19
20 In order to estimate the number of inducer cells which were necessary to induce PLNC8
21 production in *L. plantarum* NC8, a fixed inoculum of the strain NC8 (10^8 CFU/ml) was co-
22 cultured in MRS broth with inocula of increasing sizes (ranging from 10^1 to 10^7 CFU/ml)
23 of the inducer strains. The mixed cultures were incubated at 30°C for 6-8 h, and the
24 antimicrobial activity in the cell-free supernatants was quantified as described above.

Time course of PLNC8 production in co-cultures

The time course of induction of PLNC8 production by *L. plantarum* NC8 in co-culture with PLNC8-resistant (PLNC8^R) or PLNC8-sensitive (PLNC8^S) but PLNC8-inducer strains was investigated. For this, fresh MRS broth was inoculated with ca. 10⁸ CFU/ml of *L. plantarum* NC8 plus ca. 10⁷ CFU/ml of either of the inducer strains *L. lactis* MG1363 Rif^r or *P. pentosaceus* FBB63 Rif^r. The co-cultures were incubated at 30°C, and samples were removed at different time intervals. Viable cells were differentially enumerated by spreading the previously diluted samples with a Spiral System (Interscience, Saint-Nom-La Bretèche, France) on MRS agar (for *L. plantarum* NC8) or MRS agar containing 30 µg/ml of rifampin (Sigma) (for *L. lactis* MG1363 or *P. pentosaceus* FBB63). Finally, PLNC8 activity in the samples was quantified as described above.

Partial characterization of the PLNC8 inducer

Overnight broth cultures of the inducer strains (Table 1) were divided in two aliquots: one of them was centrifuged (10,000 × g, 10 min, 4°C), and filtered through a 0.22-µm-pore-size Millex-GV filter (Millipore); the other part was incubated at 55°C for 1 h or autoclaved (121°C) for 15 min. Aliquots (50 µl) of the various cell-free supernatants or heat-treated cultures were added to fresh MRS broth containing ca. 10⁸ CFU/ml of an overnight culture of *L. plantarum* NC8. The mixtures were incubated at 30°C for 6-8 h, and the PLNC8 activity quantified as described above. As controls, overnight broth pure cultures of the inducer strains were used for induction of PLNC8 production in *L. plantarum* NC8.

To test for sensitivity of the inducer to proteolytic enzymes, the heat-treated cultures showing inducing activity were adjusted to pH 7.5 with 5 M NaOH, and then treated with proteinase K (Sigma) at 1 U/ml (final concentration). Mixtures were incubated at 37°C for 1 h, and then heated at 100°C for 10 min to stop the reaction. Finally, the samples were examined for induction of PLNC8 as described above. As a control, proteinase K was inactivated by heating at 100°C for 10 min prior to be added to the heat-treated inducer cultures.

Finally, MRS broth was placed inside a dialysis tube (MW cutoff of 14.0 kDa) and then inoculated with ca. 10^7 CFU/ml of either of the inducer strains *L. lactis* MG1363 or *P. pentosaceus* FBB63. Each dialysis tubing was submerged into an Erlenmeyer flask containing MRS broth previously inoculated with ca. 10^8 CFU/ml of *L. plantarum* NC8 and then incubated at 30°C for 6-8 h. Cell-free supernatants from both separated cultures were tested for PLNC8 activity.

Partial characterization of the PLNC8 autoinducer

Cell-free supernatants from *L. lactis* MG1363- or *P. pentosaceus* FBB63-*L. plantarum* NC8 co-cultures (Fig. 1A) showing PLNC8 activity were tested for their ability to induce bacteriocin production in *L. plantarum* NC8 as described in Figure 1B. For this, 20 µl of each cell-free supernatant was added to fresh MRS broth (1 ml) containing ca. 10^8 CFU/ml of an overnight culture of *L. plantarum* NC8 and incubated at 30°C for 6-8 h (Fig. 1B). Bacteriocin activity in the cell-free supernatants (named CFS₁, Fig. 1B) was quantified as described above and expressed as autoinducing units (AIU) per ml (AIU/ml). One AIU was arbitrarily defined as the amount of autoinducer in the cell-free supernatants, called hereafter PLNC8-autoinducer, which promoted production by *L. plantarum* NC8 of one

1 BU of PLNC8. This was expressed as the reciprocal of the highest dilution exhibiting
2 induction of PLNC8 production enough for 50% inhibition of the indicator strain per ml
3 (AIU/ml). These supernatants were further used for induction of PLNC8 production in
4 fresh *L. plantarum* NC8 cultures as described above (Fig. 1B).

5 To test for heat sensitivity of the inducer, samples (1 ml) of cell-free supernatants
6 containing 1,280 BU/ml of PLNC8 were incubated at 100°C for intervals up to 60 min, or
7 autoclaved (121°C) for 15 min, and the remaining activity was quantified. Sensitivity to
8 proteinase K of the PLNC8 autoinducer was carried out as described above, and the
9 remaining AIU/ml was quantified. In addition, supernatants containing autoinducing
10 activity were also placed inside a dialysis tube and then processed as described above.

11 Finally, in order to investigate whether or not PLNC8 itself was involved in the
12 autoinducing activity, the purified peptides PLNC8 α and PLNC8 β generated from PLNC8
13 by C₂-C₁₈ reverse-phase chromatography (Maldonado et al. 2003) or mixtures of them
14 were used in autoinduction assays. For this, 0.01 μ M of PLNC8 α , 0.16 μ M of PLNC8 β or
15 mixtures of both peptides (0.01 μ M of PLNC8 α plus 0.16 μ M of PLNC8 β) were added to
16 fresh MRS broth (1 ml) containing ca. 10⁸ CFU/ml of *L. plantarum* NC8, incubated at
17 30°C for 8 h, and then the bacteriocin quantified. As a control, both individual and mixed
18 peptides were assayed by the drop diffusion test for inhibitory activity against *L. plantarum*
19 128/2.

20

Results

Induction of PLNC8 in *L. plantarum* NC8 co-cultures

The influence of pH (Ahn and Stiles 1990; Barefoot et al. 1994b; Jiménez-Díaz et al. 1993; Kaiser and Montville 1993; Kleerebezem et al. 1997), temperature (Diep et al. 2000; Leal-Sánchez et al. 2002; Mørtvedt-Abildgaard et al. 1995; Tichaczek et al. 1992), and growth conditions (Biswas et al. 1991; de Vuyst et al. 1996; Gutowski-Ecked et al. 1994; Jiménez-Díaz et al. 1993; Leal-Sánchez et al. 2002), previously described as enhancers of bacteriocin production by LAB, was studied. None of these factors appeared to promote production of PLNC8 by *L. plantarum* NC8 growing alone. However, when *L. plantarum* NC8 was co-cultured with several gram-positive strains, production of PLNC8 was induced by forty-one out of eighty-two strains tested (Table 1). These strains belonged to various genera, including *Bacillus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Listeria*, *Pediococcus*, *Staphylococcus* and *Streptococcus*, and none of them exhibited any antimicrobial activity when they were cultivated as single cultures.

The inhibitory spectrum of the cell-free supernatants from four different co-cultures was found to be the same regardless of the identity of the strain used to induce bacteriocin production by *L. plantarum* NC8 (Table 1). This could indicate that most probably the same bacteriocin, PLNC8, was produced in all cases, for this bacteriocin was indeed purified in a previous work after induction with two of these strains (Maldonado et al. 2003). On the other hand, the inducer and sensitivity phenotypes are not linked, for the four types of phenotypic combinations could be observed among the strains tested (Table 1). Only 6% of the strains tested were able to induce PLNC8 production but they were sensitive to PLNC8, and included representatives from various species belonging to the

Lactobacillus or *Pediococcus* genera. Although none of the *L. plantarum* strains were able to induce production of PLNC8, most of them were sensitive to PLNC8 (80%).

Time course of PLNC8 production by *L. plantarum* NC8 in selected co-cultures

The time courses of bacteriocin production as well as its effect on the inducer population are shown in Figure 2. Although different in total amount, the shape of the curve showing PLNC8 production was very similar in both co-cultures (Fig. 2B and 2D, respectively). Thus, the PLNC8 activity could be detected in the supernatants after 3 h of co-culture with either inducer strain, with a maximum reached after 8 (Fig. 2B) or 5 h (Fig. 2D) of incubation, respectively.

As shown in Figure 2A and 2C, the growth of the *L. plantarum* NC8 population was not affected by the co-culture with the inducer strains. Also, the *L. lactis* MG1363 population was not substantially affected by the co-culture with *L. plantarum* NC8, although a decrease in the cell number (CFU/ml) by two orders of magnitude was observed after 24 h (Fig. 2A). However, the cell number of the FBB63 strain was shown to decrease early when the PLNC8 activity was detected in the culture medium, and declined by about four log units after 24 h of incubation (Fig. 2C). These differences are most probably due to the PLNC8^S phenotype of *P. pentosaceus* FBB63.

Determination of the number of inducer cells necessary to promote PLNC8 production by *L. plantarum* NC8

A minimum inoculum of 10^4 CFU/ml of *L. lactis* MG1363 was necessary to induce PLNC8 production by *L. plantarum* NC8 (40 BU/ml). The bacteriocin production increased up to a maximum (2,560 BU/ml, for 10^6 CFU/ml of this inducer strain and

conditions) and then remained constant when higher inoculum sizes were used. Similar results were obtained with other inducing bacteria (data not shown).

Partial characterization of the PLNC8 inducer

Cell-free supernatants from single broth cultures of any of the strains listed in Table 1 were not able to induce PLNC8 production by pure cultures of *L. plantarum* NC8, thus suggesting that the inducing ability was not mediated by any metabolite secreted by the inducer strains into the culture medium. When the strain NC8 was co-cultured with inducer strains but separated by a dialysis membrane, no PLNC8 activity was detected in the *L. plantarum* NC8 cell-free supernatants.

Autoclaving of the inducer cultures prior to the addition to cultures of strain NC8 did not induce PLNC8 production except for the inducer strains *L. brevis* LB9, *L. sake* NCFB 2714, *L. lactis* MG1363, *L. lactis* MG1614, *L. lactis* CNRZ 105 or *L. lactis* CNRZ 489. However, induction of PLNC8 was completely abolished in all six cases after treatment with proteinase K.

On the other hand, some inducer strains killed by heat treatment (55°C, 1 h) retained complete inducing ability. Obviously, living cells are not necessary for induction of PLNC8 production by *L. plantarum* NC8.

Partial characterization of the PLNC8 autoinducer

Figure 2 shows the time course of PLNC8-autoinducer production by *L. plantarum* NC8 when it was co-cultured with *L. lactis* MG1363 (Fig. 2B) or *P. pentosaceus* FBB63 (Fig. 2D). Differences in total amount of autoinducer at each time interval can be observed. Co-

culture with *L. lactis* MG1363 appears to promote higher amounts of autoinducer, which was virtually constant after 5 h of incubation (Fig. 2B). In contrast, *P. pentosaceus* FBB63 promoted less PLNC8 autoinducer and it reached a maximum earlier (2-3 h) (Fig. 2D).

Cell-free supernatants from *L. lactis* MG1363- or *P. pentosaceus* FBB63-*L. plantarum* NC8 co-cultures showing PLNC8 activity (Fig. 1A) were examined for consecutive inducing activity on fresh broth cultures of *L. plantarum* NC8 (Fig. 1B). Addition of 2-5% of those supernatants (Fig. 1B, 1st induction) was found to promote bacteriocin production by *L. plantarum* NC8 pure cultures (Fig. 3). Further additions of cell-free supernatants from these newly PLNC8-containing *L. plantarum* NC8 cultures (Fig. 1B, CFS₂) to other *L. plantarum* NC8 pure cultures resulted also in induction of bacteriocin production. Nevertheless, bacteriocin activity in these consecutive supernatants decreased (from 640 in CFS₁ to 80 BU/ml in CFS₂), and was completely lost in the third culture (0 BU/ml in CFS₃). On the other hand, the autoinducing activity of the primary, exogenously induced PLNC8-containing cell-free supernatants (640 AIU/ml in CFS) substantially decreased after the first culture (80 AIU/ml in CFS₁), and was finally lost after the second one (0 AIU/ml in CFS₂) (Fig. 3). On the other hand, when increasing amounts of these PLNC8-containing cell-free supernatants (CFS in Fig. 1A) were added to *L. plantarum* NC8 pure cultures they promoted increasing amounts of PLNC8 in the new autoinduced *L. plantarum* NC8 culture supernatants, up to a maximum of 640 BU/ml. In summary, both the bacteriocin and autoinducing activities obtained upon induction of *L. plantarum* NC8 with cell-free supernatants from co-cultures (in the absence of inducer strains) were much lower than those obtained after induction with the inducer strains themselves.

The PLNC8-autoinducing activity in the cell-free supernatants from *L. lactis* MG1363- or *P. pentosaceus* FBB63-*L. plantarum* NC8 co-cultures was resistant to heat, being

1 completely stable even after autoclaving (Fig. 4). Nevertheless, it was sensitive to
2 proteinase K, thus indicating its proteinaceous nature. In contrast, PLNC8 activity was
3 partially lost after the heat treatments (Fig. 4). On the other hand, neither purified PLNC8 α
4 or PLNC8 β peptides nor mixtures of them were able to induce production of PLNC8 by *L.*
5 *plantarum* NC8 in the absence of inducer bacteria.

6 Finally, when the strain NC8 was grown in flasks in which dialysis tubing containing
7 co-culture cell-free supernatants showing autoinducing activity were placed, PLNC8
8 activity was detected in the *L. plantarum* NC8 cell-free supernatant.

9

Discussion

In this report, *L. plantarum* NC8 was shown to be unable to produce PLNC8 if grown alone in liquid cultures under the environmental conditions tested. However, co-culture with several Gram-positive bacteria resulted in production of PLNC8. Induction by such external stimulus was not the only way for *L. plantarum* NC8 to produce bacteriocin. The same stimulus also induced in strain NC8 production of a proteinaceous, heat-resistant compound. That compound was shown to induce production of PLNC8 by *L. plantarum* NC8 in the absence of a second, inducing bacterium. Obviously, strain NC8 possesses a mechanism of autoinduction for PLNC8 production. These results indicate that most probably production of bacteriocin by *L. plantarum* NC8 is a well regulated process.

Enhancement of bacteriocin production in LAB by co-culture with other Gram-positive bacteria has been previously addressed (Barefoot et al. 1994a; Barefoot et al. 1994b; Sip et al. 1998). However, in the case of *L. plantarum* NC8, bacteriocin production appeared to be switched off in the absence of an inducer strain which is apparently obligatory for PLNC8 production. It is presently unknown how the inducer bacteria and *L. plantarum* NC8 interact to finally result in PLNC8 production. However, results presented here suggest that a proteinaceous substance from the inducer strains is involved. This substance did not appear to be released to the culture medium. Otherwise, cell-free supernatants or dialyzed cultures from the inducer bacteria would have been able to induce production of PLNC8. On the other hand, living inducer cells were not strictly necessary for induction of PLNC8, for heat-killed inducer cells from some strains were able to do it. However, it is not yet known whether this inducer substance is a single protein or a cell envelope structure including protein components. This protein or cell envelope structure is probably shared by all of the inducing strains.

1 In addition to PLNC8 production, the external stimulus provided by certain Gram-
2 positive bacteria also promotes production of a PLNC8-autoinducing activity by *L.*
3 *plantarum* NC8, here termed PLNC8-autoinducer. Based on the present findings, this
4 autoinducer can be differentiated from PLNC8. First, neither PLNC8 nor its individual
5 peptides PLNC8 α or PLNC8 β were able to induce PLNC8 production in *L. plantarum*
6 NC8. Second, whereas PLNC8 was sensitive to autoclaving (retained only 12.5% of the
7 initial inhibitory activity), the autoinducer activity remained unchanged (Fig. 4). And third,
8 it is known that autoinducers are effective at concentrations as low as 10^{-14} M or less (Nes
9 et al. 1996; Nes and Eijsink 1999). In our case, no autoinducing activity was detected after
10 a second induction, whereas bacteriocin activity was still detected (Fig.1B and Fig. 3).

11 In the case of *L. plantarum* NC8, it has been demonstrated that the presence of other
12 bacteria in the same environment triggers bacteriocin production. From an ecological point
13 of view, such regulatory mechanism could be part of a defence strategy which includes
14 sensing of other bacteria and a response to these by production of bacteriocin. As soon as
15 the inducing cells have disappeared from the culture medium or their population is below a
16 certain threshold, bacteriocin production can no longer be maintained. The inducing
17 bacteria also activate the production of an autoinducer molecule in *L. plantarum* NC8
18 which apparently promotes both bacteriocin and of the autoinducer itself. However, the
19 amounts of bacteriocin and autoinducer detected in that case were apparently much lower
20 than those detected after exposure to the inducing bacteria. Hence, the stimulus provided
21 by the inducer strains seems to be primarily necessary for bacteriocin production in *L.*
22 *plantarum* NC8. On the other hand, the presence of an autoinducing activity suggests the
23 existence of a quorum sensing mechanism which regulates bacteriocin production the
24 population of *L. plantarum* NC8. Actually, the presence of two direct repeats upstream of
25 the putative -35 region of *plNC8B*, the gene encoding PLNC8 β (Maldonado et al. 2003)

- 1 which agrees with the consensus sequence and structure of promoters of class II
- 2 bacteriocins regulated by autoinduction suggests a similar mechanism.
- 3

Acknowledgments

We thank L. Axelsson (MATFORSK, Norwegian Food Research Institute, Osloveien, Norway) for providing the *L. plantarum* NC8 strain.

This work was supported by the Spanish Government through MCYT project AGL2000-1611-CO3-01. A.M. was the recipient of a grant from MCYT, Spain.

References

- Ahn C, Stiles ME (1990) Plasmid-associated bacteriocin production by a strain of *Carnobacterium piscicola* from meat. *Appl Environ Microbiol* 56:2503-2510
- Aukrust T, Blom H (1992) Transformation of *Lactobacillus* strains used in meat and vegetable fermentations. *Food Res Int* 25:253-261
- Axelsson L, Holck A (1995) The genes involved in production of and immunity to sakacin A, a bacteriocin from *Lactobacillus sake* Lb706. *J Bacteriol* 177:2125-2137
- Barefoot SF, Grinstead GA (1993) Bacteriocins for dairy propionibacteria and inducible bacteriocins of lactic acid bacteria. In: Hoover DG, Steenson LR (eds) *Bacteriocins of lactic acid bacteria*. Academic Press, Inc, New York, NY, pp 219-231
- Barefoot SF, Chen YR, Hughes TA, Bodine AB, Shearer MY, Hughes MD (1994a) Identification and purification of a protein that induces production of the *Lactobacillus acidophilus* bacteriocin lactacin B. *Appl Environ Microbiol* 60:3522-3528
- Barefoot SF, Nettles CG, Chen YR (1994b) Lactacin B, a bacteriocin produced by *Lactobacillus acidophilus*. In: de Vuyst L, Vandamme EJ (eds) *Bacteriocins of lactic acid bacteria: microbiology, genetics and applications*. Blackie Academic and Professional, London, United Kingdom, pp 353-376
- Biswas S R, Ray P, Johnson MC, Ray B (1991) Influence of growth conditions on the production of a bacteriocin, pediocin AcH, by *Pediococcus acidilacti* AcH. *Appl Environ Microbiol* 57:1265-1267

- Brurberg M B, Nes IF, Eijsink VGH (1997) Pheromone-induced production of antimicrobial peptides in *Lactobacillus*. *Mol Microbiol* 26:347-360
- de Vuyst, L (1994) Bacteriocins and bacteriocin-like substances from *Lactobacillus*. In: de Vuyst L, Vandamme EJ (eds) Bacteriocins of lactic acid bacteria: microbiology, genetics and applications. Blackie Academic and Professional, London, United Kingdom, pp 319-329
- de Vuyst L, Vandamme EJ (1994) Antimicrobial potential of lactic acid bacteria. In: de Vuyst L, Vandamme EJ (eds) Bacteriocins of lactic acid bacteria: microbiology, genetics and applications. Blackie Academic and Professional, London, United Kingdom, pp 91-142
- de Vuyst L, Callewaert R, Crabbe K (1996) Primary metabolite kinetics of bacteriocin biosynthesis by *Lactobacillus amylovorus* and evidence for stimulation of bacteriocin production under unfavourable growth conditions. *Microbiology* 142:817-827
- Diep DB, Håvarstein LS, Nissen-Meyer J, Nes IF (1994) The gene encoding plantaricin A, a bacteriocin from *Lactobacillus plantarum* C11, is located on the same transcription unit as an agr-like regulatory system. *Appl Environ Microbiol* 60:160-166
- Diep DB, Håvarstein LS, Nes IF (1995) A bacteriocin-like peptide induces bacteriocin synthesis in *Lactobacillus plantarum* C11. *Mol Microbiol* 18:631-639
- Diep DB, Håvarstein LS, Nes IF (1996) Characterization of the locus responsible for the bacteriocin production in *Lactobacillus plantarum* C11. *J Bacteriol* 178:4472-4483

- Diep DB, Axelsson L, Grefslis C, Nes IF (2000) The synthesis of the bacteriocin sakacin A is a temperature-sensitive process regulated by a pheromone peptide through a three-component regulatory system. *Microbiology* 146:2155-2160
- Dykes GA (1995) Bacteriocins: ecological and evolutionary significance. *Tree* 10:186-189
- Dykes GA, Hastings JW (1997) Selection and fitness in bacteriocin-producing bacteria. *Proc R Soc Lon B* 264:683-687
- Eijsink VGH, Brurberg MB, Middelhoven PH, Nes IF (1996) Induction of bacteriocin production in *Lactobacillus sake* by a secreted peptide. *J Bacteriol* 178:2232-2237
- Floriano B, Ruiz-Barba JL, Jiménez-Díaz R (1998) Purification and genetic characterization of enterocin I from *Enterococcus faecium* 6T1a, a novel antilisterial plasmid-encoded bacteriocin which does not belong to the pediocin family of bacteriocins. *Appl Environ Microbiol* 64:4883-4890
- Flynn S, van Sinderen D, Thornton GM, Holo H, Nes IF, Collins JK (2002) Characterization of the genetic locus responsible for the production of ABP-118, a novel bacteriocin produced by the probiotic bacterium *Lactobacillus salivarius* subsp. *salivarius* UCC118. *Microbiology* 148:973-984
- Franz CMAP, van Belkum MJ, Worobo RW, Vederas J, Stiles ME (2000) Characterization of the genetic locus responsible for production and immunity of carnobacteriocin A: the immunity gene confers cross-protection to enterocin B. *Microbiology* 146:621-631

- Geis A, Singh J, Teuber M (1983) Potential of lactic streptococci to produce bacteriocin. *Appl Environ Microbiol* 45:205-211
- Gutowski-Ecked Z, Klein C, Siegers K, Bohm K, Hammelmann M, Entian KD (1994) Growth phase-dependent regulation and membrane localization of Spa B, a protein involved in biosynthesis of the lantibiotic subtilin. *Appl Environ Microbiol* 60:1-11
- Jack R, Tagg JR, Ray B (1995) Bacteriocins of Gram-positive bacteria. *Microbiol Rev* 59:171-200
- Jiménez-Díaz R, Rios-Sánchez RM, Desmazeaud M, Ruiz-Barba JL, Piard JC (1993) Plantaricin S and T, two new bacteriocins produced by *Lactobacillus plantarum* LPCO10 isolated from a green olive fermentation. *Appl Environ Microbiol* 59:1416-1424
- Jiménez-Díaz R, Ruiz-Barba JL, Cathcart DP, Holo H, Nes IF, Sletten KH, Warner PJ (1995) Purification and partial amino acid sequence of plantaricin S, a bacteriocin produced by *Lactobacillus plantarum* LPCO10, the activity of which depends on the complementary action of two peptides. *Appl Environ Microbiol* 61:4459-4463
- Kaiser AL, Montville TJ (1993) The influence of pH and growth rate on production of the bacteriocin, bavaricin MN, in batch and continuous fermentations. *J Appl Bacteriol* 75:536-540
- Klaenhammer TR (1993) Genetics of bacteriocins produced by lactic acid bacteria. *FEMS Microbiol Rev* 12:39-86

- Kleerebezem M, Quadri LEN, Kuipers O.P, de Vos WM (1997) Quorum sensing by peptide pheromones and two-component signal-transduction systems in gram-positive bacteria. *Mol Microbiol* 24:895-904
- Kleerebezem M, Quadri LEN (2001) Peptide pheromone-dependent regulation of antimicrobial peptide production in Gram-positive bacteria: a case of multicellular behavior. *Peptides* 22:1579-1596
- Kleerebezem M, Kuipers OP, de Vos WM, Stiles ME, Quadri LEN (2001) A two-component signal transduction cascade in *Carnobacterium piscicola* LV17B: two signaling peptides and one sensor-transmitter. *Peptides* 22:1597-1601
- Klein C, Kaletta C, Entian KD (1993) Biosynthesis of the lantibiotic subtilin is regulated by a histidine kinase/response regulator system. *Appl Environ Microbiol* 59:296-303
- Kuipers OP, Beerthuyzen MB, de Ruyter PGGA, Luesink EJ, de Vos WM (1995) Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. *J Biol Chem* 270:27299-27304
- Kuipers OP, de Ruyter PGGA, Kleerebezem M, de Vos WM (1998) Quorum sensing-controlled gene expression in lactic acid bacteria. *J Biotechnol* 64:15-21
- Leal-Sánchez MV, Jiménez-Díaz R, Maldonado-Barragán A, Garrido-Fernández A, Ruiz-Barba JL (2002) Optimization of bacteriocin production by batch fermentation of *Lactobacillus plantarum* LPCO10. *Appl Environ Microbiol* 68:4465-4471

- Maldonado A, Ruiz-Barba JL, Jiménez-Díaz R (2003) Purification and genetic characterization of plantaricin NC8, a novel coculture-inducible two-peptide bacteriocin from *Lactobacillus plantarum* NC8. *Appl Environ Microbiol* 69:383-389
- Moll GN, van den Akker E, Hauge HH, Nissen-Meyer J, Nes IF, Konings WN, Driessen AJM (1999) Complementary and overlapping selectivity of the two-peptide bacteriocins plantaricin EF and JK. *J Bacteriol* 181:4848-4852
- Mørtvedt-Abildgaard CI, Nissen-Meyer J, Jelle B, Grenov B, Skaugen M, Nes IF (1995) Production and pH-dependent bactericidal activity of lactococin S, a lantibiotic from *Lactobacillus sake* L45. *Appl Environ Microbiol* 61:175-179
- Nes IF, Diep DB, Håvarstein LS, Brurberg MB, Eijsink VGH, Holo H (1996) Biosynthesis of bacteriocins in lactic acid bacteria. *Antonie van Leeuwenhoek* 70:113-128
- Nes IF, Eijsink VGH (1999) Regulation of group II peptide bacteriocin synthesis by quorum sensing mechanisms. In: Dunny GM, Winans SC (eds) *Cell-cell signaling in bacteria*. American Society for Microbiology Press, Washington, DC, pp 175-192
- Nilsen T, Nes IF, Holo H (1997) An exported inducer peptide regulates bacteriocin production in *Enterococcus faecium* CTC492. *J Bacteriol* 180:1848-1854
- Quadri LEN, Kleerebezem M, Kuipers OP, de Vos WM, Roy KL, Vederas JC, Stiles ME (1997) Characterization of a locus from *Carnobacterium piscicola* LV17B involved in bacteriocin production and immunity: evidence for global inducer-mediated transcriptional regulation. *J Bacteriol* 179:6163-6171

- Riley MA (1998) Molecular mechanisms of bacteriocin evolution. *Annu Rev Genet* 32:255-278
- Ruiz-Barba JL, Cathcart DP, Warner PJ, Jiménez-Díaz R (1994) Use of *Lactobacillus plantarum* LPCO10, a bacteriocin producer, as a starter culture in spanish-style green olive fermentations. *Appl Environ Microbiol* 60:2059-2064
- Saucier L, Poon A, Stiles ME (1995) Induction of bacteriocin in *Carnobacterium piscicola* LV17. *J Appl Bacteriol* 78:684-690
- Saucier L, Paradkar AS, Frost LS, Jensen SE, Stiles ME (1997) Transcriptional analysis and regulation of carnobacteriocin production in *Carnobacterium piscicola* LV17. *Gene* 188:271-277
- Sip A, Grajek W, Boyaval P (1998) Enhancement of bacteriocin production by *Carnobacterium divergens* AS7 in the presence of a bacteriocin-sensitive strain *Carnobacterium piscicola*. *Int Food Microbiol* 42:63-69
- Tagg JR, Dajani AS, Wannamaker LW (1976) Bacteriocins of gram-positive bacteria. *Bacteriol Rev* 40:722-756
- Tichaczek PS, Nissen-Meyer J, Nes IF, Vogel RF, Hammes WP (1992) Characterization of the bacteriocins curvacin A from *Lactobacillus curvatus* LTH1174 and sakacin P from *L. sake* LTH673. *System Appl Microbiol* 15:460-468

Table 1. Induction of plantaricin NC8 (PLNC8) production by *L. plantarum* NC8 in co-culture with gram-positive bacterial strains, and inhibitory spectrum of the induced bacteriocin.

Bacterial species	Source ^a	Induction of PLNC8 production in strain NC8 ^b	Sensitivity to induced PLNC8 from co-culture CFSS ^c
<i>Bacillus cereus</i> ATCC 9139	TNO	+	R
<i>Enterococcus faecalis</i> EF1	TNO	+	R
<i>E. faecalis</i> JH2- 2	INRA	-	R
<i>E. faecalis</i> OG1-X	UG	+	R
<i>E. faecalis</i> BM 4100 WT	TNO	+	R
<i>E. faecalis</i> CNRZ 135	INRA	+	R
<i>E. faecalis</i> CNRZ 136	INRA	+	R
<i>E. faecalis</i> CNRZ 137	INRA	+	R
<i>E. faecium</i> LP6T1a	Our collection	+	R
<i>E. faecium</i> LP6T1a-20 ^d	Our collection	+	R
<i>Lactobacillus acidophilus</i> ATCC 4356	TNO	-	R
<i>L. acidophilus</i> NCDO 1748	NCDO	+	R
<i>L. brevis</i> LB9	UV	+	R
<i>L. bulgaricus</i> ATCC 11842	TNO	+	R
<i>L. casei</i> ATCC 334	TNO	-	R
<i>L. casei</i> NCDO 161	NCDO	+	R
<i>L. casei</i> NCDO 393	CIT	-	R
<i>L. curvatus</i> NCFB 2739	TNO	-	R
<i>L. fermentum</i> ATCC 9338	TNO	+	S
<i>L. fermentum</i> ATCC 14933	ATCC	-	S
<i>L. fermentum</i> NCDO 1750	NCDO	+	S
<i>L. helveticus</i> ATCC 15009	TNO	+	R
<i>L. hilgardii</i> LB76	UV	+	S
<i>L. pentosus</i> ATCC 8041	CECT	-	S
<i>L. plantarum</i> ATCC 8014	ATCC	-	S
<i>L. plantarum</i> BOM1	Our collection	-	S
<i>L. plantarum</i> LB6	UV	-	S
<i>L. plantarum</i> LPC1	Our collection	-	S
<i>L. plantarum</i> LPC13	Our collection	-	S
<i>L. plantarum</i> LPC2	Our collection	-	S
<i>L. plantarum</i> LPD1	Our collection	-	S
<i>L. plantarum</i> LPD5	Our collection	-	S
<i>L. plantarum</i> LPD7	Our collection	-	S
<i>L. plantarum</i> LPE1	Our collection	-	R
<i>L. plantarum</i> LPE5	Our collection	-	S
<i>L. plantarum</i> LPP1	Our collection	-	S

Table 1. continued (a)

Bacterial species	Source ^a	Induction of PLNC8 production in strain NC8 ^b	Sensitivity to induced PLNC8 from co-culture CFSs ^c
<i>L. plantarum</i> LPP5	Our collection	-	R
<i>L. plantarum</i> LPS1	Our collection	-	S
<i>L. plantarum</i> LPS5	Our collection	-	S
<i>L. plantarum</i> LPS21	Our collection	-	S
<i>L. plantarum</i> LPS26	Our collection	-	S
<i>L. plantarum</i> LPS29	Our collection	-	S
<i>L. plantarum</i> LPT 70/3	Our collection	-	R
<i>L. plantarum</i> NCDO 1193	CIT	-	R
<i>L. plantarum</i> 37A	Our collection	-	S
<i>L. plantarum</i> 55-1 ^d	Our collection	-	R
<i>L. plantarum</i> 15N/1	Our collection	-	S
<i>L. plantarum</i> 128/2	Our collection	-	S
<i>L. plantarum</i> 144/1	Our collection	-	S
<i>L. reuteri</i> DSM 20016	TNO	+	R
<i>L. sake</i> NCFB 2714	TNO	+	S
<i>L. salivarius</i> NCFB 2747	TNO	+	R
<i>Lactococcus lactis</i> IL1441	INRA	+	R
<i>L. lactis</i> IL1403	INRA	+	R
<i>L. lactis</i> MG1363	CIT	+	R
<i>L. lactis</i> MG1614	CIT	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 105	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 106	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 113	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 114	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 117	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 158	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 163	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 485	INRA	+	S
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 489	INRA	+	R
<i>L. lactis</i> subsp. <i>cremoris</i> CNRZ 511	INRA	+	R
<i>Leuconostoc cremoris</i> DB1275	TNO	-	R
<i>L. mesenteroides</i> 32	INRA	+	R
<i>L. mesenteroides</i> 33	INRA	+	R
<i>Listeria innocua</i> BL 86/26	TNO	+	R
<i>L. monocytogenes</i> NCTC 7973	FVM	-	R
<i>L. monocytogenes</i> LI5 sv1/2	FVMN	-	R
<i>L. monocytogenes</i> NCTC 5105	FVM	-	R

Table 1. continued (b)

Bacterial species	Source ^a	Induction of PLNC8 production in strain NC8 ^b	Sensitivity to induced PLNC8 from co-culture CFSs ^c
<i>L. monocytogenes</i> LI1 sv4	FVM	-	R
<i>L. monocytogenes</i> Scott A	FVM	-	R
<i>Pediococcus damnosus</i> NCDO 1832	CIT	-	R
<i>P. parvulus</i> P339	UV	-	R
<i>P. pentosaceus</i> FBB63	TNO	+	S
<i>P. pentosaceus</i> PC1	TNO	+	R
<i>P. pentosaceus</i> P56	UV	-	R
<i>Staphylococcus carnosus</i> MC1	TNO	+	R
<i>Streptococcus thermophilus</i> ST20	TNO	+	R
<i>S. thermophilus</i> ST112	TNO	+	R

^aAbbreviations: ATCC, American Type Culture Collection (Rockville, Md.); CECT, Colección Española de Cultivos Tipo (Universidad de Valencia, Burjasot, Spain); CIT, Cranfield Institute of Technology (United Kingdom); DSM, DMSZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (Braunschweig, Germany); FVM, Facultad de Veterinaria, Universidad Complutense (Madrid, Spain); INRA, Institut National de la Recherche Agronomique (Jouy-en-Josas, France); NCDO, National Collection of Dairy Organisms (Reading, United Kingdom); NCFB, National Collection of Food Bacteria, c/o NCIMB Ltd. (Aberdeen, Scotland, United Kingdom); NCTC, National Collection of Type Cultures (Central Public Health Laboratory, London, United Kingdom); TNO, Nutrition and Food Research (Zeist, The Netherlands); UT, Universität Tübingen (Tübingen, Germany); UV, Universidad de Valencia (Burjasot, Spain); UG, Universidad de Granada (Granada, Spain).

^bPLNC8 determined in cell-free supernatants (CFSs) from mixed cultures of *L. plantarum* NC8 plus the tested strain was assayed by the agar drop diffusion test (Jiménez-Díaz et al. 1993) with *L. plantarum* 128/2 as the indicator strain (Maldonado et al. 2003). Symbols: +, induction of PLNC8; -, not induction.

^cSensitivity of the listed species to the PLNC8 produced by *L. plantarum* NC8 when it was co-cultured either with *L. lactis* MG1363, *P. pentosaceus* FBB63, *L. salivarius* NCFB 2747, or *L. mesenteroides* 33. Symbols: S, sensitive; R, resistant.

^dNon-bacteriocin producing strains derived from bacteriocin producing ones: *E. faecium* LP6T1a-20 from the enterocin I producer *E. faecium* LP6T1a (Floriano et al. 1998), and *L. plantarum* 55-1 from the plantaricins S and T producer *L. plantarum* LPCO10 (Jiménez-Díaz et al. 1993).

Legends of the Figures

Figure 1. The procedure for induction of PLNC8 production by *L. plantarum* NC8. Percentages (v/v) in parentheses indicate proportion of inoculum or cell-free supernatant in total (final) volume.

Figure 2. Growth (log CFU/ml), induced PLNC8 activity (log BU/ml), and PLNC8-autoinducing activity (log AIU/ml) in *L. plantarum* NC8-*L. lactis* MG1363 (panels A and B) or *L. plantarum* NC8-*P. pentosaceus* FBB63 co-cultures (panels C and D). Symbols: in A and C, *L. plantarum* NC8 in single (●) and in co-cultures (○); *L. lactis* MG1363 in single (▲) and in co-cultures (△); and *P. pentosaceus* FBB63 in single (◆) and in co-cultures (◇). In B and D, PLNC8 (■) and PLNC8-autoinducing activities (□) in the cell-free supernatants.

Figure 3. PLNC8 and PLNC8-autoinducing activities in consecutive cultures of *L. plantarum* NC8. CFS, cell-free supernatant from co-culture of *L. plantarum* NC8 and *L. lactis* MG1363; CFS₁, *L. plantarum* NC8 culture induced with a cell-free supernatant from CFS; and CFS₂, *L. plantarum* NC8 culture induced with a cell-free supernatant from CFS₁. Neither PLNC8 nor PLNC8-autoinducing activities were detected in CFS₃ (see Figure 1B).

Figure 4. Sensitivity to heat of co-culture induced *L. plantarum* NC8 cell-free supernatants showing PLNC8 activity.

A) Induction of PLNC8 production by co-culture

B) Induction of PLNC8 production by cell-free supernatants

Figure 2

Figure 3

Figure 4