



*J. Plankton Res.* (2015) 37(4): 834–850. First published online July 4, 2015 doi:10.1093/plankt/fbv055

# Larval fish assemblage structure in the surface layer of the northwestern Mediterranean under contrasting oceanographic scenarios

ITZIAR ÁLVAREZ<sup>1\*</sup>, JOSÉ MARÍA RODRÍGUEZ<sup>2</sup>, IGNACIO A. CATALÁN<sup>1</sup>, MANUEL HIDALGO<sup>3</sup>, DIEGO ÁLVAREZ-BERASTEGUI<sup>4</sup>, ROSA BALBÍN<sup>3</sup>, ALBERTO APARICIO-GONZALEZ<sup>3</sup> AND FRANCISCO ALEMANY<sup>3</sup>

<sup>1</sup>INSTITUT MEDITERRANI D'ESTUDIS AVANÇATS (IMEDEA), MIQUEL MARQUÉS 21, 07190 ESPORLES, I. BALEARIC SPAIN, <sup>2</sup>INSTITUTO ESPAÑOL DE OCEANOGRÁFIA, CENTRO OCEANOGRÁFICO DE GIJÓN, AVDA. PRÍNCIPE DE ASTURIAS 70BIS, 33212 GIJÓN, ASTURIAS, SPAIN, <sup>3</sup>INSTITUTO ESPAÑOL DE OCEANOGRÁFIA, CENTRO OCEANOGRÁFICO DE BALEARES, MOLL DE PONENT SN, 07014 PALMA DE MALLORCA, SPAIN AND <sup>4</sup>SOCIB, BALEARIC ISLANDS COASTAL OBSERVING AND FORECASTING SYSTEM, PARCIBIT, ED. NAORTE, BLOC A 2<sup>º</sup> P. PTA. 3, PALMA DE MALLORCA, SPAIN

\*CORRESPONDING AUTHOR: itziar@imedea.uib-csic.es

Received December 15, 2014; accepted June 14, 2015

Corresponding editor: Marja Koski

During the summer, the relative influence of resident Atlantic Waters (AW) and new AW largely drives the mesoscale dynamics around the Balearic Islands (NW Mediterranean). Two principal summer hydrographic scenarios were identified in the region, differentiated by the relative position of the density front between new and resident AW within the archipelago and its associated mesoscale activity. In this study, we investigated how those early summer mesoscale scenarios influence larval fish assemblages, by analyzing data from two cruises representative of these two scenarios (2004 and 2005). Redundancy analysis was used to assess the variance in the larval fish assemblage that could be significantly explained by the most parsimonious combination of available environmental variables in both years. While depth was the most important variable in explaining the larval fish assemblage structure variability observed under both scenarios, indicators of mesoscale activity (dynamic height, geostrophic velocity) contributed significantly to understanding the dynamics of the larval fish community. Mesoscale activity was higher in summer 2004, leading to higher larval fish abundances and zooplankton biomass and lower larval fish diversity than in the unusually warm summer 2005, which showed lower mesoscale activity. The larval assemblage dynamics are discussed in terms of extrinsic and species-specific factors.

**KEYWORDS:** mesoscale circulation; dynamic height; fish larvae; spawning grounds; spatial distribution; NW Mediterranean