

Far beyond the Sun – II. Probing the stellar magnetism of the young Sun ι Horologii from the photosphere to its corona

E. M. Amazo-Gómez¹,^{1*} J. D. Alvarado-Gómez¹,¹ K. Poppenhäger^{1,2},^{1,2} G. A. J. Hussain,^{3,4}
 B. E. Wood,⁵ J. J. Drake,⁶ J.-D. do Nascimento, Jr.,^{6,7} F. Anthony,⁷ J. Sanz-Forcada,⁸ B. Stelzer,⁹
 F. Del Sordo^{10,11}, M. Damasso¹², S. Redfield¹³, J. F. Donati¹⁴, P. C. König,^{4,15} G. Hébrard^{15,16}
 and P. A. Miles-Páez⁴

¹Leibniz-Institut für Astrophysik Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

²Universität Potsdam, Institut für Physik und Astronomie, Karl-Liebknecht-Straße 24/25, 14476 Potsdam-Golm, Germany

³European Space Agency, Keplerlaan 1, 2201 AZ Noordwijk, Netherlands

⁴European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748 Garching bei München, Germany

⁵Naval Research Laboratory, Space Science Division, Washington, DC 20375, USA

⁶Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge MA 02138, USA

⁷Universidade Federal do Rio Grande do Norte (UFRN), Dep. de Física, 59078-970 Natal, RN, Brazil

⁸Centro de Astrobiología (CSIC-INTA), ESAC Campus, Camino Bajo del Castillo, E-28692 Villanueva de la Cañada, Madrid, Spain

⁹Eberhard Karls Universität, Institut für Astronomie und Astrophysik, Sand 1, 72076 Tübingen, Germany

¹⁰INAF – Osservatorio Astrofisico di Catania, via Santa Sofia, 78 Catania, Italy

¹¹Institute of Space Sciences (ICE-CSIC), Campus UAB, Carrer de Can Magrans s/n, 08193 Barcelona, Spain

¹²INAF – Osservatorio di Torino, Via Osservatorio, 20, 10025 Pino Torinese TO, Italy

¹³Astronomy Department and Van Vleck Observatory, Wesleyan University, Middletown, CT 06459-0123, USA

¹⁴CNRS-IRAP, 14, avenue Edouard Belin, F-31400 Toulouse, France

¹⁵Institut d’astrophysique de Paris, UMR7095 CNRS, Sorbonne Université, 98 bis boulevard Arago, 75014 Paris, France

¹⁶Observatoire de Haute-Provence, CNRS, Université d’Aix-Marseille, 04870 Saint-Michel-l’Observatoire, France

Accepted 2023 July 5. Received 2023 July 5; in original form 2022 October 21

ABSTRACT

A comprehensive multiwavelength campaign has been carried out to probe stellar activity and variability in the young Sun-like star ι -Horologii. We present the results from long-term spectropolarimetric monitoring of the system by using the ultra-stable spectropolarimeter/velocimeter HARPS at the ESO 3.6-m telescope. Additionally, we included high-precision photometry from the NASA *Transiting Exoplanet Survey Satellite* (*TESS*) and observations in the far- and near-ultraviolet spectral regions using the STIS instrument on the NASA/ESA *Hubble Space Telescope* (*HST*). The high-quality data set allows a robust characterization of the star’s rotation period, as well as a probe of the variability using a range of spectroscopic and photometric activity proxies. By analysing the gradient of the power spectra (GPS) in the *TESS* light curves, we constrained the faculae-to-spot driver ratio ($S_{\text{fac}}/S_{\text{spot}}$) to 0.510 ± 0.023 , which indicates that the stellar surface is spot dominated during the time of the observations. We compared the photospheric activity properties derived from the GPS method with a magnetic field map of the star derived using Zeeman–Doppler imaging (ZDI) from simultaneous spectropolarimetric data for the first time. Different stellar activity proxies enable a more complete interpretation of the observed variability. For example, we observed enhanced emission in the *HST* transition line diagnostics C IV and C III, suggesting a flaring event. From the analysis of *TESS* data acquired simultaneously with the *HST* data, we investigate the photometric variability at the precise moment that the emission increased and derive correlations between different observables, probing the star from its photosphere to its corona.

Key words: stars: activity – stars: chromosphere – stars: coronae – stars: magnetic fields – stars: photosphere – stars: solar-type.

1 INTRODUCTION

An accepted understanding among solar and stellar scientists is that the 11-yr variations in global activity observed in the Sun are coherent and connected with the 22-yr magnetic cycle. These cycles have been extensively studied and correlated with a synchronized behaviour of

the solar atmospheric layers (i.e. photosphere, chromosphere, and corona; Hathaway 2015; Li, Xu & Feng 2022). To gain a better understanding of the magnetic properties of other stars, researchers have combined multiple techniques to overcome the challenges arising from the limited availability of high-quality data, as compared to the solar counterpart (see review by Reiners 2012). Techniques such as the Zeeman Doppler Imaging (ZDI; Donati et al. 1997; Hussain et al. 2000; Piskunov & Kochukhov 2002), allow us to investigate the stellar large-scale polarity. Moreover, dynamo simulation studies

* E-mail: eamazogomez@aip.de, emamazog@unal.edu.co

emphasize the significance of considering essential parameters, such as stellar inclination, rotation, and observation phase coverage, in order to improve the recovery of more accurate ZDI magnetic diagnostics (see Hackman et al. 2023).

By implementing the ZDI technique, large-scale polarity and activity modulations, including magnetic reversals, have been detected only in a few stars (e.g. Fares et al. 2009; Jeffers et al. 2017; Boro Saikia et al. 2018, 2022). These studies have also identified some of them that exhibit solar-like cyclic behaviour, such as 61 Cyg A (Boro Saikia et al. 2016). However, for most stars, the time-scale for magnetic field reversals is much shorter. Furthermore, these rapid reversals do not show a resemblance to the observed variability in chromospheric activity, unlike the case of the Sun (e.g. Donati et al. 2008; Fares et al. 2009; Morgenthaler et al. 2012).

This situation becomes particularly challenging when studying other indicators of stellar magnetism, such as coronal activity, as monitoring X-ray emissions over the entire duration of the cycle is difficult (see table 3 in Wargelin et al. 2017). There is not enough available stellar data, and further research is required to gain a better understanding of the relationship between magnetic field behaviour and activity variations in stars, similar to what we have for the Sun. This could involve supporting more long-term observational campaigns, as presented in this work, to study the coronal activity and polarimetric behaviour of stars over extended time-scales.

As mentioned, combining multiple instruments and techniques to monitor the magnetic variability of stars different from the Sun is very challenging, but still proven to be an effective way to render a robust description of them. Apart from the Sun, there are just a few systems that have been observed over multiple wavelengths and followed up regularly, (e.g. α Cen A, ϵ Eri, ξ Bootis, etc.; Drake & Smith 1993; Laming, Drake & Widing 1996; Hatzes et al. 2000; Drake et al. 2000; Pagano et al. 2004; Benedict et al. 2006; Janson et al. 2008; Ness & Jordan 2008; Dumusque et al. 2011, 2012; Robrade, Schmitt & Favata 2012; Anglada-Escudé & Butler 2012; Hatzes 2013; Jeffers et al. 2014, 2017; Boro Saikia et al. 2016, 2018, 2022; Coffaro et al. 2020).

ι Horologii (ι Hor, HD 17051, HR 810) is one of the few systems with a robust set of multiwavelength records covering a period of years (see Baliunas et al. 1995; Saar & Osten 1997; Valenti & Fischer 2005; Barnes 2007; Ramírez et al. 2017). The star has an estimated age of about ~ 625 Myr (see Vauclair et al. 2008) and similar characteristics to the young Sun (i.e. F8V-G0V; ~ 6080 K; $R_* \sim 1.16 R_\odot$; Bruntt et al. 2010, see Table 1). The star hosts a giant Jupiter-like planet at approximately 1 au (see Kürster et al. 2000; Zechmeister et al. 2013). ι Hor is more active than the Sun, e.g. $\log L_X/L_{\text{bol}} = -4.9$ on average (Sanz-Forcada, Stelzer & Metcalfe 2013; Alvarado-Gómez et al. 2018), and has one of the shortest chromospheric and coronal activity cycles known to date (~ 1.6 yr; Metcalfe et al. 2010; Sanz-Forcada et al. 2011; Sanz-Forcada et al. 2013; Sanz-Forcada et al. 2019). Besides, a magnetic activity cycle of ~ 2.1 yr was described in Alvarado-Gómez et al. (2018, hereafter Paper I, 2019a). The cycles are analysed in this paper after extended observations, evidencing the possibility of a double-cycle behaviour.

Given the resemblance between ι Hor and the young Sun, its short magnetic and activity cycles (close to 10 times shorter than the 11-yr solar cycle), the star represents a particularly interesting target to monitor in order to get a better insight into stellar magnetism generation and enable us to test the solar activity paradigm (i.e. whether it is or not correct to apply the solar activity analogy to the star). Long-term observations, as performed for ι Hor, benefit the development of more accurate models of stellar dynamos, which are critical to our understanding of magnetism throughout the main

Table 1. Fundamental properties of ι Hor.

Parameter	Value	Reference
Spectral Type	F8V – G0V	Bruntt et al. (2010)
T_{eff} [K]	6080 ± 80	Bruntt et al. (2010)
$\log(g)$	4.399 ± 0.022	Bruntt et al. (2010)
R_* [R_\odot]	1.16 ± 0.04	Bruntt et al. (2010)
M_* [M_\odot]	1.23 ± 0.12	Bruntt et al. (2010)
$v \sin i$ [km s^{-1}]	6.0 ± 0.5	Alvarado-Gómez et al. (2018)
P_{rot} [d] ^{spec}	$7.70^{+0.18}_{-0.67}$	Alvarado-Gómez et al. (2018)
$\overline{P}_{\text{rot}}$ [d] ^{photo}	$7.39^{+0.28}_{-0.16}$	This work
$\langle v_R \rangle$ [km s^{-1}] ^a	16.927 ± 0.001	This work
$\langle \log(L_X) \rangle$	28.78 ± 0.08	Sanz-Forcada et al. (2013)
Age [Myr] ^b	~ 625	Vauclair et al. (2008)

Note. Summary of the main properties of ι Hor. The (spect) and (photo) refer to the spectroscopic and photometric estimations of the rotation period values, respectively. ^a Average value from the 6 multi-epoch HARPSpol observations reported in this work and Alvarado-Gómez et al. (2018). ^b Age derived from HARPS asteroseismology. This value falls in between the estimates from gyrochronology (~ 740 Myr; Barnes 2007), and from the level of X-ray (~ 500 Myr; Sanz-Forcada et al. 2011).

sequence: the magnetism driving the star’s angular momentum loss and causing its magnetic activity to decrease over time (Wright et al. 2011, 2018; Reiners & Mohanty 2012; Reiners, Schüssler & Passegger 2014; Wright & Drake 2016; Reiners et al. 2022). The rate at which activity, and hence high-energy radiation and stellar wind decrease, is ultimately critical for rotational and possibly magnetic dynamo evolution (Garraffo et al. 2018).

Here, we show how a multiwavelength analysis allows probing activity across different atmospheric layers. By compiling the information contained in a set of activity indicators, it is possible to build a global picture of the star’s magnetic activity state. We argue that, due to its proximity, brightness, and youth, ι Hor is a keystone star to build on and test the solar-stellar analogy. In this paper, we continue the follow-up to the star as part of our intensive monitoring of ι Hor under the *Far beyond the Sun* campaign (P.I. J.D. Alvarado-Gómez). In Paper I, we described the first half of the data acquired, 3 ESO HARPSpol observational periods. Here, we cluster together the entire set of ESO HARPSpol data (6 observational periods) and simultaneous observations of the star obtained by *TESS* and *HST* telescopes.

1.1 Data exploration

Under the *Far beyond the Sun* campaign, the ι -Hor system has been intensively monitored over six semesters, during ESO observing periods P₉₆ to P₁₀₁ (between 2015 October to 2018 September). This allowed us to collect 199 data points using the spectropolarimetric mode of the HARPS instrument (HARPSpol) at the ESO 3.6-m telescope at La Silla Observatory (see Mayor et al. 2003; Piskunov et al. 2011).

In addition to the spectropolarimetric data, we analysed photometric time series obtained by the *TESS* telescope (Ricker et al. 2015). We describe in Section 2 the complete spectropolarimetric data set gathered for this project. We outline the analysis of our 2.95-yr HARPSpol follow-up, adding three additional semesters observing ι Hor to those presented in Paper I. We used Sector 2 and Sector 3, for a total data compilation of 54 d (2018 August 23–September 20, 2018 September 20–October 17, respectively). We also looked at sectors 29 and 30 (2020 August and September, respectively) obtaining similar results as described in Section 3. In




Figure 1. Top panel: long term follow up of S_{HK} value from the Ca II H & K core lines. Grey dots are the original scattered data before the smoothing averaging over the rotational period modulation. Blue points are in detail analysed in Metcalfe et al. (2010, M + 2010), Flores et al. (2017, F + 2017), and Alvarado-Gómez et al. (2018, AG + 2018). X-rays observations in red-filled rhomboids for ι -Horologii reported by Sanz-Forcada et al. (2013, 2019, SF + 2013, 2019). The purple-filled squares are new data presented in this work. As well as the Olive dashed line which corresponds to *TESS* and *HST* simultaneous observations. Bottom panel: observed minus calculated residuals of the double period fitting.

addition, we retrieve the surface faculae to spot ratio, $S_{\text{fac}}/S_{\text{spot}}$, by applying the GPS method over *TESS* photometric time series. We complemented our collection of data for ι Hor with acquired NUV and FUV spectra using the Space Telescope Imaging Spectrograph (STIS) instrument (see Woodgate et al. 1986; Gull & et al. 1986; Woodruff, Ludtke & Woodgate 1990) on board the *Hubble Space Telescope* observatory. We analyse two separate visits of 3 orbits each (one in the NUV and, two in the FUV regions), yielding 6 separate exposures for the target. Three observations were taken on 2018 September 03 over cycle 25 (P.I. J.D. Alvarado-Gómez, *HST* proposal ID 15299) and three on 2019 August 01 during cycle 26 (P.I. J.D. Alvarado-Gómez, *HST* proposal ID 15512). The STIS/*HST* observations were obtained using the Multi-Anode Microchannel Array (MAMA) detector by the E140M (FUV) and E230H (NUV) gratings. A more detailed description of the *HST* observations is presented in Section 4. Section 5 contains a multitechnique (Zeeman-Doppler Imaging, spectropolarimetry, photometry), multiwavelength (FUV, NUV, visible) comparative analysis carried out in one epoch consisting of simultaneous observations by HARPSpol, *TESS*, and *HST*. We present the main discussion and conclusions in Section 6. The data availability is described in Section 7.

2 HARPS SPECTROPOLARIMETRY

The HARPS instrument on the ESO 3.6-m telescope at the La Silla Observatory is a high-precision ultrastable échelle spectrograph, and velocimeter. It can attain a precision of 0.97 m s^{-1} over a spectral range of 378 to 691 nm, and has a spectroscopic resolution of $R = 120\,000$ (see Mayor et al. 2003; Pepe et al. 2005). In addition, the instrument offers a spectropolarimetric mode that enables observation of circularly and linearly polarized signatures across the entire spectral range (see Piskunov et al. 2011). We used the spectropolarimetric mode of the instrument over six consecutive semesters, from ESO period P_{96} to P_{101} (as shown with vertical dashed lines in Fig. 1). The three first semesters (P_{96} – P_{98}) were analysed in Paper I. In this work, we complete the analysis by adding

98 new spectropolarimetric data points collected over the final three semesters of the campaign (P_{99} – P_{101}).

Each observation sequence consisted of roughly one-hour exposures, divided into four sub-exposures between which the half-wave Fresnel rhombs were rotated to different angles to construct the circular and null polarization profiles. Homogeneously, as in Paper I, the data reduction was carried out using LIBRE-ESPRIT (Donati et al. 1997). The reduced data products consisted of unpolarized intensity profiles and circularly polarized Stokes-V signatures. Diagnostic null (N) spectra serve as checks for instrumental noise and artefacts where combining the individual sub-exposures helps to cancel out possible polarization coming from the instrument. These spectra were created by applying the ratio method (see Bagnulo et al. 2009).

Making use of the instrument’s high resolution and spectropolarimetric observational mode, we analysed the radial velocities (RVs). We extracted information about the stellar activity from indicators such as the Ca II S-index S_{HK} , and the H α emission as index $I_{\text{H}\alpha}$. We also probed the stellar magnetic behaviour by measuring the effect of the longitudinal magnetic field B_{ℓ} over the spectra. We analyse the information obtained from the HARPSpol data set in the following subsections.

2.1 Ca II H & K index

In the chromospheres of Sun-like stars, the source functions of ionized metals are generally dominated by collisional processes. In this context, the collisionally dominated Ca II H & K Fraunhofer lines and, the emission observed specifically in the line cores, turned out to be a good indicator of activity in this atmospheric layer (see Thomas 1957; Hall 2008). The enhancement of the line cores, which has been demonstrated to form in the high chromosphere, is a response to the presence of magnetically driven chromospheric heating and temperature rise (see Eberhard & Schwarzschild 1913; Thomas 1957; Wilson 1963; Skumanich 1972; Vaughan, Preston & Wilson 1978; Wilson 1978; Hall 2008). The Ca II S-index, S_{HK} , was introduced by Vaughan et al. (1978) as a consistent way to analyse the fluxes in the

Table 2. Two-period model for the S_{HK} activity evolution of ι -Hor.

Parameter	Value
<i>1st Component</i>	
Period (P_1)	547.6 ± 3.0 d (1.499 ± 0.012 yr)
Semi-amplitude (A_1)	0.0141 ± 0.0010
Phase (Φ_1)	4.73 ± 0.16 rad
<i>2nd Component</i>	
Period (P_2)	401.0 ± 7.3 d (1.097 ± 0.023 yr)
Semi-amplitude (A_2)	0.0155 ± 0.0008
Phase (Φ_2)	3.56 ± 0.04 rad
<i>Offsets $\langle S \rangle_{\text{SET}}$</i>	
CASLEO-REOSC (30) ^a	0.2323 ± 0.0013
SMARTS-RC Spec (143)	0.2389 ± 0.0010
HARPS PH3 (66)	0.2438 ± 0.0011
HARPSpol (199)	0.2443 ± 0.0005
<i>Statistics</i>	
Reduced χ_r^2	2.33
Data points	439
$\langle S \rangle_{\text{ALL}}$	0.234 ± 0.011

^aNumber of data points for a given instrument: REOSC at CASLEO, RC Spec at SMART and, HARPS at the 3.6 m ESO telescope at La Silla.

cores of Ca II H & K lines in the Sun, and was quickly extended to Sun-like stars. The S_{HK} index is a dimensionless ratio that measures the enhanced emission of the Ca II H & K lines, centred at 396.85 and 393.37 nm, by comparing the emission flux in the line cores with that in the nearby continuum. We follow the approach and relations showed in Wright et al. (2004):

$$S_{\text{HK}} = \beta \left(\frac{F_{\text{H}} + F_{\text{K}}}{F_{\text{R}} + F_{\text{V}}} \right). \quad (1)$$

Here, F_{H} and F_{K} are the fluxes measured within 0.218 nm wide triangular bandpasses centred on the line cores at 396.8469 and 393.3663 nm, while F_{R} and F_{V} are the fluxes measured in two rectangular 2 nm wide bandpasses centred on the continuum on either side of the HK lines at 400.107 and 390.107 nm, respectively. The β calibration factor depends on the implemented instrument. We used in this work the β calibration coefficient for HARPS obtained by Alvarado-Gómez et al. (2015):

$$\beta_{\text{HARPS}} = 15.39 \pm 0.65, \text{ with an offset of } 0.058.$$

We estimated a percentile error of the S_{HK} (eS_{HK}) using the signal to noise of the spectral order corresponding to the H & K lines, as

$$eS_{\text{HK}} = \frac{1}{S/N}(S_{\text{HK}}). \quad (2)$$

In Paper I, we analysed the S_{HK} variability from the first half of the campaign (P96 to P98, blue points in Fig. 1), incorporating into the analysis archive HARPS spectra, as well as, values obtained by SMARTS RC SPEC instrument reported in Metcalfe et al. (2010), and, data by CASLEO/REOSC (see Flores et al. 2017). The new S_{HK} values obtained during the ESO periods P99 to P101 are analysed here and listed in the journal of observations (see the appendix section).

The top panel of Fig. 1 shows the compilation of the four different data sets utilized for a multiparameter least-squares curve-fitting and the adjusted zero-level normalization (see, Table 2).

We used the SYSTEMIC2 console package (Meschiari et al. 2009) to re-analyse the entire data set, including the data from the second half of our HARPSpol campaign. The total data compilation comprises 439 S_{HK} values, which we used to perform a bootstrapping random periodicity analysis with 10^4 trials, and allowing periods between $p_{\text{min}} = 2.0$ d and $p_{\text{max}} = 10^4$ d within a 10^5 sampling grid. We removed the Earth's translational period of 365.25 d. We




Figure 2. Periodicity analysis on the long-term S_{HK} variations of ι -Hor (Fig. 1). Lomb–Scargle (LS) periodograms are recovering two main significant peaks above the noise level (LSP ~ 40) $P_1 = 547.65 \pm 3\text{d}$ (1.49 ± 0.01 yr) and $P_2 = 401.01 \pm 7\text{d}$ (1.09 ± 0.02 yr).

analysed the persistent periodicities in the long-term activity and found that a single activity cycle cannot explain the full data set. Instead, the superposition of two sinusoidal functions provided a much better fit to the data (10 free parameters, see Table 2). The offsets in Table 2 are the adjustment for detrend and normalizing the different instrument sets of S_{HK} values. Additionally, we accounted for the scatter attributed to the stellar rotation (grey dots in Fig. 1) by binning the instrument-specific data to the estimated rotation period of 7.8 d obtained by applying the GPS method (see Section 3.1.1). We observed a reduced scatter in the amplitude of the time series, as well as a considerable improvement of the reduced χ_r^2 of from 20.02 with the original data to 2.33 by fitting the averaged by-period time series (cyan line in Fig. 1). However, we see that the phase and period of the fitted functions are not affected by the binning procedure, indicating the robustness of the solution. From the periodogram analysis (see Fig. 2), we found two superimposed periods: $P_1 = 547.65 \pm 3\text{d}$ (1.49 ± 0.01 yr) and $P_2 = 401.01 \pm 7\text{d}$ (1.09 ± 0.02 yr).

In contrast to the X-ray cycle of 588.5 d (1.6 yr) determined by Sanz-Forcada et al. (2013) and confirmed in Sanz-Forcada et al. (2019), the total S_{HK} data showed a double periodicity; one of 1.49 yr, closer to the periodic X-ray signal, and a secondary signal of 1.09 yr. In the bottom panel of Fig. 1, we show the residuals from the optimized χ square fit, resulting in a double-period curve. We observe that the residuals are not totally flat, even with this optimal fit to the S_{HK} time series. Meanwhile, the section of S_{HK} data plot as purple points in Fig. 1 are showing the modulation of the index over about one activity cycle synchronized with the X-ray observations with *XMM-Newton* telescope presented by Sanz-Forcada et al. (2019). A possible reconciliation between having different cycles estimated by X-rays and chromospheric indexes could be due to a superposition of two predominant cycle modes (as has been suggested in Paper I and also more recently in the S_{HK} cycle of κ Ceti by Boro; Saikia et al. 2022). In Paper I was also proposed the possibility of a geometrical effect given by the misalignment of the observed features in the low chromosphere and the corona, where features located near the limbo edge can be traced in the corona but less clearly in the photosphere or chromosphere.

A comparison between the Ca II H & K line cores during minimum (P99–A, blue profile) and maximum activity periods (P100–C, red profile) is shown in Fig. 3. The ι Hor Ca II H & K line profile cores have a clear strong emission at both activity minimum and maximum periods. At activity maximum ι Hor has higher levels of emission in the H & K cores at each epoch of observation, likely indicating




Figure 3. Ca II H & K Fraunhofer core lines (left and right, respectively) for the set of observations during the minimum period of activity in P_{99–A} (blue background plot) and comparison with the set of observations during maximum activity period in P_{100–C} (red overplotted line), see comparative Ca II S-index S_{HK} values in Fig. 1 for reference. The black spectrum at the bottom of both panels corresponds to the solar Ca II H & K at activity minimum, as obtained by the HARPS instrument on 2017 April 12.

a higher degree of coverage by chromospheric active regions at a range of longitudes.

Similarly than performed for the S_{HK} index, we analysed the time series of H α emission as index $I_{\text{H}\alpha}$ and, the longitudinal component of the magnetic field B_{ℓ} in the following sections, see Fig. 4 in Sections 2.2 and 2.3.

2.2 H α emission as index $I_{\text{H}\alpha}$

The H α line centred at 656.2801 nm provides another commonly used activity proxy of cool stars. The fluxes from the line wings and core trace emission from the upper photosphere and middle chromosphere, respectively. Multiple stellar features such as flares, prominences, spicules, filaments, etc. contribute to the emission of the H α line. For example, the core of the H α line, in particular its width and line shift, is an indicator of photospheric hot intergranular walls, known as the network or faculae if closer to a concentrated magnetic region. This line is also a tracer of the plages (see Labonte 1986). Similarly, the line has been used for tracing structures in rapidly rotating young convective stars, using ‘slingshot prominences’ which appear as transient absorption features in H α (see Jardine et al. 2020).

We calculated the H α activity indicator $I_{\text{H}\alpha}$ by integrating the flux over a 0.36-nm bandpass around the H α line core at 656.2801 nm (called $F_{\text{H}\alpha}$), and dividing $F_{\text{H}\alpha}$ by the flux integrated over two continuum regions of 0.22-nm width on the red and the blue side of the H α line core: C_{B} at 655.885 nm, and C_{R} at 656.730 nm, as defined in Gizis, Reid & Hawley (2002), Wright et al. (2004), Marsden et al. (2014), see Fig. 5

$$I_{\text{H}\alpha} = \frac{F_{\text{H}\alpha}}{C_{\text{B}} + C_{\text{R}}}. \quad (3)$$

The relative positions and intensities of telluric to stellar lines, from one spectrum to another, can change due to different atmospheric conditions such as air mass and humidity during the observations. Those Earth atmospheric lines around the H α region have been identified by using the Rowland table of ‘The solar spectrum’ (Moore, Minnaert & Houtgast 1966). The effects of the telluric lines inside the considered core and wing regions of H α at about 655.8, 656.1, 656.4, and 656.9 nm (see Table B1) are corrected by subtracting the area under the line profile and applying a cubic-spline fitting. We used a mask of width W (in mÅ) to correct the spectrum, as shown coloured in red in Fig. 5.

We compare H α line profiles from activity minimum and maximum (ESO P_{99–A} and P_{100–C}) in Fig. 5. We do not observe a substantial difference in the line core for the different activity regimens, instead, we observe some slight excess towards the blue, C_{B} and red, C_{R} wing regions of the line that will make the biggest impact on the resultant indicator.

Following Fig. 6 one can say that the H α index, $I_{\text{H}\alpha}$, is correlated with the chromospheric Ca II H & K index S_{HK} , which is quantified by a moderate positive Pearson correlation of $\rho_{S_{\text{HK}}, I_{\text{H}\alpha}} = 0.82$. The correlation is very broad, given the high scatter on the H α index values. The values for $I_{\text{H}\alpha}$ obtained during ESO P₉₉ to P₁₀₁ are reported in Table C1.

2.3 Longitudinal magnetic field

Circularly polarized light gives us a direct estimation of the magnetic field along the line-of-sight, B_{ℓ} , unlike the indirect information obtained through the activity indicators discussed thus far (S_{HK} , $I_{\text{H}\alpha}$, and L_X). However, circularly polarized (Stokes V) signatures in Sun-like stars typically have relative amplitudes of 0.1 per cent.




Figure 4. Long term evolution of the H α index in ι Hor. Grey dots are the original scattered data before the smoothing averaging over the rotational period. Blue open circles during observation periods P₉₆ to P₉₈ are reported in Paper I. Purple-filled squares are the 98 values obtained during the observation periods P₉₉ to P₁₀₁. The cyan line corresponds to the reduced optimized χ -square fit to the S_{HK} data set, resulting in a double-period curve. The red line is the 1.6-yr fit from X-rays reported in Sanz-Forcada et al. (2013, 2019).




Figure 5. H α lines for the set of observations during the period of minimum activity in P_{99-A} (black background plot) and comparison with the maximum activity period in P_{100-C} (golden over-plot). The corrections for telluric lines are shown in red; see Fig. 1 for reference.

It is therefore not possible to obtain magnetic field measurements with a sufficiently high level of signal-to-background noise (S/N) in individual photospheric lines. We employ the multiline technique of least squares deconvolution (LSD) to produce a mean Stokes V signature with the co-added S/N of thousands of lines in order to robustly detect and analyse Zeeman signatures from stars like ι Hor (Donati et al. 1997; Kochukhov, Makaganiuk & Piskunov 2010).

As described in the first paper of this study, a photospheric line list, tailored to the stellar properties of ι Hor, is required to retrieve

the LSD profiles on each night. This spectral line list, containing information on line depths, rest wavelengths, and Landé factors, is generated with the aid of the Vienna atomic line database (VALD3; Ryabchikova et al. 2015; Pakhomov, Piskunov & Ryabchikova 2017). The mask computed corresponds to $T_{\text{eff}} = 6000$ K, $\log(g) = 4.5$ and microturbulent velocity of 1.0 km s⁻¹. Literature values of T_{eff} , $\log(g)$ are in Table 1. The microturbulent velocity of 1.04 km s⁻¹ is used as reference (see Bruntt et al. 2010). A depth threshold of 0.05 in normalized units, was used to retrieve the line list from the data base. The line mask was then refined following the method discussed in Alvarado-Gómez et al. (2015), matching the mask line depths to a high S/N HARPSpol spectrum of ι Hor (acquired during the first epoch of observations in 2015 October). The mask was then cleaned from very broad lines (e.g. Ca H&K, H α) and line blends that would result in large deviations from the self-similarity assumption required by the LSD procedure (see Donati et al. 1997; Kochukhov et al. 2010). The final optimized line mask for ι Hor had 8834 entries within the HARPSpol spectral range. This optimized line mask was used to retrieve nightly LSD profiles within a range of velocities between -5.0 and 45.4 km s⁻¹, with a velocity spacing appropriate for the HARPS spectrograph (~ 0.8 km s⁻¹). The typical S/N in the retrieved LSD Stokes V profiles varied between 5×10^4 and 8×10^4 , depending on the S/N of the original spectrum (see Tables C1 1 and C1 2). An example LSD profile is presented in Fig. 7.

Once we obtain the LSD profiles, we employ the centre of gravity technique (COG technique is a commonly used method to determine the centre of a photospheric line profile. At sufficiently high spectral resolution, it can be used to determine the radial velocity of the star by measuring the centre of the Stokes intensity profile in velocity space, see Uitenbroek 2003) to measure the radial velocity and compute the disc-integrated longitudinal magnetic field.

In the weak-field approximation regime, the longitudinal component of the magnetic field B_{ℓ} , is linearly proportional to the amplitude of the Stokes V parameter. In this regime, the Stokes V amplitude can be computed as follows

$$\frac{V}{I} = -g_{\text{eff}} C_z \lambda^2 \frac{dI}{I d\lambda} B_{\ell}, \quad (4)$$




Figure 6. Activity relation between $I_{H\alpha}$ and the S_{HK} index from the Ca II H & K cores. Left- and right-hand panels are coloured with B_ℓ and $|B_\ell|$, respectively.




Figure 7. Extracted LSD profile for ι Hor from the HARPSpol observations acquired in 08.09.2018. The different panels show the intensity (I), circular polarization (Stokes V), and diagnostic null (N) profiles, normalized to the continuum intensity (I_c). Both, LSD Stokes V and N , have been enhanced by a constant factor (5×10^3) for visualization purposes.

where V is the amplitude of the Stokes V profile, I is the intensity at a central wavelength λ , $C_z = 467 \text{ nm}^{-1} \text{ G}^{-1}$, and g_{eff} is the effective Landé factor (a dimensionless quantity describing the magnetic sensitivity, or response of the line, to the magnetic field).

The LSD Stokes $I(v)$ and $V(v)$ profiles represent the continuum intensity and circularly polarized profiles respectively. B_ℓ can be

estimated (in Gauss) from the following expression:

$$B_\ell = -714 \frac{\int v V(v) dv}{\lambda_0 \bar{g} \int [I_c - I(v)] dv}, \quad (5)$$

In our HARPSpol data set, the central wavelength is $\lambda_0 \simeq 509 \text{ nm}$, the mean Landé factor for the integrated LSD Stokes I $\bar{g} \simeq 1.198$, and v is the velocity in km s^{-1} .

The main source of uncertainty in B_ℓ , σ_{B_ℓ} , comes from the velocity limits over which the flux is integrated into equation (5), and the propagation of errors from the LSD technique (see Marsden et al. 2014). We used a velocity range between 5.4 and 30.2 km s^{-1} , which maximizes the B_ℓ/σ_{B_ℓ} ratio and is sufficient to encompass the entire profile. We check for instrumental noise by applying LSD and COG techniques to the null spectra and compared them with the B_ℓ measurements. This enables us to evaluate whether our B_ℓ values have contributions from instrumental polarization. The resulting values of B_ℓ , σ_{B_ℓ} , and N_ℓ , are listed in columns 10–12 of Table C1.

We look for quantified correlations between the different activity indicators S_{HK} , $I_{H\alpha}$, B_ℓ and the absolute value of the longitudinal magnetic field $|B_\ell|$. The last one, even though suffering from cancellation effects, allows us to compare the magnitude of the radial field with the activity indicators (see Figs 6 and 8). We observe a clear correlation between the S_{HK} and $I_{H\alpha}$ indicators, expressed with a Pearson coefficient of $\rho_{(S_{HK}, I_{H\alpha})} = 0.82$, as described in Section 2.2. Such a correlation is not evident between B_ℓ and the activity indexes. For example, the coefficient of correlation between B_ℓ and S_{HK} is quantified by a very weak positive Pearson correlation of $\rho_{(B_\ell, S_{HK})} = 0.025$. In the case of B_ℓ and $I_{H\alpha}$, the coefficient is $\rho_{(B_\ell, I_{H\alpha})} = 0.046$. For the values obtained between B_ℓ and $|B_\ell|$, we do observe a slightly higher anticorrelation ($\rho_{(B_\ell, |B_\ell|)} = -0.26$). The compilation of correlation coefficients between the different indicators of activity, and the RVs calculated in the following section are organized in Table 5.




Figure 8. Long term follow up of B_ℓ in i Hor. Grey dots are the original scattered data before the smoothing averaging over the rotational period. Blue open circles during observation periods P₉₆ to P₉₈ are reported in Paper I. Purple-filled squares are the 98 values obtained during the observation periods P₉₉ to P₁₀₁. The cyan line shows the optimized reduced χ -square fit to the S_{HK} data set, resulting in a double-period curve. The red curve is the 1.6-yr cycle from X-rays reported in Sanz-Forcada et al. (2013, 2019).

2.4 Radial velocity evolution

For the analysis of the radial velocity (RV) evolution presented in this section, we included pipeline-processed spectroscopic HARPS PH3 archival data¹ acquired between 2003 November and 2016 December (60 spectra), together with the HARPSpol data set acquired between 2015 October and 2018 September (199 spectra), resulting in 259 data points for the radial velocity time series.

2.4.1 Least-Squares deconvolution and bisector profiles

The RVs are measured from the LSD profiles. As outlined in Paper I, the RV of each star was measured by extracting the LSD Stokes I profile and measuring the centroid from a least-squares Gaussian fit to the data. We found that these measurements were very consistent with RVs derived from ISPEC, using the fitting procedure of the cross-correlation function from the HARPS/SOPHIE G2 line mask with typical RV errors of 2 m s^{-1} . To measure the asymmetry of the line profile, which could be due to physical mechanisms related to the stellar activity (e.g. starspots), we also computed the bisector as defined in Santerne et al. (2015), and the value of BIS , i.e. the velocity span between the average of the top and the bottom parts:

$$BIS = BIS_{\text{top}} - BIS_{\text{bottom}}, \quad (6)$$

where BIS_{top} is the average of the bisector between 60 and 90 per cent of the total contrast in the flux of the line profile and BIS_{bottom} is the average of the bisector between 10 and 40 per cent of the total contrast, where 100 per cent corresponds to the continuum and 0 per cent to the minimum of the line profile.

Across the whole time series, the value of the BIS does not exceed 143 m s^{-1} , with an average value of $\overline{BIS} \approx 92.1 \text{ m s}^{-1}$, which represents about 82 per cent of the overall measured RV variation. The fact that the BIS value has the same order of magnitude than the RV variation indicates that the asymmetry of the LSD profile induced by the stellar activity cannot be neglected. By applying the technique

to three different values of the spectral binning of the LSD profile in velocity-space, $\delta v = 0.8, 0.4$ and 0.2 km s^{-1} , we confirmed that the choice of δv does not affect the calculation of the BIS value. The results shown here are based on the BIS values derived for $\delta v = 0.8 \text{ km s}^{-1}$.

In Paper I, Alvarado-Gómez et al. (2018) first noted a hint of an *activity gradient* in the RV-subtracted residuals, which we can further demonstrate with this more complete data set (see *left-hand* and *middle* panels of Fig. 9). The stellar activity levels, as traced by the chromospheric spectral indices, follow the distribution of RV residuals, with the (O–C) values tending to be positive during higher levels of Ca II H & K S_{HK} and the $I_{\text{H}\alpha}$, and negative (O–C) values corresponding to lower activity levels. On the other hand, the disintegrated longitudinal magnetic field B_ℓ (Fig. 9, *right*), tracing the larger scale activity *jitter*, shows no apparent coherence with the RV variations, confirming that B_ℓ does not trace the global activity level in the same way that the chromospheric activity indicators do. While this dependence found in the residuals confirms the presence of activity-related *jitter*, the upper panels in Fig. 9 show no such dependence on activity, confirming that the RV variability in i Hor is caused by the presence of a planet.

3 TESS PHOTOMETRY

Building on the legacy of the Kepler mission, the *Transiting Exoplanet Survey Satellite* (TESS) (Ricker et al. 2015) – also a planet-hunting mission – has enhanced our ability to explore the photometric variability of about 200 000 bright nearby stars across the whole sky in an area 400 times larger than that covered by the Kepler mission (Borucki et al. 2010). The TESS mission has allowed us in this work to analyse simultaneously high-resolution photometric, spectropolarimetric and UV observations for i Hor. In this section, we analysed the i Hor light curves obtained by the TESS mission for two consecutive sectors, S2 and S3, from 2018 August 23 to 2018 October 17. We also pre-analysed sectors S29 and S30, but given the high level of noise present in those sectors, we focus here on data from S2 and S3, which were also acquired closer to the dates of our simultaneous *HST* NUV/FUV and spectropolarimetric observations.

¹http://archive.eso.org/wdb/wdb/adp/phase3_main/form




Figure 9. RV variations of ι Hor determined from the combined HARPS PH3 and HARPSpol data sets. The phase and orbital solution (solid line) have been fitted using the parameters reported by Zechmeister et al. (2013). The residuals are shown in the bottom panel. The colour codings denote the corresponding values of S_{HK} (left), $I_{H\alpha}$ (middle), and B_ℓ (right) for each observation (when available).

TESS light curves were obtained from the High-Level Science Products (HLSP) on the Mikulski Archive for Space Telescopes (MAST). In order to improve the analysis for the *TESS* data set, we stitched, normalized, detrended, and reduced the noise on sectors S2 and S3 following the procedure as described in de Almeida, Anthony & do Nascimento (2020), Anthony et al. (2022). In this process, we derived the *TESS* long-cadence (30 and 10-min cadence) light curves for ι Hor by performing the photometric extraction from the full-frame images (FFI) data. First, we downloaded a 35×35 pixel cutout image centred on our target using the *TESScut* tool on MAST servers (Brasseur et al. 2019). Next, we applied a constructed custom mask aperture that includes each pixel from the target star and extracted the data from the observed fields `TIME [BJD]`, `FLUX [e-/s]`, and `FLUX_ERR [e-/s]`. Then, we constructed vectors with observing times, measured fluxes, and flux uncertainties, respectively. Following the same steps, we also estimate the background signal considering the surrounding pixels of the object where the point-spread function appears to be zeroed. Before detrending the light curve, we performed a background subtraction. The resulting light curve is divided by the median and converted into normalized fluxes.

3.1 Faculae to spot ratio

Starspots and faculae imprint characteristic patterns on photometric time series, diminishing or enhancing the observed stellar brightness, while transiting the stellar disc. In this section, we focus on recovering a quantitative description of the stellar surface through the facular-to-spot ratio ($S_{\text{fac}}/S_{\text{spot}}$), by implementing the gradient of the wavelet power spectra method (GPS; see Shapiro et al. 2020; Amazo-Gómez et al. 2020a, b). Precise information on the stellar rotation period is required in order to properly analyse the stellar surface and the $S_{\text{fac}}/S_{\text{spot}}$. Consequently, we implement independent methods to retrieve the stellar rotation period and compare our findings with previous analysis in the literature.

3.1.1 Rotation Period

The stellar rotation period is key for determining a number of stellar properties and important for recovering an estimate of the distribution of features over the stellar surface. In order to obtain a better value of this parameter, we applied seven different methods to recover photometric periodicities related to the stellar rotation period.

Table 3. Rotation period analysis from different independent methods.

Method	Prot [d]	Error [d]
GP (S2)	6.64	+0.28 −0.09
GP (S3)	7.01	+0.16 −0.09
QP-GP (S2 + S3)	7.23	0.20
GLS (S2 + S3)	7.73	0.21
ACF (S2 + S3)	7.69	0.23
PS (S2 + S3)	7.94	0.32
GPS (S2 + S3)	7.78	0.18

We used quasi-periodic Gaussian-process regression (QP-GP; see Damasso & Del Sordo 2017; Foreman-Mackey et al. 2017, 2018; Barros et al. 2020), the Generalized Lomb–Scargle periodogram (GLS; see Zechmeister & Kürster 2009), the Autocorrelation Function (ACF; see McQuillan, Aigrain & Mazeh 2013; McQuillan, Mazeh & Aigrain 2014; Santos et al. 2019), Wavelet Power Spectra (PS; see Torrence & Compo 1998; García et al. 2014; Santos et al. 2019), and the Gradient of Power Spectra (GPS; e.g. see Shapiro et al. 2020; Amazo-Gómez et al. 2020a, b; Amazo-Gómez 2021). We applied those different methods (QP-GP, GLS, ACF, PS, and GPS) to analyse the combined, normalized, and stitched sectors 2 and 3 on *TESS* photometry (from 2018 August 23 to 2018 October 17, see Table 3). Additionally, we applied the GP method independently for sectors, S2 and S3. We do not observe strong deviations from any of the methods when applied to S2 and/or S3. More detailed descriptions of the different methods that we used can be found in Appendix A.

In Table 3, we compile the results from the methods applied and show the results from the GLS, ACF, PS, and GPS methods in Fig. A2. The rotational modulation from the *TESS* light curves is consistently recovered by all seven of the implemented methods. The five methods applied to the normalized and stitched S2 + S3 LCs gave us an estimate of the rotation period ranging from 7.23 to 7.94 d. Those values obtained are in agreement with the 7.718 ± 0.007 d obtained for the combined S2 + S3 LCs analysed in Sanz-Forcada et al. (2019). The two additional methods applied independently to S2 and S3 retrieved values between 6.64 to 7.48 d for S2, and 7.01 to 7.24 d for S3. We noted that this range of periods is lower than the average period of 8.43 ± 0.02 d for S2 and 7.95 ± 0.02 d for S3 previously recovered from the GLS analysis of the pipeline-reduced




Figure 10. In blue the GPS α -Factor of 400 modelled faculae to spot ratio ($S_{\text{fac}}/S_{\text{spot}}$) adapted from Shapiro et al. (2020). The α -Factor is proportional to the high-frequency inflection point found in the gradient of the power spectra, GPS, and inverse to the rotation period, as α -Factor = $\text{HFIP}/P_{\text{rot}}$. The Sun is located in the transition region between the branches of spot-dominated (left) and faculae-dominated stellar surfaces (right). In Amazo-Gómez et al. (2020a), we found an $\alpha_{\odot} = 0.158$ which corresponds to a $S_{\text{fac}}/S_{\text{spot}} \sim 3$. The pink cross shows the position of ι Hor and demonstrates that it is located in the spot-dominated branch of the diagram with a $\alpha_{\iota} = 0.209 \pm 0.011$ and a $S_{\text{fac}}/S_{\text{spot}} = 0.510 \pm 0.023$.

TESS light curves in Sanz-Forcada et al. (2019). We considered that this discrepancy in the analysis of the individual sectors might be due to the different light curve extraction procedures. In Sanz-Forcada et al. (2019), light curves were extracted after testing different apertures constructed from a summed image of all cadences from a particular sector and choosing the aperture for which the light curve showed the lowest standard deviation. As described earlier, the light curves we use here were extracted after subtracting the background signal measured from surrounding pixels where the point spread function from the target should be zero.

3.1.2 Gradient of Power Spectra, GPS method

Recent studies have suggested that our Sun exhibits particular characteristics that differentiate it from even its closest stellar analogues. For example, Metcalfe, Egeland & van Saders (2016) suggest that Sun-like stars in the middle of their main sequence life could start showing hints of a dynamo shutdown, and transition towards a different dynamo regime. Reinhold et al. (2020) found that the solar variability seems to be an outlier if compared against stars with similar temperature, age, and rotation periods (when available for those slowly rotating stars).

Further analysis by Shapiro et al. (2020) and Amazo-Gómez (2021) showed that the rotation period of stars with complex light curves, as for the solar case, can be reliably determined by implementing a novel technique based on the profile of the gradient of power spectra (GPS). In Amazo-Gómez et al. (2020a), the rotation modulation is detected at all solar activity levels, where other methods have failed. By characterizing the particular shape generated by facular (M-like shape) or spot (V-like shape) transits, recorded in the total solar irradiance (TSI) and compared simultaneously with MDI observations, it was possible to infer whether the stellar surface was dominated by facular or spot regions. The new method developed in Shapiro et al. (2020); Amazo-Gómez et al. (2020a, b);

Amazo-Gómez (2021) allows us not just to infer, but to quantify the degree of spot- or faculae-dominance on the stellar surface based on solar and stellar light curves. The quantification is made through the ratio between the bright and dark features, $S_{\text{fac}}/S_{\text{spot}}$. Through the application of the GPS method to both solar-like LC simulations and TSI observations, it has been found that Sun-like stars exhibit distribution across three distinct regimes. Those regimes characterize either the dominance by the presence of spots or faculae on the stellar light curves and, stars that seem to be in a transition between the two branches. Interestingly, a detailed characterization of the solar brightness variations found that the Sun lies in the middle of the branch between the spot- and faculae-dominant regimes. The entire methodology is described, tested and applied in Shapiro et al. (2020); Amazo-Gómez (2021).

From the stellar photometric brightness variations of ι Hor, we derived the GPS α -factor and compared this against the faculae-to-spot driver ratio ($S_{\text{fac}}/S_{\text{spot}}$) for 400 modelled light curves analysed in Shapiro et al. (2020, see Fig. 10). The α -factor is proportional to the high-frequency inflection point ($\text{HFIP} = 1.63$ d) found in the GPS, and inversely proportional to the rotation period. It is expressed as: $\alpha - F = \text{HFIP}/P_{\text{rot}}$. We found that ι Hor has a high alpha-factor $\alpha - F = 0.209 \pm 0.011$ (cf. 0.158 ± 0.007 for the Sun). As shown in Fig. 10, ι Hor (pink cross) is located in the spot-dominated branch of the diagram of α -Factor versus $S_{\text{fac}}/S_{\text{spot}}$. This indicates that the stellar surface is spot dominated, with a facular-to-spot ratio ($S_{\text{fac}}/S_{\text{spot}}$) of 0.510 ± 0.023 .

4 HST/STIS DATA

In the following paragraphs, we discuss spectroscopic observations of ι Hor acquired during *HST* cycles 25 and 26 in the far- and near-ultraviolet (FUV & NUV) regions. We used the STIS/*HST* instrument with the E140M grating and FUV-MAMA detector, and E230H grating with NUV-MAMA detector, covering 1150–

1700 Å and 1620–3100 Å, respectively². We acquired data over six orbits during cycles 25 and 26, three orbits in each cycle. The orbits during cycle 25 were acquired in 2018 September 03, and on 2019 August 01 for cycle 26. The E230H NUV-MAMA observations setup had an exposure of 2019 s and, the E140M FUV-MAMA involved two exposures of 3141 s. The spectra were reduced by the pipeline calibration code CALSTIS from the STIS science software package.

4.1 Lines in the FUV

We performed a comparison of the relative amplitudes of selected spectral lines between our STIS/*HST* stellar observations and solar UV spectra of a quiet and active region acquired by the Solar Ultraviolet Measurements of Emitted Radiation (SUMER) on board the Solar and Heliospheric Observatory (SOHO). The solar comparison data were of a quiet region on 1997 April 20, and the active region AR-NOAA8487 on 1999 March 18 (see Curdt et al. 2001; Curdt, Landi & Feldman 2004) and are available at Werner Curdt's homepage.

We used two Gaussians functions, one thin and one thick, to fit the flux lines, such as S II, S III, NV Si IV, C IV, C III, O IV, which are emission lines with broad wings, see Figs 11 to D1, and D2. The narrow line fits the core of the line profiles and may correspond to a turbulent wave dissipation or possible Alfvén wave heating mechanism (see Wood, Linsky & Ayres 1997). The second broader Gaussian is used to fit the wings, in analogy with solar observations, this may be caused by flare heating effects or, as suggested by Peter (2001), due to magnetoacoustic waves travelling through the transition region to the corona. In particular, C IV λ 1548.2 Å line is characteristic of the transition region, with an approximate temperature of 10^5 K. We find that the stellar spectra in the C IV region are comparable with a typical observation of a solar active region (green line in Fig. 12). Comparing the four spectra acquired during Cycles 25 and 26 (see dotted lines, left-hand panel of Fig. 12) we observe an enhancement of the emission of S III, C IV, C III, and O IV during the second FUV orbit of Cycle 25 (C25-II) which are detected as a higher amplitude in the line cores and broadening of the wings. The enhanced emission in the analysed lines suggests a possible flaring event. The total flux from the combined narrow and broad fitted Gaussian profile fits are shown in Table 4. Although we noticed an increase in flux in the FUV lines during C25, we could not find any clear evidence of a flare in the *TESS*-LC data that we analysed. However, there was a slight rise in flux at the start of C25-II, which can be seen in the zoomed-in bottom right-hand panel of Fig. A3.

4.2 Searching for an Astrosphere around *ι* Hor

The original objective of our *HST* program was to measure the stellar mass-loss rate of *ι* Hor employing the astrospheric technique developed by Wood et al. (2002, 2005a). This procedure is based on the detection of the Hydrogen wall emerging from the interaction between the stellar wind and the local interstellar medium (ISM). This absorption has been observed for a number of stars and is one of the few ways to detect mass-loss from low-mass stars (Wood et al. 2005b, 2021). The astrospheric H I signature (captured in the blue wing of the Lyman α line) is highly blended with ISM absorption. Therefore, searching for astrospheric absorption involves analysing the ISM properties observed toward the star as well.

²We shift from nm to Å in this section in order to facilitate comparisons with other studies analysing STIS/*HST* spectra.

Fig. 13 shows Fe II and Lyman α absorption observed towards *ι* Hor. The Fe II lines are the best available lines for studying the ISM velocity structure. Two well-separated velocity components are seen toward *ι* Hor. The two Fe II lines with rest wavelengths of 2586.650 and 2600.173 Å are fitted simultaneously, using procedures extensively used in the past (e.g. Redfield & Linsky 2008). Each absorption feature is defined by three parameters: a central wavelength (or velocity), a column density, and a Doppler parameter associated with the width of the line. The two Fe II velocity components have velocities of $v = 7.4$ and $v = 15.9$ km s⁻¹, logarithmic column densities (in cm⁻² units) of $\log N_{\text{Fe II}} = 12.53$ and $\log N_{\text{Fe II}} = 13.15$, and Doppler parameters of $b_{\text{Fe II}} = 1.94$ and $b_{\text{Fe II}} = 3.09$ km s⁻¹, respectively.

For Lyman α , the H I absorption is highly saturated and very broad (see Fig. 13). This is true for even the nearest stars with the lowest ISM column densities, but *ι* Hor seems a particularly extreme case for such a nearby star. Normally, narrow absorption from interstellar deuterium (D I) is observed to the left of the broad H I absorption, but for *ι* Hor even the D I absorption is saturated and the H I absorption is so broad that it is almost completely blended with D I. We fit the D I and H I absorption simultaneously, once again using procedures more extensively described elsewhere (e.g. Wood et al. 2005a, 2021). Although the two ISM components seen in Fe II are not separable in the much broader D I and H I absorption, we consider the two components in the Lyman α fit, constrained by the Fe II fit. In particular, we force the velocity separation and column density ratio to be the same, and for simplicity, we simply assume the two components have identical Doppler parameters. With these constraints, we measure an H I Doppler parameter of $b_H = 11.1$ km s⁻¹, and for the two components we measure H I column densities of $\log N_H = 18.09$ and $\log N_H = 18.72$, respectively.

Our ISM-only analysis fits the Lyman α data quite well, as shown in Fig. 13. We therefore see no evidence of astrospheric absorption toward *ι* Hor that would allow us to say something about the strength of its stellar wind. The ISM H I column density is much too high, and the resulting Lyman α absorption is too broad to detect the astrospheric absorption signature. The total ISM H I column density of $\log N_H = 18.81$ is in fact the third highest known within 25 pc (Wood et al. 2005a, 2021). The only higher values are toward HD 82558 ($d = 18.3$ pc, $\log N_H = 19.05$) and HD 203244 ($d = 20.8$ pc, $\log N_H = 18.82$).

4.3 Mg II H & K line

The wings, peaks, and cores of the spectral line Mg II H & K are considered reliable indicators for recovering activity information from the stellar photosphere, chromosphere, and near transition region. These lines are located in the near-ultraviolet (NUV) spectral range [$k = 2795.528$ Å and $h = 2802.704$ Å]. It is known that the line is sensitive to the strength of the magnetic field and the hydrodynamics conditions, helping to determine atmosphere conditions at a certain region. For example, in the Sun, the profile features at the base of the core line (see in Fig. 14, KW and HW) are formed in the photosphere (high 600 km, mid photosphere), while the core peaks are formed in the middle chromosphere (KPV and HPV at about high 1200 km and the KPR and HPR at about 1550 km, mid chromosphere) being very useful for calculating the velocities of inflows or outflows in the specific region of the atmosphere. The deep profiles between the core peaks (KC, HC) are known to be very sensitive and saturate during flaring events, they are formed in the upper chromosphere around 2200 km, about 200 km below the transition region (see Pereira et al. 2013; Leenaarts et al. 2013a, b; Schmit et al. 2015). We put in




Figure 11. Si III, Ly α , O V, N V, Si II and C II lines of t Hor at four *HST* visits compared with solar active (green) and quiet regions (yellow). Lines observed during visits I and II for cycle 25 are coloured in purple and blue, respectively. Lines observed during visits I and II for cycle 26 are coloured in red and rose, respectively. See additional UV lines in Section D of the appendix.




Figure 12. A comparison of C III and C IV smoothed lines obtained for *ι* Hor at four different epochs in *HST* Cycles 25 and 26 and active (green) and quiet (yellow) regions on the Sun. Lines observed during visits I and II for cycle 25 are coloured in purple and blue, respectively. Lines observed during visits I and II for cycle 26 are coloured in red and rose, respectively. A clear enhancement is seen in *ι* Hor during the second visit in Cycle 25, an indication of a possible flaring event.

Table 4. Measured flux densities for C III, C IV, Si IV, and O IV lines for visits I and II during Cycles 25 and 26 of STIS/*HST* observations compared with solar active and quiet regions (called AR and QR, respectively) observed by SUMMER/SOHO.

Flux density $10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$	C25-I	C25-II	C26-I	C26-II	Sun AR	Sun QR
$F_{\text{CIV}} 1548.2 \text{ \AA}$	9.47	13.09	8.79	9.48	4.31	1.46
$F_{\text{CIII}} 1175.7 \text{ \AA}$	4.53	6.55	3.98	4.43	1.83	1.74
$F_{\text{SiIV}} 1402.7 \text{ \AA}$	4.17	4.25	3.81	4.19	3.15	–
$F_{\text{OIV}} 1401.1 \text{ \AA}$	0.43	0.48	0.37	0.44	–	–

the stellar context these solar features to analyse the photospheric, chromospheric, and near transition region layers of *ι* Hor.

The relative difference between Mg II fluxes for the NUV orbits of cycles 25 and 26 suggests increased activity during cycle 25, as is also observed for the FUV lines. Even though higher activity was expected in cycle 26 than in cycle 25, following the predicted cycle trends from Ca and X-rays observed in Fig. 1, we observed a slightly enhanced flux in the NUV and FUV regions during cycle 25. We suggest the enhanced flux during cycle 25 could be associated with the presence of an active region (AR) and with elevated chromospheric activity (see characteristic AR shape during simultaneous *HST*-C25 observations in Fig. 15). Unfortunately, the time of observation of the Mg Line, during the first orbit of *HST*, was not simultaneously performed with the observation of the enhanced C Lines, during the third orbit of *HST*, C25II, as shown in Fig A3. Therefore, we do not observe saturation in the cores of Mg II H & K lines which indicates that the flaring event, suggested by the enhanced C IV and S IV lines in the FUV during the orbit II in C25, occurred after the first NUV orbit in C25. The relative intensity between the core peaks of Mg II H & K lines, are more prominent and easier to analyse than H & K cores of

Table 5. Pearson correlation coefficients between the different activity all indicators. Top: coefficients calculated taking the entire 199 data points obtained by HARPSpol. Bottom: coefficients calculated taking the 11 simultaneous data points obtained by HARPSpol.

	S_{HK}	$I_{\text{H}\alpha}$	B_{ℓ}	$ B_{\ell} $	RV
All data set:					
S_{HK}	1.00	0.82	0.025	−0.034	−0.015
$I_{\text{H}\alpha}$	0.82	1.00	0.046	−0.021	−0.056
B_{ℓ}	0.025	0.046	1.00	−0.26	0.042
$ B_{\ell} $	−0.034	−0.021	−0.26	1.00	0.045
RV	−0.015	−0.056	0.042	0.045	1.00
Simultaneous Obs.:					
S_{HK}	1.00	0.57	−0.095	0.28	−0.26
$I_{\text{H}\alpha}$	0.57	1.00	0.36	−0.059	−0.40
B_{ℓ}	−0.095	0.36	1.00	−0.34	0.32
$ B_{\ell} $	0.28	−0.059	−0.34	1.00	0.184
RV	−0.26	−0.40	0.32	0.184	1.00

Ca II in C25 and C26. Following Avrett (2013), the relative difference between Mg II H & K core lines indicates an averaged stellar outflow.

5 SIMULTANEOUS OBSERVATIONS

In this section, we focus on the data acquired simultaneously during the campaign. These correspond to the ESO P101C HARPSpol spectropolarimetric run (2018 August–2018 September), the Sector 28 observations from *TESS* (starting on 2018 August 23 until 2018 September 20 and overlapping from August 29 to September 16), and the coordinated Cycle 25 *HST*/STIS observations (2018 September 03). This combined data set allows us to compare the various manifestations of the magnetism of *ι* Hor, at different layers of the stellar atmosphere, on a sub-rotational period time-scale.




Figure 13. Fits to ISM absorption lines observed by *HST* toward ι Hor with two Fe II lines shown on the left, and the H I + D I Lyman α line shown on the right. The Fe II spectra are shown on a heliocentric velocity scale. The data are shown as red histograms. The thin black solid lines above the absorption are the assumed stellar emission background. Most of the stellar H I Lyman α emission is absorbed by the ISM. Two distinct Fe II velocity components are observed, and the combined absorption of the fit is shown both before (dotted lines) and after (thick solid lines) instrumental smoothing. The two components are too blended to be resolved in the Lyman α spectrum, but the Lyman α fit includes both components, with parameters constrained by the Fe II fit. The individual components are shown as dotted lines, and their combined absorption after instrumental smoothing is the thick solid line that fits the data.




Figure 14. Mg II H & K profiles observed by *HST*/STIS for the NUV single orbit during cycles 25 and 26. We describe the different features of the Mg II H & K line as from the wavelengths on the violet side of the H & K lines, we denote the minimum of the base wing as HWV and KWV, the first peak HPV and KPV. The core of the lines HC and KC. The second peak is located on the red side with respect to the core, or central depression of the line, as HPR and KPR. Finally, the minimum of the wing base is located on the red side with respect to the line core as HKR and KWR.

The different panels of Fig. 15 show the temporal evolution of various observables during this particular run. This includes photometry from *TESS*, as well as chromospheric/photospheric activity indicators (S_{HK} , $I_{H\alpha}$), radial velocity (RV), and longitudinal

magnetic field measurements (B_{ℓ}) from the HARPSpol spectra. We also indicate the specific date of our *HST*/STIS visit probing the transition region and corona. While a detailed Zeeman-Doppler Imaging (ZDI) analysis of ι Hor will be presented in a forthcoming




Figure 15. Simultaneous observations of *ι* Hor. The top panel shows the normalized light-curve section [1362.5 to 1379.5 BTJD d] obtained during S2 by the *TESS* satellite. The following panels show the simultaneous longitudinal magnetic field B_l , the $H\alpha_{\text{index}}$, the flux of calcium line cores, Ca II H & K, expressed in terms of the Ca II H & K S_{HK} , and radial velocities RVs. All values obtained from observations by the ESO/HARPSpol instrument for the EPOCH-P101C under the *Far beyond the Sun* campaign. RV error values are in the order of 2 m s^{-1} , which makes it difficult to resolve in the scale of Km s^{-1} , go to Table. C1 2 to see the RV error and S/N values per observation night. The green-bisque dash lines indicate 3 simultaneous observations of the NUV & FUV regions obtained by the *HST* STIS/NUV-MAMA instrument, with E230H and E140M gratings, respectively.

paper, we complement the aforementioned time series description with the corresponding ZDI magnetic field reconstruction associated with this epoch (Figs 16 and 17). The ZDI reconstruction follows the implementation described by Hussain et al. (2000), Hussain, Jardine & Collier Cameron (2001), Hussain et al. (2002), assuming a $P_{\text{rot}} = 7.7\text{ d}$ (Section 3.1.1) and an inclination angle of 60° . Despite the relatively good phase coverage of this epoch, it was not sufficient for an estimate of the differential rotation profile of the star. Therefore, solid-body rotation was assumed for the ZDI map shown in Figs 16. A Milne–Eddington local line profile was used which was tailored to a high S/N Stokes I spectrum of the star, together with a linear limb-darkening law coefficient set to 0.74 (appropriate for the spectral type of *ι* Hor and the bandpass of HARPSpol, see Gray 2005). The code uses a maximum-entropy regularization approach by Skilling & Bryan (1984) to guarantee the uniqueness of the solution. The local Stokes V line profile model is obtained from the Stokes I Milne–Eddington profile under the weak-field approximation (i.e. $V(\lambda) \propto \partial I / \partial \lambda$). Following the methodology described in Alvarado-Gómez et al. (2015), a grid of maximum-entropy ZDI solutions, with their corresponding fits to the LSD Stokes V profiles, is constructed from which the optimal reduced $\chi^2_{\text{ZDI}} = 1.20$ is obtained (see Fig. 17). The model is fitted to 704 data points in total, corresponding to 64 points per LSD Stokes V profile of each of the 11 consecutive nights of observation. The final map is presented in Fig 16, displaying the three components of the large-scale magnetic field (Radial: colour-scale, Meridional, and Azimuthal: vectors) in a latitude–longitude Mercator projection. The phase coverage is represented

by tick marks in the upper x-axis, labelled by the corresponding observing night of the run as indicated in the bottom panel of Fig. 15.

As can be seen by comparing Figs 15 and 16, the first night of observations revealed a predominant negative polarity in the large-scale field (see panel for B_l) associated with the visible pole of the star. The next three nights were lost owing to bad weather conditions in La Silla, which unfortunately coincided with our coordinated *HST*/STIS visit. Still, the photometric variability displayed an increase of ~ 0.2 per cent with respect to the normalized level, which could be associated with the presence of some flaring activity over those phases as registered in the FUV/NUV spectra (Section 4). That region of the ZDI map (retrieved mostly with the information provided one rotation later by nights 7 and 8) reveals relatively weak magnetic fields towards lower latitudes ($\lesssim 10\text{ G}$) with the appearance of a positive polarity region registered during the second observed night. From there and for the following 5 nights both activity indicators, S_{HK} and $I_{\text{H}\alpha}$, slowly increase to reach a maximum on night 7 of the run. After a small gap, the *TESS* light curve shows a dip at the 0.1 per cent around night 6, indicating the presence of spots at that particular phase. This agrees with the distribution obtained on this region of the ZDI map, which shows the strongest low-latitude mixed-polarity regions of the reconstruction, reaching up to 20 G in magnitude (nights 5 and 6). Such local changes in the field polarity are clearly visible in the sinusoidal behaviour of B_l during those observed phases (Fig. 15, second panel).

From the beginning of the run, the RV remains relatively stable, except for night 7 (2018 September 10 to 11) where it goes below the




Figure 16. Zeeman doppler imaging (ZDI) reconstruction of the large-scale magnetic field of ι Hor based on HARPSpol data acquired from 2018 September 02 to 16 (P101C). The colour scale shows the radial component of the magnetic field (B_R) in Gauss, while vectors trace the azimuthal (B_A) and meridional (B_M) fields normalized in size to their magnitude $(B_A^2 + B_M^2)^{1/2} = 17.7$ G. Two consecutive rotations were employed to retrieve the ZDI map, whose phase coverage is indicated by the black and red tick marks in the upper x-axis, labelled by the night of acquisition during the run (see Fig. 15). The horizontal dashed line corresponds to the visibility limit imposed by the inclination of the star ($\sim 60^\circ$).




Figure 17. Sequence of LSD Stokes V (left) and N (right) profiles used to reconstruct the ZDI map presented in Fig. 16. Observations (black symbols with errors), fits provided by the ZDI magnetic field distribution (red), and the achieved optimal reduced $\chi_{\text{ZDI}}^2 = 1.20$ are shown. The data in both panels have been enhanced by a constant factor (5×10^3) for visualization purposes. We include the observing phase (ϕ) for each night, taking the first night of this epoch as $\phi = 0.0$ (BJD = 2458 363.854 83) and using a $P_{\text{ROT}} = 7.7$ d. The blue line in the right-hand panel indicates the zero level.

long-term average given in Table 1. As can be seen from Table C1 and Fig. 15, this could be mainly due to the relatively higher noise in the spectrum of that night, resulting also in larger uncertainties for all the derived quantities. However, the presence of a spot group around nights 5–6 (2018 September 8 to 9), as can be contrasted with *TESS* photometry, could have also contributed to this relatively large drift in the RV.

Bad weather prevented observations on September 11–12, over which the photometry showed a relatively stable behaviour close to the nominal level. After this gap, the RV appears slightly lower during nights 9 to 11 than at the beginning of the run but still within the uncertainty limits for its long-term average. In contrast, B_ℓ , S_{HK} , and $I_{\text{H}\alpha}$ display very similar behaviour as during nights 3 to 5, indicating a relatively stable magnetic and activity configuration on the stellar surface over the course of the two observed rotations. One example of this is the polarity flip and relatively quick variation of B_ℓ during

these three nights (by roughly 6 G), which precede by approximately half a day a dip in the photometric light curve indicating the presence of a starspot group coming into view (see top panel of Fig. 15 around BTJD 1369.5 and 1378.5). This combined data set illustrates that while the large-scale field remains stable over two rotation periods (as demonstrated both in the B_ℓ values and in our combined map) the changing light curve and RV values hint at substantial continuous flux emergence happening on smaller spatial scales but still sufficient to modify the brightness of the star and the shape of the intensity profile.

6 DISCUSSION AND CONCLUSIONS

As presented in Paper I, ι Hor constitutes the closest located star to the intersection between the cycle branches emerging in the $P_{\text{rot}}-P_{\text{cyc}}$ diagram (Brandenburg, Saar & Turpin 1998; Böhm-Vitense 2007; Metcalfe et al. 2016). Given that the active (A) and inactive (I) cycle

periods are similar, a beating pattern is expected in the temporal evolution of different activity indicators. This is indeed observed in *ι* Hor, where the intense spectropolarimetric monitoring of the star, combined with archival observations, allowed us to refine and retrieve $P_{\text{cyc}}^A = 1.499 \pm 0.012$ and $P_{\text{cyc}}^I = 1.097 \pm 0.023$ yr. In addition, our analysis of the rotation period of *ι* Hor (Section 3) moves the star even closer to the active–inactive branches intersection, with a P_{rot} close to 8 d (refined from previous estimates at $P_{\text{rot}} > 8$ d, see e.g. Metcalfe et al. 2010). Note also that, for sparsely sampled observations (or broader binning of the available data), the beat period of $P_{\text{beat}} \simeq 4.49$ yr will dominate over the shortest periodicity (see e.g. Flores et al. 2017). Similarly, the ~ 1.97 yr period identified in Paper I was not recovered in the analysis of the whole data set, mainly due to the tighter constraints on shorter time-scales placed by the 98 additional spectropolarimetric observations to the 2-component fit (see Table 2).

The location of *ι* Hor in the spot-dominated regime over rotation time-scales ($S_{\text{fac}}/S_{\text{spot}} \sim 0.51$) is in line with its higher level of activity with respect to the Sun, reflecting its youth and relative rapid rotation. The Sun is closer to the faculae-dominated stars (towards the bottom right in Fig. 10) close to finalizing its transition from the spot-dominated (left) to the faculae-dominated stars (right). Amazo-Gómez et al. (2020a) found that the Sun displays a roughly constant $S_{\text{fac}}/S_{\text{spot}} \sim 3$ along its activity cycle. It is therefore expected that *ι* Hor would also have a stable $S_{\text{fac}}/S_{\text{spot}}$ value across its activity cycles, although further observations would be needed to confirm this.

As discussed in Section 4.2, our search for an astrospheric signal from *ι* Hor was hampered by the extremely large interstellar absorption along that line of sight (resulting in the third largest known column density within 25 pc at $\log N_{\text{H}} = 18.81$). Given the coronal activity levels of the star ($F_{\text{X}} \simeq 3.5\text{--}10.0 \times 10^5 \text{ erg s}^{-1} \text{ cm}^{-2}$), and following the mass loss–activity relation proposed for Sun-like stars ($\dot{M}_{\star} \propto F_{\text{X}}^{1.34}$, Wood et al. 2005b), the stellar wind mass-loss rate of *ι* Hor is expected to somewhere between the range $\dot{M}_{\star} \sim 9\text{--}80 \dot{M}_{\odot}$ over the course of its activity cycle. Note, however, that in periods of high activity, the X-ray flux of *ι* Hor approaches the observed break in the mass-loss–activity relation ($F_{\text{X}} \sim 10^6 \text{ erg s}^{-1} \text{ cm}^{-2}$), so that the actual \dot{M}_{\star} value could be much lower than the expectation. The results for the FUV/NUV line fluxes described in Sections 4.1 and 4.3, the emission measure analysis presented by Sanz-Forcada et al. (2019), and the different ZDI maps retrieved for the star, will be used in a forthcoming study to investigate numerically the stellar wind and planetary environment of this system (e.g. Alvarado-Gómez et al. 2016a, b).

In this study, we analysed different atmospheric layers of *ι* Hor by probing activity indicators from the photosphere to the lower corona. For this analysis, we have performed a detailed multiwavelength and multitechnique characterization of the star.

(i) We analyse the long-term variability in magnetic activity sensitive diagnostics derived using HARPS data acquired both from archival spectroscopic data and HARPS spectropolarimetric data from our programme, completing the observations and analysis started in Paper I. For the activity diagnostics, we use high signal-to-noise LSD profiles to derive time series measurements of photospheric activity, including the longitudinal magnetic field (B_{ℓ}). For the chromospheric activity, we quantify variability in the Ca II H & K and H α emission profiles using activity indices S_{HK} and $I_{\text{H}\alpha}$. The S_{HK} indices from our HARPSpol data were supplemented with archival HARPS data, spanning ten years in total.

(ii) A period analysis of the full S_{HK} data set revealed two dominant periods, indicating the presence of two overlapping chro-

mospheric activity cycles, with periods of 547.65 ± 3 d (1.49 y) and 401.01 ± 7 d (1.09 y), respectively. Sanz-Forcada et al. (2013, 2019) reported a 588.5 d (1.6 yr) cycle in both X-ray and Ca II H & K but on including the last 1.5-yr Ca II H & K data from our campaign, we found that the dominant periodicity in the chromospheric cycle (1.49-yr) is close aligned with the 1.6-yr X-ray cycle. This is relatively similar to the Sun, in which chromospheric and coronal activity cycles are aligned. The slight difference may be due to a geometrical/inclination effect given by the misalignment of the observed features in the low chromosphere and the corona, where features located near the limbo edge can be traced in the corona but less clearly in the photosphere or chromosphere. As one hemisphere of the star is obscured, if the active regions migration over the course of the activity cycle spans a wider range of latitudes than on the Sun, this could result in the double period that we find for *ι* Hor. Alternatively, the overlapping chromospheric cycles may be due to activity on the different hemispheres evolving differently.

(iii) We find a relatively strong correlation between the two chromospheric activity indicators, S_{HK} and $I_{\text{H}\alpha}$ ($\rho_{S_{\text{HK}}, I_{\text{H}\alpha}} = 0.82$). We noted no correlation between S_{HK} and longitudinal magnetic field B_{ℓ} , (Pearson correlation coefficient $\rho_{B_{\ell}, S_{\text{HK}}} = 0.025$). This is likely a consequence of the different length scales being probed – B_{ℓ} probes the large-scale photospheric magnetic field whereas S_{HK} is sensitive to disc-integrated chromospheric activity, including small-scale active regions.

(iv) We apply seven different techniques to evaluate *ι* Hor’s rotation period using *TESS* data and find a broad degree of agreement within the range 6.6 to 7.9 d. We can refine these measurements further using Stokes V spectra when optimizing stellar parameters for Zeeman–Doppler imaging – as the large-scale field does not vary as quickly as photometry (see Fig. 15) this enables a more robust measurement of the stellar rotation period.

(v) We implemented the novel Gradient of Power Spectra analysis (GPS) over time series of brightness variations. We estimated the stellar rotation periodicity of 7.78 ± 0.18 from magnetic features modulation for *ι* Hor. Furthermore, we performed a characterization of the ratio between dark spots and bright faculae features by analysing the location of gradients in the high-frequency tail of the power spectra. We analyse the location of the star in the context of the faculae to spot branching exposed in Shapiro et al. (2020); Amazo-Gómez et al. (2020a, b); Amazo-Gómez (2021) and compared it with the solar value. We derive a facular-to-spot ratio of 0.510 ± 0.023 for *ι* Hor, considerably more spot-dominated than the Sun which has $S_{\text{fac}}/S_{\text{spot}} \sim 3$ throughout its activity cycle.

(vi) We present a multiwavelength analysis of *ι* Hor bringing together optical photometry and spectropolarimetry with FUV and NUV *HST*/*STIS* spectra. The variability in the B_{ℓ} over the 17-night span of the data set is consistent with rotational modulation caused by the presence of unevenly distributed active regions that remain relatively stable over the timespan of observations. Similarly, the S_{HK} and $I_{\text{H}\alpha}$ indices do not show any significant change beyond rotational modulation – at least within the noise level of these measurements. In contrast, the *TESS* light curve clearly shows variability which cannot be explained by rotational modulation alone, indicating the emergence of new flux. There is indication that the RV also shows some evolution from one rotation cycle to the next but the effect is at the $2\text{--}3\sigma$ level so more precise measurements would be needed to confirm definitively to what extent the flux emergence that is detected in the *TESS* light curve could affect RV measurements.

(vii) We analysed and compare four different *HST*/*STIS* spectra observed during cycles 25 and 26. We observed a clearly enhanced

flux during the second orbit of C25 which suggests the possible association with a flaring event. In the near- and far- ultraviolet (NUV and FUV, respectively), we analysed the inter-combination lines such as C III, C IV, S IV, or O IV, which can be good tracers for characterizing the high chromosphere and coronal transition region. Those activity indicators trace the influence of the stellar magnetic field as it emerges from the stellar interior and weaves its way through the different atmospheric layers of the star.

The analysis here presented of ι Hor data allowed for the recovery of the ZDI maps for the entire campaign, which is part of a forthcoming publication. In addition, those maps have been used to model the geometry of magnetic regions in the stellar surface that extends into the chromosphere, corona, and beyond (e.g. Cohen & Drake 2014; Alvarado-Gómez et al. 2019b; Garraffo et al. 2022). This has been done by applying a detailed 3D magnetohydrodynamics (MHD) code (BATS-R-US; Powell et al. 1999), originally developed and validated for the solar wind and corona (e.g. Sokolov et al. 2013; van der Holst et al. 2014). This numerical approach includes all the relevant physics for calculating a coronal/wind model, having the surface magnetic field maps as a driver of a steady-state solution for the star. Coronal heating and stellar wind acceleration are self-consistently calculated via Alfvén wave turbulence dissipation in addition to radiative cooling and electron heat conduction (see Alvarado-Gómez et al. 2022). Models for the different phases of the magnetic cycle are generated and compared directly with the observations. The analysed observations are also used to predict the conditions experienced by the exoplanet during its orbit and through the magnetic cycle of its parent star. Moreover, the impact on habitability conditions due to the host-star magnetic field (and its evolution) is also studied in detail and will be published after the current manuscript.

ACKNOWLEDGEMENTS

We thank the referee for careful reading of the manuscript and constructive comments which improved the original version of the manuscript. Based on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere under ESO programmes 096.D-0257, 097.D-0420, 098.D-0187, 099.D-0236, 0100.D-0535, and 0101.D-0465. Data were obtained from the ESO Science Archive Facility under request number jalvarad.212739. This work has made use of the VALD data base, operated at Uppsala University, the Institute of Astronomy RAS in Moscow, and the University of Vienna. This research has made use of the SIMBAD data base, operated at CDS, Strasbourg, France. This research has made use of NASA's Astrophysics Data System. EMAG and JDAG were partially supported by *HST* GO-15299 and GO-15512 grants. EMAG and KP acknowledge support from the German *Leibniz-Gemeinschaft* under project number P67/2018. FDS acknowledges support from the Marie Skłodowska-Curie Actions (MSCA, grant agreement 101030103). JSF acknowledges support from the Agencia Estatal de Investigación of the Spanish Ministerio de Ciencia, Innovación y Universidades through grant PID2019-109522GB-C51.

DATA AVAILABILITY

The data underlying this article were accessed from different large-scale facilities. The derived data from this research will be shared upon reasonable request to the corresponding author. For the analysis of the activity indicators, longitudinal magnetic field, and radial velocity evolution presented we

included pipeline-processed spectroscopic HARPS PH3, http://archive.eso.org/wdb/wdb/adp/phase3_main/form and the HARPS-Polarimetry pipeline processed data query form <http://archive.eso.org/wdb/wdb/eso/repro/form>. *TESS* light curves and, *HST*/STIS data can be obtained from the High-Level Science Products (HLSP) on the Mikulski Archive for Space Telescopes (MAST, <http://archive.stsci.edu/>). The solar comparison data were of a quiet region on 1997 April 20, and the active region AR-NOAA8487 on 1999 March 18 (see Curdt et al. 2001, 2004) and are available at Werner Curdt's homepage <https://www2.mps.mpg.de/homes/curdt/>. Optimistic and conservative estimates place the inner boundary of the habitable zone (HZ) of ι Hor at 0.94 and 1.20 au, respectively available at <http://www.hzgallery.org> (Kane & Gelino 2012).

REFERENCES

- Agarwal S., Del Sordo F., Wettlaufer J. S., 2017, *AJ*, 153, 12
- Alvarado-Gómez J. D. et al., 2015, *A&A*, 582, A38
- Alvarado-Gómez J. D., Hussain G. A. J., Cohen O., Drake J. J., Garraffo C., Grunhut J., Gombosi T. I., 2016a, *A&A*, 588, A28
- Alvarado-Gómez J. D., Hussain G. A. J., Cohen O., Drake J. J., Garraffo C., Grunhut J., Gombosi T. I., 2016b, *A&A*, 594, A95
- Alvarado-Gómez J. D. et al., 2018, *MNRAS*, 473, 4326
- Alvarado-Gómez J. D. et al., 2019a, in *Solar Heliospheric and Interplanetary Environment (SHINE 2019)*, p. 110
- Alvarado-Gómez J. D., Garraffo C., Drake J. J., Brown B. P., Oishi J. S., Moschou S. P., Cohen O., 2019b, *ApJ*, 875, L12
- Alvarado-Gómez J. D. et al., 2022, *ApJ*, 928, 147
- Amazo-Gómez E. M., 2021, *Understanding the Brightness Variations of Sun-like Stars on Timescales of Stellar Rotation*. Georg-August-Universität Göttingen
- Amazo-Gómez E. M., Shapiro A. I., Solanki S. K., Krivova N. A., Kopp G., Reinhold T., Oshagh M., Reiners A., 2020a, *A&A*, 636, A69
- Amazo-Gómez E. M., Shapiro A. I., Solanki S. K., Kopp G., Oshagh M., Reinhold T., Reiners A., 2020b, *A&A*, 642, A225
- Anglada-Escudé G., Butler R. P., 2012, *ApJS*, 200, 15
- Anthony F. et al., 2022, *AJ*, 163, 257
- Avrett E. H., Landi E., McKillop S., 2013, *ApJ*, 779, 155
- Bagnulo S., Landolfi M., Landstreet J. D., Landi Degl'Innocenti E., Fossati L., Sterzik M., 2009, *PASP*, 121, 993
- Baliunas S. L. et al., 1995, *ApJ*, 438, 269
- Barnes S. A., 2007, *ApJ*, 669, 1167
- Barros S. C. C., Demangeon O., Díaz R. F., Cabrera J., Santos N. C., Faria J. P., Pereira F., 2020, *A&A*, 634, A75
- Benedict G. F. et al., 2006, *AJ*, 132, 2206
- Böhm-Vitense E., 2007, *ApJ*, 657, 486
- Boro Saikia S. et al., 2016, *A&A*, 594, A29
- Boro Saikia S. et al., 2018, *A&A*, 620, L11
- Boro Saikia S. et al., 2022, *A&A*, 658, A16
- Borucki W. J. et al., 2010, *Sci.*, 327, 977
- Brandenburg A., Saar S. H., Turpin C. R., 1998, *ApJ*, 498, L51
- Brasseur C. E., Phillip C., Fleming S. W., Mullally S. E., White R. L., 2019, *Astrophysics Source Code Library*, record ascl:1905.007
- Bruntt H. et al., 2010, *MNRAS*, 405, 1907
- Coffaro M. et al., 2020, *A&A*, 636, A49
- Cohen O., Drake J. J., 2014, *ApJ*, 783, 55
- Curdt W., Brekke P., Feldman U., Wilhelm K., Dwivedi B. N., Schühle U., Lemaire P., 2001, *A&A*, 375, 591
- Curdt W., Landi E., Feldman U., 2004, *A&A*, 427, 1045
- Damasso M., Del Sordo F., 2017, *A&A*, 599, A126
- de Almeida L., Anthony F., do Nascimento J. D., 2020, in *Proc. XLIII SAB Annual Meeting*, p. 80
- Donati J.-F., Semel M., Carter B. D., Rees D. E., Collier Cameron A., 1997, *MNRAS*, 291, 658
- Donati J. F. et al., 2008, *MNRAS*, 385, 1179

- Drake J. J., Smith G., 1993, *ApJ*, 412, 797
- Drake J. J., Peres G., Orlando S., Laming J. M., Maggio A., 2000, *ApJ*, 545, 1074
- Dumusque X., Santos N. C., Udry S., Lovis C., Bonfils X., 2011, *A&A*, 527, A82
- Dumusque X. et al., 2012, *Nature*, 491, 207
- Eberhard G., Schwarzschild K., 1913, *ApJ*, 38
- Fares R. et al., 2009, *MNRAS*, 398, 1383
- Farge M., 1992, *Annu. Rev. Fluid Mech.*, 24, 395
- Flores M. G., Buccino A. P., Saffe C. E., Mauas P. J. D., 2017, *MNRAS*, 464, 4299
- Foreman-Mackey D., 2018, *Res. Notes Am. Astron. Soc.*, 2, 31
- Foreman-Mackey D., Agol E., Ambikasaran S., Angus R., 2017, *AJ*, 154, 220
- Foreman-Mackey D., Czekala I., Agol E., Luger R., Barclay T., 2019, *dfm/exoplanet: exoplanet v0.2.4*, Zenodo
- García R. A. et al., 2009, *A&A*, 506, 41
- García R. A. et al., 2014, *A&A*, 572, A34
- Garraffo C. et al., 2018, *ApJ*, 862, 90
- Garraffo C., Alvarado-Gómez J. D., Cohen O., Drake J. J., 2022, *ApJ*, 941, L8
- Gizis J. E., Reid I. N., Hawley S. L., 2002, *AJ*, 123, 3356
- Gray D. F., 2005, *PASP*, 117, 711
- Gull T. R. et al., 1986, in Rolfe E. J., Wilson R., eds, *ESA Special Publication Vol. 263, New Insights in Astrophysics. Eight Years of UV Astronomy with IUE*. p. 653
- Hackman T., Kochukhov O., Viviani M., Warnecke J., Korpi-Lagg M. J., Lehtinen J. J., 2023, preprint ([arXiv:2306.07838](https://arxiv.org/abs/2306.07838))
- Hall J. C., 2008, *Living Rev. Sol. Phys.*, 5, 2
- Hathaway D. H., 2015, *Living Rev. Sol. Phys.*, 12, 4
- Hatzes A. P., 2013, *ApJ*, 770, 133
- Hatzes A. P. et al., 2000, *ApJ*, 544, L145
- Hussain G. A. J., Donati J.-F., Collier Cameron A., Barnes J. R., 2000, *MNRAS*, 318, 961
- Hussain G. A. J., Jardine M., Collier Cameron A., 2001, *MNRAS*, 322, 681
- Hussain G. A. J., van Ballegoijen A. A., Jardine M., Collier Cameron A., 2002, *ApJ*, 575, 1078
- Janson M., Reffert S., Brandner W., Henning T., Lenzen R., Hippler S., 2008, *A&A*, 488, 771
- Jardine M., Collier Cameron A., Donati J. F., Hussain G. A. J., 2020, *MNRAS*, 491, 4076
- Jeffers S. V., Petit P., Marsden S. C., Morin J., Donati J.-F., Folsom C. P., 2014, *A&A*, 569, A79
- Jeffers S. V., Boro Saikia S., Barnes J. R., Petit P., Marsden S. C., Jardine M. M., Vidotto A. A., *BCool Collaboration*, 2017, *MNRAS*, 471, L96
- Kane S. R., Gelino D. M., 2012, *PASP*, 124, 323
- Kochukhov O., Makaganiuk V., Piskunov N., 2010, *A&A*, 524, A5
- Kürster M., Endl M., Els S., Hatzes A. P., Cochran W. D., Döbereiner S., Dennerl K., 2000, *A&A*, 353, L33
- Labonte B. J., 1986, *ApJS*, 62, 241
- Laming J. M., Drake J. J., Widing K. G., 1996, *ApJ*, 462, 948
- Leenaarts J., Pereira T. M. D., Carlsson M., Uitenbroek H., De Pontieu B., 2013a, *ApJ*, 772, 89
- Leenaarts J., Pereira T. M. D., Carlsson M., Uitenbroek H., De Pontieu B., 2013b, *ApJ*, 772, 90
- Li K. J., Xu J. C., Feng W., 2022, *Sci. Rep.*, 12, 15877
- Lomb N. R., 1976, *Ap&SS*, 39, 447
- Marsden S. C. et al., 2014, *MNRAS*, 444, 3517
- Mayor M. et al., 2003, *The Messenger*, 114, 20
- McQuillan A., Aigrain S., Mazeh T., 2013, *MNRAS*, 432, 1203
- McQuillan A., Mazeh T., Aigrain S., 2014, *ApJS*, 211, 24
- Meschiari S., Wolf A. S., Rivera E., Laughlin G., Vogt S., Butler P., 2009, *PASP*, 121, 1016
- Metcalfe T. S., Basu S., Henry T. J., Soderblom D. R., Judge P. G., Knölker M., Mathur S., Rempel M., 2010, *ApJ*, 723, L213
- Metcalfe T. S., Egeland R., van Saders J., 2016, *ApJ*, 826, L2
- Moore C. E., Minnaert M. G. J., Houtgast J., 1966, *The Solar Spectrum 2935 Å to 8770 Å*. National Bureau of Standards Monograph. US Government Printing Office, Washington
- Morgenthaler A. et al., 2012, *A&A*, 540, A138
- Ness J.-U., Jordan C., 2008, *MNRAS*, 385, 1691
- Pagano I., Linsky J. L., Valenti J., Duncan D. K., 2004, *A&A*, 415, 331
- Pakhomov Y., Piskunov N., Ryabchikova T., 2017, in Balega Y. Y., Kudryavtsev D. O., Romanyuk I. I., Yakunin I. A., eds, *ASP Conf. Ser. Vol. 510, Stars: From Collapse to Collapse*. Astron. Soc. Pac., San Francisco, p. 518
- Pepe F. et al., 2005, *The Messenger*, 120, 22
- Pereira T. M. D., Leenaarts J., De Pontieu B., Carlsson M., Uitenbroek H., 2013, *ApJ*, 778, 143
- Peter H., 2001, *A&A*, 374, 1108
- Piskunov N., Kochukhov O., 2002, *A&A*, 381, 736
- Piskunov N. et al., 2011, *The Messenger*, 143, 7
- Powell K. G., Roe P. L., Linde T. J., Gombosi T. I., De Zeeuw D. L., 1999, *J. Comput. Phys.*, 154, 284
- Ramírez I., Yong D., Gutiérrez E., Endl M., Lambert D. L., Do Nascimento J. D. J., 2017, *ApJ*, 850, 80
- Redfield S., Linsky J. L., 2008, *ApJ*, 673, 283
- Reiners A., 2012, *Living Rev. Sol. Phys.*, 9, 1
- Reiners A., Mohanty S., 2012, *ApJ*, 746, 43
- Reiners A., Schüssler M., Passegger V. M., 2014, *ApJ*, 794, 144
- Reiners A. et al., 2022, *A&A*, 662, A41
- Reinhold T., Shapiro A. I., Solanki S. K., Montet B. T., Krivova N. A., Cameron R. H., Amazo-Gómez E. M., 2020, *Science*, 368, 518
- Ricker G. R. et al., 2015, *J. Astron. Telesc. Instrum. Sys.*, 1, 014003
- Robrade J., Schmitt J. H. M. M., Favata F., 2012, *A&A*, 543, A84
- Ryabchikova T., Piskunov N., Kurucz R. L., Stempels H. C., Heiter U., Pakhomov Y., Barklem P. S., 2015, *Phys. Scr*, 90, 054005
- Saar S. H., Osten R. A., 1997, *MNRAS*, 284, 803
- Santerne A. et al., 2015, *MNRAS*, 451, 2337
- Santos A. R. G., García R. A., Mathur S., Bugnet L., van Saders J. L., Metcalfe T. S., Simonian G. V. A., Pinsonneault M. H., 2019, *ApJS*, 244, 21
- Sanz-Forcada J., Micela G., Ribas I., Pollock A. M. T., Eiroa C., Velasco A., Solano E., García-Álvarez D., 2011, *A&A*, 532, A6
- Sanz-Forcada J., Stelzer B., Metcalfe T. S., 2013, *A&A*, 553, L6
- Sanz-Forcada J., Stelzer B., Coffaro M., Raetz S., Alvarado-Gómez J. D., 2019, *A&A*, 631, A45
- Scargle J. D., 1982, *ApJ*, 263, 835
- Schmit D., Bryans P., De Pontieu B., McIntosh S., Leenaarts J., Carlsson M., 2015, *ApJ*, 811, 127
- Shapiro A. I., Amazo-Gómez E. M., Krivova N. A., Solanki S. K., 2020, *A&A*, 633, A32
- Skilling J., Bryan R. K., 1984, *MNRAS*, 211, 111
- Skumanich A., 1972, *ApJ*, 171, 565
- Sokolov I. V. et al., 2013, *ApJ*, 764, 23
- Thomas R. N., 1957, *ApJ*, 125, 260
- Torrence C., Compo G. P., 1998, *Bull. Am. Meteorol. Soc.*, 79, 61
- Uitenbroek H., 2003, *ApJ*, 592, 1225
- Valenti J. A., Fischer D. A., 2005, *ApJS*, 159, 141
- van der Holst B., Sokolov I. V., Meng X., Jin M., Manchester IV W. B., Tóth G., Gombosi T. I., 2014, *ApJ*, 782, 81
- Vauclair S., Laymand M., Bouchy F., Vauclair G., Hui Bon Hoa A., Charpinet S., Bazot M., 2008, *A&A*, 482, L5
- Vaughan A. H., Preston G. W., Wilson O. C., 1978, *PASP*, 90, 267
- Wargelin B. J., Saar S. H., Pojmański G., Drake J. J., Kashyap V. L., 2017, *MNRAS*, 464, 3281
- Wilson O. C., 1963, *ApJ*, 138, 832
- Wilson O. C., 1978, *ApJ*, 226, 379
- Wood B. E., Linsky J. L., Ayres T. R., 1997, *ApJ*, 478, 745
- Wood B. E., Müller H.-R., Zank G. P., Linsky J. L., 2002, *ApJ*, 574, 412
- Wood B. E., Redfield S., Linsky J. L., Müller H.-R., Zank G. P., 2005a, *ApJS*, 159, 118
- Wood B. E., Müller H.-R., Zank G. P., Linsky J. L., Redfield S., 2005b, *ApJ*, 628, L143

- Wood B. E. et al., 2021, *ApJ*, 915, 37
 Woodgate B. E. et al., 1986, in Crawford D. L., ed., *Proc. SPIE Conf. Ser. Vol. 627, Instrumentation in astronomy VI*. SPIE, Bellingham, p. 350
 Woodruff R. A., Ludtke C. W., Woodgate B. E., 1990, in *Bulletin of the American Astronomical Society*. p. 1283
 Wright N. J., Drake J. J., 2016, *Nature*, 535, 526
 Wright J. T., Marcy G. W., Butler R. P., Vogt S. S., 2004, *ApJS*, 152, 261
 Wright N. J., Drake J. J., Mamajek E. E., Henry G. W., 2011, *ApJ*, 743, 48
 Wright N. J., Newton E. R., Williams P. K. G., Drake J. J., Yadav R. K., 2018, *MNRAS*, 479, 2351
 Zechmeister M., Kürster M., 2009, *A&A*, 496, 577
 Zechmeister M. et al., 2013, *A&A*, 552, A78
 Zhou Y., Leung Y., 2010, *J. Stat. Mech.: Theory Exp.*, 2010, 06021

APPENDIX A: ROTATION PERIOD MEASUREMENTS - A COMPARISON OF MULTIPLE TECHNIQUES

As noted in the main paper, we have applied seven methods to measure photometric periodicities related to the stellar rotation period: Gaussian-process regression (GP), Generalized Lomb–Scargle periodogram (GLS), Autocorrelation Function (ACF), Wavelet Power Spectra (PS), and the Gradient of Power Spectra (GPS). In Figs A1 and A2, we show the results from four of these methods: GLS, ACF, PS, and GPS.

Our GP approach makes use of a quasi-periodic kernel (see equation 1 in Damasso & Del Sordo 2017) applied to sectors 2 and 3 of the *TESS* data for ι Hor. We quantified the rotation period of the star (parameter Θ) and a time-scale related to the correlation time of the fluctuations (λ), which should account for the average lifetime of active regions on the surface of the star. We used a rather large prior on the rotation period Θ and λ , allowing the system to analyse values of Θ between 3 and 9 d and correlations spanning a time-scale covering the whole S2 sector. Our results show that the correlation time-scale should be shorter than, or of the order of, the rotation period. In the S2 analysis, we obtained a rotation $\Theta = 6.64^{+0.28}_{-0.09}$ d, $\lambda = 4.7 \pm 0.3$ d, while for S3 we obtained $\Theta = 7.01^{+0.16}_{-0.09}$ d, $\lambda = 7.7 \pm 0.7$ d.

For comparison, we also performed a quasi-periodic GP modelling using a simple covariance function (see equation 3 in Barros




Figure A1. Rotation period analysis for ι Hor. The upper panel shows detrended and stitched *TESS* light curves from sectors 2, 3, 29, and 30. The PDCSAP light curve (e open symbols) clearly shows more variability than when extracting the light curve from full frame images, (FFI, orange, and blue LC for sectors 2–3 and 29–30, respectively). Bottom panel left: Lomb–Scargle periodogram analysis. Bottom panel right: face fold FFI extracted LC for 9.6 and 7.6 d.




Figure A2. The ι Hor rotation period analysis for the normalized and stitched *TESS* light curves from Sectors 2 and 3, from 2018 August 23 to 2018 October 17 in panel (A). Panels (B) and (C) show the corresponding Generalized Lomb–Scargle (GLS) periodogram and autocorrelation function (ACF), respectively. panel (D) shows the global wavelet power spectrum, PS, calculated with the 6th degree Paul wavelet. Panel (E) shows the gradient of the power spectrum plotted in panel (D). Red and blue dotted lines in panel (E) indicate the location of the high and low-frequency inflection points location, respectively.

et al. 2020), constructed using the PYTHON *exoplanet* package (Foreman-Mackey et al. 2019) with the following properties:

$$k(\tau) = \frac{B}{2+C} e^{-\tau/L} \left(\cos\left(\frac{2\pi\tau}{P_{\text{rot}}}\right) + (1+C) \right) + \sigma^2 \delta_{ij}, \quad (\text{A1})$$

where P_{rot} corresponds to the rotation period of the star, L is the length-scale of exponential decay, and $B > 0$, $C > 0$, and $L > 0$. We chose large prior normal distributions for each hyperparameter, specifically for the rotation period and the amplitude, letting the model test values of $\log P_{\text{rot}}$ within $\mathcal{N}(1.97, 0.5)$ (corresponding to 3.2–90 d) and of $\log B$ and $\log C$ within $\mathcal{N}(-12.31, 5)$ and $\mathcal{N}(-4.47, 5)$, respectively. The central values μ of the chosen prior normal distributions were primarily set by a quick optimization test. Then, combined with a Markov chain Monte-Carlo fit, the equivalent estimate of the peak-to-peak amplitude of the oscillation is represented by $2B/(2+C)$ and is valued at $3.23 \cdot 10^{-3}$. By using this kernel, the GP approach gives a fair estimation of the amplitude of the activity *jitter* in addition to a consistent determination of the rotation period for $P_{\text{rot}} = 7.23 \pm 0.20$ d compared to the QP kernel described above.

We also employed a Multifractional approach called a Multifractional Temporally Weighted Detrended Fluctuation Analysis (Zhou & Leung 2010; Agarwal, Del Sordo & Wettlaufer 2017). This technique is optimized for detecting relevant time-scales and employs a model-free approach, also giving insight into the noise properties (see section 2 of Agarwal et al. 2017, for a detailed explanation). By analysing the S2 and S3 data, we found that the data are dominated by white noise on short time-scales (up to about 1 h), while between 1 h to about 4 d, they show red noise.

It was possible to detect both time-scales associated with these changes in noisy behaviours, and time-scales associated with periodic features. In the S2 data, we observe a typical time of 4.1 d associated




Figure A3. Zoom into the S2 *TESS*-LC section [from 1362.5 to 1379.5 BTJD d] during 3 different *STIS/HST* visits for Cycle 25. The 3 *HST* orbits are organized for observing just ones the NUV region around the Mg II H & K line, and two visits for observing the FUV region nearby the Ly α line. *ι* Hor.

with the end of the red noise phase, and periodic dynamics at 7.48 d. In S3, we found time-scales of 4.41 and 7.25 d corresponding to the end of the red noise phase and clear periodic behaviour, respectively. The uncertainties on all of these measurements are approximately 0.15 d. When analysing the combined data, we also detected strong periodicity over 21.2 d.

We also applied the Lomb–Scargle periodogram (see Lomb 1976; Scargle 1982) to the S2 + S3 data for *ι* Hor, specifically the Generalized Lomb–Scargle periodogram (GLS) version v1.03, using the formalism given by Zechmeister & Kürster (2009). In panel (b) of Fig. A2 we show the output from GLS, giving a rotation period estimate of about 7.76 d.

Panel (c) of Fig. A2 shows the result from the autocorrelation function applied to S2 + S3 data and resulting in a rotation period of 7.69 d. This is based on estimating the degree of self-similarity in the light curve, see McQuillan et al. (2014).

The results from our application of a wavelet power spectra transform are shown in panel (d) of Fig. A2. This method is optimal for a time series with a non-stationary signal at many different frequencies. It has previously been employed for determining stellar rotation periods by García et al. (2009). We use the power spectra transform analysis based on the 6th order Paul wavelet, (see Farge 1992; Torrence & Compo 1998) and recover a rotation period of 7.94 days.

Finally, we apply the GPS method to the normalized and stitched S2 + S3 LCs. In Shapiro et al. (2020), Amazo-Gómez et al. (2020a, 2020b) they found that the gradient of the wavelet power spectra transform, and in particular, the position of the high-frequency inflection point (i.e. the inflection point, IP, with the highest frequency) is connected to the stellar rotation period. The rotation period is determined from the profile of the high-frequency tail of the smoothed wavelet power spectrum (using a Paul wavelet, order 6), i.e. its part in between about a day and a quarter of the rotation

period. In particular, they identify the point where the gradient of the power spectrum (GPS) plotted in log–log scale (i.e. $d \ln P(\nu) / (d \ln \nu)$, where P is the power-spectral density and ν is frequency) reaches its maximum value. That point corresponds to the inflection point, i.e. the concavity of the power spectrum, plotted in log–log scale, changes sign there. It was shown that the period corresponding to the inflection point is connected to the rotation period of a star by a calibration factor which is a function of stellar effective temperature, metallicity, and activity. They validated their new method against Kepler light curves and the total solar irradiance (TSI) data from the Sun. For *ι* Hor, we recovered a rotation period of 7.78 ± 0.18 d using this method.

APPENDIX B: TELLURIC LINES IN H α

Removed telluric lines from H α are coloured in red in Fig. 5. See the Rowland table of The solar spectrum book Moore et al. (1966), where the equivalent width values are taken.

Table B1. Removed telluric lines from H α are coloured in red in Fig. 5. See the Rowland table of The solar spectrum book Moore et al. (1966), where the equivalent width values are taken.

λ [nm]	W [mÅ]	λ [nm]	W [mÅ]
655.7857	1.5	656.4061	4.5
655.8149	7.0	656.4206	14.0
655.8650	1.5	656.5545	3.0
655.8955	1.5	656.7850	2.0
656.0555	22.0	656.8806	3.5
656.1097	5.0	656.8900	5.0
656.3521	4.5		

APPENDIX C: JOURNAL OF OBSERVATIONS

Table C1. Journal of observations (columns 1–4) and measurements for each night (columns 5–12).

BJD (+ 2400000.) [days]	S/N I [@551 nm]	S/N V [@551 nm]	Airmass	RV [m s ⁻¹]	S _{HK}	eS _{HK}	I _{Hα}	eI _{Hα}	B _{ℓ} [G]	$\sigma_{B_{\ell}}$ [G]	N _{ℓ} [G]
ESO EPOCH P99A											
57934.90224	710	676	1.340	16909.2 ± 2.5	0.2270	0.0060	0.51123	0.00139	-3.14	0.61	0.11
57935.91343	765	742	1.281	16902.6 ± 2.3	0.2254	0.0056	0.51165	0.00128	-2.67	0.56	0.26
57936.90045	592	572	1.324	16892.9 ± 2.7	0.2260	0.0074	0.51122	0.00164	-2.78	0.73	-1.09
57937.90757	739	706	1.283	16904.5 ± 2.4	0.2266	0.0058	0.51103	0.00134	-1.42	0.58	0.46
57938.89985	264	260	1.303	16907.0 ± 3.9	0.2092	0.0173	0.51171	0.00370	-2.27	1.63	1.75
57942.91940	803	765	1.198	16919.2 ± 2.3	0.2280	0.0052	0.51075	0.00124	-2.68	0.54	0.19
57943.92830	675	639	1.168	16911.4 ± 2.5	0.2246	0.0059	0.51137	0.00147	-0.92	0.63	0.17
57944.89264	721	696	1.268	16894.6 ± 2.4	0.2233	0.0058	0.51157	0.00135	-0.53	0.60	0.93
57945.92100	855	826	1.174	16903.8 ± 2.2	0.2239	0.0049	0.51153	0.00116	-1.16	0.50	-0.07
57946.91205	654	631	1.189	16928.3 ± 2.5	0.2259	0.0063	0.51178	0.00150	-2.54	0.66	-0.14
57948.89648	660	642	1.219	16905.9 ± 2.7	0.2312	0.0063	0.51168	0.00148	-1.33	0.66	0.01
57949.91318	335	309	1.167	16916.8 ± 3.8	0.2160	0.0121	0.51136	0.00293	-3.49	1.26	-0.45
ESO EPOCH P99B											
57961.89812	833	806	1.132	16934.1 ± 2.3	0.2363	0.0051	0.51221	0.00119	-0.48	0.52	-0.33
57962.90690	352	226	1.113	16951.4 ± 3.5	0.2303	0.0125	0.51293	0.00283	2.27	2.07	-2.26
57964.91010	964	932	1.102	16941.0 ± 2.1	0.2325	0.0044	0.51163	0.00103	-1.19	0.45	-0.10
57967.90436	350	335	1.099	16931.2 ± 3.6	0.2284	0.0122	0.51144	0.00279	3.97	1.22	-0.13
57968.93869	589	545	1.076	16949.8 ± 2.7	0.2325	0.0073	0.51203	0.00165	1.31	0.73	0.69
57970.92374	577	554	1.079	16959.7 ± 2.8	0.2413	0.0070	0.51274	0.00171	-0.58	0.74	1.01
57971.93274	517	500	1.076	16960.8 ± 2.9	0.2343	0.0068	0.51196	0.00198	-1.67	0.79	0.01
57972.92529	537	526	1.077	16942.7 ± 2.9	0.2322	0.0078	0.51162	0.00182	-2.93	0.80	1.17
57973.92259	466	421	1.077	16941.5 ± 3.1	0.2326	0.0095	0.51255	0.00208	-2.53	0.95	1.36
57974.93909	577	561	1.078	16942.5 ± 2.7	0.2345	0.0071	0.51214	0.00171	-1.44	0.73	1.35
ESO EPOCH P99C											
58013.88198	703	680	1.123	16991.1 ± 2.4	0.2344	0.0060	0.51107	0.00141	0.47	0.62	0.91
58014.88046	601	589	1.125	17005.1 ± 2.6	0.2430	0.0070	0.51187	0.00163	-2.54	0.71	0.35
58017.75162	638	603	1.123	16996.7 ± 2.5	0.2394	0.0068	0.51277	0.00151	-0.10	0.67	-0.35
58019.83871	836	809	1.090	16991.8 ± 2.2	0.2364	0.0052	0.51156	0.00117	-0.29	0.52	1.13
58020.84732	591	569	1.102	16987.1 ± 2.7	0.2336	0.0065	0.51121	0.00169	0.48	0.71	0.37
58021.87703	656	630	1.155	16985.8 ± 2.5	0.2360	0.0064	0.51123	0.00149	-0.73	0.66	-0.22
58022.79833	551	526	1.076	17001.7 ± 2.9	0.2442	0.0069	0.51303	0.00182	-1.19	0.76	-1.38
58023.82878	632	602	1.090	17000.1 ± 2.6	0.2439	0.0060	0.51329	0.00159	-2.25	0.66	-0.39
58024.76855	380	337	1.084	16999.0 ± 3.4	0.2457	0.0109	0.51331	0.00258	1.48	1.21	2.79
58025.83057	334	300	1.098	16992.1 ± 3.6	0.2458	0.0149	0.51305	0.00288	0.88	1.39	-0.44
58026.80399	561	524	1.079	17016.1 ± 2.8	0.2437	0.0068	0.51313	0.00178	0.76	0.75	-0.67
58027.80589	502	481	1.082	16994.7 ± 2.9	0.2392	0.0090	0.51246	0.00190	0.86	0.87	-0.64
58028.79881	575	545	1.079	16978.9 ± 2.8	0.2379	0.0077	0.51152	0.00168	-0.08	0.75	-0.57
ESO EPOCH P100A											
58069.60736	847	816	1.125	16946.7 ± 2.3	0.2488	0.0044	0.51332	0.00120	-5.46	0.51	-0.72
58070.60682	883	854	1.122	16923.2 ± 2.3	0.2420	0.0044	0.51232	0.00113	0.39	0.48	0.09
58071.64337	1021	998	1.081	16929.5 ± 2.1	0.2393	0.0037	0.51192	0.00101	0.59	0.42	-0.34
58073.63701	1117	1048	1.082	16929.7 ± 2.0	0.2393	0.0036	0.51190	0.00090	-0.67	0.39	0.43
58074.66443	958	925	1.078	16939.8 ± 2.2	0.2427	0.0043	0.51352	0.00104	-2.58	0.45	1.11
58075.64065	989	948	1.078	16950.3 ± 2.2	0.2492	0.0041	0.51411	0.00101	-1.44	0.44	0.49
58076.65331	643	603	1.075	16927.5 ± 2.7	0.2479	0.0065	0.51334	0.00154	-1.75	0.68	0.15
58077.65193	1067	1010	1.076	16936.1 ± 2.1	0.2408	0.0039	0.51213	0.00093	-0.47	0.41	0.35
58078.66084	1111	1049	1.079	16918.6 ± 2.0	0.2378	0.0038	0.51184	0.00091	1.50	0.39	-0.18
58079.65272	873	848	1.078	16928.3 ± 2.3	0.2380	0.0046	0.51227	0.00114	-1.30	0.49	-0.25
58081.65863	1070	1042	1.083	16935.0 ± 2.0	0.2476	0.0040	0.51312	0.00093	-0.42	0.41	-0.13
ESO EPOCH P100B											
58108.56994	715	679	1.076	16863.0 ± 2.7	0.2466	0.0055	0.51485	0.00142	2.19	0.60	-0.25
58109.62286	828	785	1.132	16900.3 ± 2.4	0.2381	0.0049	0.51406	0.00117	-0.21	0.51	-0.06
58110.61173	792	753	1.118	16926.9 ± 2.3	0.2433	0.0051	0.51472	0.00123	-2.55	0.54	0.10
58111.64169	681	654	1.185	16940.7 ± 2.6	0.2512	0.0058	0.51679	0.00143	-3.99	0.63	-0.19
58112.63716	874	838	1.180	16949.6 ± 2.2	0.2612	0.0050	0.51840	0.00111	-0.39	0.50	0.03
58113.60393	985	925	1.119	16907.1 ± 2.2	0.2657	0.0045	0.51839	0.00099	-0.51	0.44	0.28
58114.58488	1052	999	1.098	16927.0 ± 2.2	0.2612	0.0040	0.51754	0.00094	-3.63	0.41	-0.32

Table C1 – *continued*

BJD (+ 2400000.) [days]	S/N I [@551 nm]	S/N V [@551 nm]	Airmass	RV [m s ⁻¹]	S _{HK}	eS _{HK}	I _{Hα}	eI _{Hα}	B _ℓ [G]	σ _{B_ℓ} [G]	N _ℓ [G]
58115.62666	959	924	1.175	16897.0 ± 2.2	0.2505	0.0045	0.51575	0.00103	1.09	0.45	-0.13
58116.58632	1161	1114	1.105	16868.6 ± 2.1	0.2421	0.0040	0.51461	0.00085	1.98	0.38	0.20
58117.61202	984	957	1.154	16895.8 ± 2.2	0.2428	0.0048	0.51346	0.00098	1.07	0.45	-0.39
58118.59907	1081	1026	1.134	16920.1 ± 2.1	0.2475	0.0041	0.51431	0.00091	-1.12	0.41	0.34
58119.54663	973	939	1.079	16949.8 ± 2.1	0.2597	0.0042	0.51619	0.00103	-1.64	0.44	0.31
58121.53995	700	679	1.079	16911.3 ± 2.6	0.2654	0.0056	0.51698	0.00140	-2.95	0.61	0.67
58122.55886	1116	1093	1.094	16922.4 ± 2.1	0.2611	0.0038	0.51667	0.00089	-2.55	0.39	-0.18
58151.56156	916	865	1.286	16914.6 ± 2.3	0.2638	0.0054	0.51728	0.00102	-1.50	0.50	0.16
58152.55135	905	874	1.257	16905.5 ± 2.3	0.2655	0.0052	0.51696	0.00107	0.22	0.48	-0.74
58153.53730	991	962	1.220	16919.4 ± 2.1	0.2600	0.0050	0.51604	0.00097	0.84	0.44	-0.18
58155.53830	878	833	1.241	16898.9 ± 2.3	0.2583	0.0052	0.51460	0.00113	-2.17	0.49	-0.28
58156.54132	668	637	1.262	16916.4 ± 2.6	0.2608	0.0068	0.51515	0.00143	-5.35	0.65	0.03
58157.52308	804	780	1.211	16914.1 ± 2.3	0.2672	0.0059	0.51698	0.00118	-1.54	0.55	0.09
58158.52441	891	862	1.223	16889.7 ± 2.4	0.2689	0.0055	0.51670	0.00107	1.27	0.50	0.39
58159.52273	857	835	1.226	16908.2 ± 2.3	0.2642	0.0054	0.51601	0.00112	0.70	0.51	-0.30
58160.52493	720	695	1.243	16911.7 ± 2.5	0.2639	0.0060	0.51597	0.00133	-0.96	0.60	0.17
58161.53424	757	740	1.287	16883.6 ± 2.6	0.2654	0.0059	0.51629	0.00126	-0.94	0.59	-0.17
58162.51359	994	952	1.224	16918.8 ± 2.2	0.2638	0.0049	0.51632	0.00097	-1.16	0.44	-0.01
ESO EPOCH P101A											
58292.90558	703	774	1.423	17031.2 ± 2.5	0.2445	0.0054	0.52289	0.00137	0.39	0.62	0.08
58293.90560	441	487	1.408	17060.5 ± 3.1	0.2470	0.0085	0.52396	0.00214	-2.19	0.98	0.39
58294.90798	499	544	1.381	17051.8 ± 3.0	0.2499	0.0069	0.52424	0.00196	-2.48	0.84	-1.06
58295.91032	745	821	1.355	17053.0 ± 2.4	0.2542	0.0047	0.52429	0.00133	-1.29	0.56	0.79
58296.90910	711	781	1.349	17077.6 ± 2.5	0.2534	0.0052	0.52485	0.00136	0.84	0.60	-0.32
58297.88170	756	839	1.487	17063.0 ± 2.3	0.2561	0.0049	0.52527	0.00129	-0.33	0.56	0.54
58298.90085	683	762	1.363	17030.8 ± 2.5	0.2469	0.0054	0.52453	0.00141	-1.18	0.62	-0.78
58306.92059	477	512	1.204	17052.7 ± 3.0	0.2413	0.0072	0.52327	0.00204	-1.69	0.87	-0.29
ESO EPOCH P101B											
58328.87595	699	663	1.167	17001.0 ± 2.5	0.2422	0.0052	0.51407	0.00138	0.78	0.61	-0.30
58329.87836	755	748	1.155	16996.3 ± 2.4	0.2387	0.0048	0.51288	0.00128	-2.05	0.56	0.12
58331.86236	656	629	1.181	16992.6 ± 2.6	0.2375	0.0048	0.51252	0.00156	-0.52	0.61	-0.74
58332.79295	691	684	1.454	16988.7 ± 2.6	0.2365	0.0055	0.51318	0.00137	-1.75	0.61	-0.92
58333.86852	641	595	1.153	16999.2 ± 2.6	0.2389	0.0056	0.51332	0.00153	-3.71	0.65	-0.31
58335.87337	864	826	1.134	16999.9 ± 2.3	0.2433	0.0041	0.51384	0.00114	0.98	0.48	-0.44
58336.85067	827	809	1.176	17010.6 ± 2.2	0.2377	0.0045	0.51248	0.00117	-1.49	0.51	0.83
58341.86137	687	653	1.127	16997.7 ± 2.5	0.2441	0.0050	0.51385	0.00144	-1.72	0.60	-0.32
ESO EPOCH P101C											
58363.85483	899	876	1.077	16962.5 ± 2.1	0.2374	0.0039	0.51192	0.00111	-2.61	0.46	0.48
58366.78328	850	810	1.145	16962.7 ± 2.3	0.2332	0.0042	0.51127	0.00118	-0.65	0.49	-0.15
58367.76771	987	935	1.174	16956.2 ± 2.0	0.2372	0.0037	0.51216	0.00101	-4.21	0.42	-1.09
58368.79676	944	898	1.113	16947.1 ± 2.1	0.2395	0.0038	0.51228	0.00106	-1.03	0.44	0.84
58369.86552	1047	991	1.079	16961.4 ± 2.0	0.2369	0.0037	0.51220	0.00094	3.32	0.40	0.22
58370.86078	1065	1027	1.079	16966.8 ± 2.0	0.2435	0.0036	0.51266	0.00093	2.06	0.39	0.13
58371.87136	758	729	1.086	16931.6 ± 2.8	0.2432	0.0071	0.51296	0.00182	-2.07	0.79	-0.22
58372.85848	847	809	1.081	16963.7 ± 2.2	0.2428	0.0045	0.51184	0.00115	-3.29	0.50	-0.22
58375.84499	676	643	1.079	16941.4 ± 2.5	0.2427	0.0057	0.51188	0.00140	-3.99	0.63	-0.17
58376.84750	741	716	1.081	16934.3 ± 2.4	0.2383	0.0051	0.51247	0.00129	-0.85	0.57	-0.85
58377.74118	663	642	1.173	16958.0 ± 2.5	0.2405	0.0055	0.51277	0.00144	2.26	0.63	-0.29

APPENDIX D: ADDITIONAL HST LINES




Figure D1. Si IV spectra of ι Hor compared with the solar active (green) and quiet regions (yellow). Enhanced emission is seen in ι Hor during the second visit in Cycle 25 though to a lesser extent than in the C III and C IV lines.




Figure D2. O IV lines of ι Hor at four epochs compared with solar active (green) and quiet regions (yellow) as in the previous plots of Si IV, C III and C IV. While enhancement is clearly seen in the O IV emission during the observation marked C25-II, the shape of O IV 1401.0 Å also appears to change.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.