

Comparative Analysis of the Volatile Fraction of Fruit Juice from Different *Citrus* Species

M. Carmen González-Mas^{1*}, José Luis Rambla^{2,3}, M. Carmen Alamar³, Abelardo Gutiérrez³, Antonio Granell²

1 Centro de Citricultura, Instituto Valenciano de Investigaciones Agrarias (IVIA), Moncada, Valencia, Spain, **2** Instituto de Biología Molecular y Celular de Plantas (IBMCP), CSIC - Universidad Politécnica de Valencia, Valencia, Spain, **3** Centro de Agrolingeniería, Instituto Valenciano de Investigaciones Agrarias (IVIA), Moncada, Valencia, Spain

Abstract

The volatile composition of fruit from four *Citrus* varieties (Powell Navel orange, Clemenules mandarine, and Fortune mandarine and Chandler pummelo) covering four different species has been studied. Over one hundred compounds were profiled after HS-SPME-GC-MS analysis, including 27 esters, 23 aldehydes, 21 alcohols, 13 monoterpene hydrocarbons, 10 ketones, 5 sesquiterpene hydrocarbons, 4 monoterpene cyclic ethers, 4 furans, and 2 aromatic hydrocarbons, which were all confirmed with standards. The differences in the volatile profile among juices of these varieties were essentially quantitative and only a few compounds were found exclusively in a single variety, mainly in Chandler. The volatile profile however was able to differentiate all four varieties and revealed complex interactions between them including the participation in the same biosynthetic pathway. Some compounds (6 esters, 2 ketones, 1 furan and 2 aromatic hydrocarbons) had never been reported earlier in *Citrus* juices. This volatile profiling platform for *Citrus* juice by HS-SPME-GC-MS and the interrelationship detected among the volatiles can be used as a roadmap for future breeding or biotechnological applications.

Citation: González-Mas MC, Rambla JL, Alamar MC, Gutiérrez A, Granell A (2011) Comparative Analysis of the Volatile Fraction of Fruit Juice from Different *Citrus* Species. PLoS ONE 6(7): e22016. doi:10.1371/journal.pone.0022016

Editor: Pierandrea Temussi, Università di Napoli Federico II, Italy

Received: March 7, 2011; **Accepted:** June 12, 2011; **Published:** July 19, 2011

Copyright: © 2011 Gonzalez-Mas et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work has been supported through the project GVPRE/2008/164 of Conselleria d'Educació of Valencian Community and the project RTA2007-00029-C02-01: Sensogest (INIA). M.C. González and A. Gutierrez are contracted by the Fundación Agroalimed (Conselleria d'Agricultura of VC). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: gonzalez_mde@gva.es

These authors contributed equally to this work.

Introduction

Developing powerful platforms for volatile analysis is a prerequisite for further insights into the volatiles biosynthetic pathways and also in the identification of the genetic and environmental effects in volatile production [1,2]. This information is relevant in the frame of current breeding programs in *Citrus* which are directed to respond to the market demand for quality fruits and are also important for the biotechnology of fruit and fruit derived product. One of the main characteristics of *Citrus* fruit quality is defined by the aroma of fruit juice. The aroma of a fresh juice is the product of a complex combination of several odour components that include esters, aldehydes, alcohols, ketones and hydrocarbons, which are collectively defined as volatile organic compounds or VOCs [3–6]. Headspace extraction coupled to GC-MS is at present the method of choice for most of the volatile analysis in food/flavour chemistry [7,8] and particularly in *Citrus* [9–17], having displaced former methods that involved complex sample preparation and large amounts of solvents [18–20]. Some studies on the compositional analysis of *Citrus* juice aroma have been described which used dynamic and static headspace extraction [21,22]. Different types of fibers have been used for *Citrus* juice analysis by HS-SPME [10,13,14,23] but the one with three components: DVB/CAR/PDMS (divinylbenzene/ carboxen/polydimethylsiloxane) is the most widely used, because of

its ability to extract a larger number of VOCs than other fibers [15,17,24,25].

So far, almost all the studies on the aroma of *Citrus* juices had been conducted on orange juice, normally using one or at most two varieties. The fragmented information available together with the different techniques and fibers used, complicates the comparison of VOCs profiles between different *Citrus* varieties present in the literature [5,10,15,16,18,19,21,26–28]. In contrast to oranges, only few studies have been conducted on mandarin [10,11] and grapefruit aroma juices [4,12,29]. No studies have been performed for the volatiles in the juice of pummelo, and only one comparative study has been reported comparing mandarin and orange juices [3]. In this paper we describe the optimization of a VOCs capture/profiling method for *Citrus* and the characterization of the volatile profile for the juice of four *Citrus* varieties: Powell Navel summer orange, Clemenules clementine mandarine, and Fortune mandarine and Chandler pummelo hybrids. All four varieties are used as parentals in order to obtain new hybrids in breeding programs and at the same time they are themselves important varieties for fresh market in the world [30]. This is the first time that different varieties corresponding to different species are analysed in parallel using the same analytical technique and therefore enable us to describe both the volatile fraction in the juice and the variability in the volatile profile between the materials analysed.

It is also the first time that the volatiles in the juice of pummelo are described.

Materials and Methods

Citrus juice

Mature fruits at optimal ripening stage [31], were collected in 2007 from trees of Powell Navel Late sweet orange (*Citrus sinensis* (L.) Osb.), Clementines (*Citrus clementine* Hort. ex Tan.), and two *Citrus* hybrids: Fortune (*C. clementine* x *C. tangerine*) and Chandler pummelo (*C. grandis* x *C. grandis*) varieties. All trees were grown in the same orchard and subjected to homogeneous cultural conditions, in order to reduce environmental effects on the volatile profile. The experimental orchard is located at the Experimental Station of *Instituto Valenciano de Investigaciones Agrarias*, Moncada, Valencia, Spain, under a mediterranean climate (averages rainfall of 515.8 mm and temperature of 15.2°C for 2007). In all cases, three biological replicate samples for each variety were obtained, each one representing at least four different fruits each. Fruit juice was obtained using a hand extractor, in order to avoid squeezing of the flavedo and to prevent contamination of the juice with peel components. After that, 10 mL aliquots of each sample were placed in 22 mL crimp cap headspace vials and kept frozen at -20°C until analyzed. Two aliquots of 10 mL corresponding to technical replicates of each sample were analyzed. The total number of analysis was 24 (3 biological samples x 2 technical replicates for the 4 varieties).

HS-SPME extraction conditions

Right before analysis, samples were thawed at 20°C for ten minutes and then were subjected to headspace solid phase micro-extraction (HS-SPME). Extraction was carried out using 10 mL of sample into a 22 mL crimp cap headspace vial. A 50/30 µm DVB/CAR/PDMS (Supelco, Bellefonte, PA, USA) fiber was used for all the analysis. Pre-incubation and extraction times were 10 and 20 min, respectively. A temperature of 50°C was selected for pre-incubation and extraction because it allowed the detection of a higher number of VOCs than when 30°C was used. Desorption was performed for 1 min at 250°C in splitless mode.

Gas chromatography–mass spectrometry conditions

VOCs trapped on the fiber were analysed by GC-MS using an autosampler COMBI PAL CTC Analytics (Zwingen, Switzerland), a 6890N GC Agilent Technologies (Santa Clara, CA, USA) and a 5975B Inert XL MSD Agilent, equipped with an Agilent J&W Scientific DB-5 ms fused silica capillary column (5%-phenyl-95%-dimethylpolysiloxane as stationary phase, 60 m length, 0.25 mm i.d., and 1 µm thickness film). Oven temperature conditions were 40°C for 2 min, 5°C/min ramp until 250°C and then held isothermally at 250°C for 5 min. Helium was used as carrier gas at 1.2 mL/min constant flow. Mass/z detection was obtained by an Agilent mass spectrometer operating in the EI mode (ionization energy, 70 eV; source temperature 230°C). Data acquisition was performed in scanning mode (mass range m/z 35–220; seven scans per second). Chromatograms and spectra were recorded and processed using the Enhanced ChemStation software for GC-MS (Agilent).

Compound identification

Compound identification was based both on the comparison between the MS for each putative compound with those of the NIST 2005 Mass Spectral library and also with the match to our GC retention time and Mass Spectra custom library which have been generated using commercially available compounds.

Compounds used as reference were of analytical grade and purchased from Sigma-Aldrich Química (Madrid, Spain), except for 2-carene, thymol and ledene, which were obtained from Extrasynthese (Genay, France). In addition to the commercial compounds, seven esters (methyl pentanoate, ethyl pentanoate, methyl heptanoate, ethyl heptanoate, methyl octanoate, ethyl octanoate, and ethyl nonanoate) were synthesized in our laboratory by acid-catalyzed esterification from analytical grade reagents. For that, 10 µL of the corresponding acid (pentanoic acid, heptanoic acid, octanoic acid, or nonanoic acid, supplied by Sigma-Aldrich) was added to 1 mL of the corresponding alcohol (methanol, ethanol) with 10 µL of H₂SO₄ 96%, and incubated at 40°C overnight. After that, a small amount of sodium carbonate was added and incubated at 4°C for 24 hours, to neutralize any remaining acid. The solution was centrifuged and the supernatant used as a ≈1% standard solution of the ester in the respective alcohol. Also, 1 mL of either 100 ppb or of 1 ppm standard solutions was analyzed in the same conditions as the samples. Only those compounds/peaks confirmed by both mass spectrum and retention time in each and every chromatogram were considered. For relative quantification, the peak area was integrated from the extracted ion chromatogram corresponding to a specific ion previously selected for each compound. A mixture of extracts representing the four varieties analysed was injected regularly as part of the injection series and was used as a reference for correction for temporal variation and fiber aging. Finally, corrected results for each compound were expressed as relative ratios to the average level present in Chandler juice. When a compound was not detected in Chandler, the ratio was calculated to a variety that contained it as indicated in Table 1.

Statistical analysis

For both Principal Component Analysis (PCA) and Hierarchical Cluster Analysis, the complete dataset including all replicates was considered. For both type of analysis, the ratio of the signal relative to that of the average in the four varieties was log 2 transformed. For PCA, the program SIMCA-P version 11 (Umetrics, Umea, Sweden) was used with the centered data. For the Hierarchical Cluster Analysis, the program Acuity 4.0 (Axon Instruments) was used, with the distance measures based on the Pearson correlation. Pearson correlation coefficients were calculated with the SPSS version 15.0 software (SPSS Inc., Chicago, USA). Data from the correlation matrix was represented as a heatmap by means of the Acuity 4.0 program.

Results and Discussion

Table 1 lists the VOCs detected in our HS-SPME-GC-MS platform and the relative levels for the four varieties analyzed. A total of 109 compounds have been identified: 27 esters (19 aliphatic and 8 monoterpene acetates), 23 aldehydes (18 aliphatic, 4 monoterpene and 1 norcarotenoid), 21 alcohols (12 aliphatic and 9 monoterpene), 13 monoterpene hydrocarbons, 10 ketones (8 aliphatic, 1 norcarotenoid and 1 monoterpene), 5 sesquiterpene hydrocarbons, 4 monoterpene cyclic ethers, 4 furans and 2 aromatic hydrocarbons. It is important to note that although more than 300 VOCs have been reported in other *Citrus* juice [26], some of them have been identified only tentatively [16,18,19,24]. To unequivocally assign chemical names to the compounds in our dataset, we have used analytical grade commercial compounds. Those compounds that were putatively identified by their mass spectra but were not confirmed with the commercial standard were not included in our dataset.

Table 1. Relative levels (fold changes) of VOCs detected in juices of four *Citrus* varieties.

Code	Cluster	Volatile Organic Compound	Family Code/ Number	Retention Time (min)	Specific Ion (m/z)	Chandler	Clemenules	Fortune	Powell
1	A1	2-carene	Mt hd/1	24.05	93	1±0.25	-	traces	traces
2	A1	(Z)-linalool oxide ^{a,b}	Alc/1	26.47	111	1±0.24	-	traces	-
3	A1	(E)-linalool oxide ^{a,b}	Alc/2	27.00	111	1±0.22	-	-	-
4	A1	(E,E)-2,4-nonadienal	Ald/1	31.10	81	1±0.31	-	-	-
5	A1	(Z)-ocimene	Mt hd/2	24.87	93	1±0.61	-	-	-
6	A1	p-cymene	Mt hd/3	24.91	119	1±0.85	-	-	-
7	A1	β-caryophyllene	Sqt/1	37.87	133	1±0.57	-	-	traces
8	A1	nootkatone ^c	Ket/1	47.87	121	1±0.33b	-	0.02±0.01a	0.02±0.01a
9	A1	α-humulene	Sqt/2	38.82	80	1±0.64b	traces	0.04±0.01a	0.03±0.01a
10	A1	β-pinene	Mt hd/4	23.46	93	1±0.88b	0.11±0.01a	0.17±0.03a	0.21±0.12a
11	A1	1-pentanol	Alc/3	14.60	42	1±0.48b	0.26±0.13a	0.11±0.02a	0.20±0.11a
12	A1	1-hexanol	Alc/4	18.55	56	1±0.70b	0.12±0.02a	0.22±0.18a	0.87±0.32b
13	A1	methyl hexanoate	Est/1	20.68	74	1±0.46c	-	0.07±0.02a	0.44±0.20b
14	A1	pseudocumene ^e	Ar/1	23.85	105	1±0.18b	0.35±0.21a	0.24±0.01a	1.11±0.43b
15	A1	ethyl hexanoate	Est/2	23.37	88	1±0.66b	0.17±0.06a	0.62±0.11ab	2.47±1.57c
16	A2	(E,E)-2,4-decadienal	Ald/2	34.17	81	1±0.68c	0.91±0.76bc	0.24±0.08ab	0.05±0.02a
17	A2	6-methyl-5-hepten-2-one ^d	Ket/2	23.03	108	1±0.40b	0.46±0.05a	0.12±0.02a	-
18	A2	octanal	Ald/3	23.77	57	1±0.22c	0.83±0.19b	0.39±0.05a	0.22±0.06a
19	A2	heptanal	Ald/4	20.00	70	1±0.26c	0.53±0.04b	0.42±0.01b	0.08±0.06a
20	A2	nonanal	Ald/5	27.30	57	1±0.15c	0.71±0.15b	0.60±0.06b	0.15±0.27a
21	A2	2,3-pentanedione	Ket/3	11.91	100	1±0.28d	0.79±0.15c	0.53±0.16b	0.07±0.04a
22	A2	(E)-2-octenal	Ald/6	25.79	70	1±0.24c	0.67±0.37b	0.51±0.15b	0.13±0.08a
23	A2	(E)-2-nonenal	Ald/7	29.21	70	1±0.28b	1.07±0.09b	0.98±0.20b	0.18±0.09a
24	A2	(E)-2-heptenal	Ald/8	22.14	83	1±0.26b	0.29±0.12a	0.23±0.05a	0.08±0.04a
25	A2	1-octen-3-one	Ket/4	22.82	70	1±0.46b	0.15±0.09a	0.19±0.07a	0.10±0.05a
26	A2	1-octen-3-ol	Alc/5	22.85	57	1±0.32b	0.11±0.02a	0.14±0.03a	0.08±0.04a
27	A2	hexanal	Ald/9	16.01	56	1±0.22b	0.22±0.03a	0.23±0.02a	0.14±0.07a
28	A2	pentanal	Ald/10	12.04	58	1±0.24b	0.19±0.02a	0.23±0.04a	0.14±0.09a
29	A2	2-pentylfuran	Fur/1	23.36	138	1±0.19b	0.27±0.01a	0.29±0.05a	0.14±0.12a
30	A2	1-heptanol	Alc/6	22.39	70	1±0.55b	0.38±0.08a	0.58±0.32a	0.31±0.05a
31	A2	α-copaene	Sqt/3	36.41	119	1±0.58b	0.22±0.05a	0.55±0.06a	0.35±0.02a
32	A2	valencene	Sqt/4	39.69	133	1±0.56b	0.15±0.02a	1.16±0.32b	0.91±0.17b
33	A2	bornyl acetate ^a	Est/3	33.48	121	1±0.87a	0.46±0.18a	0.73±0.13a	0.50±0.27a
34	A2	ethyl heptanoate	Est/4	26.82	88	1±0.67b	0.36±0.18a	1.76±0.31c	0.69±0.29ab
35	A2	propyl acetate ^{e*}	Est/5	12.52	61	-	-	1±0.53	-
36	B	2-ethylfuran	Fur/2	12.17	81	1±0.21b	2.41±0.32d	1.22±0.24c	0.26±0.08a
37	B	2-methylfuran	Fur/3	8.84	82	1±0.25b	1.62±0.20c	1.13±0.29b	0.40±0.09a
38	B	(E)-2-pentenal	Ald/11	14.30	83	1±0.45b	4.09±0.80c	1.45±0.25b	0.41±0.07a
39	B	1-penten-3-one	Ket/5	11.57	55	1±0.52a	9.49±2.20c	2.45±0.59b	0.53±0.11a
40	B	ethyl propanoate	Est/6	12.47	57	1±1.08a	20.09±3.13b	22.36±7.40b	-
41	B	ethyl 2-methylbutanoate ^f	Est/7	17.87	102	-	1±0.38b	0.16±0.04a	-
42	B	3-pentanone ^e	Ket/6	11.97	57	1±0.63b	2.55±0.57c	0.59±0.19ab	0.40±0.14a
43	B	1-penten-3-ol	Alc/7	11.46	57	1±0.64a	2.05±1.38b	0.54±0.14a	0.71±0.37a
44	B	β-citronellal ^{a,ff}	Ald/12	28.92	69	-	1±1.53	-	-
45	B	eucalyptol (1,8-cineole)	Mt cyc ether/1	25.38	154	1±0.62a	6.56±1.71c	0.30±0.06a	2.79±1.07b
46	B	decanal	Ald/13	30.59	57	1±0.19a	4.51±1.67b	0.78±0.22a	1.46±1.24a
47	B	(Z)-3-hexenal	Ald/14	15.92	69	1±0.89ab	1.54±0.32b	0.44±0.14a	0.81±0.54ab
48	B	(E)-2-hexenal	Ald/15	18.17	83	1±0.70a	3.92±0.81c	1.02±0.16ab	1.68±0.83b

Table 1. Cont.

Code	Cluster	Volatile Organic Compound	Family Code/ Number	Retention Time (min)	Specific Ion (m/z)	Chandler	Clemenules	Fortune	Powell
49	B	geranylacetone ^d	Ket/7	37.62	43	1±0.26b	4.73±0.36d	0.58±0.14a	1.65±0.67c
50	B	β-cyclocitral ^d	Ald/16	31.63	137	1±0.42a	6.45±0.60b	0.78±0.13a	0.92±0.12a
51	B	β-ionone ^d	Ket/8	38.90	177	1±0.44a	11.34±0.76c	0.88±0.18a	1.53±0.24b
52	C1	nerol ^a	Alc/8	31.28	93	1±0.47ab	0.44±0.19a	1.86±2.07b	5.43±2.18c
53	C1	hexyl acetate	Est/8	23.86	56	1±0.86a	0.46±0.10a	0.98±0.22a	3.91±0.65b
54	C1	methyl nonanoate [*]	Est/9	30.91	74	1±0.51a	0.45±0.17a	1.02±0.11a	6.04±1.81b
55	C1	neryl acetate ^a	Est/10	34.97	69	1±0.55a	0.20±0.07a	1.90±0.19a	15.53±5.87b
56	C1	(Z)-3-hexen-1-ol	Alc/9	18.16	82	1±0.92a	0.50±0.11a	0.33±0.09a	4.75±1.27b
57	C1	methyl octanoate [*]	Est/11	27.74	74	1±0.49b	0.41±0.05a	0.44±0.08ab	4.60±1.39c
58	C1	geranial ^a	Ald/17	32.61	69	1±0.41a	0.76±0.17a	0.39±0.11a	14.62±12.11b
59	C1	geraniol	Alc/10	31.95	69	1±0.63a	0.78±0.12a	1.50±0.47a	10.88±6.14b
60	C1	heptyl acetate ^g	Est/12	27.27	43	-	-	-	1±0.27
61	C1	methyl decanoate [*]	Est/13	33.94	74	1±0.82a	3.78±1.48a	3.14±0.92a	72.44±42.64b
62	C1	1-nonanol	Alc/11	29.36	70	1±0.37a	1.38±0.57a	0.82±0.30a	12.04±10.54b
63	C1	undecanal	Ald/18	33.67	57	1±0.18ab	1.23±0.19b	0.76±0.09a	1.21±0.63b
64	C1	(E)-2-hexen-1-ol ^f	Alc/12	18.48	57	-	1±0.75a	-	2.72±0.80b
65	C1	1-decanol ^f	Alc/13	32.50	115	-	1±0.40a	0.03±0.01a	5.10±2.13b
66	C1	1-octanol	Alc/14	26.00	56	1±0.33a	6.38±1.83a	2.18±0.26a	59.82±50.05b
67	C1	3-carene	Mt hd/5	24.43	93	1±0.56a	14.10±8.38b	1.39±0.64a	78.68±4.74c
68	C2	β-citronellol ^a	Alc/15	31.15	81	1±0.53a	1.47±0.39a	4.53±0.45c	2.21±0.16b
69	C2	ethyl acetate	Est/14	9.18	61	1±0.42a	5.28±1.55b	40.71±7.12c	3.70±1.37b
70	C2	(Z)-carveol ^a	Alc/16	31.84	109	1±0.49a	4.77±1.76b	6.91±0.85c	4.85±1.98b
71	C2	(E)-carveol ^a	Alc/17	31.41	109	1±0.36a	2.91±0.19b	5.35±0.71c	2.49±0.35b
72	C2	carvone ^a	Ket/9	32.31	82	1±0.68a	12.09±3.13c	20.89±1.02d	5.75±0.13b
73	C2	linalool ^a	Alc/18	27.15	93	1±0.69a	15.73±1.35c	26.77±1.66d	10.05±0.85b
74	C2	ethanol	Alc/19	5.64	45	1±0.59a	10.91±2.39c	6.05±1.64b	4.95±2.33b
75	C2	acetaldehyde	Ald/19	4.77	43	1±0.25a	3.07±0.84b	2.75±0.22b	2.96±0.49b
76	C2	dodecanal	Ald/20	36.55	57	1±2.35a	77.62±21.86d	20.38±8.79b	58.44±39.38c
77	C2	3-methylfuran [*]	Fur/4	9.16	82	1±0.32a	4.25±1.60c	2.91±0.94b	2.51±0.33b
78	C2	(E)-limonene oxide ^h	Mt cyc ether/2	28.84	94	1±0.77a	7.53±2.60b	6.81±4.59b	5.98±3.14b
79	C2	(Z)-limonene oxide ^h	Mt cyc ether/3	28.77	67	1±1.15a	2.88±0.82b	3.47±0.14b	3.74±0.50b
80	C3	camphene	Mt hd/6	22.44	93	1±1.04a	1.37±0.17ab	1.95±0.20b	3.31±1.69c
81	C3	terpinolene	Mt hd/7	27.05	121	1±0.40a	3.19±0.18b	4.78±0.33c	6.11±1.01d
82	C3	limonene	Mt hd/8	25.11	108	1±0.25a	2.23±0.12b	2.46±0.10b	2.93±0.42c
83	C3	α-pinene	Mt hd/9	21.67	93	1±0.41a	6.17±0.92b	9.17±1.18b	18.56±9.66c
84	C3	myrcene	Mt hd/10	23.28	91	1±0.42a	2.54±0.37b	3.21±0.28b	5.58±1.78c
85	C3	α-phellandrene	Mt hd/11	24.29	93	1±0.41a	3.51±0.41b	5.64±0.58b	14.03±8.35c
86	C3	α-terpineol ^a	Alc/20	30.75	59	1±0.59a	3.78±0.25b	5.67±0.69c	6.45±3.56c
87	C3	γ-terpinene	Mt hd/12	26.03	93	1±0.61a	2.97±0.44b	5.16±0.85c	5.34±2.62c
88	C3	terpinen-4-ol ^a	Alc/21	30.38	93	1±0.77a	5.65±1.41b	15.86±2.59d	10.98±5.85c
89	C3	neral ^a	Ald/21	31.81	84	1±0.55a	4.28±1.43b	6.21±0.69c	7.32±1.45d
90	C3	perillaldehyde ^a	Ald/22	33.43	68	1±0.29a	8.81±2.43b	15.19±5.28c	25.48±2.64d
91	C3	α-terpinene	Mt hd/13	24.65	121	1±0.52a	3.41±1.43b	4.75±1.80c	6.19±1.61d
92	C3	ethyl nonanoate [*]	Est/15	33.05	88	1±0.91a	0.41±0.15a	5.59±1.51b	6.88±1.16c
93	C3	geranyl acetate ^a	Est/16	35.50	69	1±0.96a	1.19±0.34a	14.09±1.39b	28.56±12.68c
94	C3	(Z)-carvyl acetate ^a	Est/17	34.50	84	1±0.46a	0.54±0.10a	4.27±1.58b	6.92±1.84b
95	C3	(E)-carvyl acetate ^a	Est/18	35.30	84	1±1.17a	0.62±0.15a	7.36±4.56b	4.79±0.55b
96	C3	citronellyl acetate ^a	Est/19	34.69	95	1±0.62a	0.94±0.31a	47.41±4.01c	19.52±3.25b
97	C3	styrene [*]	Ar/2	20.01	104	1±0.30a	2.41±0.91a	2.90±0.68a	15.10±5.92b
98	C3	ethyl octanoate	Est/20	30.04	88	1±0.55a	1.28±0.37a	4.05±0.81b	16.63±2.00c

Table 1. Cont.

Code	Cluster	Volatile Organic Compound	Family Code/ Number	Retention Time (min)	Specific Ion (m/z)	Chandler	Clemenules	Fortune	Powell
99	C3	nonyl acetate	Est/21	33.47	98	1±2.35a	4.88±6.60a	22.54±6.43a	304.38±77.26b
100	C3	ethyl butanoate	Est/22	15.88	88	1±2.036a	17.58±4.13b	27.11±5.07b	113.73±54.23c
101	C3	decyl acetate	Est/23	36.29	70	1±1.51a	48.18±18.61b	68.70±11.40c	553.34±30.06d
102	C3	α -terpinyl acetate ^{a,f}	Est/24	35.10	121	-	1±0.50a	1.47±0.34b	9.61±0.72c
103	C3	linalyl acetate ^a	Est/25	31.82	93	1±0.47a	7.09±4.24a	7.29±0.92a	83.29±21.54b
104	C3	ethyl decanoate	Est/26	35.91	88	1±0.78a	5.84±1.75ab	16.43±3.04b	62.91±32.20c
105	C3	octyl acetate	Est/27	30.47	70	1±0.53a	18.86±8.93a	64.08±0.30b	526.00±86.22c
106		3-methylbutanal	Ald/23	10.66	58	-	traces	-	-
107		1,4-cineole	Mt cyc ether/4	24.54	111	-	traces	-	traces
108		β -farnesene	Sqt/5	37.73	120	-	-	-	traces
109		γ -dodecalactone [*]	Ket/10	43.71	85	traces	-	-	-

Data are normalized to the mean values in the Chandler variety, unless otherwise indicated. Mean corresponding to n=6 values. Means followed by different letters in the same row are significantly different ($p < 0.05$) by Duncan's test. Family Code: Ald: Aldehyde; Ket: Ketone; Alc: Alcohol; Est: Ester; Fur: Furan; Mt hd: Monoterpene hydrocarbon; Sqt: Sesquiterpene; Ar: Aromatic hydrocarbon; Mt cyc ether: Monoterpene cyclic ether.

^aMonoterpene derived compound.

^bIn addition to the alcohol group, it has a tetrahydrofuran group.

^cSesquiterpene compound.

^dNorcarotenoid compound.

^eData normalized to the mean abundance in Fortune variety.

^fData normalized to the mean abundance in Clemenules variety.

^gData normalized to the mean abundance in Powell variety.

^hIts cyclic ether group is an epoxy group.

^{*}Compound reported for the first time in a Citrus juice.

doi:10.1371/journal.pone.0022016.t001

As a result eleven compounds out of a total of 109 are described here for the first time in the juice of Citrus species (6 esters, 2 ketones, 1 furan and 2 aromatic hydrocarbons) (Table 1); the remaining compounds have been described previously in Citrus juice samples [11,16,17,24,26,32–34]. Almost all the detected compounds showed dramatic changes in the levels of accumulation in at least one of the four varieties (see Table 1). To better understand the usefulness of the volatile profile to define and distinguish the four Citrus varieties, a principal component analysis (PCA) was performed. Figure 1 shows that the first two principal components explain almost 80% of the variance, and clearly separate all four varieties from one another. The first component, explaining 54% of the variance, mainly separates Chandler pummelo from all the other varieties and to a lesser extent also Powell orange from both Clemenules and Fortune. The second component explains about 25% of the variance and clearly separates Clemenules from Powell and Chandler, while Fortune would be intermediate. Finally, the third component (Figure S1) essentially separates Fortune from the rest, and the analysis of the loading plots should reveal the part of the volatile profile which is characteristic of Fortune, and is responsible of roughly 13% of the total variance. These three components together explain as much as 92% of the total variance in the dataset.

Analysis of the loadings plot reveals the compounds responsible of the separation between samples (Figure 2). The most relevant for the first component is a group of mostly terpenic compounds (β -caryophyllene, (*Z*)-ocimene, (*E,E*)-2,4-nonadienal, (*Z*)- and (*E*)-linalool oxides, *p*-cymene) which is almost exclusive of Chandler pummelo, and the compound octyl acetate, a metabolite present at relatively very high levels in Powell. The second component is defined by a group of compounds, mostly esters, with contrasting relative levels between Clemenules and Powell. The most relevant compounds contributing to the separation of Fortune from the

other varieties are revealed by the loadings plot corresponding to the third component (Figure S2), and include propyl acetate, citronellyl acetate and ethyl acetate with higher levels in Fortune, and (*E*)-2-hexen-1-ol, eucalyptol, 3-carene and 1-decanol with lower levels in this variety.

A hierarchical cluster analysis confirmed that Clemenules and Fortune presented the most similar volatile profile, while Chandler pummelo exhibited the most differential profile of them all (Figure 3). According to the pattern of VOCs presented by these four varieties, volatile compounds can be organized in three clusters, named A, B and C, with some sub-clusters (named A1, A2, C1, C2 and C3). It is therefore revealed that clusters of VOCs with differential accumulation levels rather than a few individual compounds are responsible for the separation between varieties. For the sake of clarity, compounds in Table 1 are displayed according to the same order than in the hierarchical cluster.

Correlation analysis of the volatile compounds was also performed, in order to assess how these metabolites were related to each other. When compared to the hierarchical cluster analysis, results are basically consistent. Basically, highly positively correlated volatiles were grouped in the same cluster, and compounds in distant clusters tend to show negative or non-significant correlations (Figure 4, Table S1). When descending to the metabolite to metabolite level, it can be observed a general pattern of high positive correlations of ester compounds to both their alcoholic precursor and other structurally similar esters. This suggests that the levels of these compounds, which show up to 500-fold variations between varieties, could be regulated both by enzymatic activity (by means of relatively specific alcohol acyl transferases) and by substrate availability. A strong negative correlation between ester and aldehyde levels is also observed. This also suggests an important role for alcohol dehydrogenase enzymes activity in the differences detected between the volatile




Figure 1. Principal Component Analysis score plot (t[1] vs t[2]) for the first and second principal components.
doi:10.1371/journal.pone.0022016.g001

profiles of Chandler, otherwise basically rich in sesquiterpenes and aliphatic aldehydes, and the other varieties with a volatile profile with higher abundance of alcohols and esters.

Compounds in the cluster A are present at higher levels in Chandler pummelo than in any of the other three varieties studied. Compounds which are basically exclusive of Chandler belong to sub-cluster A1 and include mostly monoterpene hydrocarbons and derivatives such as 2-carene, (*Z*)-linalool oxide, (*E*)-linalool oxide, (*Z*)-ocimene, *p*-cymene, and also (*E,E*)-2,4-nonadienal and nootkatone. Among the compounds in sub-cluster A1, 2-carene had only been identified so far in pummelo peel oil [35] and the sesquiterpene nootkatone has been frequently described in grapefruit juice [4] but rarely in other Citrus juices [18]; the remaining compounds in this subcluster have been identified also in Citrus juices [3,17,26]. Sub-cluster A2 includes aliphatic aldehydes from five to nine carbon atoms, and some olefinic aldehydes such as (*E*)-2-heptenal, (*E*)-2-octenal, (*E*)-2-nonenal, and (*E,E*)-2,4-decadienal, all of which have been described to provide herbal, fruity and floral aroma to Citrus juices [26,32]. This sub-cluster also includes the compound 2-pentylfuran, reported previously only in tangerine [34], but identified in all four of our varieties in this paper. Cluster A included the only four sesquiterpenes unambiguously identified in our analysis: β -caryophyllene, nootkatone, α -copaene and valencene (β -farnesene was only detected at the level of traces in Powell), all of which had been previously reported in Citrus juices [17,26]. However, the chromatograms of all varieties, and most notably those of

Chandler, presented a large number of unidentified sesquiterpenes (as could be inferred from their MS spectra) which corresponded to the most abundant peaks eluting between 35 and 41 min (Figure S3). The close similarity of the mass spectra of many sesquiterpenes and the lack of standards makes this identification difficult, as it requires the use of purification steps and additional analytical techniques (such as NMR, and chemical synthesis) in order to identify their exact molecular structures. Therefore, although noted here, we did not include them in our approach.

Cluster B is defined by the compounds more abundantly found in Clemenules than in any of the other three varieties. These include a set of highly correlated carotenoid derivatives probably by the action of carotenoid cleavage dioxygenases: β -cyclocitral, β -ionone and geranylacetone (Figure 4, Table S1), and 3-pentanone, a ketone reported here for the first time in a *Citrus* juice. Other compounds in cluster B have also been previously described in *Citrus* juice [34] and they include 1-penten-3-one, 2-ethylfuran, 2-methylfuran, eucalyptol and the aldehydes (*E*)-2-pentenal, decanal, (*Z*)-3-hexenal, (*E*)-2-hexenal, and finally β -citronellal, which in our analysis was only detected in Clemenules.

Sub-cluster C1 includes compounds found more abundant in Powell than in the other three varieties. The monoterpene 3-carene and the esters methyl octanoate, methyl decanoate and heptyl acetate are the most important (heptyl acetate is exclusive of Powell variety). Methyl octanoate and methyl decanoate had never been described in *Citrus* juice, although the presence of many other esters had been previously reported in *Citrus* [16,17,26].





Figure 2. Principal Component Analysis loading plot (p[1] vs p[2]) for the first and second principal components. Each number corresponds to a particular volatile compound, as indicated in Table 1. doi:10.1371/journal.pone.0022016.g002

Sub-cluster C2 includes most of the compounds which accumulated generally to higher levels in Fortune than in other varieties, such as linalool or β -citronellol.

Finally, sub-cluster C3 includes compounds which are present in smaller quantities in Chandler than in the other varieties studied. Included in this sub-cluster are monoterpene hydrocarbons such as α -phellandrene, limonene or γ -terpinene, all of which are generally described in *Citrus* juices [26]. Also neral and perillaldehyde aldehydes, and 3-methylfuran (the only one of the four furans detected here that had never been described in *Citrus* juice before) were less abundant in Chandler than in the other three varieties. Some furan compounds are considered to be originated from lipid oxidation [36], but our results suggest independent metabolic pathways for the synthesis of 2- and 3-alkyl furans. This is based in 2-methylfuran showing a very strong positive correlation to 2-ethylfuran and also to 2-pentylfuran in our samples, while no significant correlation was found to 3-methylfuran. Moreover, the majority of compounds included in this sub-cluster showed the highest levels in Powell variety, as it is the case for monoterpenes limonene, α -phellandrene and α -pinene, monoterpene acetates, aliphatic esters octyl-, nonyl-, and decyl acetate, ethyl octanoate, ethyl nonanoate and ethyl decanoate (ethyl nonanoate never been described in *Citrus* literature before) and the aromatic hydrocarbon styrene. Styrene and pseudocumene (other aromatic hydrocarbon synonymous of 1,2,4-trimethylbenzene) have been identified in all our four

varieties for the first time. Of these two, only styrene have been described previously in *Citrus* commercial juices [21] but not pseudocumene although a compound with a similar structure, 1,4-diethylbenzene, have been reported previously in tangerine juice [37].

Some volatile compounds commonly described in *Citrus* juices failed to be detected in our study (Table S2). Thus, no volatile acids were detected in the juices analyzed; in fact it is known that the contribution of the acids to the total aroma of the orange juice is very limited [24]. In addition, some esters usually described in the *Citrus* juice, such as methyl butanoate [3,14,15,28], ethyl 3-hydroxyhexanoate [5,16,28,33], or methyl *o*-(methylamino)benzoate [17] were not identified in our samples. Moreover some alcohols such as the aliphatic alcohols 2- and 3-methylbutanol [3] or the monoterpene alcohol borneol and sesquiterpene alcohols β -eudesmol and α -bisabolol [12] described in previous *Citrus* analysis were not found in our samples. Vanillin was not found in our samples either, although it has been described in many other studies in *Citrus* juices [5,27], although this compound usually appears in juices that have undergone degradation due to exposure to high temperature [26]. This is also the case for some aldehydes identified in *Citrus* aroma, such as cuminaldehyde or (*E*)-2-undecenal [4,13], or some C₁₃-norisoprenoids such as β -damascenone or α -ionone identified previously in orange juice [24]. Overall lack of detection of some of those compounds in our samples could be due to these compounds not being present in our



samples because of biological/environmental variability, although we cannot discard that differences in extraction and analytical techniques used (i.e. exposure of juices to high temperatures) or misidentification of those compounds in previous reports could be the reason.

In summary, over 100 volatile compounds have been unequivocally identified for the first time in the juice of four varieties of *Citrus* using the same analytical conditions, and therefore allowing us to perform more robust comparisons. Cluster and correlation analyses indicated interesting relationships between compounds and classes of compounds revealing the existence of interesting interactions between the biosynthetic pathways. Our results revealed also that the differences in the volatile profile in *Citrus* juice are mainly quantitative, and only a few compounds are variety-specific. What appears to be specific is the profile, i.e. relative content of a set of volatiles. Thus, according to the volatile profile, the most different varieties were Chandler and Powell, while Clemenules and Fortune were intermediate and very similar to one another. In Chandler the most characteristic volatiles were principally aliphatic aldehydes, sesquiterpenes such as nootkatone and monoterpenes such as 2-carene. Powell Navel orange showed the highest levels of esters such as nonyl acetate and of monoterpenes such as 3-carene. Clemenules showed the highest levels of ketones 3-pentanone and β -ionone and Fortune showed the highest levels of some acetate esters such as ethyl and propyl acetate, this latter almost Fortune-exclusive.

Volatile profiling of *Citrus* juice by HS-SPME-GC-MS has proven therefore to be a highly valuable tool for the characterization of fruit from different varieties. The results and volatile platform described in this paper could be used as a roadmap to guide in the selection process of *Citrus* breeding programs directed to obtain new varieties with better aroma, to monitor industrial processes that may affect aroma, and also in the study of the pathways leading to volatile production in *Citrus*.

References

- Tikunov Y, Lommen A, de Vos CHR, Verhoeven HA, Bino RJ, et al. (2005) A novel approach for nontargeted data analysis for metabolomics. Large-scale profiling of tomato fruit volatiles. *Plant Phys* 139: 1125–1137.
- Tikunov YM, de Vos RCH, Paramas AMG, Hall RD, Bovy AG (2010) A Role for Differential Glycoconjugation in the Emission of Phenylpropanoid Volatiles from Tomato Fruit Discovered Using a Metabolic Data Fusion Approach. *Plant Phys* 152: 55–70.
- Moshonas GP, Shaw PE (1997) Quantitation of Volatile Constituents in Mandarin Juices and Its Use for Comparison with Orange Juices by Multivariate Analysis. *J Agric Food Chem* 45: 3968–3972.
- Buettner A, Schieberle P (2001) Evaluation of Key Aroma Compounds in Hand-Squeezed Grapefruit Juice (*Citrus paradisi Macfayden*) by Quantitation and Flavor Reconstitution Experiments. *J Agric Food Chem* 49: 1358–1363.
- Buettner A, Schieberle P (2001) Evaluation of Aroma Differences between Hand-Squeezed Juices from Valencia Late and Navel Oranges by Quantitation of Key Odorants and Flavor Reconstitution Experiments. *J Agric Food Chem* 49: 2387–2394.
- Rouseff RL, Perez-Cacho PR, Jabalpurwala F (2009) Historical Review of Citrus Flavor Research during the Past 100 Years. *J Agric Food Chem* 57: 8115–8124.
- Cavalli J-F, Fernandez X, Lizzani-Cuvellier L, Loiseau A-M (2003) Comparison of Static Headspace, Headspace Solid Phase Microextraction, Headspace Sorptive Extraction, and Direct Thermal Desorption Techniques on Chemical Composition of French Olive Oils. *J Agric Food Chem* 51: 7709–7716.
- González-Mas MC, García-Riño LM, Alfaro C, Rambla JL, Padilla AI, et al. (2009) Headspace-Based Techniques to Identify the Principal Volatile Compounds in Red Grape Cultivars. *Int J Food Sci Tech* 44: 510–518.
- Arthur CL, Pawliszyn J (1990) Solid phase microextraction with thermal desorption using fused silica optical fibers. *Anal Chem* 62: 2145–2148.
- Jia M, Howard Zhang Q, Min DB (1998) Optimization of Solid-Phase Microextraction Analysis for Headspace Flavor Compounds of Orange Juice. *J Agric Food Chem* 46: 2744–2747.
- Jordan MJ, Tillman TN, Mucci B, Laencina J (2001) Using HS-SPME to Determine the Effects of Reducing Insoluble Solids on Aromatic Composition of Orange Juice. *Lebensm-Wiss u-Technol* 34: 244–250.
- Choi H-S (2003) Characterization of Citrus unshiu (C. unshiu Marcov. forma Miyagawa-wase) Blossom Aroma by Solid-Phase Microextraction in Conjunction with an Electronic Nose. *J Agric Food Chem* 51: 418–423.
- Choi H-S, Min K-C (2004) Headspace-SPME Analysis of Citrus Hybrid, Hallabong. *Food Sci Biotechnol* 13: 126–129.
- Jordan MJ, Goodner KL, Castillo M, Laencina J (2005) Comparison of two headspace solid phase microextraction fibres for the detection of volatile chemical concentration changes due to industrial processing of orange juice. *J Sci Food Agric* 85: 1065–1071.
- Arena E, Guarerra N, Campisi S, Asmundo CN (2006) Comparison of odour active compounds detected by gas-chromatography-olfactometry between hand-squeezed juices from different orange varieties. *Food Chem* 98: 59–63.
- Qiao Y, Xie BJ, Zhang Y, Fan G, Yao XL, et al. (2008) Characterization of Aroma Active Compounds in Fruit Juice and Peel Oil of Jinchun Sweet Orange Fruit (*Citrus sinensis* (L.) Osbeck) by GC-MS and GC-O. *Molecules* 13: 1333–1344.
- Barboni T, Luro F, Chiamomanti N, Desjoberg J-M, Muselli A, et al. (2009) Volatile composition of Hybrids Citrus juices by headspace solid-phase micro extraction/gas chromatography/mass spectrometry. *Food Chem* 116: 382–390.
- Brat P, Rega B, Alter P, Reynes M, Brillouet J-M (2003) Distribution of Volatile Compounds in the Pulp, Cloud, and Serum of Freshly Squeezed Orange Juice. *J Agric Food Chem* 51: 3442–3447.
- Selli S, Cabaroglu T, Canbas A (2004) Volatile flavour components of orange juice obtained from the cv. Kozan of Turkey. *J Food Comp Anal* 17: 789–796.
- Nunez AJ, Bemelmans JMH, Maarse H (1984) Isolation methods for the volatile components of grapefruit juice-distillation and solvent-extraction methods. *Chromatographia* 18: 153–158.
- Bylaite E, Meyer AS (2006) Characterisation of volatile aroma compounds of orange juices by three dynamic and static headspace gas chromatography techniques. *Eur Food Res Technol* 222: 176–184.
- Pérez-López AJ, Carbonell-Barrachina AA (2006) Volatile odour components and sensory quality of fresh and processed mandarin juices. *J Sci Food Agric* 86: 2404–2411.

Supporting Information

Figure S1 Principal Component Analysis score plot (t[1] vs t[3]) for the first and third principal components. (TIF)

Figure S2 Principal Component Analysis loading plot (p[1] vs p[3]) for the first and third principal components. Each number corresponds to a particular volatile compound, as indicated in Table 1. (TIF)

Figure S3 Chromatograms representing each of the varieties analyzed: A, Chandler; B, Clemenules; C, Fortune; D, Powell. (TIF)

Table S1 Pearson correlation coefficients for each of the volatile compounds. Significant correlations ($p < 0.01$) are highlighted in bold. (XLS)

Table S2 Volatile organic compounds injected as standard but not identified in any juice analyzed. (DOC)

Acknowledgments

We are grateful to the *Centro de Ecología Química Agrícola* of UPV for its donation of several commercial standards. We thank Silvia Cortes and Isabel Herrero for laboratory assistance, and Dr. Guillermo Bernet for the supply of Chandler fruits. Part of the experiments was performed at the Metabolomic lab of *Instituto de Biología Molecular y Celular de Plantas*, CSIC.

Author Contributions

Conceived and designed the experiments: MCG JLR AGU. Performed the experiments: MCG JLR. Analyzed the data: MCG JLR MCA. Contributed reagents/materials/analysis tools: MCG AGU AGR. Wrote the paper: MCG JLR AGR AGU.

23. Yo SP, Lin CH (2004) Qualitative and Quantitative Composition of the Flavour Components of Taiwan Calamondin and Philippine Calamansi Fruit. *Europ J Hort Sci* 69: 117–124.
24. Mahattanatawee K, Rouseff R, Valim MF, Naim M (2005) Identification and Aroma Impact of Norisoprenoids in Orange Juice. *J Agric Food Chem* 53: 393–397.
25. Pohjanheimo TA, Sandell MA (2009) Headspace volatiles contributing to flavour and consumer liking of wellness beverages. *Food Chem* 115: 843–851.
26. Ruiz Perez-Cacho P, Rouseff RL (2008) Fresh Squeezed Orange Juice Odor: A Review. *Crit Rev Food Sci Nutr* 48: 681–695.
27. Plotto A, Margaria CA, Goodner KL, Goodrich R, Baldwin EA (2004) Odour and flavour thresholds for key aroma components in an orange juice matrix: terpenes and aldehydes. *Flavour Fragr J* 19: 491–498.
28. Plotto A, Margaria CA, Goodner KL, Baldwin EA (2008) Odour and flavour thresholds for key aroma components in an orange juice matrix: esters and miscellaneous compounds. *Flavour Fragr J* 23: 398–406.
29. Lin J, Rouseff RL, Barros S, Naim M (2002) Aroma Composition Changes in Early Season Grapefruit Juice Produced from Thermal Concentration. *J Agric Food Chem* 50: 813–819.
30. Rodrigo MJ, Zacarias L (2006) Horticultural and Quality Aspects of Citrus Fruits. In: Hui YH, Barta J, Cano MP, Gusek TW, Sidhu JS, Sinha NK, eds. *Handbook of Fruits and Fruit Processing*, First edition. Ames: Wiley-Blackwell Publishing. pp 293–296.
31. Soler Aznar J, Soler Fayos G, Forner Giner MA, Forner Valero J (2006) Criterios para identificar variedades y patrones de cítricos (in Spanish, Criteria for identifying citrus varieties and rootstocks). Valencia, Spain: Generalitat Valenciana, Conselleria d'Agricultura, Pesca i Alimentació Press. 156 p.
32. Chisholm MG, Jell JA, Cass DM Jr. (2003) Characterization of the major odorants found in the peel oil of *Citrus reticulata* Blanco cv. Clementine using gas chromatography-olfactometry. *Flavour Fragr J* 18: 275–281.
33. Moshonas MG, Shaw PE (1994) Quantitative Determination of 46 Volatile Constituents in Fresh, Unpasteurized Orange Juices Using Dynamic Headspace Gas Chromatography. *J Agric Food Chem* 42: 1525–1528.
34. Kerbiriou P, Plotto A, Goodener K, Baldwin E, Gmitter FG Jr. (2007) Distribution of Aroma Volatiles in a Population of Tangerine Hybrids. *Proc Fla State Hort Soc* 120: 267–275.
35. Sawamura M, Kuriyama T (1988) Quantitative Determination of Volatile Constituents in the Pummelo (*Citrus grandis* Osbeck forma Tosa-buntan). *J Agric Food Chem* 36: 567–569.
36. Wang Y, Kays SJ (2000) Contribution of Volatile Compounds to the Characteristic Aroma of Baked 'Jewel' Sweetpotatoes. *J Amer Hort Sci* 125: 638–643.
37. Elmaci Y, Altug T (2005) Flavor characterization of three mandarin cultivars (Satsuma, Bodrum, Clemantine) by using GC/MS and flavour profile analysis techniques. *J Food Qual* 28: 163–170.