

Virgin olive oil polyphenol hydroxytyrosol acetate inhibits *in vitro* platelet aggregation in human whole blood: comparison with hydroxytyrosol and acetylsalicylic acid

José Antonio González Correa¹, Juan Antonio López-Villodres¹, Rocío Asensi¹, José Luis Espartero², Guillermo Rodríguez-Gutiérrez³ and José Pedro De La Cruz^{1*}

¹Department of Pharmacology, Laboratorio de Investigaciones Antitrombóticas e Isquemia Tisular (LIAIT), School of Medicine, University of Málaga, Campus de Teatinos s/n, 29071 Málaga, Spain

²Departamento de Química Orgánica y Farmacéutica, Facultad de Farmacia, Universidad de Sevilla, c/ profesor García González 2, 41012 Sevilla, Spain

³Instituto de la Grasa (CSIC), Avda. Padre García Tejero 4, 41012 Sevilla, Spain

(Received 3 June 2008 – Revised 30 July 2008 – Accepted 31 July 2008 – First published online 8 September 2008)

Hydroxytyrosol acetate (HT-AC) is a polyphenol present in virgin olive oil (VOO) at a proportion similar to hydroxytyrosol (HT) (160–479 µmol/kg oil). The present study was designed to measure the *in vitro* platelet antiaggregating activity of HT-AC in human whole blood, and compare this effect with that of HT and acetylsalicylic acid (ASA). The experiments were designed according to the standard procedure to investigate the activity of ASA. HT-AC and HT inhibited platelet aggregation induced by ADP, collagen or arachidonic acid in both whole blood and platelet-rich plasma (PRP). ASA and HT-AC had a greater effect in whole blood than in PRP when ADP or collagen was used as inducer. ASA and HT-AC had a greater effect in PRP + leucocytes than in PRP alone. All three compounds inhibited platelet thromboxane B₂ and leucocyte 6-keto-prostaglandin F_{1α} (6-keto-PF_{1α}) production. The thromboxane/6-keto-PGF_{1α} inhibition ratio (as an indirect index of the prostanoid equilibrium) was 10.8 (SE 1) for HT-AC, 1.0 (SE 0.1) for HT and 3.3 (SE 0.2) for ASA. All three compounds stimulated nitric oxide production, although HT was a weaker effect. In our experiments only concentrations higher than 500 µM (HT) or 1 mM (HT-AC and ASA) inhibited 3-nitrotyrosine production. All three compounds inhibited the production of TNFα by leucocytes, with no significant differences between them. In quantitative terms HT-AC showed a greater antiplatelet aggregating activity than HT and a similar activity to that of ASA. This effect involved a decrease in platelet thromboxane synthesis and an increase in leucocyte nitric oxide production.

Hydroxytyrosol acetate: Olive oil polyphenols: Platelet aggregation: Prostanoids: Nitric oxide

In the Mediterranean diet, which has been associated with a reduced risk of CVD^(1–4), olive oil is the main source of fat. Although a variety of factors have been related with the heart-healthy effect of this diet, there is broad agreement in attributing a major role in this beneficial effect to the polyphenols in virgin olive oil (VOO)⁽⁵⁾. This effect may originate from the action of polyphenols on different biochemical pathways, or from synergism between the effects of polyphenols and fatty acids, particularly oleic acid (a MUFA)⁽⁶⁾.

The polyphenol compound present in highest concentrations in VOO is hydroxytyrosol (3,4-dihydroxyphenylethanol; HT); its concentration varies depending on the variety of olive, geographical area, ripeness of the fruit and method of olive oil pressing, among other factors⁽⁷⁾. For example, VOO from Manzanilla variety contains 2.5–3 times as much HT (479.6 µmol/kg oil) as the Hojiblanca (159.2 µmol/kg oil), Picual (176.8 µmol/kg oil), Cornicabra (164.1 µmol/kg oil) or Arbequina (179.6 µmol/kg oil) varieties⁽⁷⁾.

The effect of polyphenols on physiopathological mechanisms involved in CVD has been characterized only in part. These effects include inhibition of platelet aggregation^(8,9), inhibition of oxidation of LDL⁽¹⁰⁾, stimulation of NO production⁽¹¹⁾ and down-regulation of the expression of endothelial adhesion molecules⁽¹²⁾. Platelet aggregation is considered one of the main events in arterial thrombosis, and this makes investigating how aggregation is inhibited by polyphenols in VOO a worthwhile topic for further inquiry.

Although HT is the major polyphenol on VOO, other polyphenols are present, including hydroxytyrosol acetate (HT-AC)⁽⁷⁾. The concentration of HT-AC is approximately equal to that of HT in some olive varieties such as Arbequina, but is twice as high as that of HT in the Picual variety, and three to four times as high in the Manzanilla and Hojiblanca varieties⁽⁷⁾.

The aims of the present study were to measure the antiplatelet aggregating effect *in vitro* of HT-AC in human blood,

Abbreviations: ASA, acetylsalicylic acid; HT, hydroxytyrosol; HT-AC, hydroxytyrosol acetate; 6-keto-PF_{1α}, 6-keto-prostaglandin F_{1α}; LPS, lipopolysaccharide; PRP, platelet-rich plasma; VOO, virgin olive oil.

* **Corresponding author:** Dr J. P. De La Cruz, fax +34 952 131568, email jpcruz@uma.es

and to compare its effects to HT and acetylsalicylic acid (ASA), the most widely used drug in the world to prevent ischaemic CVD because of its antiplatelet aggregating action.

Experimental methods

Materials

Thromboxane B₂, 6-keto-prostaglandin F_{1α} (6-keto-PF_{1α}) and TNFα enzyme immunoassay kits were from Amersham International plc (Little Chalfont, UK). Nitrite/nitrate and the 3-nitrotyrosine ELISA kits and HT were obtained from Cayman Chemical (Ann Arbor, MI, USA). HT-AC was supplied by the Department of Organic Chemistry and Pharmaceutics, School of Pharmacy, University of Seville (Seville, Spain). Collagen and ADP were obtained from Menarini Diagnóstica S.A. (Barcelona, Spain). All other reagents were from Sigma Chemical Co. (St. Louis, MO, USA).

Study design

The present *in vitro* study was carried out in blood samples from healthy volunteers who had not taken any medication during the 15 d before inclusion and VOO in the diet 24 h before. Sodium citrate (3.8%) at a proportion of 1:10 was used as the anticoagulant. In each group of experiments we used six samples from different volunteers (three men and three women in each group) with a mean age of 32.8 (SE 1.3) years. The study was approved by the Clinical Assays Committee of the Hospital Regional Universitario Carlos Haya in Malaga, Spain.

Samples were used as whole blood, but in some experiments, platelet-rich plasma (PRP), erythrocytes and leucocytes were isolated from whole blood. PRP was obtained after centrifugation of whole blood at 800 g for 10 min at 20°C; all PRP samples were adjusted to 3×10^9 platelets/l. Erythrocytes were obtained from the pellet resulting from centrifugation of whole blood to obtain PRP; erythrocytes were washed twice with PBS (pH 7.4). Leucocytes were separated on a Ficoll density gradient (Hystopaque 1077 and Hystopaque 1119) and diluted (final count: 5.2 (SE 0.2) $\times 10^9$ leucocytes) in Ca²⁺ physiological saline solution (95% viability as judged by trypan blue exclusion).

Different concentrations of ASA, HT and HT-AC (10^{-7} to 10^{-3} M) were incubated at 37°C before inducers were added. All drugs were diluted in a mixture of ethanol–PBS (100 μl ethanol plus 900 μl PBS) at a proportion of 1:10.

Analytical techniques

All techniques were run in a single-blind manner, i.e. the persons who did the assays were unaware of the origin and nature of the samples.

Platelet aggregometry

Platelet aggregation was measured as electronic impedance in whole blood, PRP, PRP plus erythrocytes, and PRP plus leucocytes (500 μl sample diluted with 500 μl PBS). We used a Chrono-Log 540 aggregometer (Chrono-Log Corp., Haverton, PA, USA) with collagen (1 μg/ml), ADP (2.5 μM) and

arachidonic acid (400 μM) to induce aggregation. Drugs were incubated at 37°C for 10 min before the aggregation inducer was added, and aggregation was recorded for 10 min. Maximum intensity of aggregation was quantified as the maximum change in electronic impedance in samples without the drug or a given concentration of each drug. The concentrations of aggregating agents were chosen according to previous experiments which yielded EC₅₀ values of 1.10 (SE 0.14) μg/ml for collagen (*n* 10), 2.41 (SE 0.11) μM for ADP and 385 (SE 21) μM for arachidonic acid (*n* 10). Erythrocytes and leucocytes were added in a proportion to reach the cell counts in whole blood (4.8 (SE 0.2) $\times 10^{12}$ erythrocytes/l and 6.0 (SE 0.4) $\times 10^6$ leucocytes/l).

Platelet production of thromboxane B₂

The production of thromboxane A₂ was determined by measuring its stable metabolite thromboxane B₂. Samples of whole blood were stimulated with calcium ionophore A23187 (50 μM) at 37°C during 30 min, then they were centrifuged at 10 000 g for 3 min, and the amount of thromboxane B₂ in the supernatant was determined with an enzyme immunoassay kit.

Leucocyte production of 6-keto-prostaglandin F_{1α}

The production of prostacyclin was determined by measuring its stable metabolite 6-keto-PF_{1α}. We used samples of leucocytes (6.5 (SE 0.4) $\times 10^6$ leucocytes/l) that were incubated for 5 min at 37°C, then 50 μM-calcium ionophore A23187 was added and the samples were shaken for 30 min. The samples were centrifuged at 10 000 g for 15 min at 4°C, and were frozen at –80°C until analysis for 6-keto-PF_{1α} by enzyme immunoassay.

Leucocyte production of nitric oxide

NO production was induced with the calcium ionophore A23187 (50 μM) and measured as nitrite + nitrate concentration in the supernatant of induced leucocyte samples (6.3 (SE 0.3) $\times 10^6$ leucocytes/l) with an enzyme immunoassay kit. The experiments were designed to determine NO production with the drug under study or in ethanol–PBS solution (control sample) for 10 min at 37°C. Briefly, leucocytes were incubated for 10 min at 37°C with 200 μl buffer containing 10 mM-HEPES, 1 mM-EDTA, 1 mM-DL-dithiothreitol (DTT), 10 μg/ml leupeptin, 4.8 mM-DL-valine, 1 mM-NADPH, 1 mM-MgCl₂, 1 mM-CaCl₂ and 20 μM-L-arginine, and then with 10 μl saline or the corresponding concentration of the drugs. Basal nitrite + nitrate production was first determined; after this step calcium ionophore A23187 was added and after 30 min of incubation at 37°C the increase in nitrite + nitrate production was evaluated.

3-Nitrotyrosine and TNFα production

Blood was incubated with bacterial membrane lipopolysaccharide (LPS; 1 μg/ml) for 18 h at 37°C. The plates were then centrifuged, and the plasma was removed and frozen at –80°C. TNFα and 3-nitrotyrosine (an indicator of peroxynitrite production) were measured in the supernatant with a

commercial enzyme immunoassay kit. In each experiment a sample of blood without LPS induction was incubated, and LPS-induced TNF α and 3-nitrotyrosine production was calculated as the difference between the LPS-induced samples and non-induced samples.

Statistical analysis

The data are expressed as means and their standard errors for ten experiments. All statistical analyses were done with the Statistical Program for Social Sciences version 12.0 (SPSS Inc., Chicago, IL, USA). One-way ANOVA followed by Bonferroni transformation and unpaired Student's *t* test were used, and differences were considered significant when $P < 0.05$.

Results

HT-AC and HT inhibited platelet aggregation induced by ADP, collagen or arachidonic acid in a concentration-dependent manner in whole blood and PRP (Figs. 1 and 2). Maximum intensity of aggregation in cultures induced with ADP was 23.51 (SE 1.95) ohms in whole blood and 19.47 (SE 1.10) ohms in PRP. After induction with collagen, maximum aggregation was 21.24 (SE 1.16) ohms in whole blood and 18.97 (SE 1.15) ohms in PRP, and in cultures induced with arachidonic acid, maximum aggregation was 23.49 (SE 0.93) ohms in whole blood and 23.75 (SE 0.95) ohms in PRP. Table 1 shows the concentrations of each compound that inhibited maximum platelet aggregation in control samples (incubated without the aggregation inducer) by 50% (IC₅₀). In whole blood, the antiplatelet aggregating effect of HT was significantly weaker than that of HT-AC or ASA. Inhibition of aggregation in whole blood was similar with HT-AC and ASA except when arachidonic acid was the inducer; in these experiments the antiaggregating effect of ASA was stronger. In samples of PRP induced with ADP or collagen we found no significant differences among the three compounds, but when arachidonic acid was the inducer, the effect of HT was significantly weaker than HT-AC or ASA.

When we compared the antiaggregating effect in whole blood and PRP, we found that the effect of HT-AC and ASA was significantly stronger when aggregation was induced with ADP or collagen (Table 1); the antiaggregating effect of HT in whole blood was similar to that in PRP. In PRP incubated with leucocytes the antiaggregating effect of HT-AC and ASA increased in comparison to the effect in PRP alone (Fig. 3). In PRP incubated with erythrocytes the antiaggregating effects of HT, HT-AC and ASA were not different statistically with respect to PRP. Maximum aggregation was 21.75 (SE 1.12) ohms in whole blood, 19.00 (SE 0.56) ohms in PRP, 21.59 (SE 1.74) ohms in PRP with erythrocytes and 16.08 (SE 1.11) ohms in PRP with leucocytes.

All three compounds inhibited thromboxane B₂ and 6-keto-PF_{1 α} production in a concentration-dependent manner (Fig. 4). Table 2 shows the IC₅₀ values for these assays. The order of potency in the inhibition of these two prostanoids was ASA > HT-AC > HT. The thromboxane/prostacyclin inhibition ratio was 10.79 (SE 1.03) for HT-AC, 1.05 (SE 0.09) for HT and 3.28 (SE 0.21) for ASA. Thromboxane B₂ production in control samples was 1.29 (SE 0.09) nmol/10⁹

platelets and 6-keto-PF_{1 α} production was 1.16 (SE 0.09) nmol/10⁹ leucocytes.

All three compounds increased nitrite + nitrate production in whole blood in a concentration-dependent manner (Fig. 5), although the effect of HT was significantly weaker than that of HT-AC and ASA (Table 2). Nitrite + nitrate concentration in control samples was 1.05 \pm 0.01 μ mol/l. The production of 3-nitrotyrosine induced with LPS was significantly inhibited only by the highest concentration of the compounds we tested (1000 μ M) (Fig. 5; Table 2). The concentration of




Fig. 1. Platelet aggregation in human whole blood after incubation with hydroxytyrosol acetate (■), hydroxytyrosol (●) and acetylsalicylic acid (◆). The inducers were ADP (2.5 μ M; A), collagen (1 μ g/ml; B) or arachidonic acid (400 μ M; C). Values are means with their standard errors depicted by vertical bars.




Fig. 2. Platelet aggregation in human platelet-rich plasma after incubation with hydroxytyrosol acetate (■), hydroxytyrosol (●) and acetylsalicylic acid (◆). The inducers were ADP (2.5 µM; A), collagen (1 µg/ml; B) or arachidonic acid (400 µM; C). Values are means with their standard errors depicted by vertical bars.

3-nitrotyrosine in control samples was 39.85 (SE 3.28) nmol/l. The production of TNF α was also inhibited in a concentration-dependent manner by all three compounds (Fig. 5), with no significant differences among them (Table 2). The concentration of TNF α in control samples was 10.36 (SE 1.21) pg/ml.

Discussion

We report the first evidence of the antiplatelet aggregating effect of HT-AC in human blood. The effect of HT-AC was stronger than that of HT and similar in quantitative terms to that of ASA. Some phenolic compounds in VOO inhibit

Table 1. Concentrations of hydroxytyrosol acetate (HT-AC), hydroxytyrosol (HT) and acetylsalicylic acid (ASA) that inhibited 50% of platelet aggregation (IC₅₀), induced with 2.5 µM-ADP, 1 µg/ml collagen or 400 µM-arachidonic acid in whole-blood and platelet-rich plasma (PRP)† (Mean values with their standard errors)

	IC ₅₀ (µM)					
	HT-AC		HT		ASA	
	Mean	SE	Mean	SE	Mean	SE
ADP						
Whole blood	94.17	7.58	738†	90	87.21	6.11
PRP	543*	35	625	85	813*	6.11
Collagen						
Whole blood	26.48	1.66	195†	21	24.94	1.18
PRP	109*	9.64	109	11	281*	28
Arachidonic acid						
Whole blood	40.08	4.10	197†	20	9.28	0.81
PRP	39.57	3.25	131†	15	6.64	0.40

Mean values were significantly different from those of whole blood (unpaired Student's *t* test): **P*<0.0001.

Mean values were significantly different from those of ASA and HT-AC: †*P*<0.0001.

‡ For details of subjects and procedures, see Experimental methods.

platelet function, e.g., HT^(8,9), oleuropein⁽⁸⁾ and phenolic isochromans⁽¹³⁾. In fact, feeding VOO to rats or rabbits was found to significantly reduce collagen-induced platelet aggregation^(14,15). The present results confirm this effect of HT, but also show that HT-AC has a greater antiaggregating effect than HT, and is almost as effective as ASA in blocking platelet aggregation. The latter observation is potentially important because ASA is the most widely used drug in the world in prophylaxis against thromboembolism.

In the present section we analyse the results obtained with HT and HT-AC in relation to the mechanisms of action for ASA, as the profile of inhibition of platelet aggregation of HT-AC is similar to that of ASA (Table 1; Figs. 1 and 2).

Firstly, the main mechanism that explains the antithrombotic effect of ASA is inhibiting the synthesis of thromboxane A₂. All three compounds inhibit thromboxane synthesis in a concentration-dependent manner. It was previously shown that HT at a concentration of 400 µM in whole blood reduced collagen- or thrombin-induced thromboxane production by approximately 75%⁽⁸⁾. Interpolating the results plotted in Fig. 4 for a concentration of 400 µM of the phenolic compounds compared here shows that HT would reduce thromboxane production by 67% and HT-AC would reduce production by 86%.

Secondly, the synthesis of thromboxane A₂ can be reduced by inhibition of the enzyme cyclooxygenase (as acting ASA) or the enzyme thromboxane synthetase. The effect of HT and HT-AC cannot be attributed to the inhibition of thromboxane synthase activity, since prostacyclin production was also inhibited (Fig. 4). The present finding suggests that cyclooxygenase was probably the target of these phenols. More than 30 years ago this mechanism of action was shown in ASA⁽¹⁶⁾, and it also appears to be important in the antiaggregating effect of HT-AC. Although platelet thromboxane and leucocyte prostacyclin production are not strictly comparable, they provide an approximate idea of the effect of these compounds on the thromboxane/prostacyclin balance through the IC₅₀ prostacyclin/IC₅₀ thromboxane ratio (see Results).




Fig. 3. Maximum intensity of platelet aggregation (I_{max}) induced with 1 $\mu\text{g/ml}$ collagen in whole blood (WB), platelet-rich plasma (PRP), PRP plus erythrocytes (PRP + RBC) and PRP plus leucocytes (PRP + L) after incubation with hydroxytyrosol acetate (A), hydroxytyrosol (B) or acetylsalicylic acid (C). Values are means with their standard errors depicted by vertical bars. Mean values were significantly different from those of WB: * $P < 0.05$, ** $P < 0.001$. Mean values were significantly different from those of PRP: + $P < 0.05$, ++ $P < 0.001$.

HT-AC inhibited thromboxane production more than prostacyclin production, an effect with clear benefits with a view to preventing thrombosis. The cyclooxygenase-inhibiting mechanism plays a potentially important role, since it was recently shown that HT has no effect on other pathways involving platelets such as cAMP, at least not at a concentration of 100 μM ⁽⁹⁾. The present findings are consistent with the weaker antiaggregating effect of both phenolic compounds and ASA when aggregation was induced by ADP, since the effect of ADP is mainly the result of its ability to lower cytoplasmic concentrations of cAMP^(17,18). However, this is not to imply




Fig. 4. Whole-blood thromboxane B_2 (A) and leucocyte keto-prostaglandin $F_{1\alpha}$ (B) induced with 50 μM -calcium ionophore A23187 after incubation with hydroxytyrosol acetate (■), hydroxytyrosol (●) and acetylsalicylic acid (◆). Values are means with their standard errors depicted by vertical bars.

that inhibition of thromboxane synthesis is the only mechanism through which HT exerts its antithrombotic potential *in vivo*, since it has also been shown to inhibit the expression of adhesion molecules such as intercellular adhesion molecule-1 and vascular cell adhesion molecule-1 expressed on endothelial cells⁽¹²⁾.

Thirdly, other mechanisms can complete inhibition of cyclooxygenase produced by HT and HT-AC. This is supported in the differential antiaggregant effects in isolated platelets and in whole blood. We found that HT had the same antiaggregating effect in both media, whereas the effect of HT-AC and ASA was stronger in whole blood, albeit with some exceptions. When arachidonic acid was used as the inducer, aggregation resulted from the stimulation of platelet thromboxane A_2 synthesis⁽¹⁹⁾. The absence of differences in the antiaggregant effect of ASA, HT and HT-AC is explained by the inhibition of cyclooxygenase. The greater antiaggregating effect of ASA and HT-AC in whole blood when platelets were induced with ADP or collagen might be due to an effect on a different biochemical pathway that mediates in the relations between leucocytes, erythrocytes and platelets. In this connection it was found that the greater effect of ASA in whole blood was mainly due to the stimulation of neutrophil NO production^(18,20), a finding also reported for other salicylate derivatives⁽²¹⁾.

Because erythrocytes did not modify the antiplatelet effect of HT and HT-AC, leucocytes could probably influence this effect through a modification in its NO production. ASA

Table 2. Concentrations of hydroxytyrosol acetate (HT-AC), hydroxytyrosol (HT) and acetylsalicylic acid (ASA) that inhibited 50% (IC₅₀) calcium ionophore-induced thromboxane B₂ and 6-keto-prostaglandin F_{1α} (6-keto-PGF_{1α}), lipopolysaccharide-induced 3-nitrotyrosine and TNFα, and concentrations of these compounds that increased calcium ionophore-induced nitrite + nitrate production by 50% (EC₅₀)‡ (Mean values with their standard errors)

	HT-AC		HT		ASA	
	Mean	SE	Mean	SE	Mean	SE
IC ₅₀ (μM)						
Thromboxane B ₂	5.31	0.63	63.64*	4.84	1.37	0.09
6-Keto-PGF _{1α}	58.33	6.00	69.40	6.41	4.53†	0.22
3-Nitrotyrosine	> 1000		658	26	> 1000	
TNFα	67.37	6.91	65.48	7.64	76.81	4.91
EC ₅₀ (μM)						
Nitrites + nitrates	1.79	0.15	101*	10	5.58	0.40

Mean values were significantly different from those of ASA and HT-AC: **P*<0.0001.

Mean values were significantly different from those of HT and HT-AC: †*P*<0.0001.

‡ For details of subjects and procedures, see Experimental methods.

and the two phenolic compounds stimulated leucocyte NO production in a concentration-dependent manner (Fig. 5), although the effect of HT was smaller in quantitative terms than that of HT-AC or ASA. The reported ability of ASA to stimulate NO production was confirmed by the present finding that in PRP plus erythrocytes, the effect of ASA and HT-AC was similar to their effect in PRP alone, whereas in PRP plus leucocytes, the effect of HT-AC was similar to its effect in whole blood, as was also noted for ASA in an earlier report⁽¹⁸⁾ and in the present study (Fig. 3). HT has been shown to increase the cytoplasmic concentration of calcium and thus contribute to the increase in NO synthesis by endothelial cells^(22,23). Our assays showed that HT-AC is also able to stimulate calcium-dependent NO production in human leucocytes.

Therefore, HT could inhibit platelet function in whole blood through an inhibition of platelet thromboxane A₂ synthesis, while HT-AC could exert a greater antiaggregant effect because of the inhibition of thromboxane A₂ synthesis plus an increase in NO leucocyte production, similar to the mechanism of action of ASA.

Fourthly, the effect of ASA on the production of inflammatory mediators has been related to its action in preventing CVD^(24,25); accordingly, we tested TNFα production in leucocytes stimulated with LPS. All three compounds compared here inhibited the production of this inflammatory mediator in a concentration-dependent way, with no differences among them in the intensity of their effect. In this connection it has been shown that exposure of LPS-induced human THP-1 cells to olive vegetation water (containing approximately 200 μM-polyphenols) reduced TNFα production by 50%⁽²⁶⁾. In the present experimental procedure, the effect of both HT and HT-AC was greater compared to the effect reported by Bitler *et al.*⁽²⁶⁾ Therefore, in addition to the antiplatelet effect of HT and HT-AC, it would be necessary to consider its effect on inhibiting the production of inflammatory mediators as an additional mechanism that could fill a possible effect in the prevention of CVD, similar to ASA.

Finally, in the genesis and evolution of vascular disease oxygen reactive species play an important role. In connection with the production of NO, the presence of free radicals facilitates peroxynitrite⁽²⁷⁾, we wished to study the potential effect of free radicals on 3-nitrotyrosine formation as an

indirect indicator of peroxynitrite production⁽²⁸⁾. Despite the antioxidant effect of polyphenols in VOO^(29,30), these compounds failed to inhibit peroxynitrite production, at least under the present experimental conditions (Fig. 5). Moreover, this antioxidant effect occurs in a range of concentrations greater than the inhibition of platelet function, therefore it does not appear that the antioxidant effect of these compounds has a direct relationship with its antiplatelet effect. In addition, blood samples used for platelet aggregation do not have conditions for formation of free radicals, so the antioxidant effect of HT and HT-AC could not influence their antiplatelet effect.

The *in vivo* significance of the results presented here with regard to inhibition of platelet aggregation by VOO phenols, mainly HT and HT-AC, must be related to its pharmacokinetic characteristics. After ingestion of 40 ml VOO with a phenolic content of 366 mg/kg olive oil, maximal plasma concentration of HT reached 25.83 μg/l in man⁽³¹⁾, which represents approximately 130 μmol HT/l. However, around 95% of HT is present in plasma in conjugate forms, suggesting an extensive first pass intestinal/hepatic metabolism of the ingested primary forms⁽³²⁾. According to these data only 1–2 μmol free HT/l is present in human plasma after ingestion of VOO. Platelet aggregation curves obtained in the present study show around 20–30% inhibition, and the corresponding thromboxane B₂ production was 30–40% for HT-AC (data obtained by extrapolating the results at 1–2 μmol/l). The possible inhibition of platelet function in the presystemic circulation, as has been demonstrated for ASA⁽³³⁾, and the need to investigate the pharmacokinetic parameters of the administration of HT or HT-AC, remain to be established in order to postulate the importance of these results in the prevention of cardiovascular events.

In summary, the present results show that HT-AC, a polyphenol present in VOO, exerts an antiaggregating effect on platelets that is greater in quantitative terms than the effect of HT and similar to the effect of ASA. This effect involves a decrease in platelet thromboxane synthesis and an increase in leucocyte NO production. The present findings, together with the reported antioxidant effect of HT-AC and its ability to inhibit TNFα production, suggest the need for further studies of the effect of HT-AC on the suite of mechanisms that regulate thrombogenesis in man.




Fig. 5. Concentration of 50 µM-calcium ionophore-induced nitrites + nitrates in leucocytes (A), and 3-nitrotyrosine (B) and TNF α (C) induced with 1 µg/ml bacterial membrane lipopolysaccharide in whole blood, after incubation with hydroxytyrosol acetate (■), hydroxytyrosol (●) and acetylsalicylic acid (◆). Values are means with their standard errors depicted by vertical bars.

Acknowledgements

This study was partially supported by a grant from the Ministerio de Ciencia y Tecnología, Spain (AGL-04-7935-C03-02). None of authors have any conflict of interest. We thank A. Pino for his invaluable technical assistance and K. Shashok for translating parts of the manuscript into English. J. P. D. L. C. and J. A. G. C. designed the study and prepared the manuscript. J. A. L.-V., R. A., J. A. G. C. and J. P. D. L. C. contributed to the successful execution of the experimental work. J. L. E. and G. R.-G. contributed to the chemical preparation, synthesis and analysis of the polyphenols used in the study.

References

- Barzi F, Woodward M, Marfisi RM, Tavazzi L, Valagussa F, Marchioli R & GISSI-Prevenzione Investigators (2003) Mediterranean diet and all-causes mortality after myocardial infarction: results from the GISSI-Prevenzione trial. *Eur J Clin Nutr* **57**, 604–611.
- de Lorgeril M, Salen P, Martin JL, Mamelle N, Monjaud I, Tou-boul P & Delaye J (1996) Effect of a Mediterranean type diet on the rate of cardiovascular complications in patients with coronary artery disease. *J Am Coll Cardiol* **28**, 1103–1108.
- Knoops KT, Groot de LC, Fidanza F, Alberti-Fidanza A, Kromhout D & van Staveren WA (2006) Comparison of three different dietary scores in relation to 10-year mortality in elderly European subjects: the HALE project. *Eur J Clin Nutr* **60**, 746–755.
- Trichopoulou A, Orfanos P, Norat T, *et al.* (2005) Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *Br Med J* **330**, 991–998.
- Covas MI (2007) Olive oil and the cardiovascular system. *Pharmacol Res* **55**, 175–186.
- Perez-Jimenez F, Delgado J, Pérez-Martínez P, López-Segura F, Fuentes F, Cortés B, Lozano A & López-Miranda J (2006) Olive oil and haemostasis: a review on its healthy effects. *Public Health Nutr* **9**, 1083–1088.
- Romero C, Medina E, Vargas J, Brenes M & De Castro A (2007) *In vitro* activity of olive oil polyphenols against helicobacter pylori. *J Agric Food Chem* **55**, 680–686.
- Petroni A, Blasevich M, Salami M, Papini N, Montedoro GF & Galli C (1995) Inhibition of platelet aggregation and eicosanoid production by phenolic components of olive oil. *Thromb Res* **78**, 151–160.
- Dell'Agli M, Maschi O, Galli GV, Fagnani R, Dal Cero E, Caruso D & Boasio E (2007) Inhibition of platelet aggregation by olive oil phenols *via* cAMP-phosphodiesterase. *Br J Nutr* **99**, 945–951.
- Fitó M, Covas MI, Lamuela-Raventós RM, Vila J, Torrents L, de la Torre C & Marrugat J (2000) Protective effect of olive oil and its phenolic compounds against low density lipoprotein oxidation. *Lipids* **35**, 633–638.
- Pignatelli P, Di Santo S, Buchetti B, Sanguigni V, Brunelli A & Violi F (2006) Polyphenols enhance platelet nitric oxide by inhibiting protein kinase C-dependent NADPH oxidase activation: effect on platelet recruitment. *FASEB J* **20**, 1082–1089.
- Dell'Agli M, Fagnani R, Mitro N, *et al.* (2006) Minor components of olive oil modulate proatherogenic adhesion molecules involved in endothelial activation. *J Agric Food Chem* **54**, 3259–3264.
- Togna GI, Togna AR, Franconi M, Marra C & Guiso M (2003) Olive oil isochromans inhibit human platelet reactivity. *J Nutr* **133**, 2532–2536.
- De La Cruz JP, Villalobos MA, Carmona JA, Martín-Romero M, Smith-Agreda JM & Sánchez de la Cuesta F (2000) Anti-thrombotic potential of olive oil administration in rabbits with elevated cholesterol. *Thromb Res* **100**, 305–315.
- González-Correa JA, Muñoz-Marín J, Arrebola MM, Guerrero A, Narbona F, López-Villodres JA & De La Cruz JP (2007) Dietary virgin olive oil reduces oxidative stress and cellular damage in rat brain slices subjected to hypoxia-reoxygenation. *Lipids* **42**, 921–929.
- Vane JR (1971) Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nat New Biol* **231**, 232–235.
- De la Cruz JP, Bellido I, Camara S, Martos F & Sanchez de la Cuesta F (1986) Effects of acetylsalicylic acid on platelet aggregation in male and female whole blood: an *in vitro* study. *Scand J Haematol* **36**, 394–397.

18. De la Cruz JP, Camara S, Bellido I, Carrasco T & Sanchez de la Cuesta F (1987) Platelet aggregation in human whole blood after chronic administration of aspirin. *Thromb Res* **46**, 133–140.
19. Zucker MB & Nachmias VT (1985) Platelet activation. *Arteriosclerosis* **5**, 2–18.
20. Lopez Farre A, Caramelo C, Esteban A, Alberola ML, Millas I, Monton M & Casado S (1995) Effect of aspirin on platelet–neutrophil interaction. Role of nitric oxide and endothelin-1. *Circulation* **91**, 2080–2088.
21. Sánchez de Miguel L, Casado S, Farré J, *et al.* (1998) Comparison of *in vitro* effects of triflusal and acetylsalicylic acid on nitric oxide synthesis by human neutrophils. *Eur J Pharmacol* **343**, 57–65.
22. Visioli F, Bellosa S & Galli C (1998) Oleuropein, the bitter principle of olives, enhances nitric oxide production by mouse macrophages. *Life Sci* **62**, 541–546.
23. Visioli F & Galli C (2002) Biological properties of olive oil phytochemicals. *Crit Rev Food Sci Nutr* **42**, 209–221.
24. Shackelford RE, Alford PB, Xue Y, Thai SF, Adams DO & Pizzo S (1997) Aspirin inhibits tumor necrosis factor alpha gene expression in murine tissue macrophages. *Mol Pharmacol* **52**, 421–429.
25. Mateos-Cáceres PJ, García-Méndez A, Farré J, *et al.* (2002) Prior aspirin use in unstable angina patients with modified plasma inflammatory markers and endothelial nitric oxide synthase in neutrophils. *Eur J Clin Invest* **32**, 895–900.
26. Bitler CM, Viale TM, Damaj B & Crea R (2005) Hydrolyzed olive vegetation water in mice has anti-inflammatory activity. *J Nutr* **135**, 1475–1479.
27. Beckman JS & Koppenol WH (1996) Nitric oxide, superoxide and peroxynitrite: the good, the bad, and the ugly. *Am J Physiol Cell Physiol* **271**, C1424–C1437.
28. Tarpey MM, Wink DA & Grisham MB (2004) Methods for detection of reactive metabolites of oxygen and nitrogen: *in vitro* and *in vivo* considerations. *Am J Physiol Regul Integr Comp Physiol* **286**, R431–R444.
29. Mateos R, Dominguez MM, Espartero JL & Cert A (2003) Antioxidant effect of phenolic compounds, alpha-tocopherol, and other minor components in virgin olive oil. *J Agric Food Chem* **51**, 7170–7175.
30. Gordon MH, Paiva-Martins F & Almeida M (2001) Antioxidant activity of hydroxytyrosol acetate compared with that of other olive oil polyphenols. *J Agric Food Chem* **49**, 2480–2485.
31. Miró-Casas E, Covas MI, Farré M, Fitò M, Ortuño J, Weinbrenner T, Roset P & De La Torre R (2003) Hydroxytyrosol disposition in humans. *Clin Chem* **49**, 945–952.
32. Fitò M, De La Torre R, Farré M, Khymenetz O, Marrugat J & Covas MI (2007) Bioavailability and antioxidant effect of olive oil phenols compounds in humans: a review. *Ann Ist Super Sanita* **43**, 375–381.
33. FitzGerald GA, Lupinetti M, Charman SA & Charman WN (1991) Presystemic acetylation of platelets by aspirin: reduction in rate of drug delivery to improve biochemical selectivity for thromboxane A₂. *J Pharmacol Exp Ther* **259**, 1043–1049.