

17 β -Estradiol – A New Modulator of Neuroglobin Levels in Neurons: Role in Neuroprotection against H₂O₂-Induced Toxicity

Elisabetta De Marinis^a Paolo Ascenzi^{a,b} Marco Pellegrini^a Paola Galluzzo^a
Pamela Bulzomi^a Maria Angeles Arevalo^c Luis Miguel Garcia-Segura^c
Maria Marino^a

^aDepartment of Biology and ^bInterdepartmental Laboratory of Electron Microscopy, University Roma Tre, Rome, Italy; ^cInstituto Cajal, CSIC, Madrid, Spain

Key Words

Neuroglobin • Estrogen receptor • H₂O₂ neurotoxicity • Neuroprotection • Apoptosis

Abstract

Although discovered in 2000, neuroglobin (Ngb) functions are still uncertain. A contribution to the role played by Ngb in neurons could certainly derive from the identification of Ngb endogenous modulators. Here, we evaluate the possibility that Ngb could be regulated by 17 β -estradiol (E₂) signaling in both SK-N-BE human neuroblastoma cell line and mouse hippocampal neurons. 1 nM E₂ rapidly induced a 300% increase in Ngb levels in both models. The E₂ effect was specific, being not induced by testosterone or dihydrotestosterone. The E₂-induced Ngb increase requires estrogen receptor (ER) β , but not ER α , as evaluated by the mimetic effect of ER β -specific agonist DPN and by the blockage of E₂ effect in ER β -silenced SK-N-BE cells. Furthermore, both rapid (15 min) ER β -dependent activation of p38/MAPK and transcriptional ER β activity were required for the estrogenic regulation of Ngb. Finally, E₂ exerted a protective effect against H₂O₂-induced neuroblastoma cell death which was completely prevented in Ngb-silenced cells. Overall,

these data suggest that Ngb is part of the E₂ signaling mechanism that is activated to exert protective effects against H₂O₂-induced neurotoxicity. Copyright © 2011 S. Karger AG, Basel

Introduction

Neuroglobin (Ngb), the third member of the globin family [1], is a monomeric hexa-coordinated heme protein of 17 kDa expressed not only in neurons of the central and peripheral nervous systems, but also in the gastrointestinal tract and in endocrine organs [1–18]. Recently, Ngb has also been detected in human glioblastoma cell lines [17] and in quiescent astrocytes of the healthy seal brain [15]. Although Ngb occurs at relatively low concentrations (μ M) in a wide range of tissues, Ngb is found at relatively high concentrations in highly metabolically active cells and certain specialized cells, such as neurons of the hypothalamus, and particularly in retinal rod cells where its concentration has been estimated to be up to 100 μ M [18–21]. Ngb binds several ligands, including diatomic gaseous ligands, and displays (pseudo-)enzymatic properties [1, 3, 6, 10, 22–33]. The *P*₅₀ value for O₂ binding

to Ngb has been reported to range between 2 and 10 Torr depending on pH, temperature, and the redox state of the cell [1, 22–24, 32].

Although discovered in 2000 [1], the cell function(s) of Ngb is still controversial. Indeed, the O₂ supply by Ngb to the mitochondria of the metabolically active neurons and retinal rod cells is highly debated [1, 15, 19, 33, 34]. In vitro, Ngb has been reported to scavenge nitrogen monoxide (NO) in the presence of high O₂ levels [25, 28, 35]; however, at low O₂ conditions Ngb may react with NO₂⁻ resulting in the formation of NO [36]. Therefore, the protective role of Ngb against NO in vivo is controversial [32, 33, 37]. Although in vitro Ngb does not react with hydrogen peroxide (H₂O₂) [25, 30] and the Ngb-NO₂ adduct reacts with H₂O₂ facilitating the nitration of aromatic substrates [30], the correlation between reactive oxygen species formation/decomposition and Ngb expression in vivo is debated [15, 38]. Moreover, Ngb has been reported to interact with several proteins (see [15]). In particular, Ngb binding to the G α protein inhibits GDP dissociation, thereby protecting cells from apoptosis [39, 40]. Recently, in silico simulations indicate Ngb capability to reduce cytochrome *c* released from mitochondria suggesting its protective role against programmed cell death [19, 41–43].

Although Ngb properties are highly debated [12, 15, 19, 33, 44], it is unlikely that Ngb has so many distinct roles [15]; nevertheless, there is no doubt that Ngb is beneficial to neurons [15]. In vivo experiments, using transgenic rodents, have shown that increased levels of Ngb significantly protect both heart and brain tissues from hypoxic insult, whereas decreased Ngb levels lead to an exacerbation of tissue death [45–47]. In this way, Ngb could protect neurons from hypoxic insult by modulating the activation of the apoptotic cascade [19, 41]. A significant contribution to highlight the role played by Ngb in neuroprotection could derive from the identification of Ngb endogenous modulator(s) (e.g., hormones and neurotransmitters), but, as far as we know, no Ngb involvement in the hormone signal transduction pathways has been identified yet.

Female sex steroid hormones could represent good candidates as Ngb modulators. Indeed, in addition to their well-established role in reproductive organs, estrogens affect areas of the brain that are not primarily involved in reproduction [48]. Growing evidence documents profound effects of estrogens on learning, memory, and mood as well as neurodevelopmental and neurodegenerative processes [49, 50]. Although most studies have been conducted on females, there is mounting recognition that estrogens play important roles in the

male brain, where they can be generated from circulating testosterone by local aromatase or synthesized de novo by neurons and glia [51]. Several sources of evidence confirm that estrogens serve as neurotrophic and neuroprotective agents. Notably, 17 β -estradiol (E₂) attenuates the toxicity of the amyloid- β peptide and glutamate in a hippocampal cell line [52]. In addition, estrogen therapy in post-menopausal women is associated with decreased incidence and enhanced recovery from ischemic stroke [51]. The protective effects of estrogens have been widely reported in different types of neuronal cells against a variety of insults, including H₂O₂ [53, 54], serum deprivation [55], oxygen-glucose deprivation [56], and iron [57]. Due to myriad and often tissue-specific estrogen effects, the precise molecular events that mediate these protective actions are not fully understood. Here, we evaluate the possibility that Ngb could be part of 17 β -estradiol-induced signals and effects in neuronal cells.

Materials and Methods

Reagents

E₂, testosterone (T), 5 α -androstan-17 β -ol-3-one (dihydrotestosterone, DHT), naringenin (Nar), insulin-like growth factor 1 (IGF-1), actinomycin D (Act), cycloheximide (Cxm), Pen-Strep solution, H₂O₂, RPMI-1640 media without phenol red, and charcoal-stripped fetal calf serum, the palmitoyl acyltransferase (PAT) inhibitor 2-bromohexadecanoic acid (2-Br-palmitate; 2-Br), the protease inhibitor cocktail, and the bovine serum albumin fraction V (BSA) were purchased from Sigma-Aldrich (St. Louis, Mo., USA).

Optimem, Hank's buffer salt solution (HBSS 1 \times), Neurobasal medium, B27 serum-free supplement, and GlutaMAX-I were purchased from Gibco-BRL (Gaithersburg, Md., USA). The p38 inhibitor SB 203 580 (SB), the AKT inhibitor, and the IGF-1 receptor (IGF-1R) inhibitor picropodophyllin (PPP) were obtained from Calbiochem (San Diego, Calif., USA). The E₂ antagonist fulvestrant (ICI 182,780, ICI), the estrogen receptor (ER) α -selective agonist 4,4',4''-(4-propyl-[¹H]-pyrazole-1,3,5-triyl)trisphenol (PPT), the ER β -selective agonist 2,3-bis(4-hydroxyphenyl)propionitrile (DPN), and the ER β -selective antagonist (R,R)-5,11-diethyl-5,6,11,12-tetrahydro-2,8-chrysenediol (THC) were obtained from Tocris (Ballwin, Mo., USA). Bradford protein assay was obtained from Bio-Rad Laboratories (Hercules, Calif., USA). The human recombinant ER α and ER β were obtained by PanVera (Madison, Wisc., USA). The anti-phospho-ERK1/2, anti-AKT, anti-ER α (MC20), anti-ER β (H150), anti-caspase-3, anti-poly(ADP-ribose)polymerase (PARP), and anti-ERK1/2 antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, Calif., USA). The polyclonal anti-phospho-AKT, anti-phospho-p38, and anti-p38 antibodies were purchased from New England Biolabs (Beverly, Mass., USA). The monoclonal anti-human Ngb (13C8) was purchased from Abcam (Cambridge, UK). The anti- β -tubulin was purchased from MP Biomedical (Solon, Ohio, USA). The chemiluminescence reagent for Western blot ECL was

obtained from GE Healthcare (Little Chalfont, UK). All the other products were from Sigma-Aldrich. Analytical or reagent grade products were used without further purification.

Cells

The human SK-N-BE neuroblastoma cell line was routinely grown in air containing 5% CO₂ in modified, phenol red-free, RPMI-1640 medium containing 10% (v/v) charcoal-stripped fetal calf serum, L-glutamine (2.0 mM), Pen-Strep solution (penicillin 100 U/ml, and streptomycin 100 mg/ml). Cells were passaged every 2 days. Cells were grown to approximately 70% confluence in 6-well plates before stimulation.

Hippocampal neurons were obtained from E18 mouse embryos after isolating the hippocampus in Ca²⁺- and Mg²⁺-free HBSS 1×. Mice were treated following the guidelines of the Council of Europe Convention ETS123, recently revised as indicated in the Directive 86/609/EEC. In addition, all protocols were approved by the Institutional Animal Care and Use Committee of CSIC-Cajal Institute (Madrid, Spain). Once 8–10 embryonic hippocampi were obtained, they were finely cut, washed twice in HBSS 1× buffer, and incubated in 0.1 mg/ml trypsin solution and 1 mg/ml DNase (Roche Diagnostics GmbH, Mannheim, Germany) for 15 min at 37°C. Trypsin and DNase were then eliminated by washing 3 times, with HBSS 1×, and the cut tissue was then triturated using a siliconized pipette. Cells were counted and plated in polylysine-coated (1 mg/ml) 6-well plates containing phenol red-free neurobasal medium supplemented with 2% (v/v) B27 serum-free supplement, 0.25% (v/v) GlutaMAX-I, and 1% (v/v) penicillin/streptomycin solution. Neurons were maintained under these conditions for 3 days at 5% CO₂ and 37°C.

Cells were simultaneously treated with vehicle (ethanol/PBS 1:10, v/v) and/or E₂ (0.1–1,000 nM), PPT (0.1–100 nM), DPN (0.1–100 nM), T (0.1–1,000 nM), DHT (0.1–1,000 nM), IGF-1 (100 ng/ml), and H₂O₂ (50 μM). When indicated, the anti-estrogen ICI (1 μM), the PAT inhibitor 2-Br (10 μM), the AKT inhibitor (1 μM), the p38 inhibitor SB (5 μM), the ERβ inhibitor THC (1 μM), the IGF-1R inhibitor PPP (100 nM), and the transcription inhibitor Act (1 μg/ml) were added 30 min before E₂ or IGF-1 administration. The translational inhibitor Cxm (10 μg/ml), was added 1 h before E₂ administration.

Cell Viability

SK-N-BE cell lines were grown to 70% confluence in 6-well plates and stimulated either with vehicle or E₂ (1 nM) or THC (1 μM). After 24 h of stimulation, cells were treated either with vehicle or with H₂O₂ 50 μM for 24 h. After treatment, cells were harvested with trypsin, and counted with Beckman Coulter Model ZM electronic particle (Palo Alto, Calif., USA).

Transfection of Short Interfering RNA

SK-N-BE cells, reaching 40–60% confluence, were transfected in a serum-free condition with either Stealth RNAi™ Ngb-silencing RNA or ERβ-silencing RNA (siRNA; Invitrogen) according to the manufacturer's instructions, using oligofectamine (Invitrogen) as the transfection reagent. The sequence used for Ngb oligonucleotides was 5'-CGUGAUUGAUGCUGCAGUGACC-AAU-3'; the sequence used for ERβ oligonucleotides was 5'-GAAGAACUCUUUGCCCGGAAAUUA-3'. The mismatch sequences used as a control were 5'-UGUGAUUUAUGGUGC-AGUAACCAAC-3' and 5'-GAAUCAUCCGUGCCAAGUAG-

AUUA-3' for Ngb and ERβ si-RNA, respectively. Briefly, oligofectamine and oligonucleotides (400 and 200 pM for Ngb siRNA and ERβ siRNA, respectively) were mixed with Optimem. The mixture was incubated for 20 min at room temperature, diluted with Optimem, and added to the cell medium for 4 h at 37°C. The medium was added to cells to reach the growing conditions (i.e., 10% (v/v) serum).

To evaluate the effective silencing of Ngb and ERβ, total proteins from cells transfected with MOCK (control), with scramble (mismatch sequence, data not shown), and with Ngb or ERβ oligonucleotides were extracted 48 h after transfection, and Ngb and ERβ expression was tested by Western blot analysis using anti-Ngb and anti-ERβ antibodies.

Western Blot Assays

After stimulation, the SK-N-BE cell line and hippocampal neurons were lysed and solubilized in 0.125 M Tris, pH 6.8, containing 10% (w/v) SDS and the protease inhibitor cocktail, then the cell lysates were boiled for 2 min. Total proteins were quantified using the Bradford protein assay. Solubilized proteins (20 μg) were resolved by 7 or 15% SDS-PAGE at 100 V for 1 h at 25°C and then electrophoretically transferred to nitrocellulose for 45 min at 100 V and 4°C. The nitrocellulose was treated with 3% (w/v) BSA in 138 mM NaCl, 25 mM Tris, pH 8.0, at 25°C for 1 h and then probed overnight at 4°C either with anti-Ngb (final dilution 1:1,000) or anti-ERα MC-20 (final dilution 1:500) or anti-ERβ H-150 (final dilution 1:3,000) or anti-caspase-3 (final dilution 1:1,000) or anti-PARP (final dilution 1:500) or anti-phospho-ERK1/2 (final dilution 1:200) or anti-phospho-AKT (final dilution 1:1,000) or anti-phospho-p38 (final dilution 1:1,000). The nitrocellulose was stripped by Restore Western Blot Stripping Buffer (Pierce Chemical, Rockford, Ill., USA) for 10 min at room temperature and then probed with anti-β-tubulin (final dilution 1:1,000) to normalize total lysate. Moreover, the nitrocellulose incubated with either anti-phospho-ERK1/2 or anti-phospho-AKT or anti-phospho-p38 was stripped and probed with anti-ERK1/2 (final dilution 1:200), anti-AKT (final dilution 1:100) and anti-p38 (final dilution 1:1,000), respectively. To evidence ERα and ERβ levels, electrophoresis was performed in the presence of 5 ng of recombinant ERα and ERβ. Antibody reaction was visualized with chemiluminescence Western blot detection reagent.

Densitometric analyses were performed by ImageJ software for Windows. The densitometry quantification of protein was normalized to tubulin.

Statistical Analysis


A statistical analysis was performed by using ANOVA followed by Tukey-Kramer post-test with the GraphPad InStat3 software system for Windows. In all cases, $p < 0.05$ was considered significant.

Results

E₂ Specifically Increases Ngb Levels in Neurons

Figure 1 shows that E₂ stimulation induced a time- and dose-dependent increase in Ngb levels in SK-N-BE cells (fig. 1A, B). The E₂ (10 nM) effect on Ngb levels started 30

Fig. 1. Effect of 17 β -estradiol (E₂) on Ngb protein levels in SK-N-BE human neuroblastoma cell line (**A, B**) and in mouse hippocampal primary neurons (**C, D**). **A, C** Time-course analysis of E₂ treatment (10 nM) on Ngb levels. **B, D** E₂ dose-dependent (0.1–1,000 nM) effect on Ngb levels (24 h of stimulation). The amount of protein was normalized by comparison with tubulin levels. The data are typical Western blots of five independent experiments (top panels); densitometric analysis related to E₂ dose- and time-dependent experiments (bottom panels). Data are means \pm SD of five different experiments. $p < 0.001$ was calculated with ANOVA followed by Tukey-Kramer post-test. **A, C** a significant vs. 0 h and b vs. 0.5 h; **B, D** a vs. 0, b vs. 0.1, c vs. 1, and d vs. 10 nM.



min after stimulation being significant 1 h after stimulation and remained constant 24 h after hormone stimulation. The E₂ dose-response curve was bell-shaped with a maximum effect at physiological E₂ concentrations (i.e., 1–10 nM; 24 h stimulation).

In contrast, the male sex steroid hormone DHT and the common precursor of E₂ and DHT, T, did not modify Ngb levels at any tested concentration (data not shown), suggesting the specificity of the E₂ effect. These data were confirmed in freshly isolated mouse hippocampal neurons (fig. 1C, D). Indeed, in these primary neurons, the E₂ effect was rapid (1 h), persistent (24 h) (fig. 1C), and specific in that neither DHT nor T were able to increase Ngb levels at any tested concentration (data not shown). In addition, even in mouse hippocampal neurons, 10 nM E₂ increased Ngb levels which remained significantly higher than control cells even at higher E₂ concentrations (fig. 1D). In line with the slight differences found in the E₂ concentration to obtain the maximum effect in both

cell types, 1 and 10 nM were used in the consecutive experiments to stimulate the SK-N-BE cell line and hippocampal neurons, respectively.

ER β was necessary for E₂-induced increase in Ngb levels. The pretreatment of SK-N-BE cells with the pure E₂ antagonist, ICI, completely prevented the E₂ effect on Ngb levels (fig. 2A), suggesting an ER-mediated mechanism. As SK-N-BE cells contain high ER β and low ER α levels (fig. 2B), cells were stimulated with either the specific ER α agonist PPT or the specific ER β agonist DPN to discriminate the role of each ER isoform in the E₂-induced Ngb level increase. Only 1 and 10 nM DPN mimicked the E₂ effect on Ngb levels (fig. 2C), whereas PPT was unable to increase Ngb levels, at any concentration investigated (fig. 2D). This result was confirmed by cell pretreatment with the specific ER β inhibitor THC, which completely prevented the E₂ effect (fig. 2C). Furthermore, the decrease of ER β protein level by ER β SiRNA transfection caused an impairment of the E₂ ability to increase




Fig. 2. Impact of estrogen receptors (ER) α and β on Ng β protein expression in SK-N-BE human neuroblastoma cell line. **A** Western blot analysis of Ng β levels in cells stimulated for 24 h with either vehicle or E₂ (1 nM) and/or the ER inhibitor ICI 182,870 (ICI; 1 μ M). **B** ER isoform levels in non-stimulated cells compared to recombinant proteins (5 ng). **C** Analysis of Ng β levels in cells stimulated for 24 h with either vehicle, E₂ (1 nM), the ER β agonist 2,3-bis(4-hydroxyphenyl)propionitrile (DPN; 1–100 nM) or the ER β -selective antagonist (R,R)-5,11-diethyl-5,6,11,12-tetrahydro-2,8-chrysenediol (THC; 1 μ M). **D** Analysis of Ng β levels in cells stimulated for 24 h with either vehicle, E₂ (1 nM) or the ER α agonist 4,4',4''-(4-propyl [1H]-pyrazole-1,3,5-triyl)trisphenol (PPT; 1–100 nM). **E** Analysis of Ng β and ER β levels in cells trans-

ected with either MOCK (control) or ER β small interference mRNA (si-ER β) in the absence or presence of E₂ (1 nM). **F** Analysis of Ng β levels in cells stimulated for 24 h with either vehicle, E₂ (1 nM) or naringenin (Nar; 0.01–10 μ M). The amount of proteins was normalized by comparison with tubulin levels. The data are typical Western blots of four independent experiments; densitometric analyses related to DPN, PPT, and Nar dose-dependent experiments (**C**, **D** and **F** bottom panels). Data are means \pm SD of four different experiments. $p < 0.001$ was calculated with ANOVA followed by Tukey-Kramer post-test. **C**, **D** a significant vs. vehicle, b vs. E₂, c vs. 1, and d vs. 10 nM. **F** a significant vs. vehicle, b vs. E₂, c vs. 0.01, and d vs. 0.1 μ M.




Fig. 3. Impact of estrogen receptors (ER) α and β on Ngb protein expression in mouse hippocampal primary neurons. **A** ER isoform levels in non-stimulated cells compared to recombinant proteins (5 ng). **B** Western blot analyses of Ngb levels in cells stimulated for 24 h with either vehicle or E_2 (10 nM) and/or the ER inhibitor ICI 182,870 (ICI; 1 μ M). **C** Analyses of Ngb levels in cells stimulated for 24 h with either vehicle, E_2 (10 nM) or the ER β agonist 2,3-bis(4-hydroxyphenyl)propionitrile (DPN; 0.1–100 nM). **D** Analyses of Ngb levels in cells stimulated for 24 h with either vehicle, E_2 (10 nM) or the ER α agonist 4,4',4''-(4-propyl [3 H]-pyrazole-1,3,5-tri-

yl)trisphenol (PPT; 0.1–100 nM). **E** Analyses of Ngb levels in cells stimulated for 24 h with either vehicle, E_2 (10 nM) and/or the ER β -selective antagonist (R,R)-5,11-diethyl-5,6,11,12-tetrahydro-2,8-chrysenediol (THC; 1 μ M). The amount of proteins was normalized by comparison with tubulin levels. Typical blots of five independent experiments are shown; densitometric analyses related to DPN and PPT dose-dependent experiments (**C**, **D** bottom panels). Data are means \pm SD of five different experiments. * $p < 0.001$ was calculated with ANOVA followed by Tukey-Kramer post-test with respect to vehicle-treated samples.

Ngb levels (fig. 2E). A further confirmation of the ER β involvement in the effect of E_2 effects derives from the results obtained using the flavonoid Nar (fig. 2F). Indeed, we previously reported that this flavonoid is a partial antagonist of E_2 in the presence of ER α [58] and an E_2 mimetic in the presence of ER β [59]. Figure 2F shows that, like E_2 , 0.1 μ M Nar was sufficient to increase Ngb levels. This effect persisted at Nar high concentrations (i.e., 1 and 10 μ M). ER β was also necessary for the E_2 -induced Ngb increase in mouse primary neurons (fig. 3). In these cells, containing a similar amount of ER α and ER β

(fig. 3A), ICI prevented the E_2 effect (fig. 3B), DPN mimicked the E_2 effect (fig. 3C), whereas PPT was unable to increase Ngb levels (fig. 3D). Cell pretreatment with the specific ER β inhibitor THC further confirmed these results (fig. 3E).

Mechanism Involved in the E_2 -Induced Increase of Ngb Levels in Neurons

ERs are ligand-activated transcription factors which possess both transcriptional and extranuclear activities [48]. Thus, we evaluated the impact on the E_2 -induced




Fig. 4. Mechanisms underlying E_2 effects on Ngb protein expression in SK-N-BE human neuroblastoma cells. **A** Western blot analyses of Ngb levels in cells stimulated for 24 h with either vehicle, E_2 (1 nM) and/or the transcription inhibitor actinomycin D (Act, 1 μ g/ml). **B** Analyses of Ngb levels in cells stimulated for 24 h with either vehicle, E_2 (1 nM) and/or the translation inhibitor cycloheximide (Cxm, 10 μ g/ml). **C** Analyses of Ngb levels in cells stimulated for 24 h with either vehicle, E_2 (1 nM) and/or the pal-

mitoylacyl transferase inhibitor 2-bromohexadecanoic acid (2-Br; 10 μ M). The amount of proteins was normalized by comparison with tubulin levels. Data are representative Western blots of independent experiments (**A–C** top panels); densitometric analyses (**A–C** bottom panels). Data are means \pm SD of four different experiments. $p < 0.001$ was calculated with ANOVA followed by Tukey-Kramer post-test with respect to (*) vehicle- or (o) E_2 -treated samples.

Ngb levels of the transcription inhibitor Act, the translation inhibitor Cxm, and the ER membrane localization inhibitor 2-Br [60]. Although the Ngb promoter sequence analysis (accession No. 12,581 from Transcriptional Regulatory Element Database <http://rulai.cshl.edu/cgibin/TRED/tred.cgi?process=home>) indicates that no canonical estrogen-responsive element (ERE) is present, SK-N-BE cell pretreatment for 24 h with Act completely prevented the E_2 effect on Ngb levels (fig. 4A). Similarly, Cxm (fig. 4B) and 2-Br (fig. 4C) impaired the increase of Ngb levels induced by E_2 , suggesting that both transcriptional and extranuclear mechanisms contribute to E_2 effects. This prompted us to evaluate which signal transduction cascade was activated by E_2 in neurons. After 15 min of stimulation with 1 nM E_2 , an increase of AKT and p38 phosphorylation in SK-N-BE cells was observed (fig. 5A, B). The E_2 -induced activation of these kinases was still present 1 h after E_2 stimulation (fig. 5A, B), but only the E_2 -induced p38 phosphorylation persisted 24 h after hormone stimulation (data not shown). By 30 min after 1 nM E_2 stimulation, the ERK1/2 phosphorylation status decreased (fig. 5A, B), in agreement with previous data obtained in cortical neurons [61]. The cell pretreatment with either AKT or p38 or ERK1/2 inhibitors suggested that neither AKT nor ERK1/2 are involved in the E_2 -induced increase of Ngb levels (data not shown),

whereas p38 activation was required for both rapid (i.e., 1 h) and long term (i.e., 24 h) E_2 effects on Ngb levels (fig. 5C, D). Similarly, only p38 inhibitor prevents an E_2 -induced Ngb level increase in hippocampal neurons (data not shown). Notably, the SK-N-BE cell transfection with ER β SiRNA reduced both ER β levels and the E_2 ability to induce p38 phosphorylation (fig. 5E). On the other hand, ER β SiRNA did not impair the E_2 -induced AKT activation (data not shown), suggesting that ER α could be the molecular mediator of AKT activation in these cells.

Since it has been reported that several actions of E_2 in the nervous system involve cross-talk between ER α and the IGF-1 receptor [62], we evaluated the possibility that the ER β -dependent E_2 -induced p38 phosphorylation and the Ngb-increased levels are dependent on the ER β -IGF-1 receptor cross-talk. IGF-1 is more efficient than E_2 to activate AKT phosphorylation and cell pretreatment with PPP, the IGF-1 receptor inhibitor, strongly prevented both IGF-1 and E_2 effects on AKT activation (fig. 5F), confirming that cross-talk between the IGF-1 receptor and ERs is important for AKT activation. However, IGF-1 was unable to increase p38 phosphorylation and PPP did not prevent E_2 -induced p38 activation (fig. 5F). In addition, IGF-1 did not modify Ngb levels in SK-N-BE cells (fig. 5G) further sustaining the high specificity of the E_2 effect.




Fig. 5. Rapid signal transduction pathways activated by E₂ and impact of E₂-dependent rapid signal inhibitors on Ngb protein levels in SK-N-BE human neuroblastoma cell line. **A** Time-course analyses of phosphorylated (P) and unphosphorylated AKT, p38, and ERK1/2 in cells stimulated for 0, 15, 30, and 60 min with E₂ (1 nM). **B** Densitometric analysis related to E₂-induced AKT, p38, and ERK1/2 phosphorylation experiments. Data are means ± SD of four different experiments. * p < 0.001 was calculated with ANOVA followed by Tukey-Kramer post-test: a significant vs. 0, b vs. 15, and c vs. 30 min. **C, D** Analyses of Ngb levels in cells stimulated for 1 h (**C**) or 24 h (**D**) with either vehicle, E₂ (1 nM) and/or the p38 inhibitor SB-203580 (SB; 5 μM). **E** Analysis of Ngb, ERβ, and phosphorylated (P) and unphosphorylated p38 levels in cells

transfected with either MOCK (control) or ERβ small interference mRNA (si-ERβ) in the absence or presence of E₂ (1 nM). The amount of proteins was normalized by comparison with tubulin levels. Data are representative Western blots of three independent experiments. **F** Western blot analyses of phosphorylated (P) and unphosphorylated AKT and p38 levels in cells treated for 24 h with either vehicle, E₂ (1 nM) or IGF-1 (100 ng/ml). When indicated, cells were pretreated with the IGF-1 receptor inhibitor picropodophyllin (PPP; 100 μM). **G** Analyses of Ngb protein levels in cells treated for 24 h with either vehicle, E₂ (1 nM) or IGF-1 (100 ng/ml). The amount of proteins was normalized by comparison with tubulin levels. Data are representative Western blots of three independent experiments.




Fig. 6. E₂ effect on SK-N-BE human neuroblastoma cell line viability. **A** Cells were grown in the presence of either vehicle or E₂ (1 nM) and counted at the indicated time. Data are means ± SD of five independent experiments carried out in duplicate. **B** Cells were grown for 24 h in the presence of either vehicle, E₂ (1 nM) or E₂ in the presence of the ERβ-selective antagonist (R,R)-5,11-diethyl-5,6,11,12-tetrahydro-2,8-chrysenediol (THC; 1 μM). After 24 h, cells were stimulated with H₂O₂ 50 μM (24 h of stimulation), and counted. Data are means ± SD of five independent experiments carried out in duplicate. *p* < 0.001 was calculated with ANOVA followed by Tukey-Kramer post-test: a significant vs. Vehicle - H₂O₂, b vs. vehicle + H₂O₂, c vs. E₂ - H₂O₂, d vs. E₂ + H₂O₂, and e vs. THC + E₂ - H₂O₂. **C** Western blot analyses of caspase-3 activation and poly(ADP-ribose) polymerase (PARP) cleavage were performed on cells stimulated with either the vehicle or pretreated with E₂ (1 nM) for 24 h in the presence or absence of THC pretreatment, and then treated with H₂O₂ 50 μM (24 h of stimulation). Staurosporin (2 μM for 24 h) was used as positive control of caspase activation. The amount of proteins was normalized by comparison with tubulin level. Data are representative Western

blots of three independent experiments. **D** Western blot analysis of Ngb protein levels in cells transfected with either MOCK (control) or with Ngb small interference mRNA (si-Ngb) in the absence or presence of E₂ (1 nM). The amount of proteins was normalized by comparison with tubulin levels. Data are representative Western blots of three independent experiments. Cells transfected with either MOCK (empty bars) or si-Ngb (filled bars) were grown for 24 h in the presence of either the vehicle or E₂ (1 nM), stimulated with H₂O₂ 50 μM (24 h of stimulation), and counted. Data are means ± SD of three independent experiments carried out in duplicate. *p* < 0.001 was calculated with ANOVA followed by Tukey-Kramer post-test: a significant vs. vehicle - H₂O₂, b vs. vehicle + H₂O₂, c vs. E₂ + H₂O₂, d vs. vehicle - H₂O₂ si-Ngb, and e vs. E₂ - H₂O₂ si-Ngb. **E** Western blot analyses of caspase-3 activation and poly(ADP-ribose) polymerase (PARP) cleavage were performed on cells transfected with either MOCK or si-Ngb and stimulated with either vehicle or 50 μM H₂O₂ in the presence or absence of E₂ (1 nM) (24 h pretreatment). The amount of proteins was normalized by comparison with tubulin level. Data are representative Western blots of three independent experiments.

Ngb Involvement in E₂-Induced Protection against H₂O₂-Induced Oxidative Stress

Finally we evaluated the role played by increased levels of Ngb on E₂ effects in SK-N-BE cells. E₂ stimulation did not modify SK-N-BE cell number (fig. 6A) but, as expected [61], reduced by 50% the H₂O₂-induced decrease in cell number (fig. 6B) as well as the increase of the 17-kDa active caspase-3 subunit and the cleavage of the caspase-3 substrate PARP (fig. 6C). Staurosporin (2 μM for 24 h) was used as positive control of caspase activation (fig. 6C). Notably, the E₂-protective effect against H₂O₂-induced neuron toxicity seems to require ERβ, since the cell pretreatment with the specific ERβ inhibitor THC completely prevented E₂ effects (fig. 6B, C). In line with this result, E₂ was unable to counteract the H₂O₂-induced decrease in cell number (fig. 6D) and the activation of the pro-apoptotic cascade (i.e., caspase-3 activation and PARP cleavage) (fig. 6E) in Ngb-silenced SK-N-BE cells.

Discussion

The aim of this paper was to identify a possible endogenous modulator of Ngb. Thus, we investigated the effect of E₂, a well-known neurotrophic and neuroprotective hormone [52–57, 63–66], on Ngb expression. Our results indicated that E₂ increases Ngb levels of about 300% in both the human neuroblastoma cell line and mouse primary hippocampal neurons. Although it has been obtained with a qualitative technique (i.e., Western blot), this effect is conspicuous in that the well-known E₂ effect on cyclin D1 expression, playing a relevant role in E₂-induced cell proliferation, is only of about 50–70% [67]. In addition, the E₂-induced Ngb increase is rapid (1 h), persistent (24 h), and specific, being not mimicked by either the male sex steroid hormone DHT or by the common precursor T or by IGF-1, another well-known neuroprotective hormone. These results represent the first evidence for steroid hormone modulation of globin levels in cells. Recently, it has been reported that hemoglobin is specifically expressed in neurons, its expression being upregulated by erythropoietin and accompanied by enhanced brain oxygenation under physiologic and hypoxic conditions [68]. At the present, the relationship between hemoglobin and Ngb in neurons is still unclear. Although hemoglobin α-chains and Ngb are expressed in the same nerve cells, Ngb levels are not increased by erythropoietin [68]. It is therefore unlikely that they have a tightly linked function, e.g. in facilitated oxygen trans-

port; however, hemoglobin and Ngb could fulfill independent tasks in neurons.

It is now well known that sex steroid hormones have numerous effects on the brain throughout the lifespan, beginning during gestation and continuing on into senescence [69]. However, the inability of Ngb to react with androgens renders Ngb a new E₂ target that should be added to the variety of E₂-specific actions on the brain which include mood, locomotor activity, pain sensitivity, vulnerability to epilepsy, attentional mechanisms, and cognition [64].

The E₂ effect on Ngb levels is rapid and dose-dependent with the maximum effect at E₂ physiological concentration (i.e., 1–10 nM). Notably, the E₂ dose-response curve results in being bell-shaped. This is typical for E₂ [70], whose effects are mediated by two receptor isoforms. Accordingly, the plant-derived flavonoid Nar, which partially blocks the rapid activities of ERα [58], increased Ngb levels with a plateau at 1 μM concentration. This result suggests a functional antagonism between the activities of ERα and ERβ in neurons, as has been reported in other cell types [71].

Although human neuroblastoma cell line and mouse primary hippocampal neurons express different levels of both ER isoforms, the effect of E₂ on Ngb levels specifically requires all ERβ activities. In fact, ERβ extranuclear and genomic signals cross-talk each other to guarantee both the rapid (1 h) and the persistent (24 h) E₂ effects. In particular, the rapid (15 min) and persistent (24 h) ERβ-mediated p38 activation is required for E₂-induced Ngb increase. The E₂-dependent activation of p38, a mitogen-activated protein kinase (MAPK) family component, seems to represent a conserved pathway in ERβ-based E₂ rapid signals. Indeed, the E₂-induced ERβ-mediated activation of the p38/MAPK occurs in ERβ-transfected HeLa cells and in ERβ-containing rat myoblasts and colon adenocarcinoma cells [70, 72, 73]. This signaling pathway transduces different E₂ effects depending on the cell context. In fact, p38 activation is required for E₂-induced apoptosis of cancer cells [72, 73], for E₂-induced gene transcription [73], and for E₂-induced protection against oxidative stress in neuroblastoma cell line (present results) and in rat myoblasts [Marino, unpubl. results]. Thus, although Ngb promoter does not contain any canonical ERE, it is not surprising that the transcription inhibitor Act completely prevents the increase of the E₂-induced Ngb levels. The sequence analysis (accession No. 12,581 from Transcriptional Regulatory Element Database <http://rulai.cshl.edu/cgi-bin/TRED/tred.cgi?process=home>) indicates that several non-canonical half

ERE sites are present in the Ngf promoter along with a responsive element for other transcription factors. These results suggest that the E₂-induced Ngf transcription could be mediated by tethered interactions of ER with other transcription factors to activate gene expression (i.e., indirect genomic mechanism) [48]. Thus, the integration between extranuclear and genomic events may be required to provide plasticity for the neuronal response to E₂. The hormone rapidly induces AKT activation and ERK dephosphorylation in neuroblastoma cells. These effects are still detected in ER α -containing but ER β -silenced cells. Although these kinases are not directly involved in an E₂-induced Ngf increase, they could contribute to the E₂ effects in neurons, since AKT activation has been associated with the increase of the anti-apoptotic protein Bcl-2 and to the E₂-induced cell survival [72], while E₂ protected cortical neurons against oxidative stress by reducing H₂O₂-induced activation of ERK1/2 [61].

The main result reported here is that Ngf is part of the E₂ response to H₂O₂-induced toxicity. In fact, exposure to 50 μ M H₂O₂ induces neuroblastoma cell death (about 50%) which is accompanied by a dramatic increase in caspase-3 activation. The cell pretreatment with E₂ (1 nM) decreases cell death and reduces caspase-3 activation triggered by exposure to H₂O₂ in good accordance with the literature [52–54, 61, 74]. This E₂ effect against H₂O₂ toxicity is completely prevented by treatment with THC, ER β inhibitor, and by knocking out Ngf using small interfering RNA.

Exposure to H₂O₂ induces a robust increase of reactive oxygen species in cells (followed by oxidation of lipids, proteins, and DNA), intracellular calcium increase, glutathione depletion, mitochondria dysfunction, and caspase-3 activation followed by apoptotic cell death [74]. It has been demonstrated that E₂ exerts protective effects on several of these cellular events including potent attenuation of lipid peroxidation, attenuated ATP depletion, al-

leviated intracellular calcium elevation, ablated mitochondrial calcium loading (with the subsequent mitochondrial membrane potential maintenance), reduced caspase-3 activation, and enhanced cell survival ([74] and present results). At present, it is difficult to discriminate the role played by Ngf in each of these E₂-induced cellular outcomes because it is not yet clear if they require ER α or ER β or both receptors. Moreover, most of the E₂-protective effects have been obtained at pharmacological E₂ concentrations (0.1–10 μ M) [74]. However, it has recently shown that the elevation of human Ngf expression in neurons prior to insult with H₂O₂ enhances cell viability and results in a significant decrease in oxidative stress and an increased intracellular ATP concentration [75]. In addition, a linkage of Ngf to oxidative metabolism has been proposed [76]. These data are strongly suggestive of the involvement of Ngf in E₂-induced attenuated ATP depletion. Furthermore, in the presence of Ngf the initially released cytochrome *c* may be sequestered by the Ngf [42, 43] and the apoptotic cascade avoided [19].

In the future, different strategies (e.g., E₂-induced Ngf localization, association between Ngf and ERs, and E₂-induced ER-mediated Ngf promoter activity) will substantiate Ngf action in neurons, our laboratories being active in this field. However, from now, Ngf can be regarded as part of signals activated by E₂ to exert neuroprotective effects definitely validating the role played by Ngf as an anti-apoptotic neuroprotective globin.

Acknowledgements

The generous gift of SK-N-BE human neuroblastoma cell line from Dr. Roberto Rivabene (Institute of Public Health, ISS, Rome, Italy) and of staurosporin from Dr. Antonio Antocchia (University Roma Tre) are gratefully acknowledged. This work was supported by grants from University Roma Tre (CLAR 2009 to P.A.) and the Italian Ministry of Health (Strategico 2008 to M.M).

References

- Burmester T, Weich B, Reinhardt S, Hankeln T: A vertebrate globin expressed in brain. *Nature* 2000;407:520–523.
- Moens L, Dewilde S: Globins in the brain. *Nature* 2000;407:461–462.
- Trent JT 3rd, Watts RA, Hargrove MS: Human neuroglobin, a hexacoordinate hemoglobin that reversibly binds oxygen. *J Biol Chem* 2001;276:30106–30110.
- Pesce A, Bolognesi M, Bocedi A, Ascenzi P, Dewilde S, Moens L, Hankeln T, Burmester T: Neuroglobin and cytoglobin. Fresh blood for the vertebrate globin family. *EMBO Rep* 2002;3:1146–1151.
- Reuss S, Saaler-Reinhardt S, Weich B, Wystub S, Reuss MH, Burmester T, Hankeln T: Expression analysis of neuroglobin mRNA in rodent tissues. *Neurosci* 2002;115:645–656.
- Pesce A, Dewilde S, Nardini M, Moens L, Ascenzi P, Hankeln T, Burmester T, Bolognesi M: Human brain neuroglobin structure reveals a distinct mode of controlling oxygen affinity. *Structure* 2003;11:1087–1095.
- Wystub S, Laufs T, Schmidt M, Burmester T, Maas U, Saaler-Reinhardt S, Hankeln T, Reuss S: Localization of neuroglobin protein in the mouse brain. *Neurosci Lett* 2003;346:114–116.

- 8 Burmester T, Hankeln T: Neuroglobin: a respiratory protein of the nervous system. *News Physiol Sci* 2004;19:110–113.
- 9 Hankeln T, Wystub S, Laufs T, Schmidt M, Gerlach F, Saaler-Reinhardt S, Reuss S, Burmester T: The cellular and subcellular localization of neuroglobin and cytoglobin – a clue to their function? *IUBMB Life* 2004;56:671–679.
- 10 Vallone B, Nienhaus K, Matthes A, Brunori M, Nienhaus GU: The structure of carbon-monoxo neuroglobin reveals a heme-sliding mechanism for control of ligand affinity. *Proc Natl Acad Sci USA* 2004;101:17351–17356.
- 11 Hankeln T, Ebner B, Fuchs C, Gerlach F, Haberkamp M, Laufs TL, Roesner A, Schmidt M, Weich B, Wystub S, Saaler-Reinhardt S, Reuss S, Bolognesi M, De Sanctis D, Marden MC, Kiger L, Moens L, Dewilde S, Nevo E, Avivi A, Weber RE, Fago A, Burmester T: Neuroglobin and cytoglobin in search of their role in the vertebrate globin family. *J Inorg Biochem* 2005;99:110–119.
- 12 Brunori M, Vallone B: A globin for the brain. *FASEB J* 2006;20:2192–2197.
- 13 Nienhaus K, Nienhaus GU: Searching for neuroglobin's role in the brain. *IUBMB Life* 2007;59:490–497.
- 14 Greenberg DA, Jin K, Khan AA: Neuroglobin: an endogenous neuroprotectant. *Curr Opin Pharmacol* 2008;8:20–24.
- 15 Burmester T, Hankeln T: What is the function of neuroglobin? *J Exp Biol* 2009;212:1423–1428.
- 16 Yu Z, Fan X, Lo EH, Wang X: Neuroprotective roles and mechanisms of neuroglobin. *Neurol Res* 2009;31:122–127.
- 17 Emara M, Turner AR, Allalunis-Turner J: Hypoxic regulation of cytoglobin and neuroglobin expression in human normal and tumor tissues. *Cancer Cell Int* 2010;10:33–49.
- 18 Hundahl CA, Allen GC, Hannibal J, Kjaer K, Rehfeld JF, Dewilde S, Nyengaard JR, Kelsen J, Hay-Schmidt A: Anatomical characterization of cytoglobin and neuroglobin mRNA and protein expression in the mouse brain. *Brain Res* 2010;1331:58–73.
- 19 Fago A, Mathews AJ, Brittain T: A role for neuroglobin: resetting the trigger level for apoptosis in neuronal and retinal cells. *IUBMB Life* 2008;60:398–401.
- 20 Schmidt M, Giessl A, Laufs T, Hankeln T, Wolfrum U, Burmester T: How does the eye breathe? Evidence for neuroglobin-mediated oxygen supply in the mammalian retina. *J Biol Chem* 2003;278:1932–1935.
- 21 Bentmann A, Schmidt M, Reuss S, Wolfrum U, Hankeln T, Burmester T: Divergent distribution in vascular and avascular mammalian retinae links neuroglobin to cellular respiration. *J Biol Chem* 2005;280:20660–20665.
- 22 Dewilde S, Kiger L, Burmester T, Hankeln T, Baudin-Creuz V, Aerts T, Marden MC, Caubergs R, Moens L: Biochemical characterization and ligand binding properties of neuroglobin, a novel member of the globin family. *J Biol Chem* 2001;276:38949–38955.
- 23 Hamdane D, Kiger L, Dewilde S, Green BN, Pesce A, Uzan J, Burmester T, Hankeln T, Bolognesi M, Moens L, Marden MC: The redox state of the cell regulates the ligand binding affinity of human neuroglobin and cytoglobin. *J Biol Chem* 2003;278:51713–51721.
- 24 Fago A, Hundahl C, Malte H, Weber RE: Functional properties of neuroglobin and cytoglobin: insights into the ancestral physiological roles of globins. *IUBMB Life* 2004;56:689–696.
- 25 Herold S, Fago A, Weber RE, Dewilde S, Moens L: Reactivity studies of the Fe(III) and Fe(II)NO forms of human neuroglobin reveal a potential role against oxidative stress. *J Biol Chem* 2004;279:22841–22847.
- 26 Kiger L, Uzan J, Dewilde S, Burmester T, Hankeln T, Moens L, Hamdane D, Baudin-Creuz V, Marden M: Neuroglobin ligand binding kinetics. *IUBMB Life* 2004;56:709–719.
- 27 Trandafir F, van Doorslaer S, Dewilde S, Moens L: Temperature dependence of NO binding modes in human neuroglobin. *Biochim Biophys Acta* 2004;1702:153–161.
- 28 Brunori M, Giuffrè A, Nienhaus K, Nienhaus GU, Scandurra FM, Vallone B: Neuroglobin, nitric oxide, and oxygen: functional pathways and conformational changes. *Proc Natl Acad Sci USA* 2005;102:8483–8488.
- 29 Fago A, Mathews AJ, Dewilde S, Moens L, Brittain T: The reactions of neuroglobin with CO: evidence for two forms of the ferrous protein. *J Inorg Biochem* 2006;100:1339–1343.
- 30 Nicolis S, Monzani E, Ciaccio C, Ascenzi P, Moens L, Casella L: Reactivity and endogenous modification by nitrite and hydrogen peroxide: does human neuroglobin act only as a scavenger? *Biochem J* 2007;407:89–99.
- 31 Brittain T, Yosaatmadja Y, Henty K: The interaction of human neuroglobin with hydrogen sulphide. *IUBMB Life* 2008;60:135–138.
- 32 Giuffrè A, Moschetti T, Vallone B, Brunori M: Neuroglobin: enzymatic reduction and oxygen affinity. *Biochem Biophys Res Commun* 2008;367:893–898.
- 33 Kakar S, Hoffman FG, Storz JF, Fabian M, Hargrove MS: Structure and reactivity of hexacoordinate hemoglobins. *Biophys Chem* 2010;152:1–14.
- 34 Ostojic J, Sakaguchi DS, de Lathouder Y, Hargrove MS, Trent JT 3rd, Kwon YH, Kardouk RH, Kuehn MH, Betts DM, Grozdanic S: Neuroglobin and cytoglobin: oxygen-binding proteins in retinal neurons. *Invest Ophthalmol Vis Sci* 2006;47:1016–1023.
- 35 Herold S, Fago A: Reactions of peroxynitrite with globin proteins and their possible physiological role. *Comp Biochem Physiol A Mol Integr Physiol* 2005;142:124–129.
- 36 Petersen MG, Dewilde S, Fago A: Reactions of ferrous neuroglobin and cytoglobin with nitrite under anaerobic conditions. *J Inorg Biochem* 2008;102:1777–1782.
- 37 Jin K, Mao XO, Xie L, Khan AA, Greenberg DA: Neuroglobin protects against nitric oxide toxicity. *Neurosci Lett* 2008;430:135–137.
- 38 Fordel E, Thijs L, Moens L, Dewilde S: Neuroglobin and cytoglobin expression in mice. Evidence for a correlation with reactive oxygen species scavenging. *FEBS J* 2007;274:1312–1317.
- 39 Khan AA, Mao XO, Banwait S, DerMardrossian CM, Bokoch GM, Jin K, Greenberg DA: Regulation of hypoxic neuronal death signaling by neuroglobin. *FASEB J* 2008;22:1737–1747.
- 40 Watanabe S, Wakasugi K: Neuroprotective function of human neuroglobin is correlated with its guanine nucleotide dissociation inhibitor activity. *Biochem Biophys Res Commun* 2008;369:695–700.
- 41 Fago A, Mathews AJ, Moens L, Dewilde S, Brittain T: The reaction of neuroglobin with potential redox protein partners cytochrome *b₅* and cytochrome *c*. *FEBS Lett* 2006;580:4884–4888.
- 42 Brittain T, Skommer J, Raychaudhuri S, Birch N: An antiapoptotic neuroprotective role for neuroglobin. *Int J Mol Sci* 2010;11:2306–2321.
- 43 Raychaudhuri S, Skommer J, Henty K, Birch N, Brittain T: Neuroglobin protects nerve cells from apoptosis by inhibiting the intrinsic pathway of cell death. *Apoptosis* 2010;15:401–411.
- 44 Pesce A, De Sanctis D, Nardini M, Dewilde S, Moens L, Hankeln T, Burmester T, Ascenzi P, Bolognesi M: Reversible hexa- to penta-coordination of the heme Fe atom modulates ligand binding properties of neuroglobin and cytoglobin. *IUBMB Life* 2004;56:657–664.
- 45 Sun Y, Jin K, Peel A, Mao XO, Xie L, Greenberg DA: Neuroglobin protects the brain from experimental stroke in vivo. *Proc Natl Acad Sci USA* 2003;100:3497–3500.
- 46 Khan AA, Wang Y, Sun Y, Mao XO, Xie L, Miles E, Graboski J, Chen S, Ellerby LM, Jin K, Greenberg DA: Neuroglobin-overexpressing transgenic mice are resistant to cerebral and myocardial ischemia. *Proc Natl Acad Sci USA* 2006;103:17944–17948.
- 47 Jin K, Mao Y, Mao X, Xie L, Greenberg DA: Neuroglobin expression in ischemic stroke. *Stroke* 2010;41:557–559.
- 48 Ascenzi P, Bocedi A, Marino M: Structure-function relationship of estrogen receptor α and β : impact on human health. *Mol Aspects Med* 2006;27:299–402.
- 49 Manthey D, Behl C: From structural biochemistry to expression profiling: neuroprotective activities of estrogen. *Neuroscience* 2006;138:845–850.

- 50 Correia SC, Santos RX, Cardoso S, Carvalho C, Santos MS, Oliveira CR, Moreira PI: Effects of estrogen in the brain: is it a neuroprotective agent in Alzheimer's disease? *Curr Aging Sci* 2010;3:113-126.
- 51 Gillies GE, McArthur S: Estrogen actions in the brain and the basis for differential action in men and women: a case for sex-specific medicines. *Pharmacol Rev* 2010;62:155-198.
- 52 Behl C, Skutella T, Lezoualc'h F, Post A, Widmann M, Newton CJ, Holsboer F: Neuroprotection against oxidative stress by estrogens: structure-activity relationship. *Mol Pharmacol* 1997;51:535-541.
- 53 Sawada H, Ibi M, Kihara T, Urushitani M, Akaike A, Shimohama S: Estradiol protects mesencephalic dopaminergic neurons from oxidative stress-induced neuronal death. *J Neurosci Res* 1998;54:707-719.
- 54 Green PS, Simpkins JW: Neuroprotective effects of estrogens: potential mechanisms of action. *Int J Dev Neurosci* 2000;18:347-358.
- 55 Bae YH, Hwang JY, Kim YH, Koh JY: Antioxidative neuroprotection by estrogens in mouse cortical cultures. *J Korean Med Sci* 2000;15:327-336.
- 56 Wilson ME, Dubal DB, Wise PM: Estradiol protects against injury-induced cell death in cortical explant cultures: a role for estrogen receptors. *Brain Res* 2000;873:235-242.
- 57 Blum-Degen D, Haas M, Pohli S, Harth R, Romer W, Oettel M, Riederer P, Gotz ME: Scavestrogens protect IMR 32 cells from oxidative stress-induced cell death. *Toxicol Appl Pharmacol* 1998;152:49-55.
- 58 Galluzzo P, Ascenzi P, Bulzomi P, Marino M: The nutritional flavanone naringenin triggers antiestrogenic effects by regulating estrogen receptor α -palmitoylation. *Endocrinology* 2008;149:2567-2575.
- 59 Totta P, Acconcia F, Leone S, Cardillo I, Marino M: Mechanisms of naringenin-induced apoptotic cascade in cancer cells: involvement of estrogen receptor α and β signalling. *IUBMB Life* 2004;56:491-499.
- 60 Acconcia F, Ascenzi P, Bocedi A, Spisni E, Tomasi V, Trentalance A, Visca P, Marino M: Palmitoylation-dependent estrogen receptor α membrane localization: regulation by 17 β -estradiol. *Mol Biol Cell* 2005;16:231-237.
- 61 Numakawa Y, Matsumoto T, Yokomaku D, Taguchi T, Niki E, Hatanaka H, Kunugi H, Numakawa T: 17 β -Estradiol protects cortical neurons against oxidative stress-induced cell death through reduction in the activity of mitogen-activated protein kinase and in the accumulation of intracellular calcium. *Endocrinology* 2007;148:627-637.
- 62 Marin R, Diaz M, Alonso R, Sanz A, Arevalo MA, Garcia-Segura LM: Role of estrogen receptor α in membrane-initiated signaling in neural cells: interaction with IGF-1 receptor. *J Steroid Biochem Mol Biol* 2009;114:2-7.
- 63 Regan RF, Guo Y: Estrogens attenuate neuronal injury due to hemoglobin, chemical hypoxia and excitatory amino acids in murine cortical cultures. *Brain Res* 1997;764:133-140.
- 64 McEwen BS: Sex, stress and the hippocampus: allostasis, allostatic load and the aging process. *Neurobiol Aging* 2002;23:921-939.
- 65 Marin R, Ramirez C, Morales A, Gonzalez M, Alonso R, Diaz M: Modulation of A β -induced neurotoxicity by estrogen receptor α and other associated proteins in lipid rafts. *Steroids* 2008;73:992-996.
- 66 Vasudevan N, Pfaff DW: Non-genomic actions of estrogens and their interaction with genomic actions in the brain. *Front Neuroendocrinol* 2008;29:238-257.
- 67 Marino M, Acconcia F, Bresciani F, Weisz A, Trentalance A: Distinct nongenomic signal transduction pathways controlled by 17 β -estradiol regulate DNA synthesis and cyclin D₁ gene transcription in HepG2 cells. *Mol Biol Cell* 2002;13:3720-3729.
- 68 Schelshorn DW, Schneider A, Kuschinsky W, Weber D, Kruger C, Dittgen T, Bourgers HF, Sabouri F, Gassler N, Bach A, Maurer MH: Expression of hemoglobin in rodent neurons. *J Cereb Blood Flow Metab* 2009;29:585-595.
- 69 Swaab DF: Sexual differentiation of the brain and behavior. *Best Pract Res Clin Endocrinol Metab* 2007;21:431-444.
- 70 Galluzzo P, Rastelli C, Bulzomi P, Acconcia F, Pallottini V, Marino M: 17 β -Estradiol regulates the first steps of skeletal muscle cell differentiation via ER α -mediated signals. *Am J Physiol Cell Physiol* 2009;297:C1249-C1262.
- 71 Matthews J, Gustafsson JA: Estrogen signaling: a subtle balance between ER α and ER β . *Mol Intervent* 2003;3:281-292.
- 72 Acconcia F, Totta P, Ogawa S, Cardillo I, Inoue S, Leone S, Trentalance A, Muramatsu M, Marino M: Survival versus apoptotic 17 β -estradiol effect: role of ER α - and ER β -activated nongenomic signaling. *J Cell Physiol* 2005;203:193-201.
- 73 Galluzzo P, Caiazza F, Moreno S, Marino M: Role of ER β palmitoylation in the inhibition of human colon cancer cell proliferation. *Endocr Relat Cancer* 2007;14:153-167.
- 74 Wang X, Dykens JA, Perez E, Liu R, Yang S, Covey DF, Simpkins JW: Neuroprotective effects of 17 β -estradiol and nonfeminizing estrogens against H₂O₂ toxicity in human neuroblastoma SK-N-SH cells. *Mol Pharmacol* 2006;70:395-404.
- 75 Antao ST, Duong TT, Aran R, Witting PK: Neuroglobin overexpression in cultured human neuronal cells protects against hydrogen peroxide insult via activating phosphoinositide-3 kinase and opening the mitochondrial K(ATP) channel. *Antioxid Redox Signal* 2010;13:769-781.
- 76 Mitz SA, Reuss S, Folkow LP, Blix AS, Ramirez JM, Hankeln T, Burmester T: When the brain goes diving: glial oxidative metabolism may confer hypoxia tolerance to the seal brain. *Neurosci* 2009;163:552-560.