

# MICROCLIMATIC AND PHYSIOLOGICAL CHANGES UNDER A CENTER PIVOT SYSTEM IRRIGATING MAIZE

Yenny Urrego-Pereira<sup>a\*</sup>; José Cavero<sup>a</sup>, Eva T. Medina<sup>b</sup> and Antonio Martínez-Cob<sup>a</sup>

<sup>a</sup> *Departamento de Suelo y Agua, Estación Experimental de Aula Dei (EEAD), Consejo Superior de Investigaciones Científicas (CSIC), Avda. Montañana 1005, 50059 Zaragoza, Spain.*

<sup>b</sup> *Unidad de Suelos y Riegos (Unidad Asociada EEAD-CSIC), Centro de Investigación y Tecnología Agroalimentaria (CITA-DGA), Avda. Montañana 930, 50059 Zaragoza, Spain.*

## ABSTRACT

The microclimatic (air temperature and vapour pressure deficit (VPD)) and physiological (canopy temperature and plant transpiration) changes due to center pivot sprinkle irrigation were monitored at a commercial plot of maize (*Zea mays* L.). Two treatments were considered: a) *moist*, measurements taken at three spots on a transect when the pivot was running over it; b) *dry*, measurements taken simultaneously at a fourth spot D, 270 m apart. A total of 34 irrigation events were monitored, seven of which included plant transpiration measurements. For the *transpiration-measured* irrigation events, significant ( $P = 0.05$ ) reductions in the monitored variables for the *moist* treatment were observed for 0.6 to 2.1 h *before*, *during* and 0.5 to 2.4 h *after* the irrigation. The average decreases for the phase *during* were 1.8 to 2.1 °C for air temperature, 0.53 to

---

\* Corresponding author: Tel.: +34-976716075; Fax: +34-976716145; E-mail: [yurrego@eead.csic.es](mailto:yurrego@eead.csic.es) (Yenny Urrego-Pereira).

1 0.61 kPa for VPD, 3.1 to 3.8 °C for canopy temperature, and 0.22 to 0.28 mm h<sup>-1</sup>  
2 (30 to 36 %) for transpiration. Lower reductions were found for the phases *before*  
3 *and after*. The duration of the microclimatic changes decreased as the distance  
4 from the centre of the pivot increased (from 3.9 to 2.2 h), but the duration of the  
5 physiological changes was similar in the different pivot arm portions (≈4 h).  
6 Microclimatic and physiological changes were higher in drier and warmer days.  
7 Transpiration reduction due to irrigation was higher as closer to the center of the  
8 pivot and represented 5 to 7% of the applied water. The estimated reduction of  
9 ET represented 1.5 to 1.8% of the applied water. The reduction of transpiration  
10 and ET is positive because it represents a reduction in irrigation requirements.

11 The decrease in maize canopy temperature could be positive or negative  
12 depending on its effect on photosynthesis.

### 13 **Abbreviations**

14 VPD, vapor pressure deficit; PAP, pivot arm portion; GMT, Greenwich Mean Time;  
15 SIAR, Spanish Irrigation Advisory System; CV, coefficient of variation.

### 16 **KEYWORDS**

17 Air temperature; Vapor pressure deficit; Canopy temperature; Transpiration; Center  
18 pivot sprinkler irrigation

19

## 1 1. INTRODUCTION

2 The acreage irrigated by sprinkler irrigation systems has increased in order to better  
3 meet the crop water requirements and increase the water application efficiency and the  
4 crop yield. For instance, sprinkler irrigation systems represent about 23% of the 3.5 Mha  
5 of irrigated land in Spain. Among the different types of sprinkler irrigation systems, the  
6 center pivot offers several advantages such as the lower invest cost compared to solid-  
7 set, the high degree of automation and the high water application efficiency. Due to  
8 these factors, their use has become widespread around the world (Allen et al., 2000).  
9 Thus, center pivot is used in about 32 to 40 % of the irrigated land in several Spanish  
10 Irrigation Districts (MAGRAMA 2011). In USA, the land irrigated by center pivot has  
11 increased by more than 50% from 1986 to 1996 (Evans 2001), while it accounted for  
12 83% of the sprinkler systems on 2008, i.e. about 47% of the 22.2 Mha of irrigated land  
13 (USDA 2008).

14 Thompson et al. (1993) reported that during solid-set sprinkler irrigation a total amount  
15 of energy equivalent to 24% of the net radiation was transferred from the plant  
16 environment to the water droplets as they warmed during flight and after they impacted  
17 the canopy and soil. This leads to sprinkler irrigation water losses by evaporation during  
18 and after the irrigation. This evaporation of water modifies the crop microclimate. A  
19 decrease of air temperature and air vapor pressure deficit (VPD) has been reported  
20 (Robinson 1970; Steiner et al. 1983; Thompson et al. 1993; Tolk et al. 1995; Liu and  
21 Kang 2006a; Caverro et al. 2009). The microclimatic changes can also cause several  
22 crop responses. Howell et al. (1971) reported that during mist irrigation of peas (*Vigna*  
23 *unguiculata* (L.) Walp. ), the air and canopy temperature decreased and the leaf water

1 potential increased; in addition, a higher yield was observed as compared to the non-  
2 mist irrigated treatment. Liu and Kang (2006b) also reported decreases of canopy  
3 temperature of wheat (*Triticum aestivum* L.) of 0.3 to 2.8 °C in a sprinkler irrigated field  
4 as compared to a non-sprinkled field.

5 In maize (*Zea mays* L.), microclimatic (air temperature and air VPD) and plant  
6 physiological (canopy temperature, plant transpiration, leaf water potential) changes  
7 have been reported during sprinkler irrigation. Steiner et al. (1983) compared the  
8 microclimatic and physiological conditions of crop maize under two types of irrigation  
9 system: center pivot sprinkler and surface irrigation. They reported that long-term daily  
10 average canopy and air temperatures at the center pivot field were 1.0 °C and 1.5 °C,  
11 respectively, cooler than at the surface irrigation field. This cooling effect of the center  
12 pivot irrigation was higher during days of high evaporative demand. Using a lateral  
13 move sprinkler irrigation system, Tolk et al. (1995) observed that during daytime  
14 irrigations the VPD decreased about 1.4 KPa, the canopy temperature decreased about  
15 5.3°C, and the transpiration rates decreased by 32%. Finally, Cavero et al. (2009)  
16 reported that during daytime solid-set sprinkler irrigation the air temperature and the  
17 VPD (measured at 0.5 m below crop canopy height) decreased between 3.3 and 4.4 °C,  
18 and between 1.0 and 1.2 KPa, respectively; these decreases were lower when  
19 monitored at higher measurement heights. Cavero et al. (2009) also reported that  
20 canopy temperature decreased between 4 to 6 °C, the crop transpiration rate was  
21 reduced by 58%, and leaf water potential increased from values of -1.2 to -1.4 MPa to  
22 values of -0.52 to -0.57 MPa. In general, these studies report that these microclimatic  
23 and plant physiological changes during the sprinkler irrigation event only last for a few  
24 hours after the irrigation finishes.

1 These microclimatic and physiological changes affect the efficiency of the sprinkler  
2 irrigation application (Tolk et al., 1995; Martínez-Cob et al., 2008), so they are relevant  
3 to the modeling of sprinkler irrigation efficiency (Zhao et al., 2012). It is important to  
4 gather information of those changes under different irrigation systems and  
5 meteorological, crop and land conditions so a larger database can be obtained to test  
6 those models under different scenarios.

7 In center pivots due to its rotation movement, the water application rates and irrigation  
8 duration vary along the pivot arm portions. As the abovementioned microclimatic and  
9 physiological changes are the consequence of the evaporation water lost during and  
10 after the irrigation and those losses depend on the application rates and irrigation  
11 duration, the microclimatic and physiological changes could be different along the  
12 different pivot arm portions and they could also be different compared to other sprinkler  
13 irrigation systems. This variability and a detailed analysis of the microclimatic changes  
14 during the irrigation events were not addressed in previous work (Steiner et al., 1983).  
15 The goal of this work was to study the variability of the magnitude and duration of the  
16 microclimatic (air temperature and VPD) and physiological (canopy temperature and  
17 plant transpiration) changes in a center pivot sprinkler system irrigating a maize crop  
18 and how they affect the water use.

## 19 **2. MATERIALS AND METHODS**

### 20 **2.1 Experimental site**

21 The experiment was carried out from June to September 2008 in a maize commercial  
22 plot of 32.3 ha irrigated with a center pivot sprinkler irrigation system (VXP, Irrifrance,  
23 Paulhan, France), and located at Valfarta (Huesca, Spain) (41°33'N latitude and 0°07'W

1 longitude; 354 m altitude). The climate is Mediterranean semiarid with a yearly average  
2 precipitation of 400 mm and mean annual air temperature of 14.3° C.

3 Maize cultivar Pioneer PR34N44 was planted on 15 April 2008. A final plant density of  
4 68,000 plants ha<sup>-1</sup> was attained as determined by counting and averaging the number of  
5 plants within 15 sampling spots of 3.0 m<sup>2</sup> each. This sampling was performed on 6  
6 October, few days before harvest. The soil is classified as Typic Torrfluvents and the  
7 texture is silty loam. Agronomic practices (fertilization, weeds and pest control, etc.)  
8 common in the region were conducted by the owner of the commercial plot. For  
9 irrigation scheduling, the farmer obtained the maize irrigation requirements from the  
10 Spanish Irrigation Advisory System (SIAR) (MARM, 2011). The SIAR System includes a  
11 network of automated weather stations, one of them located 3 km southeast from the  
12 experimental plot. This station (thereafter the 'nearby grass station') is located over  
13 grass following the reference conditions defined by Allen et al. (1998). The SIAR  
14 System uses the average daily meteorological data recorded (air temperature and  
15 relative humidity, wind speed and direction, solar radiation and precipitation) to get daily  
16 estimates of ET<sub>o</sub>, and locally adjusted maize crop coefficients to calculate the weekly  
17 crop evapotranspiration and irrigation requirements using the FAO approach (Allen et al.  
18 1998). An irrigation efficiency of 0.85 is used to calculate the irrigation requirements.

19 The pivot arm was 322 m long and was divided into six portions of 49.4 m length each  
20 and a final overhang of 25.6 m length (Fig.1, Table 1). The main pipe had a diameter of  
21 0.163 m. The number of impact sprinklers in each pivot arm portion and the general  
22 characteristics of the center pivot system are shown in Table 1. All sprinklers had a  
23 pressure regulator (Model PSR30, Senninger Irrigation Inc., Clement, FL, USA) and  
24 were located at the top of the main pipe. A complete turn of the center pivot over the

1 whole plot lasted about 31 h. The weekly irrigation requirement was divided by the water  
2 depth applied by the pivot ( $\approx 13$  mm) to determine the number of irrigations per week.  
3 Irrigation pressure ( $P_i$ , kPa) was measured along the measurement period using  
4 pressure transducers (Model 2200/2600, Gems Basingstoke, Hampshire, UK) installed  
5 in the last sprinkler of pivot arm portions 2, 4 and 5 (Fig. 1). Each pressure transducer  
6 was placed between the pressure regulator and the sprinkler and connected to a logger  
7 (Model ES120, Dickson, Addison, IL, USA) which monitored and recorded pressure  
8 values every 5 min.

## 9 **2.2 Microclimatic and physiological changes**

10 Determining the microclimatic and physiological changes occurring during the irrigation  
11 events required simultaneous measurements at an irrigated and a non-irrigated area  
12 (i.e. an area under the same conditions than those of the irrigated area but irrigated at a  
13 different time). Thus three meteorological stations (thereafter, the experimental weather  
14 stations A, B and C) and sap flow measurement systems were installed at a transect AC  
15 located at northeast of the plot (Fig.1), approximately in the middle of the pivot arm  
16 portions 2, 4 and 5, respectively. A fourth meteorological station (thereafter, the  
17 experimental weather station D) and a sap flow measurement system were installed at  
18 spot D, 270 m far away from transect AC (Fig. 1). Two treatments were established in  
19 this field experiment: a) *moist* treatment, measurements taken at the stations A to C in  
20 the transect AC when the pivot arm was irrigating it; b) *dry* treatment, measurements  
21 simultaneously taken at the station D when the transect AC was being irrigated. At that  
22 time, about 8 to 10 h have passed since the pivot arm irrigated the spot D due to the  
23 duration of the rotating movement (counter clockwise) of the pivot (about 31 h), and the

1 distance between the dry spot D and the transect AC (270 m). This time was enough to  
2 dry out all intercepted water from plants in the area surrounding that spot D by the time  
3 the pivot arm reached the transect AC. For this reason, that spot D was considered the  
4 dry treatment. The size of the pivot irrigation system, its speed and the localization of  
5 the different sensors allowed enough fetch for the different measurements.

6 An air temperature and relative humidity probe (HMP45C, Vaisala, Helsinki, Finland)  
7 was installed at each experimental weather station (A to D) at 2.9 m above ground.  
8 Measurement height was kept constant along the experiment. The accuracy of the  
9 probes was  $\pm 0.3^{\circ}\text{C}$  for air temperature and  $\pm 2\%$  for relative humidity. For canopy  
10 temperature measurement, an infrared thermometer (Model IRR-P, Apogee  
11 Instruments, Inc., Roseville, CA) with an accuracy of  $\pm 0.5^{\circ}\text{C}$  was also placed at three of  
12 the experimental weather stations (A, B and D) at 1.0 m above the crop canopy with an  
13 angle of  $45^{\circ}$ , oriented towards the north. In addition, a net radiometer (NR-Lite,  
14 Kipp&Zonen, Delft, The Netherlands) with an accuracy  $\pm 30 \text{ W m}^{-2}$  at  $1000 \text{ W m}^{-2}$ , and a  
15 cup anemometer (A100R, Vector Instruments, Rhyl, UK) with an accuracy of  $\pm 0.1 \text{ m s}^{-1}$   
16 were installed at 2.9 m above ground at experimental weather station D. At each  
17 experimental weather station, those variables (air temperature and relative humidity,  
18 canopy temperature, and, at spot D, net radiation and windspeed) were monitored  
19 continuously every 10 s and their average values were recorded every 5 min by a  
20 datalogger (model CR10X at experimental weather stations A and B, model CR23X at  
21 experimental weather stations C and D, Campbell Scientific, Inc. Logan, UT, USA). The  
22 VPD was calculated from the recorded data of air temperature and relative humidity,  
23 following the methodology described by Allen et al. (1998). These meteorological data

1 were also used for a direct estimation of maize evapotranspiration (ET) at each spot A,  
2 B, C and D as described in section 2.3.

3 The transpiration rates were determined from sap flow measurements using the heat  
4 balance method (Baker and van Bavel, 1987; Weibel and Boersma, 1995; Van Bavel  
5 2005). This method was chosen because it had been previously used on maize in  
6 similar studies to this (Tolk et al. 1995; Martínez-Cob et al. 2008). Next to each  
7 meteorological station, a Flow4 datalogger (Dynamax, Houston, TX, USA) was installed  
8 to monitor, log and process data collected by four sap gauges SGB19 (Dynamax) each  
9 of them installed on a plant. Readings were taken every 10 min. The sap gauges were  
10 moved to a another set of four plants within the same area of the field on July 25 and 14  
11 August of 2008 to avoid any possible damage to the plants (Van Bavel 2005). Each  
12 gauge had a soft foam collar surrounding the electronics. In addition, once installed in  
13 the plant, each gauge was surrounded by a weather shield (aluminium bubble foil) such  
14 it held a cylindrical shape. The aluminium top shield was secured using insulation tape.  
15 The shield kept out water and prevented radiation from affecting readings (Van Bavel,  
16 2005). Following this author, the datalogger was set to apply a continuous average  
17 voltage of 4.0 V while the heater resistance of the different gauges varied between 58.9  
18 to 64.6  $\Omega$ . Van Bavel (2005) thoroughly describes the elements of the gauges, the  
19 electronics, the recorded variables and the equations used to process them to obtain  
20 transpiration rates at each gauge. The 10-min transpiration rates at each measurement  
21 spot were determined as the average of those obtained from the four sampled plants  
22 per spot. These values were determined in  $\text{g h}^{-1}$  and transformed into  $\text{mm h}^{-1}$  using the  
23 average number of plants  $\text{m}^{-2}$  measured at each spot ( $6.8 \text{ plants m}^{-2}$ ). Unlike the 5-min  
24 averages of air temperature and VPD and canopy temperature that were recorded

1 continuously along the experiment, the 10-min transpiration rates were only recorded for  
2 specific irrigation events due to limitations of the memory of the dataloggers used. For  
3 the abovementioned time scan (10 min), the datalogger's memory could only hold 24 h  
4 data so the values from 3 hours before the pivot arm passed over the transect AC until  
5 at least 6 hour after passing were recorded. Those specific irrigation events were  
6 monitored in situ, in general once per week.

7 We considered two set of data for the different variables (temperature and VPD of the  
8 air, canopy temperature and plant transpiration):

9 a) *Transpiration-measured* irrigation events: the seven irrigation events for which  
10 plant transpiration rate was measured and we were in situ to observe when the  
11 center pivot was passing over the transect AC.

12 b) *Remaining* irrigation events: the 27 irrigation events for which transpiration rate  
13 was not measured and we were not in situ to observe when the center pivot was  
14 passing over the transect AC.

15 A *transpiration-measured* irrigation event was established as the time ( $t_{ir}$ ) that took the  
16 pivot to run over a distance L of 18 m, 9 m either side of the transect AC. This value of 9  
17 m was established by visual inspection of the moistening radius of the pivot sprinklers.  
18 For each irrigation event, the value of  $t_{ir}$  was different for each monitored pivot arm  
19 portion (2, 4 and 5). For each *transpiration-measured* irrigation event the 5-min irrigation  
20 pressure values were averaged for the time  $t_{ir}$ . These average irrigation pressure values  
21 ( $P_i$ , kPa) were used to calculate the gross water depth applied ( $I_s$ , mm) at each  
22 monitored pivot arm portion. The following expression, derived from the Torricelli  
23 equation (Norman et al. 1990), was used:

$$1 \quad I_s = \frac{0.00035 \pi c_d P i^{0.5} (d_L^2 + d_s^2) 3600 t_g}{A_s} \quad (1)$$

2 where:  $c_d$  is the discharge coefficient, 0.98 (Martínez-Cob et al., 2008);  $d_L$  is the large  
 3 nozzle diameter (mm);  $d_s$  is the small nozzle diameter (mm);  $t_g$  is the time (h) to  
 4 complete a turn;  $A_s$  is the surface area (m<sup>2</sup>) irrigated by the sprinklers of pivot arm  
 5 portion. The corresponding surface area for pivot arm portions 2, 4 and 5 were 23,177,  
 6 53,887 and 69,242 m<sup>2</sup>, respectively.

7 The time  $t_g$  was determined for each pivot arm portion as follows:

$$8 \quad t_g = \frac{2\pi \cdot r}{\omega} \quad (2)$$

9 where:  $r$  is the radius at the end of the evaluated pivot arm portion (m);  $\omega$  is the angular  
 10 speed of the pivot (m h<sup>-1</sup>) computed from the values of  $t_{ir}$ .

11 The *remaining* irrigation events were not identified in situ. They were defined as those  
 12 periods for which differences between the two treatments (*dry* and *moist*), for each pivot  
 13 arm portion and variable (temperature and VPD of the air, and canopy temperature),  
 14 were higher than the accuracy of measuring instruments, and the evolution of the 5-min  
 15 values of these variables was similar to that observed during the *transpiration-measured*  
 16 irrigation events. Only those *remaining* irrigation events identified for daytime periods  
 17 (between 8:00 and 18:00 h Greenwich Mean Time (GMT)) were selected.

18 The half-hour values of wind speed and direction, solar radiation, air temperature, and  
 19 relative humidity recorded by the 'nearby grass station' were used to characterize the

1 general standard meteorological conditions in the area during the different irrigation  
2 events.

### 3 **2.3. Estimation of maize evapotranspiration**

4 It is expected that the maize evapotranspiration is also affected by the microclimatic and  
5 plant physiological changes occurring before, during and after the irrigation events. The  
6 FAO Penman-Monteith equation (Allen et al., 1998) only describes a particular  
7 application of the Penman-Monteith equation, that for calculation of reference  
8 evapotranspiration (ET for an hypothetical, grass-like crop, 0.12 m high and with a fixed  
9 surface resistance of 70 s m<sup>-1</sup>). However, the Penman-Monteith equation can be used  
10 for the direct calculation (i.e. without using crop coefficients and reference ET) of any  
11 crop evapotranspiration as the surface and aerodynamic resistance values required in  
12 these computations are crop specific (Allen et al., 1996, 1998). Therefore, this equation  
13 was chosen for estimation of maize evapotranspiration for each *transpiration-measured*  
14 irrigation event at the four spots A to D using the corresponding 5-min averages air  
15 temperature and relative humidity recorded at the experimental weather stations during  
16 the different phases identified for transpiration changes. Thus three sets of moist maize  
17 ET (for pivot arm portions 2, 4 and 5) and one set of dry maize ET were computed  
18 following Allen et al. (1996):

$$19 \quad ET_{\text{corn}} = \frac{300}{10^6 \lambda} \frac{\Delta(R_n - G) + \rho_a c_p \text{VPD} / r_a}{\Delta + \gamma \left(1 + \frac{r_c}{r_a}\right)} \quad (3)$$

20 where:  $\lambda$ , latent heat of vaporization (MJ Kg<sup>-1</sup>);  $R_n$ , net radiation (W m<sup>-2</sup>);  $G$ , soil heat flux  
21 (W m<sup>-2</sup>);  $\Delta$ , slope of the saturation vapour pressure curve versus the temperature (kPa

1 °C<sup>-1</sup>);  $\rho_a$ , air density (Kg m<sup>-3</sup>);  $c_p$ , specific heat of the air (J Kg<sup>-1</sup> °C<sup>-1</sup>); VPD, vapour  
 2 pressure deficit (kPa);  $r_a$ , aerodynamic resistance of maize (s m<sup>-1</sup>);  $r_c$ , bulk stomatal  
 3 (canopy) resistance of maize (s m<sup>-1</sup>);  $\gamma$ , psychrometric constant (kPa °C<sup>-1</sup>).

4 The variables  $\lambda$ ,  $\Delta$ ,  $\rho_a$ ,  $\gamma$ , and  $c_p$  were estimated from the measured air temperature and  
 5 relative humidity following standard procedures described by Allen et al. (1998). Note  
 6 that the estimated values of these parameters were different at the four spots A to D as  
 7 the corresponding values of air temperature and relative humidity were used. G was  
 8 estimated from net radiation following Allen et al. (1996):

$$9 \quad G = 0.4 e^{-0.5LAI} R_n \quad (4)$$

10 where LAI is the daily leaf area index estimated from measured crop height as  
 11 suggested by Allen et al. (1996). As the crop height at the four spots A to D was quite  
 12 similar, only a single set of LAI values was estimated and used.

13 The aerodynamic resistance  $r_a$  (s m<sup>-1</sup>) to vapour transfer was estimated following Allen  
 14 et al. (1996):

$$15 \quad r_a = \frac{\left[ \ln\left(\frac{z_u - d}{z_{0m}}\right) \right] \left[ \ln\left(\frac{z_h - d}{z_{0h}}\right) \right]}{k^2 u_{zu}} \quad (5)$$

16 where:  $u_{zu}$  is the wind speed (m s<sup>-1</sup>) measured at a height  $z_u$  at spot D, it was assumed  
 17 that wind speed was not affected by irrigation at the transect A-C;  $k$  is the von Karman's  
 18 constant (0.41);  $z_u$  and  $z_h$  are the measurement heights (m) above ground of wind  
 19 speed, and air temperature and relative humidity, respectively; and  $d$ ,  $z_{0m}$ , and  $z_{0h}$  (all  
 20 three in m) are the zero-plane displacement and the roughness lengths for momentum

1 and heat transfer, respectively, estimated (daily) as a function of crop height ( $h_c$ ) and  
 2 LAI following Farahani and Bausch (1995) and Kjelgaard et al. (1994):

$$3 \quad d = 1.1 h_c \ln[1 + (c_d \text{LAI})^{1/4}] \quad (6)$$

$$4 \quad z_{om} = 0.3 h_c (1 - d/h_c) \quad (7)$$

5 and,

$$6 \quad z_{oh} = 0.2 z_{om} \quad (8)$$

7 where  $c_d$  is the mean drag coefficient for individual leaves (0.07). Eq. (6) was chosen as  
 8 the product ( $c_d \text{LAI}$ ) was above 0.2 (Farahani and Bausch 1995) due to the LAI values  
 9 around 4.0 estimated during the monitoring period as crop height was about 2.5 m. The  
 10 same roughness parameters,  $d$ ,  $z_{om}$  and  $z_{oh}$ , were used at the four spots A to D as the  
 11 average crop height was similar.

12 The bulk canopy resistance ( $\text{s m}^{-1}$ ),  $r_c$ , was estimated following Farahani and Bausch  
 13 (1995):

$$14 \quad r_c = \left\{ c_0 \text{LAI} + \frac{c_1}{c_2 C} \ln \left[ \frac{1 + c_2 C R_s}{1 + c_2 C R_s \exp(-C \text{LAI})} \right] \right\}^{-1} \quad (9)$$

15 where:  $R_s$  is the incoming solar radiation ( $\text{W m}^{-2}$ );  $c_0$  is the minimum stomatal  
 16 conductance ( $0.0005 \text{ m s}^{-1}$ );  $c_1$  and  $c_2$  are constants defined as  $c_1 = 3.2\text{E-}5 \text{ m s}^{-1}$  and  $c_2$   
 17  $= 5.7\text{E-}5 \text{ m s}^{-1}$ ; and  $C$  is the light extinction coefficient, assumed to be 0.50 as  
 18 suggested by Cavero et al. (1999, 2000) for similar crop and climatic conditions to those  
 19 in this work.  $R_s$  was that measured at the 'nearby grass station' and it was assumed to  
 20 be the same along the pivot surface area.

1 Net radiation was only measured at spot D. Though net short-wave radiation should not  
 2 be affected by irrigated, net long-wave radiation could be. Therefore, the net radiation  
 3 (in  $W\ m^{-2}$ ) at the four spots A to D was estimated as described in Allen et al. (1996,  
 4 1998):

$$5\ R_n = (1 - \alpha)R_s - \sigma T_{mk}^4 \left( 0.34 - 0.14\sqrt{e_a} \right) \left( 1.35 \frac{R_s}{R_{sol}} - 0.35 \right) \quad (10)$$

6 where:  $\alpha$  is albedo (assumed to be equal to 0.23);  $\sigma$  is the Stefan-Boltzmann constant  
 7 ( $2.04E-10\ MJ\ K^{-4}\ m^{-2}\ h^{-1}$ );  $T_{mk}$  is the mean air temperature at the corresponding 5-min  
 8 period and spot A to D ( $^{\circ}K$ );  $e_a$  is actual vapour pressure (kPa) estimated from air  
 9 temperature and relative humidity at each spot A to D; and  $R_{sol}$  is the clear-sky solar  
 10 radiation ( $W\ m^{-2}$ ) estimated at clear-sky days following Allen et al. (1996, 1998).

#### 11 **2.4. Statistical analysis**

12 The 5-min averages of air temperature, air VPD and canopy temperature, and the 10-  
 13 min values of transpiration rates recorded for the *moist* treatment were compared to  
 14 those simultaneously recorded for the dry treatment for each *transpiration-measured*  
 15 irrigation event. Three phases were identified: 1) phase *before*: time period that started  
 16 when the differences between the *dry* and *moist* treatments were higher than the  
 17 accuracy of measuring instruments and finished when the pivot arm portion was 9 m  
 18 ahead of the transect AC; 2) phase *during*: time period corresponding to  $t_{ir}$ ; and 3)  
 19 phase *after*: time period that started when the arm portion had surpassed the transect  
 20 AC by 9 m and finished when the differences between treatments were lower than the  
 21 accuracy of measuring instruments. For a given irrigation event and pivot arm portion,  
 22 the duration of the phases *before* and *after* was independently established for each

1 monitored variable. Once identified the different phases for each irrigation event, the 5-  
2 min values (or 10-min values) of each variable and pivot arm portion were averaged  
3 over the time of duration of each phase. These average values obtained in the *moist*  
4 and *dry* treatments in the different irrigation events for each pivot arm portion and phase  
5 were compared with a paired t-test and a level of significance of  $P= 0.05$ .

6 For the *remaining* irrigation events, it was established a single phase integrating the  
7 phases *before*, *during* and *after* established for the *transpiration-measured* irrigated  
8 events. The 5-min values of air temperature and VPD and canopy temperature were  
9 averaged for the overall duration of each *remaining* irrigation event. The values obtained  
10 in the *moist* and *dry* treatments in the different irrigation events for each pivot arm  
11 portion were compared with a paired t-test and a level of significance of  $P= 0.05$ .

12 The values of ET obtained in the *moist* and *dry* treatments in the different irrigation  
13 events for each pivot arm portion and phase (before, during and after) were compared  
14 using a paired t-test and a level of significance of  $P= 0.05$ .

15 Linear regression analysis was used to determine the relationships between the  
16 microclimatic and physiological changes due to sprinkler irrigation and the climatic  
17 conditions.

18 The Statgraphics software was used for the analysis.

19

### 1 3. RESULTS AND DISCUSSION

#### 2 3.1 Characteristics of the irrigation events

3 The average duration of the *transpiration-measured* irrigation events decreased with the  
4 distance to the center of the pivot ranging from 1.6 h for pivot arm portion 2 to 0.5 h for  
5 pivot arm portion 5 (Table 2). All these irrigation events started between 8:25 and 10:55  
6 GMT. The average irrigation pressure in the three pivot arm portions was 197 kPa with  
7 a coefficient of variation (CV) of 3% (Table 2). This low CV value indicated a quite  
8 constant irrigation pressure during the *transpiration-measured* irrigation events. On  
9 average, the irrigation pressure in the pivot arm portion 2 was slightly higher (209 kPa)  
10 than that in the portions 4 (190 kPa) and 5 (192 kPa) (Table 2). The average applied  
11 water in the three monitored pivot arm portions was similar, 14.2 (pivot arm portion 2),  
12 13.1 (pivot arm portion 4), and 13.9 mm (pivot arm portion 5).

13 There were some differences between the average meteorological conditions recorded  
14 during the *transpiration-measured* irrigations events (Table 3). The overall mean air  
15 temperature was 27.2°C, but the individual mean temperatures ranged between 22.8 °C  
16 (13 August) and 32.5 °C (31 July). The cooler irrigation event (13 August) was also the  
17 windiest, while the hottest irrigation event (31 July) showed the highest VPD. No  
18 precipitation was recorded during the *transpiration-measured* irrigation events.

#### 19 3.2 Microclimatic changes

20 The time evolution of air temperature and VPD recorded from 2 h before until 6 h after  
21 the *transpiration-measured* irrigation event on 6 August 2008 is shown in Fig. 2. Before  
22 the irrigation event, there was a period for which there were no differences between  
23 treatments; but as the center pivot was approaching transect AC, the values recorded at

1 the *moist* treatment started to decrease compared to the dry treatment (phase *before*).  
2 For the phase *during*, that decrease became much higher. Finally, for the phase *after*,  
3 the observed reductions at the *moist* treatment, although gradually diminishing, lasted  
4 for some time until finally the values became again similar to those recorded at the *dry*  
5 treatment. In general, this time evolution was similar to that observed for all  
6 *transpiration-measured* irrigation events.

7 In general terms, the evolution of the monitored variables studied in this work during and  
8 after the *transpiration-measured* irrigation events was similar to that described in  
9 previous works (Steiner et al. 1983; Thompson et al. 1993; Tolk et al. 1995; Saadia et  
10 al. 1996; Liu and Kang 2006a; Cavero et al. 2009). However, a reduction of the air  
11 temperature and VPD before the pivot irrigated the AC transect has not been previously  
12 observed. Monteith and Unsworth (2008) indicated that the values recorded by a  
13 meteorological station are affected by the vegetation type and characteristics and the  
14 plant-atmosphere interchange within the fetch distance surrounding the station,  
15 particularly upwind the measurement spot. Roughly, the fetch distance is estimated as  
16 100 times the measurement height; thus, in this work, the fetch distance was about 290  
17 m around the station skewed to the upwind direction. Due to the rotating movement of  
18 the pivot, the nearby areas were already being moistened by irrigation as the pivot was  
19 approaching the transect AC, leading to microclimatic changes at those nearby areas,  
20 within the fetch distance of the station at that transect. Then those microclimatic  
21 changes at the nearby areas were likely causing the differences among treatments  
22 observed at the phase *before*. This effect was somewhat larger when the wind was  
23 blowing from the east as the pivot rotation was counter-clockwise. Thus, the average  
24 decrease in air temperature and VPD were about 0.7 °C and 0.21 kPa, respectively,

1 with a duration of about 0.9-1.0 h for those events for which predominant wind direction  
2 was east (with average windspeed of about  $1.9 \text{ m s}^{-1}$ ), while the average decrease in air  
3 temperature and VPD were about  $0.5 \text{ }^{\circ}\text{C}$  and  $0.15 \text{ kPa}$ , respectively, with a duration of  
4 about 0.5-0.7 h for those events for which wind was blowing from other directions (with  
5 average windspeed of about  $3.0 \text{ m s}^{-1}$ ).

6 The differences between treatments for the air temperature and VPD during the  
7 *transpiration-measured* irrigation events were significant ( $P=0.05$ ) for the three phases,  
8 *before*, *during* and *after*, and the three pivot arm portions (Tables 4 and 5). For the  
9 phase *before*, the average decreases for the *moist* treatment were  $0.5$  to  $0.7 \text{ }^{\circ}\text{C}$  ( $2.1$  to  
10  $2.8 \%$ ) for air temperature, and  $0.16$  to  $0.25 \text{ kPa}$  ( $14.2$  to  $20.6 \%$ ) for VPD of the air. The  
11 average duration of phase *before* was  $0.6$  to  $0.8 \text{ h}$  for air temperature and  $0.6$  to  $0.7 \text{ h}$   
12 for VPD of the air. For the phase *during*, the average decreases for the *moist* treatment  
13 were much higher than those for the phase *before* and amounted  $1.8$  to  $2.1 \text{ }^{\circ}\text{C}$  ( $7.1$  to  
14  $8.2\%$ ) for air temperature and  $0.53$  to  $0.61 \text{ kPa}$  ( $37.8$  to  $45.9\%$ ) for VPD of the air.  
15 Finally, for the phase *after*, the decreases for the *moist* treatment were lower than those  
16 for the phase *during* amounting  $0.8$  to  $1.3 \text{ }^{\circ}\text{C}$  ( $2.8$  to  $4.9\%$ ) for air temperature and  $0.30$   
17 to  $0.41 \text{ kPa}$  ( $14.8$  to  $26.6\%$ ) for VPD of the air.

18 The observed decreases in air temperature and VPD in this study were similar to the  
19 reductions in the long-term daily averages of air temperature and VPD due to center  
20 pivot sprinkler irrigation reported by Steiner et al. (1983). The decreases in air  
21 temperature and VPD listed on Tables 4 and 5 for the phases *during* and *after* were  
22 within the ranges reported by previous works on sprinkler irrigation with other irrigation  
23 systems (Thompson et al., 1993; Tolk et al., 1995; Saadia et al., 1996; Liu and Kang  
24 2006a; Caverro et al., 2009). The decrease in air temperature and VPD lasted about 1.3

1 h after the irrigation event finished (Tables 4 and 5), which is similar to durations reported in other works (Thompson et al., 1993; Tolk et al., 1995; Saadia et al., 1996; Caverro et al., 2009).

4 For the *remaining* irrigation events, the air temperature for the *moist* treatment significantly decreased 1.4 to 1.6 °C (5.3 to 6.0 %) on average, while the VPD of the air significantly decreased 0.46 to 0.48 kPa (24.2 to 26.2 %) on average (Table 6). These decreases were slightly higher than those observed for the *transpiration-measured* irrigation events when the phases *before*, *during* and *after* were integrated into a single period (Table 6). This slight difference between the microclimatic changes observed for the *transpiration-measured* and those for the *remaining* irrigation events was probably due to the climatic conditions during both measurement periods. As discussed later, the observed decreases in air temperature and VPD for the *moist* treatment were higher as the air temperature and VPD at the 'nearby grass station' were higher (Figs.3 and 4). Given that the *transpiration-measured* irrigation events were monitored early in the morning while the *remaining* irrigation events covered the whole daytime period, air temperature and VPD at the 'nearby grass station' were lower during the *transpiration-measured* irrigation events. Thus, changes were slightly lower in the *transpiration-measured* irrigation events.

19 The magnitude of the decreases in air temperature and VPD for the *moist* treatment was, in general terms, relatively similar between the three pivot arm portions for both *transpiration-measured* and *remaining* irrigation events (Table 6); nevertheless the decreases in VPD of the air for the phase *after* at the former irrigation events slightly increased from the center to the end of the pivot (Table 5). The main difference between the three pivot arm portions was the duration of those decreases in air temperature and

1 VPD. That duration was highly variable as indicated by the high coefficients of variation  
2 obtained (Tables 4 to 6), but, on average, the total durations of the microclimatic  
3 changes observed for the *transpiration-measured* irrigation events (when integrating the  
4 three phases *before*, *during* and *after*) were much higher at pivot arm portion 2 (the  
5 closest to the center of the pivot) than the duration at the pivot arm portion 5, the  
6 furthest from the center of the pivot. Relatively similar results were observed for the  
7 *remaining* irrigation events; the duration of the microclimatic changes at pivot arm  
8 portion 2 was about 0.6 h longer than the duration at pivot arm portion 5 (Table 6).  
9 There were no differences in the duration of the microclimatic changes between the  
10 pivot arm portions 4 and 5. This difference in the duration of the microclimatic changes  
11 was mainly due to the longer duration of the irrigation at pivot arm portion 2 (Table 5)  
12 and at a lesser extent to the longer presence of the pivot irrigating nearby areas within  
13 the fetch distance to experimental weather station A such that the microclimatic  
14 changes in those areas were also affecting to the readings of that station.

15 The average decrease in air temperature and VPD observed both in the 7 *transpiration-*  
16 *measured* irrigation events and the 27 *remaining* irrigation events was higher as the air  
17 temperature and VPD measured over grass at the 'nearby grass station' were also  
18 higher (Figs. 3 and 4). The linear regressions between the decreases in air temperature  
19 for the *moist* treatment and the air temperature at the 'nearby grass station', and  
20 between the decreases in VPD of the air for the *moist* treatment and the VPD of the air  
21 at the 'nearby grass station' were significant for the three pivot arm portions for the  
22 phase *during* at the *transpiration-measured* irrigation events (Figs. 3A and 4A) and for  
23 the whole period of microclimatic changes at the *remaining* irrigation events (Figs. 3B  
24 and 4B). The corresponding coefficients of determination ranged from 0.50 to 0.85. The

1 12 regression slopes shown in Figs. 3 and 4 were significant at  $P < 0.01$  (except one  
2 significant at  $P=0.07$ ). In general, the relationships between the microclimatic changes  
3 and the mean meteorological conditions at the 'nearby grass station' increased from  
4 pivot arm portion 2 to pivot arm portion 5. During the irrigation phase, the reduction of  
5 air temperature and VPD as the value of these variables increased in the 'nearby grass  
6 station' was greater in the outer pivot arm portion, probably due to the higher  
7 instantaneous water application rate. The relationships were stronger for the  
8 *transpiration-monitored* irrigation events (Figs. 3A and 4A) as they are calculated only  
9 for the phase *during*, while these relationships for the *remaining* irrigation events (Figs.  
10 3B and 4B) were calculated for the whole period of microclimatic changes. These  
11 weaker relationships found at the *remaining* irrigation events were due to the integration  
12 of the three phases identified for the *transpiration-measured* irrigation events. In other  
13 words, including the microclimatic changes for the phases *before* and *after* smoothes  
14 the relationship between the general climatic conditions and the microclimatic changes  
15 observed for the phase *during*.

### 16 **3.3 Physiological changes**

17 Both physiological variables studied in this work (maize canopy temperature and  
18 transpiration) showed a similar behaviour for the monitored irrigation events. The time  
19 evolution recorded from 2 h before until 6 h after the *transpiration-measured* irrigation  
20 event on 6 August 2008 is shown in Fig. 2. Before the irrigation event, there was a  
21 period for which there were no differences between treatments. As the center pivot was  
22 approaching to transect AC, the canopy temperature and maize transpiration recorded  
23 at the *moist* treatment started to decrease compared to the dry treatment (phase  
24 *before*). For the phase *during*, that decrease became much higher. Finally, for the phase

1 *after*, the observed reductions at the *moist* treatment, although gradually diminishing,  
2 lasted for some time until finally the values became again similar to those recorded at  
3 the *dry* treatment. Thus, for the *transpiration-measured* irrigation events phase *before*  
4 the canopy temperature and the transpiration rates for the *moist* treatment decreased  
5 1.0 to 1.2 °C (4.3 to 5.2%) and 0.15 to 0.19 mm h<sup>-1</sup> (23.8 to 31.7 %), respectively  
6 (Tables 7 and 8). For the phase *during*, these decreases were higher and ranged from  
7 3.1 to 3.8 °C (11.7 to 14.5 %) for canopy temperature and from 0.22 to 0.28 mm h<sup>-1</sup>  
8 (30.1 to 36.4 %) for transpiration rates. For the phase *after*, the physiological changes  
9 were smaller; thus the decreases for the *moist* treatment ranged from 1.1 to 1.4 °C (4.0  
10 to 5.2 %) for canopy temperature and 0.14 to 0.24 mm h<sup>-1</sup> (17.1 to 27.9 %) for  
11 transpiration rates (Tables 7 and 8).

12 The magnitude and duration of the canopy temperature decreases for the *moist*  
13 treatment for the *remaining* irrigation events were similar to those observed for the  
14 *transpiration-measured* irrigation events when integrating the three phases, *before*,  
15 *during* and *after* (Table 6). Again, integrating the phases *before* and *after* smoothes the  
16 canopy temperature changes observed for the phase *during*. On average, the decrease  
17 of canopy temperature for the *remaining* irrigation events was 1.8 °C, which was similar  
18 in the *transpiration-measured* irrigation events.

19 Transpiration reduction due to irrigation in the *moist* treatment ranged from 0.75 mm  
20 (22%) to 1.03 mm (30%) (Table 8), with a higher reduction as closer to the center of the  
21 pivot. This represents between 5 to 7% of the applied water. Tolk et al. (1995) working  
22 with a lateral move sprinkler irrigation system found 1.59 mm (32%) transpiration  
23 reduction, which represented 10% of applied water. Considering the time period when  
24 transpiration changes occurred, the estimated crop evapotranspiration (ET<sub>c</sub>) was also

1 reduced in the *moist* treatment (Table 9). However, the reduction of  $ET_c$  (8 to 10%) in  
2 the *moist* treatment was less than the reduction of transpiration (22 to 30%) and was  
3 similar in the different pivot arm portions. In general,  $ET_c$  reduction *before* and *after*  
4 irrigation was less than 8%, and was around 15% *during* the irrigation. The  $ET_c$   
5 reduction due to irrigation in the *moist* treatment ranged from 0.22 mm (8%) to 0.26 mm  
6 (10%) (Table 9), which represents 1.5 to 1.8% of the applied water. This reduction must  
7 be taken into account when calculating the irrigation requirements. Frost and Schwallen  
8 (1960) reported a 18% decrease of crop evapotranspiration due to sprinkler irrigation,  
9 which was greater than the 8 to 10% estimated in our work.

10 Transpiration reduction *during* irrigation was related to the decrease of air temperature  
11 and VPD, but not to the decrease of canopy temperature (Fig. 5). The strongest  
12 relationship was with the decrease of air VPD as found by others (Tolk et al. 1995; Ray  
13 et al. 2002; Yu et al. 2003). A stepwise regression analysis of the transpiration reduction  
14 versus the decrease of air temperature, of air VPD and of canopy temperature showed  
15 that the reduction of air VPD was the only variable that explained the transpiration  
16 reduction. However, for the phases *before* and *after* the smaller changes in all these  
17 variables did not allow to establish a clear relationship with the slight reduction of  
18 transpiration rate (Fig. 5).

19 Similar to the microclimatic change of air VPD due to irrigation, the transpiration rate  
20 reduction was also greater when the VPD of the air at the 'nearby grass station' was  
21 higher (drier days). Thus, Fig. 6 shows the strong relationship between the transpiration  
22 reduction due to irrigation and the VPD of the air at the 'nearby grass station'. This  
23 result agrees with previous works (Tolk et al. 1995; Martínez-Cob et al. 2008; Caverro et  
24 al. 2009). However, the reduction of canopy temperature was not related to the VPD of

1 the air but for the pivot arm portion 4 was greater when the temperature of the air at the  
2 'nearby grass station' was higher (warmer days) (Fig. 7). The strength of the relationship  
3 between transpiration reduction and the VPD of the air was higher for the pivot arm  
4 portions 4 and 5 (Fig. 6). These results suggests that physiological changes due to  
5 sprinkler irrigation in the areas furthest from the centre of the pivot were more affected  
6 by the general climatic conditions outside the plot. Likewise, the results of Figs. 3A, 4A,  
7 6 and 7 indicate that the microclimatic and physiological changes are more relevant  
8 under high evaporative demand conditions as those changes are the result of the  
9 evaporation of a portion of the applied water during the irrigation.

10 The existence of microclimatic and physiological changes before the plants receive the  
11 irrigation water had not been previously reported in detail for sprinkler irrigation systems  
12 and it was likely due to the effect of the changes occurring in the nearby areas as the  
13 pivot arm was moving towards the monitored transect. This specific behaviour of the  
14 pivot irrigation systems before the irrigation events deserves to be modelled by sprinkler  
15 irrigation efficiency models that also include the microclimatic and physiological changes  
16 due to the irrigation (Zhao et al., 2012). Thus, the results reported here can be helpful  
17 for the improvement and application of those models under different conditions and  
18 scenarios.

19 The relative decrease of the canopy temperature due to irrigation was somewhat higher  
20 than that of the air temperature, while the relative decrease of the transpiration due to  
21 irrigation was higher than that of the VPD of the air. Also, the physiological changes for  
22 the phases *before* and *after* lasted, in general, longer than the microclimatic changes  
23 (Tables 4, 5, 7 and 8). There was a slight tendency for this duration being longer for the  
24 pivot arm portions 4 and 5 with respect to pivot arm portion 2. In general terms, the

1 decreases in canopy temperature and transpiration rates for the phase *during* observed  
2 in this work were lower than those reported for solid-set sprinkler irrigation systems  
3 (Cavero et al. 2009) and lateral-move sprinkler irrigation systems (Tolk et al. 1995)  
4 irrigating also a maize crop. Cavero et al. (2009) argued that higher application rates of  
5 irrigation water increase the cooling effect on plants and thus enhance the canopy  
6 temperature decreases at the *moist* treatment. In this study, application rates for the  
7 pivot arm portion 4 (21.8 mm h<sup>-1</sup>) were much higher than those of the studies by Tolk et  
8 al. (1995) and Cavero et al. (2009), which were much closer to the application rates for  
9 the pivot arm portion 2 (8.9 mm h<sup>-1</sup>) in this study. The duration of the irrigation event  
10 was much shorter in our study than at the two abovementioned works. This shorter  
11 irrigation duration could explain why the cooling effect of the irrigation water on canopy  
12 temperature was less than in the works of Tolk et al. (1995) and Cavero et al. (2009).  
13 Fig. 2 shows that the decrease in canopy temperature progresses as the irrigation is  
14 occurring. Thus shorter irrigation durations would lead to smaller canopy temperature  
15 decreases for the *moist* treatment. In addition to the shorter duration of the irrigation  
16 event (phase *during*), the lower decreases in air temperature, VPD of the air and  
17 transpiration rates (Tables 4, 5, 7 and 8) at the *moist* treatment obtained in this work  
18 compared with those reported by Tolk et al. (1995) and Cavero et al. (2009) were also  
19 probably due to the climatic conditions during those events, which in this work were  
20 performed before solar noon when the evaporative demand and the VPD of the air are,  
21 in general, lower than those for afternoon periods when the irrigations reported by those  
22 authors were performed. Despite the shorter duration of the *transpiration-measured*  
23 irrigation events in this study, the average canopy temperature decrease for pivot arm

1 portion 4 was higher than that of pivot arm portion 2 which agrees with the influence of  
2 the application rates as suggested by Cavero et al. (2009).

3 These microclimatic and physiological changes are the consequence of the evaporation  
4 of irrigation applied water while travelling through the air and the evaporation of  
5 intercepted water by stem and leaves of the plants. The amount of intercepted water  
6 depends mainly on the architecture of the crop and in the case of maize values of 0.4 to  
7 2.7 mm have been reported (Norman and Campbell 1983; Steiner et al. 1983). Thus,  
8 the volume of water evaporated during the irrigation is usually higher than that  
9 evaporated after the irrigation. Subsequently, the microclimatic and physiological  
10 changes are usually higher for the phase *during* than those for the phase *after*.

11 The temperature and VPD of the air were measured at 0.5 m above the crop canopy  
12 while canopy temperature is measured at the crop canopy height and transpiration of  
13 the plant with the sap flow integrates the transpiration along all the plant height. Cavero  
14 et al. (2009) found that the microclimatic changes due to sprinkler irrigation (decrease of  
15 air temperature and VPD) were smaller and lasted for less time after the irrigation as the  
16 measurement height was higher. Thus, the lower height of measurement of  
17 physiological changes (canopy temperature and plant transpiration) could explain that  
18 these changes lasted longer and were greater than the microclimatic changes.

#### 19 **4. CONCLUSIONS**

- 20 • Center pivot sprinkler irrigation significantly reduced air temperature and VPD  
21 (microclimatic changes) and canopy temperature and maize transpiration rates  
22 (physiological changes). These changes occurred for some time before (about  
23 0.6 to 2.1 h), during and some time after (about 0.8 to 2.4 h) the irrigation events.

- 1       • Physiological changes lasted longer than microclimatic changes, particularly after  
2       the irrigation events, likely due to the effect of the evaporation of the intercepted  
3       water and to the higher measurement height of microclimatic changes.
- 4       • Center pivot sprinkler irrigation decreased the air temperature by 1.8 to 2.1 °C,  
5       the air VPD by 0.53 to 0.61 kPa, the canopy temperature by 3.1 to 3.8 °C and the  
6       transpiration rate by 0.22 to 0.28 mm h<sup>-1</sup>. These decreases were lower for the  
7       phases *before* and *after* and were greater in drier and warmer days.
- 8       • The duration of the microclimatic changes decreased as the distance from the  
9       centre of the pivot increased, but the duration of the physiological changes was  
10      similar in the different pivot arm portions.
- 11      • Transpiration reduction due to irrigation was higher as closer to the center of the  
12      pivot and represented between 5 to 7% of the applied water. However, the  
13      reduction of ET was similar in the different pivot arm portions and represented  
14      1.5 to 1.8% of the applied water.
- 15      • The decrease in maize canopy temperature could be positive or negative,  
16      depending on its effect on photosynthesis. The reduction of transpiration and ET  
17      must be considered positive because it represents a reduction of irrigation  
18      requirements. Whether the physiological changes will result in increased plant  
19      production should be further studied.

20

## 1 REFERENCES

- 2 Allen, R.G., Pruitt, W.O., Businger, J.A., Fritschen, L.J., Jensen, M.E., and Quinn, F.H.  
3 1996. Evaporation and Transpiration. Hydrology Handbook (Manuals and Reports  
4 on Engineering Practice No. 28). American Society of Civil Engineers (ASCE),  
5 Nueva York, EE.UU. 768 pp.
- 6 Allen, R.G., Pereira, L.S., Raes, D., and Smith, M. 1998. Crop evapotranspiration:  
7 guidelines for computing crop water requirements. FAO Irrigation and Drainage  
8 Paper 56, FAO, Rome, Italy.
- 9 Allen, R.G., Keller, J., and Martin, D. 2000. Center Pivot System Design. The Irrigation  
10 Association VA, USA. 300 p.
- 11 Baker, J.M., and Van Bavel, C.H.M. 1987. Measurement of mass flow of water in the  
12 stems of herbaceous plants. *Plant Cell & Environment*, 10: 777-782.
- 13 Cavero, J., Zaragoza, C., Suso, M.L., and Pardo, A. 1999. Competition between maize  
14 and *Datura stramonium* in an irrigated field under semi-arid conditions. *Weed*  
15 *Research*, 39: 225-240.
- 16 Cavero, J., Zaragoza, C., Bastiaans, L., Suso, M.L., and Pardo, A. 2000. The relevance  
17 of morphological plasticity in the simulation of competition between maize and  
18 *Datura stramonium*. *Weed Research*, 40: 163-180.
- 19 Cavero, J., Medina, E.T., Puig, M., and Martinez-Cob, A. 2009. Sprinkler Irrigation  
20 Changes Maize Canopy Microclimate and Crop Water Status, Transpiration, and  
21 Temperature. *Agronomy Journal*, 101: 854-864.

- 1 Evans, R.G. 2001. Center Pivot Irrigation; Research Report; USDA-Agricultural  
2 Research Service: Sidney, MT, USA.
- 3 Farahani, H.J., and Bausch, W.C.1995. Performance of evapotranspiration models for  
4 maize-bare soil to closed canopy. Transactions of the ASAE, 38: 1049-1059.
- 5 Frost, K.R., and Schwalen, H.C. 1960. Evapotranspiration during sprinkler irrigation.  
6 Transactions of the ASAE, 3: 18-20, 24.
- 7 Howell, T.A., Hiler, E.A., and Van Bavel, C.H.M. 1971. Crop response to mist irrigation.  
8 Transactions of the ASAE, 14: 906-910.
- 9 Kjelgaard, J.F., Stockle, C.O., Villar, J.M., Evans, R.G., and Campbell, G.S. 1994.  
10 Evaluating methods to estimate corn evapotranspiration from short-time interval  
11 weather data. Transactions of the ASAE, 37: 1825-1833.
- 12 Liu, H.J., and Kang, Y.H. 2006a. Effect of sprinkler irrigation on microclimate in the  
13 winter wheat field in the North China Plain. Agricultural Water Management, 84: 3-  
14 19.
- 15 Liu, H.J., and Kang, Y.H. 2006b. Regulating field microclimate using sprinkler misting  
16 under hot-dry windy conditions. Biosystems Engineering, 95: 349-358.
- 17 MAGRAMA 2011. Encuesta de superficies y rendimientos de cultivos en España  
18 (ESYRCE). Ministerio de Agricultura, Alimentación y Medio Ambiente.
- 19 MARM 2011. Sistema de Información Agroclimática para el Regadío (SIAR).  
20 [http://www.marm.es/es/agua/temas/observatorio-del-regadio-espanol/sistema-de-  
21 informacion-agroclimatica-para-el-regadio/](http://www.marm.es/es/agua/temas/observatorio-del-regadio-espanol/sistema-de-<br/>21 informacion-agroclimatica-para-el-regadio/). Spanish Ministry of Natural, Rural and  
22 Marine Environment (MARM). Data were downloaded on January 2011.

- 1 Martinez-Cob, A., Playan, E., Zapata, N., Caverro, J., Medina, E.T., and Puig, M. 2008.  
2 Contribution of evapotranspiration reduction during sprinkler irrigation to application  
3 efficiency. *Journal of Irrigation and Drainage Engineering*, 134: 745-756.
- 4 Monteith, J.L., and Unsworth, M.H. 2008. *Principles of environmental physics*, 3rd edn.  
5 Burlington, MA, USA.
- 6 Norman, J. M., and Campbell, G. 1983. Application of a plant-environment model to  
7 problems in irrigation. In: D. Hillel, (Ed.), *Advances in Irrigation*. Academic Press,  
8 New York, vol. 2, pp. 155-188.
- 9 Norman, E., Joyce, R., and Whittaker, M. 1990. *Advanced design and technology*. 3rd  
10 ed. Longman, Harlow, Essex.
- 11 Ray, J.D., Gesch, R.W., Sinclair, T.R., and Allen, L.H. 2002. The effect of vapour  
12 pressure deficit on maize transpiration response to a drying soil. *Plant Soil*, 238:  
13 113–121.
- 14 Robinson, F.E. 1970. Modifying an arid microclimate with sprinklers. *Agricultural*  
15 *Engineering*, 51: 465.
- 16 Saadia, R., Huber, L., and Lacroix , B. 1996. Using evaporative cooling to fight heat  
17 stress in corn. The potential of sprinkler irrigation to reduce air and reproductive  
18 organ temperature. *Agronomie*, 16: 465-477.
- 19 Steiner, J.L., Kanemasu, E.T., and Hasza, D. 1983. Microclimatic and crop responses to  
20 center pivot sprinkler and to surface irrigation. *Irrigation Science*, 4: 201-214.

- 1 Thompson, A.L., Gilley, J.R., and Norman, J.M. 1993. A sprinkler water droplet  
2 evaporation and plant canopy model: II. Model application. Transactions of the  
3 ASAE, 36: 743-750.
- 4 Tolk, J.A., Howell, T.A., Steiner, J.L., Krieg, D.R., and Schneider, A.D. 1995. Role of  
5 transpiration suppression by evaporation of intercepted water in improving irrigation  
6 efficiency. Irrigation Science, 16: 89-95.
- 7 USDA 2008. Farm and Ranch Irrigation Survey (2008), Vol. 3, Special Studies-Part 1 of  
8 the 2007 Census of Agriculture, AC-07-SS-1, National Agricultural Statistics Service,  
9 (Updated July 2010).
- 10 Van Bavel, M.G. 2005. Flow4 Installation and Operation Manual, Dynamax Inc.,  
11 Houston, Texas, 191 pp.
- 12 Wanjura, D.F., and Upchurch, D.R. 2000. Canopy temperature characterizations of corn  
13 and cotton water status. Transactions of the ASAE, 43:867-875.
- 14 Weibel, F.P., and Boersma, K. 1995. An improved stem heat-balance method using  
15 analog heat control. Agricultural and Forest Meteorology, 75: 191-208.
- 16 Yu, G.R., Kobayashi, T., Zhuang, J., Wang, Q.F., and Qu, L.Q. 2003. A coupled model  
17 of photosynthesis-transpiration based on the stomatal behaviour for maize (*Zea*  
18 *mays* L.) grown in the field. Plant Soil, 249: 401–416.
- 19 Zhao, W., Li, J., and Li, Y. 2012. Modeling sprinkler efficiency with consideration of  
20 microclimate modification effects. Agricultural Water Management, 161: 116-122.

21

22

1 Table 1. Main characteristics of the center pivot system.

Pivot arm portion	Distance from centre pivot	Number of sprinklers	Nozzle diameters	Spacing between sprinklers
	m		mm	m
1	0 – 48	5	2.8 - 4.8	9.3
2	48 – 98	7	4.8 - 5.4	7.0
3	98 – 147	8	4.2 - 5.8	6.2
4	147 – 197	8	4.8 - 6.0	6.2
5	197 – 246	16	4.2 - 6.0	3.1
6	246 – 295	16	4.4 - 5.6	3.1
Overhang	295 – 321	8	4.6 - 5.4	2.9

2

3

1 Table 2. Mean characteristics of the *transpiration-measured* irrigations events.

Pivot arm section	Date	Starting time <sup>a</sup>	Duration	Pressure		Applied water depth
				Mean	CV <sup>b</sup>	
		----- h -----		KPa	%	mm
2	24 July	0825	1.70	208		14.6
	31 July	0935	1.62	218	0.8	14.9
	6 August	0900	1.62	214	0.5	14.8
	13 August	0905	1.55	216	0.6	14.2
	21 August	0920	1.55	198	1.0	13.6
	28 August	0945	1.62	207	1.0	14.5
	10 September	1015	1.62	208	3.3	14.6
4	24 July	0840	0.62	193	2.2	13.6
	31 July	0955	0.60	197	1.6	13.8
	6 August	0915	0.60	194	1.9	13.6
	13 August	0920	0.57	197	1.0	13.1
	21 August	0935	0.57	177	2.4	12.4
	28 August	1000	0.60	181	1.9	13.2
	10 September	1030	0.60	194	1.6	13.7
5	24 July	0840	0.50	195	0.3	14.4
	31 July	0955	0.48	199	1.3	14.5
	6 August	0920	0.48	197	1.5	14.5
	13 August	0920	0.45	200	0.7	13.9
	21 August	0940	0.45	174	0.8	13.0
	28 August	1000	0.48	179	0.9	13.8
	10 September	1030	0.48	202	0.6	14.7

2 <sup>a</sup> Greenwich Mean Time.

3 <sup>b</sup> CV, coefficient of variation.

4

1 Table 3. Meteorological conditions during each *transpiration-measured* irrigation event,  
 2 recorded at the 'nearby grass station' of Valfarta<sup>a</sup>.

Date	Air temperature °C	Air vapor pressure deficit kPa	Wind speed m s <sup>-1</sup>	Solar radiation W m <sup>-2</sup>
24 July	28.0	1.9	2.0	742
31 July	32.5	3.6	1.5	880
6 August	29.9	2.6	1.8	772
13 August	22.8	1.7	3.5	723
21 August	27.4	1.7	2.0	780
28 August	26.6	1.6	0.9	768
10 September	27.3	1.8	1.6	651

3 <sup>a</sup> Station included in the network SIAR (Spanish Irrigation Advisory System) (MARM  
 4 2011).

5

1 Table 4. Average values of air temperature recorded in the *moist* ( $T_{MT}$ ) and *dry* ( $T_{DT}$ )  
 2 treatments before, during and after the *transpiration-measured* irrigation events at the  
 3 pivot arm portions 2, 4 and 5.

Phase	Pivot arm portion	N <sup>a</sup>	Air Temperature				Duration	
			Mean $T_{MT}$	SD <sup>b</sup>	Mean $T_{DT}$	SD	Mean	CV <sup>c</sup>
			-----( <sup>o</sup> C) -----				h	%
Before	2	6	23.8 b <sup>d</sup>	± 1.7	24.4 a	± 1.9	0.8	20
	4	4	23.4 b	± 2.4	23.9 a	± 2.5	0.6	54
	5	4	24.4 b	± 2.0	25.1 a	± 2.1	0.7	43
During	2	7	23.5 b	± 1.9	25.6 a	± 2.4	1.6	3
	4	7	23.4 b	± 1.9	25.2 a	± 2.3	0.6	11
	5	7	23.2 b	± 1.6	25.2 a	± 2.3	0.5	10
After	2	7	27.9 b	± 2.5	28.7 a	± 2.3	1.5	21
	4	7	26.3 b	± 3.2	27.4 a	± 2.8	1.3	31
	5	7	25.1 b	± 2.5	26.4 a	± 2.3	1.0	29

4 <sup>a</sup> Number of *transpiration-measured* irrigation events.

5 <sup>b</sup> Standard deviation.

6 <sup>c</sup> Coefficient of variation.

7 <sup>d</sup> For each phase and pivot arm portion the air temperature values marked with different  
 8 letters indicate that they were significantly different after a paired *t*-test ( $P = 0.05$ ).

9

1 Table 5. Average values of air vapor pressure deficit recorded in the *moist* (T<sub>MT</sub>) and *dry*  
 2 (T<sub>DT</sub>) treatments before, during and after the *transpiration-measured* irrigation events at  
 3 the pivot arm portions 2, 4 and 5.

Phase	Pivot arm portion	N <sup>a</sup>	Vapor pressure deficit				Duration	
			Mean T <sub>MT</sub>	SD <sup>b</sup>	Mean T <sub>DT</sub>	SD	Mean	CV <sup>c</sup>
			------(KPa) -----				h	%
Before	2	6	0.92 b <sup>d</sup>	± 0.37	1.11 a	± 0.45	0.7	43
	4	4	0.97 b	± 0.46	1.13 a	± 0.51	0.6	54
	5	4	1.01 b	± 0.40	1.26 a	± 0.48	0.7	43
During	2	7	0.87 b	± 0.25	1.40 a	± 0.44	1.6	3
	4	7	0.80 b	± 0.30	1.33 a	± 0.43	0.6	11
	5	7	0.72 b	± 0.20	1.33 a	± 0.44	0.5	10
After	2	7	1.72 b	± 0.51	2.02 a	± 0.51	1.4	21
	4	7	1.39 b	± 0.62	1.76 a	± 0.54	1.3	31
	5	7	1.13 b	± 0.37	1.54 a	± 0.41	1.1	30

4 <sup>a</sup> Number of *transpiration-measured* irrigation events.

5 <sup>b</sup> Standard deviation.

6 <sup>c</sup> Coefficient of variation.

7 <sup>d</sup> For each phase and pivot arm portion the air temperature values marked with different  
 8 letters indicate that they were significantly different after a paired *t*-test (P = 0.05).

9

10

1 Table 6. Average values of air temperature, air vapor pressure deficit (VPD) and canopy  
 2 temperature recorded in the *moist* ( $T_{MT}$ ) and *dry* ( $T_{DT}$ ) treatments along the different  
 3 pivot arm portions (PAP) for the 7 *transpiration-measured* irrigation events and the 27  
 4 *remaining* irrigation events.

Variable	PAP	<i>Transpiration-measured</i> irrigation events				<i>Remaining</i> irrigation events			
		$T_{MT}$	$T_{DT}$	Duration		$T_{MT}$	$T_{DT}$	Duration	
		Mean <sup>a</sup>	CV <sup>b</sup>	Mean	CV	Mean	CV		
		----- °C -----	-----	h	%	----- °C -----	-----	h	%
Air temperature	2	25.4 b <sup>c</sup>	26.7 a	3.8	9	25.0 b	26.6 a	3.0	30
	4	25.1 b	26.2 a	2.3	28	25.2 b	26.8 a	2.4	29
	5	24.5 b	25.7 a	2.0	26	25.1 b	26.5 a	2.4	33
		----- kPa -----	-----			----- kPa -----	-----		
Air VPD	2	1.21 b	1.58 a	3.6	14	1.38 b	1.84 a	3.0	30
	4	1.20 b	1.56 a	2.3	28	1.44 b	1.90 a	2.4	29
	5	1.03 b	1.43 a	2.1	23	1.35 b	1.83 a	2.4	33
		----- °C -----	-----			----- °C -----	-----		
Canopy temperature	2	23.1 b	24.9 a	4.1	30	24.2 b	26.0 a	4.0	22
	4	23.0 b	24.7 a	4.5	36	24.3 b	26.1 a	3.6	21

5 <sup>a</sup>, Mean duration.

6 <sup>b</sup>, Coefficient of variation.

7 <sup>c</sup> For each variable, pivot arm portion and irrigation type event the values marked with  
 8 different letters were significantly different after a paired *t*-test ( $P = 0.05$ ).

1 Table 7. Average values of canopy temperature recorded in the *moist* ( $T_{MT}$ ) and *dry*  
 2 ( $T_{DT}$ ) treatments before, during and after the *transpiration-measured* irrigation events at  
 3 the pivot arm portions 2 and 4.

Phase	Pivot arm portion	N <sup>a</sup>	Canopy Temperature				Duration	
			Mean $T_{MT}$	SD <sup>b</sup>	Mean $T_{DT}$	SD	Mean	CV <sup>c</sup>
			-----( <sup>o</sup> C) -----				h	%
Before	2	7	22.3 b <sup>d</sup>	± 2.7	23.3 a	± 2.8	1.8	64
	4	7	22.0 b	± 2.6	23.2 a	± 2.8	2.1	56
During	2	7	23.3 b	± 2.1	26.4 a	± 2.0	1.6	3
	4	7	22.4 b	± 1.9	26.2 a	± 2.0	0.6	11
After	2	7	26.2 b	± 2.2	27.3 a	± 1.9	0.8	84
	4	7	25.5 b	± 1.9	26.9 a	± 1.7	1.7	44

4 <sup>a</sup> Number of *transpiration-measured* irrigation events.

5 <sup>b</sup> Standard deviation.

6 <sup>c</sup> Coefficient of variation.

7 <sup>d</sup> For each phase and pivot arm portion the canopy temperature values marked with  
 8 different letters indicate that they were significantly different after a paired *t*-test (P =  
 9 0.05).

10

1 Table 8. Average values of transpiration rate recorded in the *moist* ( $T_{MT}$ ) and *dry* ( $T_{DT}$ )  
 2 treatments before, during and after the *transpiration-measured* irrigation events at the  
 3 pivot arm portions (PAP) 2, 4 and 5. The transpiration reduction duration and magnitude  
 4 is also shown.

Phase	PAP	N <sup>a</sup>	Transpiration rate				Duration		Transpiration reduction	
			$T_{MT}$	SD <sup>b</sup>	$T_{DT}$	SD	Mean	CV <sup>c</sup>	Mean	
			----- mm h <sup>-1</sup> -----				h	%	mm	%
Before	2	3	0.48 b <sup>d</sup>	± 0.15	0.63 a	± 0.14	1.0	76	0.15	24
	4	7	0.41 b	± 0.07	0.60 a	± 0.15	0.9	67	0.17	31
	5	3	0.53 b	± 0.17	0.69 a	± 0.18	1.7	60	0.27	24
During	2	7	0.48 b	± 0.08	0.75 a	± 0.15	1.6	3	0.45	36
	4	7	0.51 b	± 0.06	0.73 a	± 0.15	0.6	11	0.15	30
	5	7	0.49 b	± 0.07	0.77 a	± 0.14	0.5	10	0.14	36
After	2	7	0.62 b	± 0.16	0.86 a	± 0.11	1.8	73	0.43	28
	4	7	0.58 b	± 0.15	0.80 a	± 0.14	2.4	48	0.54	28
	5	7	0.68 b	± 0.11	0.82 a	± 0.13	2.4	63	0.34	17
All	2						3.9	43	1.03	30
	4						4.0	35	0.86	29
	5						3.6	67	0.75	22

5 <sup>a</sup> Number of *transpiration-measured* irrigation events.

6 <sup>b</sup> Standard deviation.

7 <sup>c</sup> Coefficient of variation.

8 <sup>d</sup> For each phase and pivot arm portion the transpiration rate values marked with  
 9 different letters indicate that they were significantly different after a paired *t*-test (P =  
 10 0.05).

11

12

13

14

15

16

17

18

19

20

21

22

23

1 Table 9. Average values of evapotranspiration (ET) rate estimated in the *moist* ( $T_{MT}$ )  
 2 and *dry* ( $T_{DT}$ ) treatments before, during and after the *transpiration-measured* irrigation  
 3 events at the pivot arm portions (PAP) 2, 4 and 5. The duration of periods was the same  
 4 as for transpiration. The ET reduction magnitude is also shown.

Phase	PAP	N <sup>a</sup>	ET rate				ET reduction	
			$T_{MT}$	SD <sup>b</sup>	$T_{DT}$	SD	Mean	
			----- mm h <sup>-1</sup> -----				mm	%
Before	2	3	0.56 b <sup>c</sup>	± 0.09	0.60 a	± 0.08	0.04	7.7
	4	7	0.57 b	± 0.07	0.59 a	± 0.07	0.02	3.5
	5	3	0.58 b	± 0.11	0.63 a	± 0.09	0.06	8.0
During	2	7	0.62 b	± 0.11	0.72 a	± 0.12	0.16	13.8
	4	7	0.60 b	± 0.11	0.70 a	± 0.12	0.07	14.9
	5	7	0.58 b	± 0.10	0.70 a	± 0.12	0.06	16.8
After	2	7	0.84 b	± 0.16	0.89 a	± 0.15	0.08	6.2
	4	7	0.76 b	± 0.16	0.82 a	± 0.13	0.14	8.2
	5	7	0.73 b	± 0.14	0.79 a	± 0.13	0.13	7.6
All	2						0.26	9.6
	4						0.22	7.8
	5						0.22	9.3

5 <sup>a</sup> Number of *transpiration-measured* irrigation events.

6 <sup>b</sup> Standard deviation.

7 <sup>c</sup> For each phase and pivot arm portion the ET rate values marked with different letters  
 8 indicate that they were significantly different after a paired *t*-test (P = 0.05).

9


10

11

12


13

Figure\_1  
[Click here to download high resolution image](#)




Figure\_2


[Click here to download high resolution image](#)




Figure\_3  
[Click here to download high resolution image](#)




Figure\_4  
[Click here to download high resolution image](#)



Figure\_5  
[Click here to download high resolution image](#)




Figure\_6  
[Click here to download high resolution image](#)



Figure\_7

[Click here to download high resolution image](#)



1 **FIGURE CAPTIONS**

2 Fig. 1. Overview of the center pivot sprinkler irrigation system at the commercial  
3 plot. A to C, meteorological stations and sap flow measurement systems for  
4 *moist* treatment. D, meteorological station and sap flow measurement  
5 system for *dry* treatment. Ap, pivot arm. Pr, irrigation pressure transducers.  
6 R1-6, radius at the end of each pivot arm portion. M, direction of center pivot  
7 movement.

8 Fig. 2. 5-min averages of microclimatic variables (temperature and vapour  
9 pressure deficit of the air) and physiological variables (canopy temperature  
10 and transpiration rate) monitored at the different pivot arm portions on 6  
11 August, 2008 since 2 h before until 6 h after the irrigation event. MT, *moist*  
12 treatment. DT, *dry* treatment. The vertical solid lines indicate the start and  
13 the end of the irrigation event over the transect AC. The vertical dashed  
14 lines indicate the period during which the monitored variables were different  
15 between the two treatments *before* (B), *during* (D) and *after* (A) irrigation  
16 event.

17 Fig. 3. Relationship between the decrease in air temperature due to sprinkler  
18 irrigation and the air temperature measured over grass at a nearby weather  
19 station. PAP, Pivot Arm Portions. A, the Y axis represents the decrease in  
20 air temperature for the phase *during* at the 7 *transpiration-measured*  
21 irrigation events. B, the Y axis represents the decrease in air temperature  
22 observed for the whole duration of the 27 *remaining* irrigation events.

23 Fig. 4. Relationship between the decrease in vapour pressure deficit (VPD) due  
24 to sprinkler irrigation and the VPD measured over grass at a nearby

1 standard weather station. PAP, Pivot Arm Portions. A, the Y axis represents  
2 the decrease in VPD for the phase *during* at the 7 *transpiration-measured*  
3 irrigation events. B, the Y axis represents the decrease in VPD observed for  
4 the whole duration of the 27 *remaining* irrigation events.

5 Fig. 5. Relationship between the maize transpiration reduction for the different  
6 phases (before, during and after) and the reduction of air temperature, air  
7 VPD and canopy temperature at the 7 *transpiration-measured* center pivot  
8 irrigation events.

9 Fig. 6. Relationship between the maize transpiration reduction for the phase  
10 *during* at the 7 *transpiration-measured* irrigation events at the different pivot  
11 arm portions (PAP), and the vapor pressure deficit (VPD) of the air  
12 measured at a nearby standard weather station.

13 Fig. 7. Relationship between the maize canopy temperature reduction for the  
14 phase *during* at the 7 *transpiration-measured* irrigation events at the  
15 different pivot arm portions (PAP), and the air temperature measured at a  
16 nearby standard weather station.

17