

## Determining evapotranspiration in an olive orchard in southwest Spain

J.M. Torres-Ruíz, J.E. Fernández, I.F. Girón.  
Instituto de Recursos Naturales y Agrobiología  
IRNASE-CSIC  
Apartado 1052, 41080-Sevilla  
Spain  
M.J. Martín-Palomo  
Departamento de Ciencias Agroforestales  
Universidad de Sevilla  
41013-Sevilla, Spain

R. Romero, J.A. Jiménez-Bocanegra,  
I. García Tejero  
IFAPA, Centro Las Torres-Tomejil  
Ctra. Sevilla-Cazalla, km 12.2  
41200-Alcalá del Río, Sevilla, Spain

**Keywords:** *Olea europaea*, evapotranspiration, irrigation, transpiration, modeling

### Abstract

The aim of this work was to evaluate, for an olive orchard in the Aljarafe county, the method developed by Orgaz et al. (2005) for determining the crop evapotranspiration ( $ET_c$ ). We compared the calculated  $ET_c$  ( $ET_{c\ Orgaz}$ ) values with those determined by the crop coefficient approach ( $ET_{c\ crop\ coef}$ ), as described by Fernández et al. (2006), who used coefficient values previously calibrated for our orchard conditions. In addition, we compared the tree transpiration ( $E_p$ ) values estimated with the mentioned Excel application ( $E_{p\ Orgaz}$ ) with those simulated by a transpiration model ( $E_{p\ sim}$ ) based on Penman-Monteith, validated for our orchard conditions. Results showed that the Excel application is a user-friendly tool valid for calculating reasonably accurate values of  $ET_c$  from very few easy-to-measure inputs. The crop coefficient approach does not have this limitation, but years with unusual leaf area density may lead to errors on the calculated  $ET_c$ .  $E_{p\ sim}$  is highly affected by variables difficult to measure in commercial orchards, such as the leaf area and the available soil water. In addition, processes related to leaf aging, soil temperature and recovery after drought are not included yet in the model, which affects the reliability of the  $E_{p\ sim}$  values at the end of the irrigation.

### INTRODUCTION

A correct irrigation management involves a precise determination of the crop water needs. For olive orchards, as well as for orchards of many other fruit tree species, the water crop coefficient method (Allen et al., 1998) has been widely used with that purpose (Fernández and Moreno, 1999). Still, the method has limitations derived from its empirical character, which makes difficult to extrapolate the crop coefficient ( $K_c$ ) values to orchards different from those in which the existing values have been obtained. Recently, Orgaz et al. (2005) described a more mechanistic method for calculating  $K_c$  in any type of olive orchard, based on basic features easy to quantify in commercial orchards. Later, Testi et al. (2006) and Orgaz et al. (2006) described with detail this approach, and included a new component in the calculation of  $K_c$ , accounting for the evaporation of the water intercepted by the canopy. The Orgaz et al. (2005) method was used to develop a user-friendly Excel application intended to calculate crop water requirements in commercial orchards, which was delivered with the book by Pastor et al. (2005). Our aim was to evaluate the method by Orgaz et al. (2005) in an olive orchard representative of the commercial orchards in the Aljarafe county, the main production

area for ‘Manzanilla de Sevilla’. We compared the  $ET_c$  Orgaz values with the  $ET_c$  crop coef values. Results from the soil water balance carried in the same orchard by Palomo et al (2002) show that the  $K_c$  values suggested by Fernández et al. (2006) are reasonably precise for our orchard conditions. In addition, we compared the  $E_p$  Orgaz values with the  $E_p$  sim values.

## MATERIALS AND METHODS

### Orchard characteristics

The experiments were made at ‘La Hampa’, the experimental farm of the *Instituto de Recursos Naturales y Agrobiología* (IRNASE-CSIC). The farm is located in the Aljarafe county, close to Coria del Río, at 15 km to the southwest of Seville (37° 17’ N, 6° 3’ W, 30 m a.s.l.). The olive orchard was planted in 1969 with ‘Manzanilla de Sevilla’ olive trees at 7 m × 5 m spacing. The orchard is representative of those in the area. The trees have a single trunk with two main branches from 0.7 to 1.5 m above ground. The effective depth of the root system is about 0.9 m (Fernández et al., 1991). The canopy is spherical and open at the top. Average values of canopy volume and leaf area density (LAD) at the end of the growing season were 36 m<sup>3</sup> and 1.6 m<sup>2</sup> m<sup>-3</sup>, respectively. The ground covered by the canopies was about 34%. Herbicides were used during the irrigation season to keep the soil free of weeds.

The soil is a sandy loam of about 1.6-2.0 m depth, depending on the location. The texture is quite homogeneous, with average values of 14.8% clay, 7.0% silt, 4.7% fine sand and 73.5% coarse sand. The volumetric soil water contents ( $\theta$ , cm<sup>3</sup> cm<sup>-3</sup>) at 0 MPa and -1.5 MPa are 0.33 cm<sup>3</sup> cm<sup>-3</sup> and 0.10 cm<sup>3</sup> cm<sup>-3</sup>, respectively. Field  $\theta$  values for field capacity are about 0.21 cm<sup>3</sup> cm<sup>-3</sup>. The climate is typically Mediterranean, with a mild, wet season from October to April. The rest of the year is hot and dry. The average precipitation ( $P$ ) and  $ET_o$  values in the area are 499 mm and 1165 mm, respectively (period 1971-2007). Main weather records for the irrigation season of 2007, the experimental period, are shown in Figure 1. The experimental orchard was irrigated daily to replace the crop water needs, with a lateral per row with five 3 L hour<sup>-1</sup> drippers per tree, 1 m apart. The irrigation season went from May 14 to October 2.

### Calculation of $ET_c$

We used the crop coefficient method (Allen et al., 1998) to determine daily values of  $ET_c$  as  $ET_c = K_c K_r ET_o$ . We used the  $K_c$  values determined by Fernández et al. (2006) for the orchard conditions (0.76 in May, 0.70 in June, 0.63 in July and August, 0.72 in September and 0.77 in October). The value of the coefficient related to the percentage of ground covered by the crop ( $K_r$ ), calculated after Fereres and Castel (1981), was 0.71. We used the REF-ET software (Allen 2000) and the records of the weather station next to the orchard to calculate daily  $ET_o$  values with the FAO56 Penman-Monteith equation (Allen et al., 1998).

We also calculated the  $ET_c$  values with the method described by Orgaz et al. (2005). More precisely, we used the Excel application delivered with the book by Pastor (2005). With this tool, monthly values of  $ET_c$ ,  $E_p$  and soil evaporation ( $E_s$ ), both from the dry and wetted ground areas, are easily determined. The required inputs are related to the irrigation system and management, the weather conditions and the tree size and spacing, as described above.

### Transpiration model

We used the Penman-Monteith equation to model daily  $E_p$  values in the orchard, for the whole irrigation season. We followed the approach described by Moreno et al. (1996):

$$\lambda E_{p\ sim} = f_1 \frac{sR_{n,l}}{s + \gamma(2 + g_{b,l}/g_{c,l})} + f_1 \frac{\rho c_p D_a g_{b,l}}{s + \gamma(2 + g_{b,l}/g_{c,l})} + f_s \frac{\rho c_p D_a g_{b,s}}{s + \gamma(2 + g_{b,s}/g_{c,s})} ,$$

where  $E_{p\ sim}$  is in  $s^{-1} m^{-2}$  leaf plan area;  $f_1 = A_l/A$  and  $f_s = A_s/A$ , being  $A$  ( $m^2$ ) the total (one-side) leaf area. This comprises the area of sunlit leaves ( $A_l$ ) and that of shade leaves ( $A_s$ ). The values of  $A_l$  and  $A_s$  were estimated with the RATP model (Sinoquet et al., 2001), which calculates the radiation transfer through the canopy (Diaz-Espejo et al., 2002);  $R_{n,l}$  ( $W m^{-2}$  of leaf plan area) is the net, all-wave radiation of the lit leaves (assumed to be zero for the shade leaves);  $D_a$  (Pa) is the vapour pressure deficit of the air;  $\lambda$  is the latent heat of vapourisation ( $2.454 J kg^{-1}$ );  $g_c$  is the leaf stomatal conductance ( $m s^{-1}$ ) and  $g_b$  is the leaf-canopy boundary-layer conductance ( $m s^{-1}$ ); the values of  $g_c$  were calculated as  $g_c = g_m f(PPF) f(T_l) f(D_a) f(\theta)$ , where PPF is the incident photosynthetically active photon flux,  $T_l$  is the leaf temperature and  $g_m$  is the reference stomatal conductance measured under standard conditions ( $PPF = 1600 \mu mol m^{-2} s^{-1}$ ,  $T_l = 25 \text{ }^\circ C$ ,  $D_a = 1 \text{ kPa}$ ,  $\theta$  at field capacity =  $0.21 m^3 m^{-3}$ ). The  $f$  functions, described in Diaz-Espejo et al. (2006), were validated for our orchard conditions.

## Measurements

The volumetric soil water contents ( $\theta$ ) in the soil around three representative trees were measured every 7-10 days with a Profile probe, at 0.1, 0.2, 0.3, 0.4, 0.6 and 1.0 m depths, and at 1, 2, and 3 m from the tree trunk. From the  $\theta$  values we calculated the relative extractable water (REW) of the soil as  $REW = (R - R_{min})/(R_{max} - R_{min})$ , being  $R$  (mm) the actual soil water content,  $R_{min}$  (mm) the minimum soil water content measured during the experiments, and  $R_{max}$  (mm) the soil water content at field capacity (Granier, 1987). The leaf area (LA,  $m^2$  one side) of the three trees in which  $\theta$  was measured was estimated as described by Fernández et al. (2006). Half-hour values of the main meteorological variables were continuously recorded by an automatic weather station next to the olive orchard.

## RESULTS AND DISCUSSION

Most of the 2007 irrigation season was dry and hot. The first significant precipitation amounts were recorded from mid September (Fig. 1), as usual in the area. The weather records showed a yearly value of  $P$  (411,1 mm) below average, and a yearly value of  $ET_o$  (1235.0 mm) above average.

The irrigation volumes (IA, mm) were enough to keep soil water contents close to field capacity for most of the irrigation season (Fig. 2). This surely contributed to the increase in leaf area recorded during the growing season, which extended until late August (Fig. 3). Later we observed a slight decrease, caused by the oldest leaves falling from the trees. The LA values shown in Fig. 3 are similar to those recorded by Palomo et al. (2002) during the experiments in which they evaluated the crop coefficient method for our orchard conditions.

The time course of  $ET_o$  for the whole irrigation season,  $ET_{c\ crop\ coef.}$  and  $ET_{c\ Orgaz}$ , are shown in Fig. 4. The figure also shows the  $E_{p\ sim}$ , as well as  $E_{p\ Orgaz}$ . For the whole irrigation season,  $ET_{c\ crop\ coef.}$  and  $ET_{c\ Orgaz}$  amounted to  $3413 m^3 ha^{-1}$  and  $3839 m^3 ha^{-1}$ ,

respectively. Differences were greater in July and August, in which the monthly  $ET_{c \text{ Orgaz}}$  values were 21% and 18% greater, respectively, than those of  $ET_{c \text{ crop coef.}}$ . The REW values (Fig. 2) recorded on those months suggest that the IA estimated with the crop coefficient method were reasonably correct. Therefore, the  $ET_{c \text{ Orgaz}}$  values for July and August can be too high for the orchard conditions. Pruning practices in the Aljarafe, that led to trees with low leaf area density, may account for this. The dynamics of  $ET_{c \text{ crop coef.}}$  echoed that of  $ET_o$ , as expected, showing that the crop coefficient method can be useful to control high frequency irrigation. The  $ET_{c \text{ Orgaz}}$  values varied, from month to month, according to  $ET_o$ , showing also their utility for controlling irrigation, although with a lower resolution. The  $E_{p \text{ sim}}$  values also echoed  $ET_o$ , although not always were lower than  $ET_{c \text{ crop coef.}}$ . This was especially evident late in the season (Fig. 4, bottom graph). We expected a lower precision of the model at that time of the year, since some aspects related to leaf aging, leaf gas exchange recovery after drought, or the influence of soil temperature in  $g_c$ , are not considered in the current version of the model. In addition, some disagreement between actual and  $E_{p \text{ sim}}$  values must be expected, because the transpiration model is sensitive to variables difficult to estimate precisely. This is the case of  $\theta$  and LA. Thus, the calculation of the available water it is diffculted by the high spatio-temporal variability both of  $\theta$  and root density; and, for LA, our experience shows that errors may easily amount to  $\pm 20\%$  or more. Concerning the  $E_{p \text{ Orgaz}}$  values calculated in July and August, two dry months, those amounted to 78% of  $ET_{c \text{ Orgaz}}$ , being the remaining 22% the  $E_s$ . For mid September to mid October, a wet period (Fig. 1), the estimated  $E_{p \text{ Orgaz}}$  amounted to 56% of  $ET_{c \text{ Orgaz}}$  ( $E_s = 44\%$  of  $ET_c$ , since this approach does not takes into account the water intercepted by the canopy). Testi et al. (2006) estimated, for an intensive olive orchard of similar characteristics, the annual value of  $E_s$  amounted to 35% of  $ET_c$ .

## CONCLUSIONS

For the Aljarafe conditions, where the trees are severely pruned, the  $ET_{c \text{ Orgaz}}$  values could be too high, especially at the summer months. Using properly evaluated  $K_c$  values for the orchard conditions is not enough to get accurate estimations of  $ET_c$  when using the crop coefficient approach: among other factors, variations in LA from year to year, caused by differences on the pruning intensity or the impact of fungal diseases, may lead to erroneous  $ET_c$  estimations. Cautions taken, the approach may provide reliable  $ET_c$  values with the required resolution for high frequency irrigation. The tested transpiration model is valid for research purposes, but it has limitations for determining  $E_p$  in commercial orchards.

## ACKNOWLEDGEMENTS

This work has been funded by the Spanish Ministry of Education and Science, research project No.AGL2006-04666/AGR, and by the EU, research project ref. STREP 023120.

## Literature Cited

- Allen, R. 2002. REF-ET Reference Evapotranspiration Software. Kimberly Research and Extension Center. University of Idaho. <http://www.kimberly.uidaho.edu/ref-et/>.
- Allen, R., Pereira, L.S., Raes, D. and Smith, M. 1998. Crop evapotranspiration. Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56. FAO, Rome, Italy.
- Díaz-Espejo, A., Hafidi, B., Fernández, J.E., Palomo, M.J. and Sinoquet, H. 2002. Transpiration and photosynthesis of the olive tree: a model approach. *Acta Hort* 586:457-460.
- Díaz-Espejo, A., Walcroft, A., Fernández, J.E., Hafidi, B., Palomo, M.J. and Girón I.F. 2006. Modelling photosynthesis in olive leaves under drought conditions. *Tree Physiol* 26: 1445-1456.
- Fernández, J.E., Díaz-Espejo, A., Infante, J.M., Durán, P., Palomo, M.J., Chamorro, V., Girón, I.F. and Villagarcía, L. 2006a. Water relations and gas exchange in olive trees under regulated deficit irrigation and partial rootzone drying. *Plant Soil* 284: 273-291.
- Fernández, J.E. and Moreno, F. 1999. Water use by the olive tree. *J Crop Prod.* 2: 101-162.
- Fernández, J.E., Moreno, F., Cabrera, F., Arrue, J.L., and Martín-Aranda, J. 1991. Drip irrigation, soil characteristics and the root distribution and root activity of olive trees. *Plant Soil*, 133: 239-251.
- Fereres, E. and Castel, J.R. 1981. Drip Irrigation Management. Division of Agricultural Sciences, University of California, Leaflet 21259.
- Granier, A. 1987. Evaluation of transpiration in a Douglas-fir stand by means of sap flow measurements. *Tree Physiol.* 3: 309-320.
- Moreno, F., Fernández, J.E., Clothier, B.E. and Green, S.R. 1996. Transpiration and root water uptake by olive trees. *Plant Soil* 184: 85-96.
- Orgaz, F., Testi, L., Villalobos, F.J. and Fereres, E. 2006. Water requirements of olive orchards-II: determination of crop coefficients for irrigation scheduling. *Irrig Sci.* 24: 77-84.
- Orgaz, F., Villalobos, F.J., Testi, L., Pastor, M., Hidalgo, J.C. and Fereres, E. 2005. Programación de riegos en plantaciones de olivar. Metodología para el cálculo de las necesidades de agua de riego en el olivar regado por goteo. In: *Cultivo del Olivo con Riego Localizado*. M. Pastor (Ed.). Coedición Junta de Andalucía, Consejería de Agricultura y Pesca, y Ediciones Mundi-Prensa, pp 83-137.
- Palomo, M.J., Moreno, F., Fernández, J.E., Díaz-Espejo, A. and Girón, I.F. 2002. Determining water consumption in olive orchards using the water balance approach. *Agric. Water Manage.* 55: 15-35.
- Pastor, M. (Ed.) 2005. *Cultivo del Olivo con Riego Localizado*. Coedición Junta de Andalucía, Consejería de Agricultura y Pesca, y Ediciones Mundi-Prensa, 783 pp.
- Sinoquet, H., Le Roux, X., Adam, B., Ameglio, T. and Daudet, F.A. 2001. RATP: a model for simulating the spatial distribution of radiation absorption, transpiration and photosynthesis within canopies: application to an isolated tree crown. *Plant Cell Environ* 24:395-406.
- Testi, L., Villalobos, F.J., Orgaz, F. and Fereres, E. 2006. Water requirements of olive orchards: I simulation of daily evapotranspiration for scenario analysis. *Irrig Sci* 24: 69-76.

## Figures




Fig. 1. Weather records collected during the experimental period by the water station next to the experimental olive orchard.  $R_s$  = solar global radiation,  $T_a$  = temperature of the air;  $\text{RH}_a$  = relative humidity of the air;  $P$  = precipitation amounts; DOY = day of year.




Fig. 2. Irrigation amounts (IA) supplied to the olive orchard during the experimental period. These amounts were calculated using the crop coefficient method by Allen et al. (1998) (see text for details). Also shown are the precipitation ( $P$ ) amounts recorded by the weather station next to the orchard, as well as the resulting relative extractable water (REW) values calculated from the soil water contents recorded in the orchard. DOY = day of year.




Fig. 3. Leaf area (LA) values recorded during the experimental period (mean  $\pm$  SE,  $n = 2$ ). DOY = day of year.




Fig. 4. Time courses of the potential evapotranspiration ( $ET_o$  FAO56 P-M) and crop evapotranspiration calculated both by the crop coefficient method ( $ET_c$  crop coef.) and after Orgaz et al. (2005) ( $ET_c$  Orgaz). Also show are the daily tree transpiration values simulated with our model ( $E_p$  sim) and estimated by the Orgaz et al. (2005) approach ( $E_p$  Orgaz). See text for details.