

1 **Chemical variations in the biostructures produced by soil ecosystem engineers –**
2 **Examples from the Neotropical savannas**

3

4 Juan J. Jiménez^{1,*}, Thibaud Decaëns²

5

6 ¹ *Laboratoire d'Ecologie des Sols Tropicaux, IRD, 32 Av. H. Varagnat, F-93143 Bondy,*
7 *France*

8 ² *Laboratoire d'Ecologie. UPRES-EA 1293. UFR Sciences. Université de Rouen. F-76821*
9 *Mont Saint Aignan Cedex, France*

10

11 ¹ Address during the study

12 * Corresponding author (current address):

13 Carbon Management and Sequestration Center

14 School of Environment and Natural Resources

15 The Ohio State University

16 2021 Coffey Road

17 Columbus, OH-43210

18 USA

19 Tel: +614 292 2298

20 Fax: +614 292 7432

21 *E-mail address:* jimenez.58@osu.edu

22

23 XIVth International Colloquium on Soil Zoology and Ecology

24 Session 4 – *Ecosystem engineering by soil fauna* / Oral contribution

25 **Abstract**

26

27 “Ecosystem engineers” are those organisms capable to modify physically the environment
28 by producing “biogenic” structures (BS). Termites, earthworms, ants and other large
29 macroinvertebrates produce BS with varying properties. In this study, our objective was to
30 quantify total C_{org} ($\text{g kg dry soil}^{-1}$), NH_4^+ and NO_3^- concentrations ($\mu\text{g g dry soil}^{-1}$) in
31 different parts of the BS produced by termites and ants to test the hypothesis that higher
32 concentrations are found where new building material is deposited, i.e. at the top of the BS.
33 The study was carried out in a natural savanna (NS), an introduced grass-legume pasture
34 (IP) and a gallery forest (GF) at the Carimagua Research Station in the Eastern Plains of
35 Colombia. Progressive sampling distances across the BS were used, i.e. from the top to the
36 base of the BS by using proportional distances, i.e. 20-100% for large BS and 50-100% for
37 the smallest BS, and these were compared with two types of control soil, 120% or 150% in
38 the case of large and small BS, respectively, and soil sampled 1 m away from the BS. All
39 the BS analysed had, in general, higher concentrations of nutrients than the control soil.
40 There were differences in the variables measured in the BS according to the organism that
41 produced them. The lowest values of C_{org} were observed in the BS (surface dumps)
42 deposited by fungus-growing ants (*Trachymyrmex* sp. in the NS, and *Atta laevigata* in the
43 GF), while the highest concentrations were found in the BS produced by termites in the GF,
44 where a high N concentration was also observed. Nutrient concentrations were higher in
45 general in the BS than in the control soil in all cases. However, other BS seemed not to
46 have any influence in the surrounding soil. We concluded that the activity of soil ecosystem

47 engineers increased the spatial variability of chemical parameters measured in this study.

48 The ecological significance of these differences is discussed.

49

50 *Keywords:* Soil macrofauna / Ecosystem engineers / Soil ecology / Termites / Ants /

51 Savanna

52

53 **1. Introduction**

54 By definition “ecosystem engineers” or “ecological engineers” (*sensu* [12,25])
55 are those organisms that modify physically the environment in which they live. The
56 engineer organisms do so by producing “biogenic” structures or biostructures (BS)
57 that impact in some soil processes and affect the spatial and trophic resources for one,
58 or generally more, organisms. The peculiarity of the BS is that such processes
59 continue in the absence of the organisms that created them [12,1,14]. As a result the
60 abundance and community structure of other organisms are modified without
61 establishing any direct trophic relationship [12,13].

62 Ants and termites are important regulators of soil aggregate structure as they
63 remove (ants) or ingest (termites) large amounts of soil that can be either remove it
64 from the bottom to the top soil (fungus growing ants) or egest it above or in the soil
65 profile (termites). In doing so, they form BS with constituting aggregates of different
66 sizes and characteristics, i.e. ant hills, termite mounds. These BS have varying
67 characteristics according to the species and the soil where they carry their activities
68 [17,5]. These various effects upon habitat structure are part of the numerous sources

69 of soil ecosystem heterogeneity and hence may affect soil biota diversity with
70 important functional consequences [6].

71 The BS may sometimes cover a large proportion of the soil surface. Nonetheless, there
72 is a lack of studies dealing with a description and characterization of the BS produced by
73 these invertebrates. Their description can be used to establish a functional classification of
74 these organisms in order to assess their contribution to soil processes and ecosystem
75 function. The morphology, size, abundance and physico-chemical properties of BS are a
76 previous step to evaluate their indirect effects' type and wideness in the surrounding
77 environment at a given scale [12,13]. Thus, it is necessary to describe the dynamics and
78 phenomena that occur in the BS [20]. The BS may reflect functional attributes of the
79 species producing them that are linked to the definition of ecosystem engineers. These
80 structures and the specific environment associated to them have been given the name of
81 "functional domain" [15]. These are places where specific soil processes occur at certain
82 spatial and temporal scales, so that the effects of ecosystem engineers in the ecosystem
83 through their functional domain can be quite significant [14,15]. The functional domain is
84 then a part of the soil that is influenced by a regulator that can be either biotic or abiotic.

85 The BS can be separated from the soil due to its different physico-chemical properties.
86 Decaëns et al. [9] set up a classification of engineer organisms (macroinvertebrates) in the
87 savannas from Carimagua. They demonstrated the rich diversity of BS produced by
88 ecological engineers in the soil surface of the natural savanna, i.e. i) compact structures,
89 rich in organic matter (earthworm casts), ii) soft structures, rich in organic matter (termite
90 mounds), and iii) soft granular structures poor in organic matter (ant nests). In this study,
91 however, our objective was to quantify the organic C (C_{org}), NH_4^+ and NO_3^- concentrations

92 in different parts of the BS produced by termites and ants to test the hypothesis that higher
93 concentrations are found where new building material is deposited, i.e. at the top of the BS.
94 The criterion was set up “a priori” since differences in concentration of these nutrients are
95 supposed to occur owing to the age, with the oldest part located at the bottom of the BS,
96 since these structures are normally constructed upwards.

97

98 **2. Materials and Methods**

99 *2.1. Study site*

100 The study was carried out in a natural savanna (NS), an introduced grass-legume
101 pasture (IP) and a gallery forest (GF) at the CORPOICA – CIAT Carimagua research
102 station (Figure 1), in the well-drained isohyperthermic savannas of the Eastern Plains of
103 Colombia (4° 37' N, 71°19' W and 175 m altitude). Average annual rainfall and
104 temperature are about 2,280 mm and 26 °C respectively with a dry season from December
105 to March. Soils at the study site are oxisols characterized by their acidity (pH [H₂O] = 4.5)
106 and a high Al saturation (>90%).

107 In the NS *Trachypogon vestitus* Anderss, *Paspalum pectinatum* Nees, *Axonopus aureus*
108 Beauv., *Schyzachyrium hirtiflorum* Ness, *Gymnopogon foliosus* Nees and *Hiptis conferta*
109 Pohl ex Benth. (Labiatae) are the most frequent grass species. The IP was an association of
110 *Brachiaria humidicola* Rendle with three different legumes, *Arachis pintoii* Krap & Greg,
111 *Desmodium ovalifolium* Wall. and *Stylosanthes capitata* Vog. Pasture was sown in 1993
112 and legume resown in 1996. Stocking rates for the pasture were 1 cattle ha⁻¹ (1 animal unit
113 [AU] = 250 kg live weight) in the dry season and 2 AU ha⁻¹ in the rainy period.

114 The GF where the BS were sampled was in an site called “La Reserva”, very close to
115 the Carimagua Lake (Figure 1). The dominant vegetation in the GF is constituted by several
116 tree species such as *Ficus* spp., *Dendropanax arboreux*, *Enterolobium* sp., *Jacaranda*
117 *copaia*, *Copernicia tectorum*, *Cecropia* sp. and palm forests of *Mauritia flexuosa* and *M.*
118 *minor*.

119

120 2.2. Identification of BS and morphological descriptions

121 The different plots studied were thoroughly checked and all BS found were described,
122 and the macroinvertebrates responsible for their construction identified as precisely as
123 possible (family, genus, or species). We restricted this study to the BS produced by termites
124 and ants, and did not include earthworm casts, since these were intensively studied, at least
125 for one anecic species [8,9].

126

127 2.3. Sampling procedure

128 The study was conducted in the middle of the rainy season of 1999 (August). Complete
129 BS produced by ecosystem engineers in the area were sampled. The protocol of sampling
130 procedure is indicated in Figure 2. For those BS of large size, i.e. more than 80 cm height
131 we sampled at 0, 20, 40, 60, 80 and 100% of the distance from the top to the base; due to
132 size reasons only 0, 50, and 100% sampling distance was used for the smallest BS. Two
133 types of control soil were used, (1) soil taken aside the BS, i.e., 120% and 150% for large
134 and small BS, respectively, and (2) soil taken 1 m away from the BS. A small metal
135 cylinder (5 cm Ø) was used to sample at 0-5 and 5-10 cm (Figure 2).

136 Four replicates, i.e. four BS produced by the same organism, were sampled at each site.
137 Each sample taken at different distances, i.e., 0, 20, etc., was introduced separately in
138 plastic bags and put in an ice chest to preserve further mineralisation processes and carried
139 to the laboratory. We only sampled those BS that were sufficiently represented to permit
140 the collection of enough material for laboratory determinations. The ants and termites that
141 might be found in the samples were carefully removed before preserving the samples at 4
142 °C prior to analysis. In total 380 samples were analysed.

143

144 *2.4. Chemical analysis*

145 Chemical analysis were carried out at the “Centro Internacional de Agricultura
146 Tropical” (CIAT) headquarters in Cali, so samples were sent from Carimagua in an ice
147 chest and all the samples inside plastic bags to avoid direct contact with ice. NH_4^+ and
148 NO_3^- concentrations were determined following standard techniques recommended by the
149 Tropical Soil Biology and Fertility Programme (TSBF) [2]. We used a colorimetric method
150 after acid digestion to measure total C concentrations [11] in samples that were dried at 75
151 °C for 48 h.

152

153 *2.5. Statistical treatments*

154 Data were transformed before analysis to reduce the asymmetry of the frequency
155 distribution. Normalisation of data was obtained using the Shapiro-Wilks test for normality.
156 Mean comparisons were performed with one-way ANOVA.

157

158 **3. Results**

159 *3.1 Diversity and description of BS*

160 Decaëns et al. [9] described fourteen types of BS and the invertebrates responsible for
161 their construction on the soil surface in the NS. Out of 14 BS we collected eight types of
162 BS (Table I), three epigeic ant nests (Plates 1a-c) and five types of epigeic termite domes
163 (one located above trees) (Plates 2a-c). Table II lists those macroinvertebrates identified
164 and the size of the BS, including those listed in [9]. In this paper only the two termite BS
165 sampled in the GF are described:

- 166 • *Nasutitermes* sp.1 (unidentified species)

167 The BS constructed by this termite species is an epigeic conic mound of large size
168 (60 cm height x 50 cm Ø). The surface of the mound is rough, with cemented
169 material and with colour similar to the surrounding soil (Plate 3a, from J.J.
170 Jiménez).

- 171 • *Nasutitermes* sp. 2 (unidentified species)

172 This termite constructs an arboreal BS which is located in the range of 1.70 – 3.70
173 m above soil surface. The BS is a spheric, pasteboard-like structure, and generally
174 some decomposed leaves are visible throughout the surface. There seems to be
175 some kind of specificity between this termite and the tree where nests are built. All
176 BS were found in trees belonging to the same species with about the same
177 dimensions, i.e. 5-8 cm Ø (arboreal termite mound, Plate 3b, from J.J. Jiménez).

178

179 *3.2. C_{org} , NH_4^+ , and NO_3^- in BS and control soil*

180 Concentrations of C_{org} were lowest in the BS structures produced by ants, especially
181 those deposited by *A. laevigata* and *Trachymyrmex* sp. in the NS (Figure 3a, b). In the GF
182 values of C_{org} were also rather low in the BS of *A. laevigata*, although higher than the
183 control soil (only significant for 5-10 cm in control soil 1 and for both 0-5 and 5-10 cm of
184 control soil 2). In the case of termites, similar values were obtained for *Spinitermes* sp.
185 (Figure 3a) (ca. 6, considering the BS from the NS and IP) and *Velocitermes* sp. (Figure 3b)
186 (mean values ranging from 5 to 9 throughout the BS). The highest C_{org} concentrations were
187 observed in the BS of *Nasutitermes* sp.1 in the GF (15 – 22), *Microcerotermes* sp. in the NS
188 (20 – 31), and *Nasutitermes* sp2 in the GF (37– 54).

189 Regarding NH_4^+ concentrations in the large BS the highest values were obtained in
190 *Spinitermes* sp. (Figure 4a). High NH_4^+ concentration was also obtained in the BS produced
191 by *Nasutitermes* sp1. in the GF, although these values were on average three times lower
192 than those obtained for *Spinitermes* sp. Regarding the BS produced by *A. laevigata*, NH_4^+
193 concentrations were the lowest (below $10 \mu\text{g g dry soil}^{-1}$ in all cases and distances).
194 However, when considering the entire set of BS analysed, the highest NH_4^+ concentrations
195 were obtained in the arboreal pasteboard-like BS produced by *Nasutitermes* sp2 in the GF
196 (between $1,200$ and $1,600 \mu\text{g g dry soil}^{-1}$, depending on the sampling distance) (Figure 4b).
197 There were significant differences between the BS and both types of control soil for the BS
198 produced by termites in the GF (ANOVA, $P < 0.01$).

199 Regarding NO_3^- the highest values were found in the BS produced by *Nasutitermes*
200 sp1 and *Nasutitermes* sp2 (Figure 5a, b). In general these values were ten times higher
201 (above $1,000 \mu\text{g g dry soil}^{-1}$) than those obtained for the rest of BS. Rather high values of

202 NO_3^- were also obtained in the BS produced by *A. laevigata* in the GF. The BS constructed
203 by *Spinitermes* sp. had the lowest concentrations of NO_3^- .

204 In general it was observed a decrease in all variables measured as a function of
205 sampling distance, thus revealing that the most recent material deposited in the BS
206 corresponded to the distance 0%, unless an area in the BS had to be repaired that might lead
207 to higher values in other sampling distances. The highest concentration of NH_4^+ was
208 obtained in the first sampling distance, i.e. 0% (Figure 4). Regarding NO_3^- concentrations
209 these were rather low for the entire set of small BS (Figure 5b).

210 When comparing the entire set of BS, there were significant differences in the 0%
211 sampling distance for NH_4^+ concentrations (ANOVA, $F = 37.70$; d.f. = 6; $p < 0.001$), and
212 C_{org} concentrations (ANOVA, $F = 10.14$; d.f. = 6; $p < 0.001$), but not for NO_3^-
213 concentrations ($F = 2.27$; d.f. = 6; $p > 0.05$).

214 When comparing both types of control soil, i.e. the sampling distance beside the BS,
215 120% or 150% and control soil (0-5 and 5-10 cm) only statistically significant differences
216 were found for C_{org} in the BS produced by *Microcerotermes* sp. in the NS, and *Spinitermes*
217 in both NS and IP systems (ANOVA, $P < 0.05$).

218

219 3.3. Effects of sampling site

220 Only for two species, *A. laevigata* (NS, GF and IP) and *Spinitermes* sp. (NS, IP)
221 statistical analysis between systems could be performed. No significant differences were
222 found between both NH_4^+ (ANOVA, $F = 0.09$; d.f. = 1; $p > 0.05$) and NO_3^- (ANOVA, $F =$
223 0.29 ; d.f. = 1; $p > 0.05$) concentrations in the BS produced by *Spinitermes* sp. However,

224 significant differences appeared regarding total C_{org} concentrations (ANOVA, $F = 10.12$;
225 d.f. = 1; $p < 0.005$).

226 In the case of *A. laevigata* all systems studied could be compared. There was no
227 significant differences for NH_4^+ (ANOVA, $F = 2.92$; d.f. = 2; $p > 0.05$) but differences were
228 highly significant for NO_3^- concentrations (ANOVA, $F = 321.58$; d.f. = 2; $p < 0.001$) and
229 total C_{org} (ANOVA, $F = 8.90$; d.f. = 2; $p < 0.001$).

230

231 **4. Discussion**

232 The activities of soil ecosystem engineers contribute to the variability of chemical
233 concentrations in the BS they produce, and sometimes in the surrounding environment. Our
234 results seemed to confirm the presence of a mosaic of areas with different C and N
235 concentrations. Attention must be paid when sampling these structures and preliminary
236 characterizations of chemical properties are sought.

237 The comparison of our results with those reported in other studies seems difficult since
238 the sampling methodology employed here has not been used before. However, some
239 general considerations can be taken into account. In a study conducted in the Brazilian
240 Cerrados it was found that soil organic carbon was enriched by a factor of 3.5 (90.2 g/kg
241 C_{org} in clayey Oxisols) and 11.5 (109 g/kg C_{org} in the loamy Oxisols) in the tops of the
242 epigeic mounds of termites of the genera *Armitermes* and *Dihoplotermes* [27]; these
243 termites selected fresh and partly decomposed organic matter. In our study, C_{org} , NH_4^+ and
244 NO_3^- concentrations were also higher in the BS compared to the control soil, except in the
245 BS produced by *Nasutitermes* sp2 in the gallery forest (Figure 4b), since here we compared

246 an arboreal carton-type BS with control soil, showing the differences in the composition of
247 microbial population in both types of substrates.

248 Differences in total C_{org} , NH_4^+ , and NO_3^- concentrations observed between BS and
249 the control soil are due to several reasons depending on the species and how these BS were
250 built. For instance, higher amounts of organic matter are found in faeces and salivary
251 excretions of termites [4,19]. Epigeic termite mounds are normally cemented with soil
252 particles that contain variable quantities of salivary secretions and excrements [17,18]. In
253 certain cases, the walls of the termite mounds are made of a pasteboard-like material that is
254 very rich in C_{org} , e.g. *Microcerotermes* sp. in the savanna and the arboreal nests constructed
255 by *Nasutitermes* sp2. in the GF.

256 The epigeic domes built by ants, in contrast to the structures built by termites were in
257 this study pale yellow or light orange. It is worth noticing that the BS produced by *A.*
258 *laevigata* in the pasture was of smaller size and the one from the GF was the biggest. In the
259 pasture the presence of cattle and also de control of ant populations are factors responsible
260 of this size, otherwise these BS can be quite big as usually seen in the natural ecosystems.
261 Fungus growing ants bring their depositions (soil plus fungi residues) up from the deeper
262 horizons of the soil profile (B horizon begins at 26 cm depth in these soils). Ants
263 accumulate all these residues on the soil surface. Despite the high amount of soil removed
264 by foraging ants the BS are poor in organic matter concentration and this soil does not
265 undergo significant modifications [9].

266 The highest values of NO_3^- were obtained in the BS produced by *Nasutitermes* sp1 in
267 the GF. This BS is very compacted and not only salivary secretions and faeces of termites
268 but also hydrosoluble carbohydrates of microbial origin [26] could be responsible of its

269 compact nature. This must be further tested since no correlation between the structural
270 stability of biogenic aggregates and hydrosoluble carbohydrates for the same BS has been
271 found [9]. In our case it was also common to observe green areas over the surface of the
272 BS indicating the presence of other microorganisms such as algae that may have
273 contributed to the great values of C_{org} (Figure 3d). This was also confirmed by analysing
274 the organic matter concentrations of this structure by Near Infrared Reflectance
275 Spectroscopy (NIRS) [10] and enzymology analyses [21].

276 Owing to the sampling distance 0% or the zone of the BS where the “most recently”
277 deposited material is found (confirmed by the results obtained in this study), NH_4^+ values
278 were in general highest in the BS produced by *Spinitermes* sp. (in the savanna) and
279 *Nasutitermes* sp1 and *Nasutitermes* sp2 (from the GF). These values were higher than those
280 obtained in the BS produced by another soil ecosystem engineer (an anecic earthworm) in
281 the same area [9].

282 High NH_4^+ concentration probably indicated that N mineralization was high in the
283 recently deposited material in these BS. Nitrifying microorganisms would be greatly
284 activated in the BS produced by *A. laevigata* and *Nasutitermes* sp1 in the GF. These BS are
285 the biggest found at the study site and the values obtained in this study were higher than
286 those reported by [9]. A detailed study on the role of which microbial populations are
287 enhanced or inhibited in these BS merits further efforts.

288 Termites are the most important decomposers of all invertebrates in tropical forests
289 [3]. Termites strongly influence soil organic matter and nutrient dynamics [16]. Total C_{org}
290 concentrations in this study were different according to the BS and were in general higher
291 than reported in other studies, for example, the termite mounds of *Microcerotermes*

292 *nervosus* Hill 1942 from northern Australia (5-15% organic C) [18]. In another species,
293 *Amitermes laurensis* Mjoerberg C_{org} (%) ranged from 6.9 in the upper part of the termite
294 mound to 3.9 and 1.7 in the central and lower parts, respectively [22]. In our study the
295 highest values were obtained in the termite structures produced in the GF by *Nasutitermes*
296 sp1 (ranging from 15 to 22) and *Nasutitermes* sp2 (from 37 to 54).

297 Termite effect of total soil respiration results from direct CO_2 emissions from
298 respiration of live tissues (termite and fungal tissues) and from the additional soil
299 respiration due to the stimulation of soil microbial metabolism in the processed material.
300 The high concentration of C_{org} , and NO_3^- observed in the BS produced by *Nasutitermes* sp1
301 in the GF may enhance the activity of micro-organisms in a similar way to that described
302 for ants [7].

303 In the natural savannas, fire and termites cause emissions of CO_2 , CO, CH_4 , NO, N_2O
304 to the atmosphere [10]. It has been shown that the only contribution to net emissions of
305 CH_4 is made by *Spinitermes* sp., and it is much lesser than the emissions from cattle (0.10
306 $Tg\ yr^{-1}$) or direct emissions from biomass burning (0.06 $Tg\ yr^{-1}$) [23]. All CH_4 generated by
307 subterranean termites is oxidized by soils before escaping into the atmosphere. Estimated
308 fluxes due to termites were reported as 7.2 $g\ CH_4\ ha^{-1}\ yr^{-1}$ in the NS. The integrated annual
309 CH_4 flux coming from termite mounds in the Llanos is 76 $Mg\ CH_4\ yr^{-1}$. This value is only
310 about 0.0004% of the total global emissions of 19.7 $Tg\ CH_4$ attributed to termites [24].

311 Our results also highlighted the evidence that the activities of some ecosystem
312 engineers affect the surrounding soil by significantly increasing the concentration of some
313 nutrients compared to the control soil. For example, some termites are accumulating C_{org}
314 aside the BS which is an area that would correspond to the “functional domain” of the

315 ecosystem engineer [15], although more studies are needed. A further study should be
316 considered to assess the microbial communities in different termite mounds under the
317 hypothesis of different microbial community composition within the same functional
318 domain, and whether this is the result of BS building or habitat preferences by these
319 organisms, as suggested by [7]. Besides it remains unknown which taxa of the microbial
320 community are activated in the mounds of termites in both savanna ecosystem and GF from
321 the Colombian Llanos.

322 Finally, we conclude that the activity of soil ecosystem engineers increased the spatial
323 variability of chemical parameters measured in this study. Other BS, however seemed not
324 to have any influence in the surrounding soil. Nutrient concentrations were higher in
325 general in the BS than in the control soil in all cases. This shows the fact of a nutrient
326 variability in the systems studied that may affect ecosystem processes and functional
327 diversity of micro-organisms and plants at certain scales. A model of the dynamics of
328 nutrients in the BS in these savannas and gallery forests of the Colombian “Llanos” and
329 how above-ground plant communities are affected by the activities of soil ecosystem
330 engineers should be further addressed. Besides, some unanswered questions remain, e.g.
331 which proportion of C in the mounds of leaf-foraging ants is of plant or soil-derived origin?
332 A further assessment with recently available techniques like Near Infrared Spectroscopy
333 (NIRS) can help answer these questions. It would also be necessary to initiate studies about
334 the lifespan and dynamics of break-down of these BS [8] in different environments to test
335 the hypothesis of higher functional diversity of soil ecosystem engineers and soil processes,
336 and also the dynamics of BS when the organisms that produce them are no longer present.

337

338 **5. Acknowledgements**

339 We wish to acknowledge the help of Paola Pinto in organic matter and C
340 determinations, and Neusa Asakawa for NH_4^+ and NO_3^- determinations at CIAT lab
341 facilities. Sincere thanks to Drs. Richard Thomas and Edgar Amézquita (CIAT) for their
342 technical and financial support, Stephen Lapointe for the confirmation of ant species
343 identification and Paul Eggleton (The Natural History Museum, London) for a preliminar
344 identification of termite genera. Financial support was provided by the Soil, Water and
345 Nutrient management systemwide programme of the CGIAR and also through the IBOY-
346 DIVERSITAS programme. We also greatly want to thank Clive G. Jones for useful
347 discussions held at the XIV International Colloquium of Soil Zoology and Ecology (ICSZ)
348 in Rouen (France).

349

350 **6. References**

- 351 [1] J. M. Anderson, Soil organisms as engineers: microsite modulation of macroscale
352 processes, in: C. G. Jones, J. H. Lawton, eds., *Linking Species and Ecosystems*,
353 Chapman and Hall, London, pp. 94-106, 1995.
- 354 [2] J. M. Anderson, J. Ingram (eds), *Tropical soil biology and fertility, a handbook of*
355 *methods*, CAB International, Oxford, 1993.
- 356 [3] D. E. Bignell, P. Eggleton, Termites in ecosystems, in: T. Abe, M. Higashi, D. E.
357 Bignell, eds., *Termites: Evolution, Sociality, Symbiosis, Ecology*, Kluwer
358 Academic Press, Dordrecht, 2000.
- 359 [4] H. I. J. Black, M. J. N. Okwakol, Agricultural intensification, soil biodiversity and
360 agroecosystem function in the tropics: the role of termites, *Applied Soil Ecology* 6
361 (1997) 37–53.
- 362 [5] E. Blanchart, P. Lavelle, E. Braudeau, Y. L. Bissnais, C. Valentin, Regulation of
363 soil structure by geophagous earthworm activities in humid savannas of Côte
364 d'Ivoire, *Soil Biology and Biochemistry* 29 (1997) 431-439.

- 365 [6] L. J. Brussaard, V. M Behan-Pelletier, D. E. Bignell, V. K. Brown, W. Didden, P.
366 Folgarait, C. Fragoso, D. W. Freckman, V. S. R. Gupta; T. Hattori, D. L.
367 Hawksworth, C. Klopatek, P. Lavelle, D. W. Malloch, J. Rusek, B. Söderström, J.
368 M. Tiedje, R. A. Virginia, Biodiversity and ecosystem functioning in the soil,
369 *Ambio* 26(8) (1997) 563-570.
- 370 [7] J. Dauber, V. Wolters, Microbial activity and functional diversity in the mounds of
371 three different ant species, *Soil Biology and Biochemistry* 32 (2000) 93-99.
- 372 [8] T. Decaëns, Degradation dynamics of surface earthworm casts in grasslands of the
373 Eastern Plains of Colombia, *Biology and Fertility of Soils* 32 (2000) 149-156.
- 374 [9] T. Decaëns, J. H. Galvis, E. Amézquita, Propriétés des structures produites par les
375 ingénieurs écologiques à la surface du sol d'une savane colombienne, *Compte*
376 *Rendus Academie des Sciences de Paris* 324 (2001) 465-478.
- 377 [10] M. Hedde, R. Joffre, P. Lavelle, J. J. Jiménez, T. Decaëns, Specific functional
378 signatures in soil macro-invertebrate bio-structures, *Functional Ecology* 19 (2005)
379 785-793.
- 380 [11] V. J. C. Houba, J. J. van der Lee, I. Novozamsky, J. Walinga, Soil and plant
381 analysis, in: Department of Soil Science and Plant Nutrition, ed. *Soil analysis*
382 *procedures*, 4th ed. Agricultural University, Wageningen, The Netherlands, p 4-14,
383 1988.
- 384 [12] C. G. Jones, J. H. Lawton, M. Shachak, Organisms as ecosystem engineers, *Oikos*
385 69 (1994) 373-386.
- 386 [13] C. G. Jones, J. H. Lawton, M. Shachak, Positive and negative effects of organisms
387 as physical ecosystem engineers, *Ecology* 78 (1997) 1946-1657.
- 388 [14] P. Lavelle, Faunal activities and soil processes: adaptive strategies that determine
389 ecosystem function, *Advances in Ecological Research* 27 (1997) 93-132.
- 390 [15] P. Lavelle, Functional domains in soils, *Ecological Research* 17 (2002) 441-450.
- 391 [16] P. Lavelle, A. V. Spain, *Soil Ecology*, Kluwer Academic Publishers, 2001.
- 392 [17] K. E. Lee, R. C. Foster, Soil fauna and soil structure, *Australian Journal of Soil*
393 *Research* 29 (1991) 745-775.
- 394 [18] K. E. Lee, T. G. Wood, Physical and chemical effects on soils of some Australian
395 termites, and their pedological significance, *Pedobiologia* 11 (1971) 376-409.

- 396 [19] L. A. Lobry-de-Bruyn, A. J. Conacher, The role of termites and ants in soil
397 modification: a review, *Australian Journal of Soil Research* 28 (1990) 55–93.
- 398 [20] A. Martin, J.Y.C. Marinissen, Biological and physico-chemical processes in
399 excrements of soil animals, *Geoderma* 56 (1993) 331-347.
- 400 [21] P. Mora, E. Miambi, C. Rouland, T. Decaëns, J. J. Jiménez, Functional typology of
401 structures produced by earthworms, ants and termites, *Soil Biology and*
402 *Biochemistry* 37 (2005) 1043-1048.
- 403 [22] T. Okello-Oloya, A. Spain, R. D. John, Selected chemical characteristics of the
404 mounds of two species of *Amitermes* (Isoptera, Termitinae) and their adjacent
405 surface soils from northeastern Australia. *Revue d'Ecologie et Biologie du Sol* 22
406 (1985) 291-311.
- 407 [23] M. A. Rondón, J. M. Duxbury, R. J. Thomas, Effects of Land Use Change in the
408 “Llanos of Colombia” on Fluxes of Methane and Nitrous Oxide, and on Radiative
409 Forcing of the Atmosphere, in preparation
- 410 [24] M. G. Sanderson, Biomass of termites and their emissions of methane and carbon
411 dioxide, a global data base, *Global Biogeography* 10 (1996) 543-557.
- 412 [25] N. E. Stork, P. Eggleton, Invertebrates as determinants and indicators of soil
413 quality, *American Journal of Alternative Agriculture* 7 (1992) 38-47.
- 414 [26] J. M. Tisdall, Possible role of soil microorganisms in aggregation in soils, *Plant Soil*
415 159 (1994) 115-121.
- 416 [27] W. Zech, W. Amelung, H. Neufedt, Organic matter in termite mounds of the
417 Brazilian Cerrados, in: R. J. Thomas, M. A. Ayarza (eds.) *Sustainable Land*
418 *Management for the Oxisols of the Latin American Savannas*, CIAT publication no.
419 312, 231 p, 1999.
- 420

421 **Tables**

422 Table I

423 Taxonomic position of macroinvertebrates together with a brief description of the BS produced

Macroinvertebrates				Structures				
Order	Family	Species	Feeding regime	Type	Colour	Shape	Aspect	
Ants								
Hymenoptera	Formicidae	<i>Atta laevigata</i> (F. Smith, 1858)	Fungi growers	Dome	orange	truncated cone	Mobile rubbish (loose aggregates)	Plate 1a
Hymenoptera	Formicidae	<i>Acromyrmex landolti</i> (Forel, 1885)	Fungi growers	Dome	orange	cone	Mobile rubbish	Plate 1b
Hymenoptera	Formicidae	<i>Trachymyrmex</i> sp. Forel	Fungi growers	Dome	orange	cone	Mobile rubbish	Plate 1c
Termites								
Isoptera	Termitidae	<i>Microcerotermes</i> sp. Silvestri 1901	Soil feeder	Dome	black	cylinder	Pasteboard-like material	Plate 2a
Isoptera	Termitidae	<i>Spinitermes</i> sp. Wasmann 1867	Soil feeder	Dome	black	cone	cemented material	Plate 2b
Isoptera	Termitidae	<i>Velocitermes</i> sp. Holmgren 1912	Litter feeder	Dome	grey	cone	cemented material	Plate 2c
Isoptera	Termitidae	<i>Nasutitermes</i> Dudley 1890 sp1.	Soil feeder?	Dome	grey	cone	cemented material	Plate 3a
Isoptera	Termitidae	<i>Nasutitermes</i> Dudley sp.2	Soil-litter feeder?	Arboreal	Dark brown	Sphere	Pasteboard-like material	Plate 3b

424

425

426 Table II

427 Size of the biogenic structures sampled (mean and standard deviation)

Species	Site	Dimensions (cm)	
		Height	Radius
<i>Atta laevigata</i>	NS	17.75 ± 8.26	50 ± 34.64
	GF	50 ± 16.33	33.75 ± 24.28
	IP	11 ± 8.76	29.75 ± 20.60
<i>Acromyrmex landolti</i>	NS	4 ± 0.82	27 ± 10.13
<i>Trachymyrmex</i> sp.	NS	6 ± 1.6	14.5 ± 4.4
<i>Microcerotermes</i> sp.	NS	12.5 ± 2.1	5.5 ± 0.6
<i>Spinitermes</i> sp.	NS	49.25 ± 14.2	19.75 ± 3.9
	IP	50 ± 16.8	27.25 ± 9.3
<i>Velocitermes</i> sp.	NS	13.25 ± 2.4	6.5 ± 1.3
<i>Nasutitermes</i> sp1	GF	57.5 ± 22.5	18.9 ± 8.5
<i>Nasutitermes</i> sp2	GF	57.5 ± 9.6	25 ± 5.8

428

429 **Figure legends**

430

431 Figure 1. A map of the Carimagua research station (adapted from G. Escobar)

432

433 Figure 2. Illustration of the sampling location along the distance (expressed as %) from
434 the top (0) to the base (100) of the BS. Large BS allowed the collection of more samples
435 to be taken (left) than small ones (right).

436

437 Figure 3. Concentration of C_{org} in large (a) and small (b) BS produced by some ants and
438 termites in the different systems studied. The distance from the top to the base of the BS
439 is given as percentage(100%); 120% and 150% indicate the distance outside the BS
440 (control soil 1) and CS indicates the control soil 2 (0-5 cm).

441

442 Figure 4. Concentration of NH_4^+ in large (a) and small (b) BS produced by some ants and
443 termites in the different systems studied. The distance from the top to the base of the BS
444 is given as percentage(100%); 120% and 150% indicate the distance outside the BS
445 (control soil 1) and CS indicates the control soil 2 (0-5 cm).

446

447 Figure 5. Concentration of NO_3^- in large (a) and small (b) BS produced by some ants and
448 termites in the different systems studied. The distance from the top to the base of the BS
449 is given as percentage(100%); 120% and 150% indicate the distance outside the BS
450 (control soil 1) and CS indicates the control soil 2 (0-5 cm).

451

452 **List of plates**

453

454 Plate 1. Biogenic structures produced by ants in the savanna that were analysed; a) *Atta*
455 *laevigata* Smith nest (Picture from J.J. Jiménez), b) *Acromyrmex landolti* Forel nest
456 (Picture from J.J. Jiménez), c) *Trachymyrmex* sp. (picture from T. Decaëns)


457

458 Plate 2. Biogenic structures produced by termites in the savanna that were analysed a)
459 *Microcerotermes* sp., b) *Spinitermes* sp., c) *Velocitermes* sp. (all pictures from T.
460 Decaëns).

461

462 Plate 3. Biogenic structures produced by termites in the GF that were analysed a)
463 *Nasutitermes* sp1, b) *Nasutitermes* sp2 (pictures by Jiménez).


464



465


466

467




468

469 Figure 2.




470
 471
 472
 473 Figure 3.



474
475
476
477
478

Figure 4.



480
 481
 482
 483 Figure 5.



Plate 1. BS produced by ants in the savanna.



Plate 2. BS produced by termites in the savanna (Decaëns et al. 2001).




Plate 3. BS produced by termites in the gallery forest.