Running page head: Host use pattern of *Afropinnotheres monodi*

Host use pattern of the pea crab *Afropinnotheres monodi*: potential effects on its reproductive success and geographical expansion

Pilar Drake^{1,*}, Elena Marco Herrero¹, Maria Dulce Subida^{1,2}, Alberto M. Arias¹ and José A. Cuesta¹

¹ Instituto de Ciencias Marinas de Andalucía, CSIC, Avenida República Saharaui, 2, 11519 Puerto Real, Cádiz, Spain.

Phone: +34 95683612;

Fax: +34 956834701;

² Center for Marine Conservation (Estación Costera de Investigaciones Marinas), Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile. Casilla 193, Correo 22, Santiago, C.P. 6513677, Chile.

*e-mail: pilar.drake@icman.csic.es

ABSTRACT: The successful settlement of the African pea crab *Afropinnotheres monodi* in the Bay of Cadiz was analysed to get new insights into the evolution of pea crab parasite life history traits. The pea crab lives symbiotically and with high prevalence in the bivalves *Cerastoderma glaucum* and *Mytilus galloprovincialis*, and with low prevalence in *Scrobicularia plana*. A remarkable monopolisation of host (one crab/bivalve) occurred irrespective of host species and crab demographic categories (males, hard females, soft females), probably as an optimisation of resources in small-size hosts. However, there was a clear asymmetry in host use by the different crab

categories. A 1:1 sex ratio was found in crabs harboured by *C. glaucum* and *S. plana*, with most of the female crabs being small and hard specimens, whereas crabs inhabiting *M. galloprovincialis* were primarily large reproductive females. Ovigerous females were found throughout the year in *M. galloprovincialis* and there was a strong correlation between female size and fecundity, suggesting that the females harboured by this host were the major contributors to the reproductive effort of the studied population. Conversely, most of the new crabs recruiting to the population were harboured by the remaining two host species. We hypothesise that such a generalist but asymmetrical usage of bivalve hosts by pea crabs may have clear benefits for species in geographical expansion as *A. monodi*, facilitating its dispersal to new locations and its successful settlement in sheltered systems, as the Bay of Cadiz.

KEY WORDS: Symbiotic crabs, Multiple hosts, Life cycle, *Afropinnotheres monodi*, *Cerastoderma glaucum*, *Mytilus galloprovincialis*, *Scrobicularia plana*

INTRODUCTION

As a result of natural cycles and human activity, boundaries of the geographic distribution of species are dynamic (Brown et al. 1996; Doney et al. 2012; Jones et al. 2013). The successful settlement of species on a new location depends on a suitable mechanism of dispersal and on the ability to find appropriate habitats there (Bradbury & Snelgrove 2001). In the case of symbiotic lifestyles (in its etymological sense, i.e. including parasitism, commensalism and mutualism), this means that the invasive stage of symbionts must arrive to the new location and find there a stable population of its original host and/or of a new suitable host. A symbiont that is not host-specific is

1 expected to be more successful in colonising new areas than strict host specialists.
2 Furthermore, the use of more than one host species may increase the fitness of
3 symbiotic species: symbionts may sequentially infest different hosts in order to optimize
4 the use of resources (De Bruyn et al. 2010). In the long term this can lead to a more
5 complex life cycle; however, the initial and transitory step of the adaptation (facultative
6 use of additional hosts) may favour the geographical expansion of the symbiotic guests.

7 Partnerships in marine symbiosis often include species of different size and
8 lifestyles: commonly, one small mobile species (symbiotic guest) associated to a more
9 sedentary species (symbiotic host). Due to the diverse morphology and lifestyle of
10 hosts, there is also a wide diversity of social groups among guests: in large and
11 morphologically complex hosts, several individuals of the smaller symbiotic guest
12 generally share the same host; whereas in hosts with small body size and/or low
13 morphological complexity the symbiotic partner usually lives alone (De Bruyn et al.
14 2009; Baeza & Díaz-Valdés 2011). Since the smaller symbiotic guest obtains essentially
15 refuge and food from its larger host, the adoption of a solitary lifestyle may respond to
16 the optimization of resources provided by small-size hosts (Baeza & Thiel 2003, 2007).
17 However, such solitary lifestyle causes that reproductive males and/or females must
18 temporarily leave their hosts for mating. Mating may take place inside the host of the
19 female symbiont, or in the water column, with the consequent increase in the risk of
20 predation on guests (Asama & Yamaoka 2009; Trottier & Jeffs 2012). Moreover, such
21 mating-related migrations confer further complexity to the symbiont life cycle.

22 Among marine symbionts, pinnotherid pea crabs display a wide diversity of
23 host-guest interactions. Pinnotherids may live as endo- or ectosymbionts of different
24 taxonomic groups of invertebrates, mainly in the mantle cavity of molluscs, burrows of
25 polychaetes, the integument of echinoids and the branchial sac of tunicates (Schmitt et

al. 1973; Grove & Woodin 1996; Subida et al. 2011; Baeza & Díaz-Valdés 2011; Jossart et al. 2013). As most brachyurans, pinnotherids have a complex free-living larval cycle, which represents the dispersive phase of the species (Scheltema 1986). After the settlement of megalopae, pinnotherids usually show a facultative free-living stage in both sexes (males and hard females). Thereafter, reproductive females have a last obligated symbiotic stage, during which soft females live in their host (Silas & Alagarwami 1967; Soong 1997; Becker & Türkay, 2010). In symbiotic pea crabs with a solitary lifestyle (one crab per host), males usually maximize their reproductive success by roaming among host individuals in search of receptive sedentary females (the “pure-search polygynandry of sedentary females” model proposed by Baeza & Thiel 2007). Thus, the spatio-temporal distribution of receptive females becomes crucial in determining the distribution of the remaining demographic categories (De Bruyn et al. 2009; Ocampo et al. 2012). From all the above-mentioned, pea crabs are considered an exceptionally interesting group to explore the different types of behavioural, phenotypic and genetic adaptations of symbionts, during the evolution process toward complex parasitic cycles (Grove et al. 2000; Jossart et al. 2013). Despite the increasing interest in host use and mating behaviour of pinnotherids in recent years (De Bruyn et al. 2009, 2010; Hernández et al. 2012; Ocampo et al. 2012; Jossart et al. 2013; Peiró et al. 2013), several aspects of their complex lifestyle are still unknown (Becker & Türkay 2010).

In the current scenario of rising seawater temperatures in southern Europe (Sanderson et al. 2011), new records and/or abundance increments of several African marine species in the Iberian Peninsula have been related to the warmer environmental conditions (Cabral et al. 2001; Subida et al. 2011). According to the very scarce information available, this could also be the case of the African pea crab

Afropinnotheres monodi Manning, 1993, which currently shows its northernmost populations in southern Europe (Subida et al. 2011). No information on hosts is available for the earliest African records, but European populations have been recorded living in the shell cavity of several bivalves (Subida et al. 2011). Thus, the arrival of *A. monodi* to the Gulf of Cádiz could be either due to the accidental transport of pea crabs in ships' fouling, i.e. in mussels (Apte et al. 2000), or to the larval dispersal from the North African populations. Regardless of the mechanism behind the introduction of *A. monodi* in Europe, it is reasonable to assume that the northernmost populations are probably in process of adapting to the new colonized environments. Thus, the apparently fast and undoubtedly successful settlement of the pea crab *A. monodi* in the Bay of Cadiz offers a unique opportunity to get new insights into the evolution of pea crab symbiotic life history traits.

MATERIALS AND METHODS

Study site and distribution of the bivalve hosts

All host individuals were collected within the Bay of Cadiz (southwestern Spain), an area that was originally part of the Guadalete river estuary but is now a marine marsh system as a consequence of sedimentary processes and dams. The Río San Pedro inlet, a sinuous seawater channel characterized by semidiurnal mesotides (tidal range 1 to 3.5 m), runs through the northeastern area of the marsh. The tidal current from the bay flows along the inlet, whereas the freshwater inflow is insignificant except during periods of heavy rain. The inlet shows a soft muddy bed at the inner zone and a muddy sand bed close to the river mouth (Figure 1).

A stable population of the clam *Scrobicularia plana* inhabits the intertidal mud flats of the Río San Pedro inlet, where clams burrow relatively deep into the sediment of the high tide zone. Conversely, a very high density of the cockle *Cerastoderma glaucum* was observed for some years in the muddy sand flats located at the mouth of the Río San Pedro inlet, where cockles may be found lightly burrowed (in the first ≈ 5 cm of sediment) in the mid-tidal zone; massive cockle mortality has been recorded in some summers and/or after rainy periods (Drake et al unpublished data). Crowded patches of the mussel *Mytilus galloprovincialis* can be found all around the Bay of Cadiz, attached to artificial hard structures, such as bridge pillars and concrete blocks used to protect harbour walls; as mussels usually settle at the middle-low tidal level, they remain emerged for shorter periods of time during each tidal cycle than clams and cockles.

Collection of hosts and crabs

Scrobicularia plana clams were monthly sampled for two years (June 2010 to May 2012). Although an occasional presence of the cockle *Cerastoderma glaucum* was recorded at the mouth of the Río San Pedro inlet during the first sampling year, a crowded population of cockles inhabited the study area from May 2011 to September 2012. During this period, monthly samples of cockles were collected in the same date as clams. The remarkably low abundance of pea crab reproductive females in clams and cockles, together with the concomitant high prevalence of non-reproductive pea crabs in the latter host, led us to search for an additional bivalve host in the area. In the Bay of Cadiz, *Afropinnotheres monodi* has been also collected in the bivalves *Chamelea gallina*, *Donax trunculus*, *Macra stultorum*, *Mytilus galloprovincialis*, *Spisula solida* and *Venerupis decussatus* (Drake et al unpublished data). Since reproductive females

were only found in the mussel beds of *Mytilus galloprovincialis*, this species was sampled monthly from June 2012 to May 2013. Mussel individuals were collected within the Bay of Cádiz from bridge pillars and concrete blocks of harbour protection; the demographic characteristics of the harboured pea crabs were similar in both sampling locations. During this last annual cycle, monthly samples of cockles were collected on the same date that mussel sampling took place, until the disappearance of the cockle population in October 2012; *Scrobicularia plana* clams were also occasionally sampled to corroborate that pea crab reproductive females were still extremely rare in this host.

In order to facilitate the collection of the three host bivalves in their respective habitats, clams, cockles and mussels were haphazardly sampled during the low tide of spring tides (full moon). *Scrobicularia plana* clams were collected by hand digging on the fine muddy sediment; *Cerastoderma glaucum* cockles were unburied from the sediment with a drug fork and then collected by hand; finally, mussels were removed from the hard substrates using a knife.

Host use pattern

The collected living bivalves were quickly taken to the laboratory; no crabs were observed leaving its host during transportation. In the laboratory, the shell length (SL = maximum distance along the anterior-posterior axis of the shell) of each host bivalve was measured to the nearest 0.1 mm with a dial caliper (Tesa Cal IP65). Afterwards hosts were opened and carefully inspected for the presence of symbiotic crabs.

Due to differences in shell morphology between the three studied hosts, for a same shell length the empty volume of the shell cavity, available for associated crabs,

can be very different. Thus, for each bivalve species, the empty volume of the shell cavity was estimated in a subsample of the collected individuals, as follows. After host dissection and retrieval of crabs, the soft tissues of the bivalves were placed back into the shell and each valve was filled with water. The empty volume of the shell cavity was measured as the total volume of water retained within the valves. This data was further used to construct, for each host species, the best-fitting regression models between shell length and the free volume of the shell cavity. Since volume data did not meet statistical assumptions of the parametric ANOVA test, interspecific differences in mean empty volume of host individuals harbouring *A. monodi* were assessed by using the non-parametric Kruskal Wallis H test.

All pea crabs found within each host individual were killed by freezing at (-20 °C) and then preserved and stored in ethanol (80%) for further demographic observations. The carapace width (CW = maximum cephalothorax width) of each pea crab was measured to the nearest 0.01 mm under a stereomicroscope provided with a calibrated ocular micrometer. Crabs were sexed on the basis of the presence (male) or absence (female) of gonopods. Furthermore, each female crab was classified either as hard or soft taking into account abdomen morphology, cephalotorax consistence and shape and pleopods morphology: hard females show a flattened and quadrangular cephalothorax and swimming legs, whereas the body of soft females is mostly spherical and lacks swimming legs. Lastly, soft females carrying eggs (embryos) were classified as ovigerous females.

The host use patterns of *Afropinnotheres monodi* were assessed by describing the characteristics of the crab population inside each host species, the seasonal and host-size related structure of these populations and the mean prevalence of the pea crabs in each host species. The mean prevalence of crabs was estimated as the average

prevalence values found at each sampling date. Mean carapace width of pea crab males and females from different hosts were compared using one-way ANOVA (on log-transformed data), or the Kruskal Wallis H test when data did not meet statistical assumptions of the parametric test.

The relationship between host size (SL) and crab size (CW) was assessed by means of a linear correlation analysis; the significance of the estimated Pearson correlation coefficient was used to measure the strength of this relationship. Additionally, the observed crab prevalence in each host size class was compared with the expected prevalence in the random Poisson distribution. Similarly, to test whether pea crabs tend to live solitarily, in pairs or aggregated in higher number inside the host, the frequency of occurrence of hosts without crabs and with different number of crabs was compared with the expected frequencies under a random distribution. For each host species, the observed sexual composition of pairs of crabs sharing a single host individual were compared with the expected frequencies of heterosexual and homosexual pairs in a random distribution. Also, the observed proportion of males to females was tested for deviation from a 1:1 sex ratio. In both cases, significant differences between the observed and the expected values were assessed using Chi-square tests (Zar 2010).

The fecundity of *Afropinnotheres monodi* was determined by counting all eggs carried by 108 ovigerous females. Eggs were extracted from the incubation chamber by immersing females during five minutes in an aqueous solution of sodium hypochlorite (10 g of active chlorine l^{-1}) followed by a soft agitation (adapted from Góes et al. 2005). Due to the extremely low prevalence of pea crab reproductive females in clams and cockles, most ovigerous females used for estimating fecundity were retrieved from mussels; namely, only 3 females were retrieved from *Scrobicularia plana* and 11 from

Cerastoderma glaucum cockles, whereas the remaining 94 ovigerous females were retrieved from the mussel *Mytilus galloprovincialis*. The relationship between carapace width and the number of embryos per female was fit to a power model by regression analysis. Since there was a significant effect of female size on the number of embryos, seasonal and host differences in fecundity were tested with an ANCOVA (covariate: log-transformed female CW; factors: season and host species) using the Student-Newman-Keuls a posteriori test.

Although a 5% significance level ($P = 0.05$) was considered for all statistical tests, in the multiple correlation analyses between host and crab sizes, the Bonferroni correction was used to avoid a spurious increase of significant correlations; in this case, a P of 0.005 was considered equivalent to a corrected P of 0.05 (Zar 2010).

RESULTS

Crab populations in different hosts

A total of 1557 pea crabs were collected during this study, distributed as follows: 225 in 9441 *Scrobicularia plana*, 920 in 2661 *Cerastoderma glaucum* and 412 in 868 *Mytilus galloprovincialis*. The mean prevalence of the pea crab was remarkably higher in mussels (45.9%) and cockles (41.1%) than in clams (4.1%). No megalopae of *Afropinnotheres monodi* were found in the three studied hosts. The youngest stage observed, the first crab (first stage after larval metamorphosis with $CW \leq 1\text{mm}$), was occasionally found in cockles (Table 1). The remaining ontogenetic stages of the pea crab were present in the three studied hosts, but their proportion and mean size differed between hosts (Table 1 and Figure 2). Namely, the proportion of soft females was

extremely low among the female crabs retrieved from *Scrobicularia plana* (13.3%) and *Cerastoderma glaucum* (13.1%) but extremely high in those retrieved from *Mytilus galloprovincialis* (97.4%). Moreover, the maximal size and hence the mean size of soft females were also significantly higher in crabs inhabiting mussels ($F = 190.3$, $P < 0.01$). Contrastingly, hard females represented almost 50% of the pea crabs retrieved from clams and cockles and were seldom found inside mussels; differences in the size of hard females retrieved in different hosts were moderated ($F = 2.0$, $P > 0.05$). The sex ratio of pea crabs collected in *Scrobicularia plana* (0.81) and *Cerastoderma glaucum* (1.07) did not significantly differ from 1:1 ($\chi^2 = 2.6$ and 1.07, respectively, $P > 0.05$), but a significantly higher number of females was observed in crabs obtained from *Mytilus galloprovincialis* (sex ratio = 0.14; $\chi^2 = 224.5$, $P < 0.01$) (Figure 2). Besides, the maximal size and hence the mean size of males were significantly higher in crabs retrieved from mussels ($H = 80.6$, $P < 0.01$).

The above-mentioned differences in the proportion of ontogenetic stages among hosts occurred all year round, although moderate seasonal changes could be observed (Figure 3). Minimal seasonal changes were observed for the pea crab population inhabiting cockles. Pea crab population in clams displayed monthly changes in the sex ratio lacking a clear seasonal pattern. Despite ovigerous crab females could be found all year round inside mussels, their proportion in autumn (October to December) was relatively low compared with the rest of the year. The occasional record of ovigerous females inside clams (April and June) and cockles (July and August) occurred during the period of higher reproductive activity (higher number of ovigerous females in mussels) of the pea crab (Figure 3).

Interactions between crabs

The number of pea crabs per host varied between 0 and 1 in clams and between 0 and 2 in mussels and cockles. Namely, 100% of the clams, 99.5 % of the mussels and 96.6% of the cockles harboured a single crab. Thus, crab distribution among hosts did not display a random pattern: there were more bivalves harbouring one crab and fewer bivalves without crabs or harbouring more than one crab, than those expected by chance alone. Nevertheless, the observed departure from a random pattern was statistically significant (χ^2 test, $P < 0.01$) only for the two host species with high prevalence of the pea crabs: mussels and cockles (Figure 4).

A total of 31 *Cerastoderma glaucum* harboured pairs of crabs, from which 17 (54.8%) were heterosexual pairs (15 with one hard female and one male and 2 with one soft female and one male) and 14 were homosexual pairs (10 with two males and 4 with two hard females). In mussels, of the 5 pairs of crabs sharing the same host individual, 4 were heterosexual (with one soft female and one male) and a single one was homosexual with one soft and one hard female. Taking into account the different sex ratio observed in both hosts (Figure 2), in cockles the sexual composition of the crab pairs did not differ significantly from that expected by chance alone ($\chi^2 = 2.0$; $P > 0.05$); whereas in mussels there were fewer female-female pairs than expected by chance alone ($\chi^2 = 10.1$; $P < 0.05$). However, due to the extremely low number of crab pairs found in mussels, the latter should be considered with caution.

Host size effects

The frequency distributions of crabs in host size classes were relatively similar for the three bivalves, with a mean host size of 26.8 mm in *Cerastoderma glaucum*, 27.3 mm in *Mytilus galloprovincialis* and 29.8 mm in *Scrobicularia plana* (Figure 5). Contrastingly, the corresponding empty volume of the shell cavity was significantly different among hosts: 3.3 ml in *Cerastoderma glaucum*, 7.4 ml in *Mytilus galloprovincialis* and 2.2 ml in *Scrobicularia plana* ($H = 649.3$, $P < 0.01$). Furthermore, 100% of the clams and 92.2% of the cockles harbouring crabs were individuals with an empty volume in the shell cavity lower than 5 ml, whereas only 20.8% of mussels followed this pattern (Figure 6).

Host size did not significantly affect the prevalence of crabs in *Scrobicularia plana*, with a higher number of crabs collected in the most frequent size classes. On the contrary, a smaller crab prevalence than expected from a random distribution was observed in the smallest individuals of *Cerastoderma glaucum* and *Mytilus galloprovincialis*; however such host size-related bias was only statistically significant in cockles (Table 1 and Figure 5).

Regardless of the host species, no clear host size-related distribution pattern of males and hard females of *Afropinnotheres monodi* was observed, although soft females were more frequent in larger host individuals. Nevertheless, when restricting the analysis to crabs collected inside bivalves with similar empty volume in the shell cavity (volume < 5ml), the estimated sex ratio of crabs retrieved from mussels proved to be significantly different from 1:1, with a higher prevalence of females (Figure 6).

The size of soft females, showed a highly significant positive correlation with mussel size and a moderate positive correlation with cockle size. No significant relationship was observed between host size and the size of hard female crabs,

regardless of the host species. A moderate positive correlation was also found for host size and the size of males retrieved from cockles and mussels (Table 1 and Figure 6).

The fecundity of *Afropinnotheres monodi* increased significantly with female size regardless of host species (ANCOVA, $F_{1,101} = 283.2$, $P < 0.01$) (Figure 7). After removing the effect of female size, fecundity did not differ significantly between seasons (ANCOVA, $F_{3,101} = 2.4$, $P > 0.05$), but the host effect was significant (ANCOVA, $F_{2,101} = 19.0$, $P < 0.01$), due to the lower number of eggs per female in crabs retrieved from clams (Student-Newman-Keuls test; $P < 0.01$).

DISCUSSION

From the three European species of pinnotherids currently accepted (Becker & Türkay 2010), *Pinnotheres petunculi* Hesse, 1872, *Pinnotheres pisum* (Linnaeus, 1767) and *Nepinnotheres pinnotheres* (Linnaeus, 1758), only the latter two species have been occasionally collected in the Gulf of Cadiz (Zariquiey Álvarez 1968; González-Gordillo et al. 2001; González-Gordillo et al. 2003a; López de la Rosa et al. 2002). But beside these European species, one small free-living specimen identified as *Afropinnotheres* sp., was collected in the Bay of Cadiz in 1995 (López de la Rosa et al. 2002) and, more recently, the presence of adults of *Afropinnotheres monodi* was reported in three different bivalve hosts from the Gulf of Cadiz (Subida et al. 2011). A revision of larval pinnotherid specimens previously collected in the Río San Pedro inlet (July 1991 to June 1992; June 1998 to August 1998) and in the Guadalete estuary (in spring-summer of 2006 and 2007) demonstrated that zoeae and megalopae of this African pea crab were being misidentified as *N. pinnotheres* (Drake et al. 1998; González-Gordillo et al.

2003b; Olaguer-Feliú et al. 2010). In fact, neither *P. pisum* nor *N. pinnotheres* were collected during the present study, which indicates that *A. monodi* is currently the dominant symbiotic pinnotherid in intertidal bivalve populations of the Bay of Cadiz. The expansion of the distribution range of African marine species into colder European waters is expected in the current scenario of rising seawater temperatures. Several other African crabs have their northward limit of distribution in the Gulf of Cadiz or in the Portuguese coast (Drake et al. 1998; García Raso & Manjón-Cabeza 1996). However, since at least one of the hosts of *A. monodi* is a fouling organism (i.e. mussel), its arrival to the European coasts could have occurred by its accidental transport in ships' fouling (Apte et al. 2000). Regardless of the pathway through which species coming from warmer latitudes are introduced in northern habitats, the reproductive cycles of their northernmost populations are usually determined by temperature. Namely, the breeding periods of these species in the Gulf of Cadiz are usually shorter (falling in the warmest months) than in their African populations (González-Gordillo et al. 1990; Rodríguez et al. 1997). However, larval stages (Drake et al. 1998) and ovigerous females of *A. monodi* (this study) were found all year round in the studied area, in contrast with the shorter (mainly in summer) reproductive activity of *N. pinnotheres* and *P. pisum* (Zariquiey Álvarez 1968; Becker 2010). Furthermore, the autumnal decrease of the reproductive activity of *A. monodi* observed in the studied area could disappear in the near future if the seawater temperature of the Mediterranean area keeps increasing (Sanderson et al. 2011). Likewise, *A. monodi* has an extended development, with five larval stages (Marco-Herrero et al. in prep), large enough to facilitate dispersal and colonization of new areas (Abelló et al. 2003). Taken together, all these features may have contributed to the quick and successful settlement of *A. monodi* in the Gulf of Cadiz, possibly to the detriment of *N. pinnotheres* and *P. pisum* populations.

Both males and females of *Afropinnotheres monodi* tend to live alone inside their host. Two hypotheses have been proposed to explain why the one crab per host relationship is recurrent among other pea crab species: 1) the crab releases a chemical cue that discourages conspecific guests from entering the same host; 2) a prior resident crab might show aggressive behaviour toward a posterior invading one (Bell 1984; Haines et al. 1994; Soong 1997; Takeda et al. 1997). Despite that some species of pea crabs seem to show chemoreception (Stevens 1990; Ambrosio & Brooks 2011), when *A. monodi* individuals were retrieved from their hosts and immediately placed together in an aquarium in presence of living hosts, multiple infestation by crabs was more frequent than in field observations (Drake et al. unpublished data). This finding suggests that *A. monodi* crabs cannot detect the presence of conspecifics in the host, or, in some circumstances (i.e. more guests than available hosts) the presence of conspecifics is not a limiting factor. In addition, in the few pairs of crabs observed inside the studied hosts, heterosexual pairs were not more frequent than expected by chance alone. Thus, mating in this species might take place outside hosts or, if occurring inside hosts, might last a very short period of time, after which males abandon the temporarily shared host (Hamel et al. 1999; Ocampo et al. 2012; Peiró et al. 2013). That is, the dominance of solitary guest crabs in this population of *A. monodi* suggests that the selection of the host in this species tends to be random (opportunity-related) and, consequently, the occasional pairs observed inside hosts were accidental and transitory. This host monopolisation is an efficient adaptation (more benefits than costs) to hosts with small body size and/or low morphological complexity (Baeza & Thiel 2003, 2007; Hernández et al. 2012), as the host bivalves in this study. In fact, mussels, the host that harboured mainly soft females, showed fewer female-female pairs than expected by chance. As males and hard females have significantly smaller body size (Figures 2 and 6) and

stronger swimming capacity than soft females, they may roam among host in search of empty hosts (both sexes) or receptive females (only males). In contrast, soft females seem to be totally adapted to a symbiotic lifestyle and never leave the host (De Bruyn et al. 2009; Becker et al. 2011, 2012). Thus, from the point of view of costs and benefits, host monopolization could be more relevant for reproductive females than for the remaining demographic categories when host resources (space, food and oxygen) are limiting (Baeza et al. 2002; Narvarte & Saiz 2004; Baeza & Thiel 2007; De Bruyn et al. 2009).

Like the rest of pea crabs recorded in European coasts (Becker & Türkay 2010), *Afropinnotheres monodi* is not host-specific (Subida et al. 2011) and has been found in a high number of bivalve species, resembling the pattern shown by *Pinnotheres pisum*. Indeed, in the Bay of Cadiz symbiotic males and females of this African species were found throughout the year inside the three studied hosts. However, there were clear differences in the global (total number of crabs) and specific (by demographic categories) prevalence of *A. monodi* in each host (Table 1 and Figure 2). Results of laboratory experiments, which aimed to assess host preferences in other species of pea crabs, did not always match the host preferences predicted from field observations of the same species (Ocampo et al. 2012; De Bruyn et al. 2010). Therefore, if we assume that in the field the *A. monodi* distribution on the different hosts was random, the prevalence in each host should be proportional to the corresponding host accessibility and availability. From the three studied hosts, *Scrobicularia plana* clams are the host showing less accessibility to crab guests since they bury deeply in the sediment (more difficult to be reached by crabs) of the higher intertidal zone (more time emerged). In fact *S. plana* was the host in which *A. monodi* showed the lowest prevalence. However, the varying habitat characteristics of the three hosts alone do not explain the

1 asymmetrical host usage by each demographic category observed for the studied pea
2 crab. Under the assumption that pea crabs are optimally adapted to the environment,
3 their current performance must be determined by costs and benefits associated to the
4 different behaviours (Baeza & Thiel 2007). In the case of *A. monodi*, we hypothesize
5 that the different host uses observed could be triggered by an ontogenetic change in the
6 swimming behaviours and habitat preferences of crabs. That is, if megalopae of this
7 species tend to choose the soft sediment of sheltered sand-muddy flats for their benthic
8 settlement (Olague-Feliú et al. 2010), immature males and females of *A. monodi* could
9 be more likely to infest benthic species inhabiting the sediment, as cockles and clams,
10 which are small in size (low empty space in the shell cavity) but abundant and protected
11 from predators. However, when the pea crabs grow, their swimming capability increases
12 favouring the adoption of a temporal free roaming behaviour (De Bruyn et al. 2009;
13 Jossart et al. 2013). Although the risk of predation is higher for pea crabs out of their
14 hosts, to get refuge in larger hosts as mussels is a relevant benefit (increase of resources)
15 for future ovigerous females. Assuming that males are able to distinguish between
16 female stages (Diesel 1988; De Bruyn et al. 2009; Ambrosio & Brooks 2011), they
17 could increase their mating opportunities by living near hosts infested with receptive
18 females. Indeed, *A. monodi* males were found in the 1:1 sex ratio only on the two host
19 species harbouring mainly hard/receptive females (clams and cockles), whereas soft
20 females were found mainly in mussels. An analogous asymmetrical host usage by the
21 different demographic categories has been reported for the pea crab *Fabia subquadrata*
22 in the Puget Sound (Washington, USA) area (Garth & Abbot 1980). With the available
23 information, the life history of *A. monodi* hypothesised herein is the most plausible to
24 explain the results obtained in the Bay of Cádiz. However, in the absence of a fully
25 simultaneous sampling of the studied hosts and of experimental assays demonstrating

1 the movement of pea crabs between hosts, the inferences concerning this aspect of the
2 guest's autoecology should be considered with caution.

3 As for other pea crab species (Christensen & McDermott 1958; Soong 1997;
4 Kane & Farley 2006; Ocampo et al. 2012), there was a strong correlation between host
5 size and symbiotic ovigerous female size of *Afropinnotheres monodi*, which suggests
6 that space availability within hosts is a relevant factor in determining the final size of
7 this sedentary phase (soft females) of symbiotic crabs and, consequently, of its
8 reproductive pattern. Similarly, a positive relationship between the fecundity and the
9 body size of female pea crabs, previously reported for other species (Bell & Stancyk
10 1983; Baeza & Thiel 2000; De Bruyn et al. 2010), has been observed for *A. monodi*:
11 large ovigerous females found inside mussels have much higher fecundity than the
12 scarce and small ovigerous females inhabiting clams and cockles (Figure 7).
13 Nevertheless, when the body size of female crabs was taken into account, differences in
14 size-specific fecundity were only observed between females inhabiting clams and those
15 from the remaining two host species, suggesting that the flattened morphology of the
16 shell of *Scrobicularia plana* makes this host rather unsuitable for the almost-spherical
17 ovigerous female pea crabs. That is, the size of reproductive females that use cockles as
18 a host is constrained by the relatively small size of this host, and crabs inhabiting clams
19 also show a decrease of the size-specific fecundity. Thus, even if all demographic
20 categories of *A. monodi* could be observed inside the three studied hosts, the
21 reproductive effort of this crab would still correspond mainly to females inhabiting
22 mussels. Moreover, large body size ovigerous females of *A. monodi* showed higher
23 (Bell & Stancyk 1983; Hamel et al. 1999; Baeza & Thiel 2000; De Bruyn et al. 2010) or
24 similar (Silas & Alagarwami 1967; Ocampo et al. 2012) fecundity than other pea crab
25 species. However, the input of new recruits to the adult population occurred mainly in

clams and cockles, which convert these smaller-size hosts in a relevant step to the establishment of abundant and stable populations of *A. monodi* in the studied area. According to Debruyne et al. (2010), this type of asymmetrical usage of hosts by parasitic pea crabs may be comparable with the transitory stages predicted by models of evolution of complex parasite life cycles (Parker et al. 2003). Namely, the asymmetrical usage of clams, cockles and mussels by *A. monodi* in the Gulf of Cadiz resembles the transitional stage predicted by the “downward incorporation” of a new host into the life cycle of this pea crab. In any case, the generalist but asymmetrical usage of bivalve hosts found in *A. monodi* may have clear benefits for species in geographical expansion: first, the presence of all demographic categories in the three hosts can facilitate the settlement of the species in new locations; second, the asymmetrical use of different hosts can lead to the settlement of copious and stable populations in some areas, as complex estuarine habitats in the case of *A. monodi*, which act as a massive source of pelagic larvae with high capacity of dispersal. Thus, the use of more than one host species by symbiotic species may increase its fitness and, simultaneously, shape their life cycle and evolution (De Bruyne et al. 2010). Nevertheless, the incorrect election of the definitive host by females may become a disadvantage: after metamorphosis, reproductive females become too immobile to leave the host and/or reach and enter a new one. In fact, during the study, we have collected died mussels whose valves harboured large ovigerous females of *A. monodi* in unhealthy condition or even dead.

Several species of pinnotherids worldwide parasitize commercially exploited bivalves (Stauber 1945; Christensen & McDermott 1958; Silas & Alagaraswami 1967; Sun et al. 2006). In shellfish farms, a significant loss of production has been observed even with low levels of pea crab infection (Trottier et al. 2012). Furthermore, pea crabs may rapidly colonize bivalves from a shellfish farm, and subsequently infect nearby

1 shellfish farms (Trottier & Jeffs 2012). However, considering the asymmetrical use of
2 different hosts by *A. monodi*, we expect that the strongest threats may be posed to
3 shellfish farms located in sheltered waters (bays, inlets, rías, harbours, etc.), where
4 populations of the different hosts used by this African species coexist. As the species
5 seems to be in clear northward expansion, it may become, in the near future, a
6 significant threat for European bivalve aquaculture. Thus, the confirmation of the
7 hypotheses presented in this study, by experimental essays, is of high social and
8 economic interest.

9 In conclusion, the African pea crab *Afropinnotheres monodi* seems to be a
10 symbiotic species currently in geographical expansion showing a series of features in its
11 life cycle (wide reproductive period, high fecundity, long planktonic phase, generalist
12 and asymmetric host use), that facilitate its quick and successful settlement in new
13 locations. Both characteristics make this African species particularly relevant to
14 studying the evolution of pea crab symbiotic life history traits.

17 **Aknowledgments**

18 We thank M Espigares and A Moreno for their help in field collection and sample
19 analysis, and the three anonymous reviewers, which provided us useful suggestions for
20 the elaboration of this final version. This research work was funded by the remnant
21 funds of the PAI RNM108 research group.

23 **References**

- 1 Abelló P, Visauta E, Bucci A, Demestre M (2003) Noves dades sobre l'expansió
- 2 del cranc *Percnon gibbesi* (Brachyura: Grapsidae: Plagusinae) a la Mediterrània
- 3 occidental. Boll Soc Hist Nat Balears 46:73–77.
- 4 Ambrosio LJ, Brooks WR (2011) Recognition and use of ascidian hosts, and mate
- 5 acquisition by the symbiotic pea crab *Tunicotheres moseri* (Rathbun, 1918): the role
- 6 of chemical, visual and tactile cues. Symbiosis 53:53–61
- 7 Apte S, Holland BS, Godwin LS, Gardner JPA (2000) Jumping ship: a stepping stone
- 8 event mediating transfer of non-indigenous species via a potentially unsuitable
- 9 environment. Biological Invasions 2:75–79.
- 10 Asama H, Yamaoka K (2009) Life history of the pea crab, *Pinnotheres sinensis*, in
- 11 terms of infestation in the bivalve mollusc, *Septifer virgatus*. Marine Biodiversity
- 12 Records 2:e77
- 13 Baeza JA, Thiel M (2000) Host use pattern and life history of *Liopetrolisthes mitra*, a
- 14 crab associate of the black sea urchin *Tetrapygus niger*. J Mar Biol Ass UK 80:639–
- 15 645
- 16 Baeza JA, Díaz-Valdes M (2011) The symbiotic shrimp *Ascidonia flavomaculata* lives
- 17 solitarily in the tunicate *Ascidia mentula*: implications for its mating system.
- 18 Invertebr Biol 130:351–361
- 19 Baeza JA, Thiel M (2003) Predicting territorial behaviour in symbiotic crabs using host
- 20 characteristics: a comparative study and proposal of a model. Mar Biol 142:93–100
- 21 Baeza JA, Thiel M (2007) The mating system of symbiotic crustaceans: a conceptual
- 22 model based on optimality and ecological constraints. In: Duffy JE, Thiel M (eds)
- 23 Evolutionary ecology of social and sexual systems. Crustaceans as model organisms.
- 24 Oxford University Press, Oxford, p 249–267

- 1 Baeza JA, Stotz W, Thiel M (2002) The agonistic behaviour and development of
- 2 territoriality during ontogeny of the sea anemone dwelling crab *Allopetrolisthes*
- 3 *spinifrons*. Mar Freshw Behav Physiol 35:189–202
- 4 Becker C (2010) European pea crabs - taxonomy, morphology, and host-Ecology
- 5 (Crustacea: Brachyura: Pinnotheridae). PhD thesis, Goethe-University (Frankfurt)
- 6 Becker C, Türkay M (2010) Taxonomy and morphology of European pea crabs
- 7 (Crustacea: Brachyura: Pinnotheridae). J Nat Hist 44:1555–1575
- 8 Becker C, Brandis D, Storch V (2011) Morphology of the female reproductive system
- 9 of European pea crabs (Crustacea, Decapoda, Brachyura, Pinnotheridae). J Morphol
- 10 272:12-26
- 11 Becker C, Türkay M, Brandis D (2012) The male copulatory system of European pea
- 12 crabs (Crustacea, Brachyura, Pinnotheridae). J Morphol 273:1306-1318
- 13 Bell JL (1984) Changing residence: dynamics of the symbiotic relationship between
- 14 *Dissodactylus mellitae rathbun* (Pinnotheridae) and *Mellita quinquiesperforata*
- 15 (Leske) (Echinodermata) J Exp Mar Biol Ecol 82:101-115
- 16 Bell JL, Stancyk SE (1983) Population dynamics and reproduction of *Dissodactylus*
- 17 *mellitae* (Brachyura: Pinnotheridae) on its sand dollar host *Mellita*
- 18 *quinquiesperforata* (Echinodermata). Mar Ecol Prog Ser 13:141–149
- 19 Bradbury IR, Snelgrove PVR (2001) Contrasting larval transport in demersal fish and
- 20 benthic invertebrates: the roles of behaviour and advective processes in determining
- 21 spatial pattern. Can J Fish Aquat Sci 58:811-823
- 22 Brown JH, Stevens GC, Kaufman DM (1996) The geographic range: size, shape,
- 23 boundaries, and internal structure. Annu Rev Ecol Syst 27:597–623
- 24 Cabral HN, Costa MJ, Salgado JP (2001) Does the Tagus estuary fish community
- 25 reflect environmental changes? Clim Res 18:119-126

- 1 Castañeda E, Drake P (2008) Spatio-temporal distribution of *Lekanesphaera* species in
2 relation to estuarine gradients within a temperate European estuary (SW Spain) with
3 regulated freshwater inflow. *Cienc Mar* 34:125-141
- 4 Christensen AM, McDermott JJ (1958) Life-history and biology of the oyster crab,
5 *Pinnotheres ostreum* Say. *Biol Bull Woods Hole* 114:146-179
- 6 De Bruyn C, Rigaud T, David B, De Ridder C (2009) Nature and consequences of the
7 symbiotic relationship between the crab *Dissodactylus primitivus* and its echinoid
8 host *Meoma ventricosa*. *Mar Ecol Prog Ser* 375:173–183
- 9 De Bruyn C, David B, De Ridder C, Rigaud T (2010) Asymmetric exploitation of two
10 echinoid host species by a parasitic pea crab and its consequences for the parasitic
11 life cycle. *Mar Ecol Prog Ser* 398:183–191
- 12 Diesel R (1988) Male-female association in the spider crab *Inachus phalangium*: the
13 influence of female reproductive stage and size. *J Crust Biol* 8:63-69.
- 14 Doney SC, Ruckelshaus M, Duffy JE, Barry JP, Chan F, English CA, Galindo HM,
15 Grebmeier JM, Hollowed AB, Knowlton N, Polovina J, Rabalais NN, Sydeman WJ,
16 Talley LD (2012) Climate change impacts on marine ecosystems. *Annu Rev Mar Sci*
17 4:11–37
- 18 Drake P, Arias AM, Rodríguez A (1998) Seasonal and tidal abundance patterns of
19 decapod crustacean larvae in a shallow inlet (SW Spain). *J Plankton Res* 20:585-601
- 20 Góes JM, Fransozo A, Fernández-Góes LC (2005) Fecundity of *Eriphia gonagra*
21 (Fabritius, 1781) (Crustacea, Brachyura, Xanthidae) in the Ubatuba region, São
22 Paulo Brazil. *Nauplius* 13:127-136
- 23 García Raso JE, Manjón-Cabeza E (1996) New record of *Liocarcinus mcleayi* (Barnard,
24 1947), new combination (Decapoda, Brachyura, Portunidae) from South Europe.
25 *Crustaceana* 69:84-93

- 1 Garth JS, Abbott DP (1980) Brachyura: the true crabs. In R. H. Morris, D. P. Abbott
2 and E. C. Haderlie, eds., Intertidal Invertebrates of California, Stanford Univ Press,
3 Stanford, p 594-630.
- 4 González-Gordillo JI, Cuesta JA, Pablos F (1990) Adiciones al conocimiento de los
5 crustáceos decápodos de las zonas mediolitoral e infralitoral de las costas
6 suratlánticas andaluzas (Suroeste España). 1. Brachyura. Cah Biol Mar 31:417-429
- 7 González-Gordillo JI, Dos Santos A, Rodríguez A (2001) Checklist and annotated
8 bibliography of decapod crustacean larvae from the Southwestern European coast
9 (Gibraltar Strait area). Sci Mar 65:275-305
- 10 González-Gordillo JI, Dos Santos A, Rodríguez A (2003a) Comparative seasonal and
11 spatial distribution of decapod larvae assemblages in three coastal zones off the
12 south-western Iberian Peninsula. Acta Oecologica 24:219-233
- 13 González-Gordillo JI, Arias AM, Rodríguez A, Drake P (2003b) Recruitment patterns
14 of decapod crustacean megalopae in a shallow inlet (SW Spain) related to life history
15 strategies. Est Coast Shelf Sci 56:593–607
- 16 Grove MW, Finelli CM, Wetthey DS, Woodin SA (2000) The effects of symbiotic crabs
17 on the pumping activity and growth rates of *Chaetopterus variopedatus*. J Exp Mar
18 Bio Ecol 246:31-52
- 19 Grove MW, Woodin SA (1996) Conspecific recognition and host choice in a pea crab,
20 *Pinnixa chaetoptera* (Brachyura: Pinnotheridae). Biol Bull 190:359-366
- 21 Haines CMC, Edmunds M, Pewsey AR (1994) The pea crab, *Pinnotheres pisum*
22 (Linnaeus, 1767), and its association with the common mussel, *Mytilus edulis*
23 (Linnaeus, 1758), in the Solent (UK). J Shellfish Res 13:5–10

- 1 Hamel JF, Ng PKL, Merrier A (1999) Life cycle of the pea crab *Pinnotheres halingi* sf.
2 nov., an obligate symbiont of the sea cucumber *Holothuria scabra* Jaeger. *Ophelia*
3 50:149-175
- 4 Hernández JE, Bolaños JA, Palazón JL, Hernández G, Lira C, Baeza JA (2012) The
5 enigmatic life history of the symbiotic crab *Tunicotheres moseri* (Crustacea,
6 Brachyura, Pinnotheridae): implications for its mating system and population
7 structure. *Biol Bull* 223:278–290.
- 8 Jones MC, Dye SR, Fernandes JA, Frölicher TL, Pinnegar JK (2013) Predicting the
9 Impact of Climate Change on Threatened Species in UK Waters. *PLoS ONE*
10 8:e54216
- 11 Jossart Q, David B, De Bruyn C, De Ridder C, Rigaud T, Wattier RA (2013) No
12 evidence of host specialization in a parasitic pea-crab exploiting two echinoid hosts.
13 *Mar Ecol Prog Ser* 475:167–176
- 14 Kane K, Farley GS (2006) Body size of the endosymbiotic pea crab *Tumidotheres*
15 *maculatus*: larger hosts hold larger crabs. *Gulf and Caribbean Research* 18:27–33
- 16 López de la Rosa I, García Raso JE, Rodríguez A (2002) Evolution of a decapod
17 community (Crustacea) of shallow soft bottoms with seaweeds from southern
18 Europe. *J Mar Biol Ass UK* 82:85-95
- 19 Narvarte MA, Saiz MN (2004) Effects of the pinnotherid crab *Tumidotheres maculatus*
20 on the Tehuelche scallop *Aequipecten tehuelchus* in the San Matías Gulf, Argentina.
21 *Fish Res* 67:207–214
- 22 Ocampo EH, Nuñez JD, Cledón M, Baeza JA (2012) Host specific reproductive
23 benefits, host selection behaviour and host use pattern of the pinnotherid crab
24 *Calyptraeothers garthi*. *J Exp Mar Biol Ecol* 429:36–46

- 1 Olaguer-Feliú AO, Flores AAV, Queiroga H, González-Gordillo JI (2010) Shelf and
2 estuarine transport mechanisms affecting the supply of competent larvae in a suite of
3 brachyuran crabs with different life histories. *Mar Ecol Prog Ser* 410:125–141
- 4 Parker GA, Chubb JC, Ball MA, Guy N, Roberts GN (2003) Evolution of complex life
5 cycles in helminth parasites. *Nature* 425:480–484
- 6 Parmesan C (2006) Ecological and evolutionary responses to recent climate change.
7 *Annu Rev Ecol Evol Syst* 37:637–669
- 8 Peiró DF, Baeza JA, Mantelatto FL (2013) Host-use pattern and sexual dimorphism
9 reveals the mating system of the symbiotic pea crab *Austinixa aidae* (Crustacea:
10 Brachyura: Pinnotheridae). *J Mar Biol Ass UK* 93:715–723
- 11 Prista N, Vasconcelosa RP, Costa MJ, Cabrala H (2003) The demersal fish assemblage
12 of the coastal area adjacent to the Tagus estuary (Portugal): relationships with
13 environmental conditions. *Oceanol Acta* 26:525–535
- 14 Rodríguez A, Drake P, Arias AM (1997) Reproductive periods and larval abundance
15 patterns of the crabs *Panopeus africanus* and *Uca tangeri* in a shallow inlet (SW
16 Spain). *Mar Ecol Prog Ser* 149:133–142
- 17 Sanderson MG, Hemming DL, Betts RA (2011) Regional temperature and precipitation
18 changes under high-end ($\geq 4^{\circ}\text{C}$) global warming. *Phil Trans R Soc A* 369:85–98
- 19 Scheltema RS (1986) On dispersal and planktonic larvae of benthic invertebrates: an
20 eclectic overview and summary of problems. *Bull Mar Sci* 39:290–322
- 21 Silas EG, Alagarwami K (1967) On an instance of parasitisation by the pea-crab
22 (*Pinnotheres* sp.) on the backwater clam (*Meretrix casta* (Chemnitz)) from India,
23 with a review of the work on the systematics, ecology, biology and ethology of pea
24 crabs of the genus *Pinnotheres* Latreille. *Proc Mar Biol Ass India* (symposium series
25 2) 3:1161–1227

- 1 Schmitt WL, McCain JC, Davidson ES (1973) Decapoda I, Brachyura I, family
- 2 Pinnotheridae. In: Gruner HE, Holthuis LB (eds) Crustaceorum Catalogus, 3:1-160
- 3 Soong K (1997) Some life history observations on the pea crab, *Pinnotheres*
- 4 *tsingtaoensis*, symbiotic with the bivalve mollusc, *Sanguinolaria acuta*. Crustaceana
- 5 70:855–866
- 6 Stauber L (1945) *Pinnotheres ostreum*, parasitic on the American oyster, *Ostrea*
- 7 (*Gryphaea*) *virginica*. Biol Bull 88:269–291
- 8 Stevens PM (1990) Specificity of host recognition of individuals from different host
- 9 races of symbiotic pea crabs (Decapoda: Pinnotheridae). J Exp Mar Biol Ecol
- 10 143:193–207
- 11 Subida MD, Arias AM, Drake P, García-Raso JE, Rodríguez A, Cuesta JA (2011) On
- 12 the occurrence of *Afropinnotheres monodi* Manning, 1993 (Decapoda:
- 13 Pinnotheridae) in European waters. J Crust Biol 31:367-369
- 14 Sun W, Sun S, Yuqi W, Baowen Y, Weibo S (2006) The prevalence of the pea crab,
- 15 *Pinnotheres sinensis*, and its impact on the condition of the cultured mussel, *Mytilus*
- 16 *galloprovincialis*, in Jiaonan waters (Shandong Province, China) Aquaculture
- 17 253:57-63
- 18 Takeda S, Tamura S, Washio M (1997) Relationship between the pea crab *Pinnixa*
- 19 *tumida* and its endobenthic holothurian host *Paracaudina chilensis*. Mar Ecol Prog
- 20 Ser 149:143-154
- 21 Trottier O, Jeffs AG (2012) Biological characteristics of parasitic *Nepinnotheres*
- 22 *novaezelandiae* within a *Perna canaliculus* farm. Dis Aquat Org 101:61-68
- 23 Trottier O, Walker D, Jeffs AG (2012) Impact of the parasitic pea crab *Pinnotheres*
- 24 *novaezelandiae* on aquacultured New Zealand green-lipped mussels, *Perna*
- 25 *canaliculus*. Aquaculture 344/346:23–28

- 1 Udekem D'Acoz, C (1999) Inventaire et distribution des crustacés décapodes de
- 2 l'Atlantique nord-oriental, de la Méditerranée et des eaux continentales adjacentes au
- 3 nord de 25 N. Collection Patrimoines Naturels, 40 Muséum national d'Histoire
- 4 naturelle: Paris. ISBN 2-86515-114-10. X
- 5 Zar JH (2010) Biostatistical Analysis. Prentice-Hall, Englewood Cliffs, New Jersey
- 6 Zariquiey Álvarez R (1968) Crustáceos decápodos ibéricos. Inv Pesq 32:1-236
- 7

Table 1. Mean prevalence of the pea crab *Afropinnotheres monodi* in the bivalves *Scrobicularia plana*, *Cerastoderma glaucum* and *Mytilus galloprovincialis*. For each host species and demographic crab category, host shell length range (shell cavity volume range), mean crab carapace width (carapace width range) and Pearson correlation coefficient between the crab carapace width (CW) and the host shell length (SL) are presented. Host shell lengths and crab carapace widths in mm; shell volume in ml.

	<i>Scrobicularia plana</i>	<i>Cerastoderma glaucum</i>	<i>Mytilus galloprovincialis</i>
Prevalence % (n° hosts)	4.1 (9441)	41.1 (2661)	45.9 (868)
Host shell length (Vol)			
Male	22.0-37.5 (1.0-4.0)	15.1-36.4 (0.8-7.2)	14.0-44.3- (1.3-24.5)
Hard Female	17.1-36.8 (0.5-3.8)	14.1-35.1 (0.7-6.6)	13.1-37.2 (1.1-16.7)
Soft Female	27.0-35.5 (1.7-3.5)	21.8-35.2 (2.0-6.6)	14.6-49.8 (1.4-35.5)
Crab carapace width			
Male	2.93 (1.80-3.80)	2.88 (1.35-7.33)	4.66 (2.01-9.83)
Hard Female	2.88 (1.39-4.65)	2.78 (0.65-6.83)	3.29 (2.27-5.50)
Soft Female	4.41 (3.70-5.53)	5.61 (3.77-8.67)	9.29 (3.33-13.67)
CW-SL correlation			
Male	0.13 (ns)	0.21 (*)	0.37 (*)
Hard Female	0.21 (ns)	0.04(ns)	0.18 (ns)
Soft Female	0.04 (ns)	0.34 (*)	0.63 (**)

Correlation significance level: ns, $P > 0.05$; *, $P < 0.05$; **, $P < 0.01$

Figure Legends

Fig. 1. Sampling sites of the bivalve host populations harbouring the pea crab *Afropinnotheres monodi* in the Bay of Cadiz (SW Iberian Peninsula).

Fig. 2. Population structure (size and demographic category), of the pea crab *Afropinnotheres monodi* in the three studied hosts. SF: soft female; HF: hard female; M: male. Sex ratio of pea crabs harboured by the mussel *Mytilus galloprovincialis* was significantly different from 1:1.

Fig. 3. Seasonal pattern in the occurrence of each demographic category of the pea crab *Afropinnotheres monodi* in the three host species. OF: ovigerous soft female; SF: non-ovigerous soft female; HF: hard female; M: male. Numbers on x-axis (in parentheses): numbers of pea crabs collected. Dates below host species names (in parentheses): sampling periods of the corresponding host.

Fig. 4 Population distribution of the pea crab *Afropinnotheres monodi* in the three host species. Observed frequency of occurrence of crabs in *Cerastoderma edule* and *Mytilus galloprovincialis* differs significantly from the expected in a random distribution.

Fig. 5 Percentage of occurrence of each demographic category of the pea crab *Afropinnotheres monodi* in each host size class (upper row) and frequency distribution of pea crabs in the different size classes of each host species *Scrobicularia plana*, *Cerastoderma edule* and *Mytilus galloprovincialis* (lower row).

Fig. 6 For each host species and pea crab demographic category, relationship between the shell length/shell volume cavity of host and the carapace width (CW) of the pea crab *Afropinnotheres monodi*. Below, relationship between shell length (SL) and shell volume cavity (SV) of the hosts *Scrobicularia plana* (square), *Cerastoderma edule* (triangle) and *Mytilus galloprovincialis* (circle). Sex ratio of *Afropinnotheres monodi* retrieved from small hosts (shell volume cavity < 5 ml) was significantly different from 1:1 in the mussel *Mytilus galloprovincialis*.

Fig. 7. Relationship between the crab carapace width (CW) and the number of eggs (No) carried by ovigerous female of *Afropinnotheres monodi* retrieved from the bivalves *Scrobicularia plana* (square), *Cerastoderma edule* (triangle) and *Mytilus galloprovincialis* (circle). The filling colour of symbols indicates the season of host and pea crab collection (open: spring; light-grey: summer; black: autumn; dark-grey: winter).

Figure 1

1
2 Figure 2

1
2 Figure 3

Figure 4

1

2
Figure 5

1

2
3
4

Figure 6

1
2 Figure 7
3