

Hybrid Quantization of Inflationary Universes

Guillermo A. Mena Marugán

Instituto de Estructura de la Materia, CSIC
(Mikel Fernández-Méndez,
Javier Olmedo & José Velhinho)

JARRAMPLAS
23 March 2013



CSIC

CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

The model




- We consider **perturbed** FRW universes filled with a **massive** scalar field.
- The scalar field is minimally coupled.
- The model can generate inflation.

- The most interesting case is flat spatial topology. It is also the simplest.
- The effects of **spatial curvature** can be studied by considering, e.g., spherical topology.
- We assume **compact** spatial sections.

The model

It's been well studied, even in LQC, though...

- Anomalies: Incorporate quantum effects, not the starting point for quantization.
- Effective dynamics: Needs a true derivation.



- **Approximations:** As few as possible. Should be derived or at least checked for consistency.

- In many cases these checks are only internal, within the approximated description.

Perturbations about flat FRW

- Truncation at quadratic order in the action.
- Includes backreaction at that order.
- Tests the validity of less refined truncations and provides the way to develop **approximation** methods, controlling their range of application.

Hybrid approach

Effects of quantum geometry are only accounted for in the background

- Successfully applied in Gowdy cosmologies.
- In those cases there is no truncation. This is no drawback (think of the harmonic oscillator).
- In the present case, we only deal with the quadratically perturbed model.

Uniqueness of the Fock description

- Infinite **ambiguity** in selecting a Fock representation in QFT in curved space-times.
- This can be restricted by appealing to *background symmetries*.
- Typically this is not sufficient in non-stationarity.
- Proposal: demand the **UNITARITY** of the quantum evolution.
The conventional interpretation of QM is guaranteed.
This goes beyond the viewpoint of algebraic quantizations.
- There is a natural ambiguity in the **separation of the background** from the field.
In cosmology, this introduces time-dependent canonical field transformations.
- Remarkably, symmetry invariance and dynamical unitarity select a **UNIQUE canonical pair** and a UNIQUE Fock representation for their CCR's.

Uniqueness of the Fock description




Uniqueness of the Fock description



- Recent works **DO NOT incorporate** the correct scaling (**AA&N**). This affects the quantum description, and in particular the *effective* approaches therein derived.
- Moreover, one can even consider non-local canonical transformations, respecting the decoupling of field modes.

The **UNIQUENESS** of the quantization, up to unitary equivalence, is guaranteed.

Loop Quantum FRW Cosmology




- Avoids the Big Bang.
- Specific **proposal** such that:
 - Evolution can be defined even without ideal clocks (massless field).
 - The WdW limit is unambiguous in each superselection sector.
 - It is optimal for numerical computation.
- Control of changes of densitization in the scalar constraint.
The lapse function is **not a function on phase space**.

Classical system: FRW

- Massive scalar field minimally coupled to a compact, flat FRW universe.

Geometry: $A_a^i = c^0 e_a^i (2\pi)^{-1}; \quad E_i^a = p \sqrt{0} e^0 e_i^a (2\pi)^{-2}. \quad \{c, p\} = 8\pi G \gamma / 3.$



$$a^2 = e^{2\alpha} = [p] (2\pi \sigma)^{-2}; \quad \pi_\alpha = -pc (\gamma 8\pi^3 \sigma^2)^{-1}. \quad \sigma^2 = G (6\pi^2)^{-1}.$$

Matter:

$$\varphi = (2\pi)^{3/2} \sigma \phi; \quad \pi_\varphi = (2\pi)^{-3/2} \sigma^{-1} \pi_\phi.$$

Hamiltonian constraint:

$$C_0 = -\frac{6}{\gamma^2} \sqrt{|p|} c^2 + \frac{8\pi G}{V} (\pi_\phi^2 + m^2 V^2 \phi^2).$$

$$V = [p]^{3/2}.$$



Classical system: Modes

- We expand inhomogeneities in a (real) **Fourier basis**:

$$Q_{\vec{n},+} = \frac{1}{2\pi^{3/2}} \cos \vec{n} \cdot \vec{\theta}, \quad Q_{\vec{n},-} = \frac{1}{2\pi^{3/2}} \sin \vec{n} \cdot \vec{\theta}. \quad \vec{n} \in \mathbb{Z}^3, \quad n_1 \geq 0.$$

- The basis is **orthonormal**, and we exclude the zero mode in the expansions.
- These functions are eigenmodes of the Laplace-Beltrami operator of the standard flat metric on the three-torus, with eigenvalue

$$-\omega_n^2 = -\vec{n} \cdot \vec{n}.$$

- We only consider **scalar perturbations**: decoupled from vector and tensor perturbations at dominant order.

Classical system: Inhomogeneities

- Mode expansion of the inhomogeneities:

$$h_{ij} = (\sigma e^\alpha)^2 \left[{}^0 h_{ij} + 2\epsilon (2\pi)^{3/2} \sum \left\{ a_{\vec{n},\pm}(t) Q_{\vec{n},\pm} {}^0 h_{ij} + b_{\vec{n},\pm}(t) \left(\frac{3}{\omega_n^2} (Q_{\vec{n},\pm})_{,ij} + Q_{\vec{n},\pm} {}^0 h_{ij} \right) \right\} \right],$$

$$N = \sigma N_0(t) \left[1 + \epsilon (2\pi)^{3/2} \sum g_{\vec{n},\pm}(t) Q_{\vec{n},\pm} \right],$$

$$N_i = \epsilon (2\pi)^{3/2} \sigma^2 e^\alpha \sum \frac{k_{\vec{n},\pm}(t)}{\omega_n} (Q_{\vec{n},\pm})_{,i},$$

$$\Phi = \frac{1}{\sigma} \left[\frac{\varphi(t)}{(2\pi)^{3/2}} + \epsilon \sum f_{\vec{n},\pm}(t) Q_{\vec{n},\pm} \right].$$

The corrections *include* in principle higher-order perturbations.

Classical system: Action

- Truncating the action at **quadratic** order in perturbations, one obtains:

$$H = \frac{N_0 \sigma}{16\pi G} C_0 + \epsilon^2 \sum \left(N_0 H_2^{\vec{n}, \pm} + N_0 g_{\vec{n}, \pm} H_1^{\vec{n}, \pm} + k_{\vec{n}, \pm} \tilde{H}_1^{\vec{n}, \pm} \right),$$

$$\begin{aligned} H_2^{\vec{n}, \pm} 2e^{3\alpha} = & -\pi_{a_{\vec{n}, \pm}}^2 + \pi_{b_{\vec{n}, \pm}}^2 + \pi_{f_{\vec{n}, \pm}}^2 + 2\pi_\alpha \left(a_{\vec{n}, \pm} \pi_{a_{\vec{n}, \pm}} + 4b_{\vec{n}, \pm} \pi_{b_{\vec{n}, \pm}} \right) - 6\pi_\varphi a_{\vec{n}, \pm} \pi_{f_{\vec{n}, \pm}} \\ & + \pi_\alpha^2 \left(\frac{1}{2} a_{\vec{n}, \pm}^2 + 10b_{\vec{n}, \pm}^2 \right) + \pi_\varphi^2 \left(\frac{15}{2} a_{\vec{n}, \pm}^2 + 6b_{\vec{n}, \pm}^2 \right) - \frac{e^{4\alpha}}{3} \left[\omega_n^2 a_{\vec{n}, \pm}^2 + (\omega_n^2 - 18)b_{\vec{n}, \pm}^2 \right] \\ & + e^{4\alpha} \omega_n^2 \left[f_{\vec{n}, \pm}^2 - \frac{2}{3} a_{\vec{n}, \pm} b_{\vec{n}, \pm} \right] + e^{6\alpha} m^2 \sigma^2 \left[\varphi^2 \left(\frac{3}{2} a_{\vec{n}, \pm}^2 + 6b_{\vec{n}, \pm}^2 \right) + 6\varphi a_{\vec{n}, \pm} f_{\vec{n}, \pm} + f_{\vec{n}, \pm}^2 \right], \end{aligned}$$

$$\begin{aligned} H_1^{\vec{n}, \pm} 2e^{3\alpha} = & 2\pi_\varphi \pi_{f_{\vec{n}, \pm}} - 2\pi_\alpha \pi_{a_{\vec{n}, \pm}} - (\pi_\alpha^2 + 3\pi_\varphi^2) a_{\vec{n}, \pm} - \frac{2}{3} e^{4\alpha} \omega_n^2 (a_{\vec{n}, \pm} + b_{\vec{n}, \pm}) \\ & + e^{6\alpha} m^2 \sigma^2 \varphi (3\varphi a_{\vec{n}, \pm} + 2f_{\vec{n}, \pm}) \end{aligned}$$

$$\tilde{H}_1^{\vec{n}, \pm} 3e^\alpha = \pi_{b_{\vec{n}, \pm}} - \pi_{a_{\vec{n}, \pm}} + \pi_\alpha (a_{\vec{n}, \pm} + 4b_{\vec{n}, \pm}) + 3\pi_\varphi f_{\vec{n}, \pm}.$$

Longitudinal gauge



- We can adopt **longitudinal gauge** by imposing:

$$\pi_{a_{\vec{n},\pm}} - \pi_{\alpha} a_{\vec{n},\pm} - 3\pi_{\varphi} f_{\vec{n},\pm} = 0, \quad b_{\vec{n},\pm} = 0.$$

- This removes the constraints *linear* in perturbations.

$$\pi_{b_{\vec{n},\pm}} = 0, \quad a_{\vec{n},\pm} = 3 \frac{\pi_{\varphi} \pi_{f_{\vec{n},\pm}} + (e^{6\alpha} m^2 \sigma^2 \varphi - 3\pi_{\alpha} \pi_{\varphi}) f_{\vec{n},\pm}}{9\pi_{\varphi}^2 + \omega_n^2 e^{4\alpha}}.$$

- Together with dynamical stability, this fixes $g_{\vec{n},\pm} = -a_{\vec{n},\pm}$, $k_{\vec{n},\pm} = 0$.

The shift vanishes, and the spatial metric is proportional to ${}^0 h_{ij}$.

Longitudinal gauge: Reduction



- After **REDUCTION**, a canonical set is:

$$\bar{\varphi} = \varphi + 3 \sum a_{\bar{n}, \pm} f_{\bar{n}, \pm}, \quad \pi_{\bar{\varphi}} = \pi_{\varphi},$$


$$\bar{\alpha} = \alpha + \frac{1}{2} \sum (a_{\bar{n}, \pm}^2 + f_{\bar{n}, \pm}^2), \quad \pi_{\bar{\alpha}} = \pi_{\alpha} - \sum f_{\bar{n}, \pm} (\pi_{f_{\bar{n}, \pm}} - 3\pi_{\varphi} a_{\bar{n}, \pm} - \pi_{\alpha} f_{\bar{n}, \pm}),$$

$$\bar{f}_{\bar{n}, \pm} = e^{\alpha} f_{\bar{n}, \pm}, \quad \pi_{\bar{f}_{\bar{n}, \pm}} = e^{-\alpha} (\pi_{f_{\bar{n}, \pm}} - 3\pi_{\varphi} a_{\bar{n}, \pm} - \pi_{\alpha} f_{\bar{n}, \pm}).$$

The genuine background variables are corrected with **quadratic** perturbations.

We have already **scaled** the matter field variables.

Longitudinal gauge: Dynamics



- The modes of the scaled matter field satisfy a quasi-KG equation with time-dependent mass:

$$\ddot{\bar{f}}_{\vec{n},\pm} + r_n \dot{\bar{f}}_{\vec{n},\pm} + (\omega_n^2 + s + s_n) \bar{f}_{\vec{n},\pm} = 0,$$
$$\pi_{\bar{f}_{\vec{n},\pm}} = (1 + p_n) \dot{\bar{f}}_{\vec{n},\pm} + q_n \bar{f}_{\vec{n},\pm},$$

$$s = m^2 \sigma^2 e^{2\bar{\alpha}} - \frac{e^{-4\bar{\alpha}}}{2} \left(\pi_{\bar{\alpha}}^2 + 21 \pi_{\bar{\varphi}}^2 + 3 e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi}^2 \right).$$

r_n, s_n, p_n, q_n are of order ω_n^{-2} .

- For any given background, there exists a **UNIQUE** Fock quantization with the symmetry of the three-torus and unitary dynamics.
- The system can be put in the form of a KG field with time-dependent mass by means of a **mode-dependent** canonical quantization, varying in time.
- This transformation is **unitarily** implementable in the privileged quantization.

Longitudinal gauge: Hamiltonian

- The remaining **Hamiltonian constraint** reads:

$$H = \frac{N_0 \sigma}{16\pi G} C_0 + \epsilon^2 N_0 \sum H_2^{\vec{n}, \pm}, \quad H_2^{\vec{n}, \pm} 2e^{\bar{\alpha}} = \bar{E}_{\bar{f}\bar{f}} \bar{f}_{\vec{n}, \pm}^2 + \bar{E}_{\bar{f}\pi} \bar{f}_{\vec{n}, \pm} \pi_{\bar{f}_{\vec{n}, \pm}} + \bar{E}_{\pi\pi} \pi_{\bar{f}_{\vec{n}, \pm}}^2,$$

$$\bar{E}_{\bar{f}\bar{f}}^n = \omega_n^2 + e^{2\bar{\alpha}} m^2 \sigma^2 - \frac{e^{-4\bar{\alpha}}}{2} \left(\pi_{\bar{\alpha}}^2 + 15 \pi_{\bar{\varphi}}^2 + 3 e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi}^2 \right) - \frac{3}{\omega_n^2} e^{-8\bar{\alpha}} \left(e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi} - 2 \pi_{\bar{\alpha}} \pi_{\bar{\varphi}} \right)^2.$$

$$\bar{E}_{\bar{f}\pi}^n = -\frac{3}{\omega_n^2} e^{-6\bar{\alpha}} \pi_{\bar{\varphi}} \left(e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi} - 2 \pi_{\bar{\alpha}} \pi_{\bar{\varphi}} \right), \quad \bar{E}_{\pi\pi}^n = 1 - \frac{3}{\omega_n^2} e^{-4\bar{\alpha}} \pi_{\bar{\varphi}}^2.$$

The corrections in cyan are of order ω_n^{-2} .

Longitudinal gauge: Metric (at linear order)


$$h_{ij} = (\sigma e^{\bar{\alpha}})^0 h_{ij} \left[1 + \epsilon 2 (2\pi)^{3/2} \sum a_{\vec{n}, \pm} Q_{\vec{n}, \pm} \right],$$

$$N = \sigma N_0 \left(1 - \epsilon (2\pi)^{3/2} \sum a_{\vec{n}, \pm} Q_{\vec{n}, \pm} \right), \quad N_i = 0,$$

$$a_{\vec{n}, \pm} = \frac{3}{\omega_n^2} e^{-3\bar{\alpha}} \left[\pi_{\bar{\varphi}} \pi_{\bar{f}_{\vec{n}, \pm}} + e^{-2\bar{\alpha}} \left(e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi} - 2\pi_{\bar{\alpha}} \pi_{\bar{\varphi}} \right) \bar{f}_{\vec{n}, \pm} \right],$$

$$\Phi = \frac{1}{\sigma} \left(\frac{\bar{\varphi}}{(2\pi)^{3/2}} + \epsilon e^{-\bar{\alpha}} \sum \bar{f}_{\vec{n}, \pm} Q_{\vec{n}, \pm} \right).$$

Gauge invariants



- The **Mukhanov-Sasaki** modes and their momenta have the expression:

$$v_{\vec{n},\pm} = A_n \bar{f}_{\vec{n},\pm} + B_n \pi_{\bar{f}_{\vec{n},\pm}}, \quad \pi_{v_{\vec{n},\pm}} = \dot{v}_{\vec{n},\pm} = F_n \bar{f}_{\vec{n},\pm} + G_n \pi_{\bar{f}_{\vec{n},\pm}},$$

$$A_n = 1 + \frac{3e^{-4\bar{\alpha}} \pi_{\bar{\varphi}}}{\omega_n^2 \pi_{\bar{\alpha}}} \left(e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi} - 2\pi_{\bar{\alpha}} \pi_{\bar{\varphi}} \right), \quad B_n = \frac{3e^{-2\bar{\alpha}} \pi_{\bar{\varphi}}^2}{\omega_n^2 \pi_{\bar{\alpha}}},$$

$$F_n = -\frac{3e^{-2\bar{\alpha}} \pi_{\bar{\varphi}}^2}{\pi_{\bar{\alpha}}} - \frac{3e^{-6\bar{\alpha}}}{\omega_n^2 \pi_{\bar{\alpha}}} \left[e^{12\bar{\alpha}} m^4 \sigma^4 \bar{\varphi}^2 - \frac{e^{6\bar{\alpha}} \pi_{\bar{\varphi}}}{2\pi_{\bar{\alpha}}} m^2 \sigma^2 \bar{\varphi} \left(5\pi_{\bar{\alpha}}^2 - 3\pi_{\bar{\varphi}}^2 + 3e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi}^2 \right) \right] \\ - \frac{3e^{-6\bar{\alpha}} \pi_{\bar{\varphi}}^2}{2\omega_n^2 \pi_{\bar{\alpha}}} \left(11\pi_{\bar{\alpha}} - 15\pi_{\bar{\varphi}}^2 - 3e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi}^2 \right),$$

$$G_n = 1 + \frac{3e^{-4\bar{\alpha}} \pi_{\bar{\varphi}}}{2\omega_n^2 \pi_{\bar{\alpha}}} \left[-2e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi} + \frac{\pi_{\bar{\varphi}}}{\pi_{\bar{\alpha}}} \left(\pi_{\bar{\alpha}}^2 - 3\pi_{\bar{\varphi}}^2 + 3e^{6\bar{\alpha}} m^2 \sigma^2 \bar{\varphi}^2 \right) \right].$$

- If we construct annihilation and creation variables with these invariants (for zero mass), the Bogoliubov transformation, which is **mode dependent**, is **UNITARY** in the privileged Fock quantization .

Robustness under gauge fixing

- Similar results are obtained in the gauge of flat spatial sections $a_{\vec{n}\pm} = b_{\vec{n},\pm} = 0$.
- Moreover, the same **symplectic** structure for **gauge invariants** is obtained.

Quantization: Homogeneous sector

- We quantize the homogeneous sector with standard loop techniques, using improved dynamics and the MMO proposal.
- In the volume basis $\{|v\rangle; v \in \mathbb{R}\}$, with $\hat{V} = |\hat{p}|^{3/2}$,

$$\hat{N}_{\bar{\mu}}|v\rangle = |v+1\rangle, \quad \hat{p}|v\rangle = \text{sgn}(v)(2\pi\gamma G\hbar\sqrt{\Delta}|v|)^{2/3}|v\rangle.$$

- The kinematic Hilbert space is $H_{kin}^{FRW-LQC} \otimes H_{kin}^{matt}$.
- The inverse volume is regularized as usual.

$$\widehat{\left[\frac{1}{V}\right]} = \widehat{\left[\frac{1}{\sqrt{|p|}}\right]}^3, \quad \widehat{\left[\frac{1}{\sqrt{|p|}}\right]} = \frac{3}{4\pi\gamma G\hbar\sqrt{\Delta}} \widehat{\text{sgn}(p)} \sqrt{|\hat{p}|} \left(\hat{N}_{-\bar{\mu}} \sqrt{|\hat{p}|} \hat{N}_{\bar{\mu}} - \hat{N}_{\bar{\mu}} \sqrt{|\hat{p}|} \hat{N}_{-\bar{\mu}} \right).$$

Quantization: Homogeneous Hamiltonian

- After **decoupling the zero-volume** state, we change densitization for the *FRW* constraint:

$$\hat{C}_0 = \left[\frac{1}{V} \right]^{1/2} \hat{C}_0 \left[\frac{1}{V} \right]^{1/2} .$$

$$\hat{C}_0 = -\frac{6}{\gamma^2} \hat{\Omega}_0^2 + 8\pi G \left(\hat{\pi}_\phi^2 + m^2 \hat{\phi}^2 \hat{V}^2 \right).$$

- The gravitational part, with the **MMO proposal**, is:

$$\hat{\Omega}_0 = \frac{1}{4i\sqrt{\Delta}} \hat{V}^{1/2} \left[\overline{\text{sgn}(p)} \left(\hat{N}_{2\bar{\mu}} - \hat{N}_{-2\bar{\mu}} \right) + \left(\hat{N}_{2\bar{\mu}} - \hat{N}_{-2\bar{\mu}} \right) \overline{\text{sgn}(p)} \right] \hat{V}^{1/2} .$$

Takes into account the triad orientation (manifest in anisotropic scenarios).

- This operator has the generic form


$$\hat{\Omega}_0^2 |v\rangle = f_+(v) |v+4\rangle + f(v) |v\rangle + f_-(v) |v-4\rangle .$$

Quantization: Superselection

- $\hat{\Omega}_0^2$ can be seen as a difference operator.

$$\hat{\Omega}_0^2|v\rangle = f_+(v)|v+4\rangle + f(v)|v\rangle + f_-(v)|v-4\rangle.$$

- The real function $f_+(v)$ ($f_-(v)$) vanishes in the **interval** $[-4,0]$ ($[0,4]$).
- The operator preserves the **superselection** sectors $\mathcal{L}_{\pm\epsilon}^{(4)} := \{\pm(\epsilon + 4n), n \in \mathbb{N}\}$




- This operator is selfadjoint in those sectors. Its eigenfunctions are real, and determined by their value at the **minimum volume** $\epsilon \in (0,4]$.

Quantization: Homogeneous states

- **Solutions** to the constraint are determined, e.g., by their initial values at minimum volume.
- If the scalar field serves as a clock, an alternate possibility is to give the value at a section of constant field. This is not always possible.

- The space of *physical* states can be identified, e. g., with $L^2(\mathbb{R}, d\phi)$.



Fock and hybrid quantizations

- We quantize the **rescaled inhomogeneous modes** using annihilation and creation variables constructed from our canonical variables and zero mass.
- We obtain a **Fock space** \mathcal{F} , with basis of *n-particle* states:

$$\left\{ |N\rangle = |N_{(1,0,0),+}, N_{(1,0,0),-}, \dots\rangle; \quad N_{\vec{n},\pm} \in \mathbb{N}, \quad \sum N_{\vec{n},\pm} < \infty \right\}.$$

- We proceed to a hybrid quantization, with Hilbert space

$$H_{kin}^{FRW-LQC} \otimes H_{kin}^{matt} \otimes \mathcal{F}.$$

- The Hamiltonian constraint is **not trivial**.

Quantum Hamiltonian of the perturbations

- We quantize the quadratic contribution of the perturbations to the Hamiltonian adapting the quantization **proposals of the homogeneous sector** and using a symmetric factor ordering:
 - ☆ We **symmetrize** products of the type $\hat{\phi} \hat{\pi}_\phi$.
 - ☆ We take a **symmetric geometric** factor ordering $V^k A \rightarrow \hat{V}^{k/2} \hat{A} \hat{V}^{k/2}$.
 - ☆ We adopt the **LQC** representation $(cp)^{2m} \rightarrow [\hat{\Omega}_0^2]^m$.
 - ☆ In order to **preserve the FRW superselection sectors**, we adopt the prescription $(cp)^{2m+1} \rightarrow [\hat{\Omega}_0^2]^{m/2} \hat{\Lambda}_0 [\hat{\Omega}_0^2]^{m/2}$, where

$$\hat{\Lambda}_0 = -\frac{i}{8\sqrt{\Delta}} \hat{V}^{1/2} \left[\widehat{\text{sgn}(p)} (\hat{N}_{4\bar{\mu}} - \hat{N}_{-4\bar{\mu}}) + (\hat{N}_{4\bar{\mu}} - \hat{N}_{-4\bar{\mu}}) \widehat{\text{sgn}(p)} \right] \hat{V}^{1/2}.$$

The situation is similar to that found with the Hubble parameter in LQC.

Quantum Hamiltonian of the perturbations

- With the FRW densitization:

$$\hat{H}_2^{\bar{n},\pm} = \frac{\sigma}{16\pi G} \left[\frac{1}{V} \right]^{1/2} \hat{C}_2^{\bar{n},\pm} \left[\frac{1}{V} \right]^{1/2}.$$

$$\hat{C}_2^{\bar{n},\pm} = 6(2\pi)^4 \sigma^2 \left[2\omega_n \left[\frac{1}{V} \right]^{-2/3} + \frac{\hat{Y}^-}{\omega_n} + \frac{\hat{Z}}{\omega_n^3} \right] \hat{N}_{\bar{n},\pm} + 4\pi G \left[\left(\frac{\hat{Y}^+}{\omega_n} + \frac{\hat{Z}}{\omega_n^3} \right) \hat{X}_{\bar{n},\pm}^+ + \frac{3i\sigma^2 \hat{W}}{\omega_n^2} \hat{X}_{\bar{n},\pm}^- \right],$$

$$\hat{N}_{\bar{n},\pm} = \hat{a}_{f_{\bar{n},\pm}^-}^\dagger \hat{a}_{f_{\bar{n},\pm}^-}, \quad \hat{X}_{\bar{n},\pm}^\pm = \left(\hat{a}_{f_{\bar{n},\pm}^-}^\dagger \right)^2 \pm \left(\hat{a}_{f_{\bar{n},\pm}^-} \right)^2,$$

$$\hat{Y}^\pm = \frac{m^2}{(2\pi)^2} - \pi \sigma^2 \left[\frac{1}{V} \right]^{1/3} \left(\frac{1}{\gamma^2 (2\pi)^3 \sigma^2} \hat{\Omega}_0^2 + 3(5 \pm 2) \hat{\pi}_\phi^2 + 3m^2 \hat{V}^2 \hat{\phi}^2 \right) \left[\frac{1}{V} \right]^{1/3},$$

$$\hat{Z} = -\frac{3\sigma^2}{2\pi} \left[\frac{1}{V} \right] \left(\frac{2}{\gamma} \hat{\Lambda}_0 \hat{\pi}_\phi + m^2 \hat{V}^2 \hat{\phi} \right)^2 \left[\frac{1}{V} \right],$$

$$\hat{W} = -\left[\frac{1}{V} \right]^{2/3} \left(\frac{4}{\gamma} \hat{\Lambda}_0 \hat{\pi}_\phi^2 + m^2 \hat{V}^2 (\hat{\phi} \hat{\pi}_\phi + \hat{\pi}_\phi \hat{\phi}) \right) \left[\frac{1}{V} \right]^{2/3}.$$

Solutions to the constraint

- If the matter field serves as a **clock**:

$$\hat{C}_0 + \epsilon^2 \left(\sum \hat{C}_2^{\vec{n}, \pm} \right) = 0.$$

$$\langle \Psi | \hat{\pi}_\phi = \frac{1}{\sqrt{8\pi G}} \langle \Psi | \left[\hat{\Theta}_0^2 - \epsilon^2 \left(\sum \hat{C}_2^{\vec{n}, \pm} \right)^\dagger \right]^{1/2} \approx \frac{1}{\sqrt{8\pi G}} \langle \Psi | \left[\hat{\Theta}_0 - \frac{\epsilon^2}{2} \hat{\Theta}_0^{-1} \left(\sum \hat{C}_2^{\vec{n}, \pm} \right)^\dagger \right],$$

$$\hat{\Theta}_0^2 = P \left(8\pi G \hat{\pi}_\phi^2 - \hat{C}_0 \right).$$

- We can pass to an **interaction picture** and use a Born-Oppenheimer-like approximation.
- This can be done even without the above *perturbative expansion*.
- This leads to a sort of **effective** QFT for the inhomogeneities.

Physical states

- An alternate **perturbative** scheme:

$$|\Psi\rangle = |\Psi^{(0)}\rangle + \epsilon^2 |\Psi^{(2)}\rangle \dots$$

- **FRW solution:** $|\Psi^{(0)}\rangle \hat{C}_0 = 0,$

$$\hat{C}_0 = -\frac{6}{\gamma^2} \hat{\Omega}_0^2 + 8\pi G (\hat{\pi}_\phi^2 + m^2 \hat{\phi}^2 \hat{V}^2).$$

- **Evolution of the perturbations:**

$$|\Psi^{(2)}\rangle \hat{C}_0 = -|\Psi^{(0)}\rangle \left(\sum \hat{C}_2^{\vec{n}, \pm} \right)^\dagger.$$

- Solutions are characterized by their initial data at **minimum volume**.
- From these data we arrive, e.g., at the **physical Hilbert space** $H_{kin}^{matt} \otimes \mathcal{F}$.

Conclusions



- We have considered a perturbed FRW universe with a **massive** scalar field.
- Two **approximations**:
 - ☆ The action has been truncated to second order in the perturbations.
 - ☆ A hybrid quantization scheme has been adopted.
- **First** complete **quantization** of a model with inflation within LQC ($k=1$).
- **Backreaction** has been included.

Conclusion



- For quantum simulations, the FRW prescription is **optimal**.
- Opposite to the situation in other analyses, the inhomogeneities have **UNITARY** dynamics in an (*effective*) QFT approximation.
- **No internal time** (matter clock) is needed. If a matter clock is available, one can obtain the inhomogeneities evolution adopting an **interaction picture**.
- Generally, one can construct quantum states perturbatively from data at **minimum volume**. This allows one to get a **physical Hilbert space**.