Immunoassays for trifloxystrobin analysis.

Part II. Assay development and application to residue

determination in food

Josep V. Mercader,^{b,1} Rosario López-Moreno,^{a,1} Francesc A. Esteve-Turrillas,^b

Antonio Abad-Somovilla,^{a,*} Antonio Abad-Fuentes,^{b,**}

^a *Department of Biotechnology, Institute of Agrochemistry and Food Technology, Consejo Superior de Investigaciones Científicas (IATA-CSIC), Agustí Escardino 7, 46980 Paterna, València, Spain*

^b *Department of Organic Chemistry, Universitat de València, Doctor Moliner 50, 46100 Burjassot, València, Spain*

¹ These authors contributed equally to this work

* Corresponding author. Tel.: +34-963544509; fax: +34-963544328

** Corresponding author. Tel.: +34-963900022; fax: +34-963636301.

E-mail addresses: antonio.abad@uv.es (A. Abad-Somovilla), aabad@iata.csic.es (A. Abad-Fuentes).

21   **Abstract**

22   Immunochemical assays constitute complementary analytical methods for small organic
23   molecule determination. We herein describe the characterization and optimization of two
24   competitive enzyme-linked immunosorbent assays in different formats using monoclonal
25   antibodies to the Quinone outside inhibitor (QoI) fungicide trifloxystrobin. Antibody
26   selectivity was evaluated using a variety of agrochemicals and the main trifloxystrobin
27   metabolite. Acceptable tolerance of the immunoassay to methanol, ethanol, and
28   acetonitrile was observed in all cases, whereas a dissimilar influence of buffer pH and ionic
29   strength was found. Moreover, the influence of Tween 20 over the analytical parameters
30   was studied. The limits of detection of the optimized assays were below $0.1 \mu\text{g L}^{-1}$.
31   Excellent recoveries, even at $10 \mu\text{g kg}^{-1}$, were obtained when strawberry, tomato, and
32   cucumber samples spiked with trifloxystrobin were analyzed. Finally, statistical agreement
33   was found between immunoassay and reference chromatographic results using blind-
34   spiked and in-field treated samples.

35

1. Introduction

Trifloxystrobin is a modern agrochemical exhibiting rapid and potent fungicidal activity and possessing a particular mode of action that confers much lower toxicity to humans and the environment than other traditional antimycotic substances. Due to this fact, its use has rapidly grown after it was introduced in the market in the late 1990s – global sales reached 450 M€ in 2010 (Bayer, 2010). Trifloxystrobin belongs to the strobilurin family of new-generation agrochemicals. It has been registered in more than 80 countries, and nowadays it is commercialized by Bayer CropScience under different trade names (Flint, Delaro, Madison, etc.) as the only active ingredient or formulated together with other fungicides. Although some cases of natural or acquired resistance of fungal pathogens like *Ventura inaequalis* or *Fusarium graminearum* towards trifloxystrobin have been recently reported (Frederick & Cox, 2012; Dubos, Pasquali, Pogoda, Hoffmann, & Beyer, 2011), trifloxystrobin is widely employed to fight fungal diseases in a variety of crops such as cereals, strawberries, oranges, grapes, tomatoes, cucumbers, etc. Metabolism in plants is low but complex, with the demethylated derivative being the most relevant metabolite (Fig. 1), yet trifloxystrobin accounts for more than 95% of the residue. Consequently, most regulatory agencies just include the pesticide itself as the only analytical target in samples from plant origin, even though particular legislations also take into account the demethylated metabolite (Health Canada, 2012). In the EU, trifloxystrobin was approved as low risk active substance for plant protection in 2003. European and US maximum residue limits (MRL) for trifloxystrobin range from 0.05 to 30 mg kg⁻¹ in authorized crops. According to the 2010 EU report on pesticide residues in food

(EFSA, 2013), this fungicide was one of the most frequently found pesticides – between 2% and 5% of the analyzed food samples contained trifloxystrobin residues at or below the MRL – and it was among the most common combinations in samples with multiple residues.

During the last decade, an effort has been put in order to develop analytical methods for trifloxystrobin analysis in foodstuffs. The first approach was described in 2002 and comprised liquid chromatography coupled to tandem mass spectrometry (MS) with electrospray ionization (Taylor, Hunter, Hunter, Lindsay, & Le Bouhellec, 2002). Recently, multiresidue analysis including trifloxystrobin was addressed by high-performance liquid chromatography (HPLC) or gas chromatography (GC) with different detection systems (Abreu, Correia, Herbert, Santos, & Alves, 2005; Campillo, Vinas, Aguinaga, Ferez, & Hernandez-Cordoba, 2010; Likas, Tsiropoulos, & Miliadis, 2007; Mastovska, Dorweiler, Lehotay, Wegscheid, & Szpylka, 2010; Sannino, Bolzoni, & Bandini, 2004; Schurek, Vaclavik, Hooijerink, Lacina, Poustka, Sharman, et al., 2008). Alternatively, for low cost, in-situ, and/or rapid applications in samples with limited number of target analytes, immunochemical methods can be an attractive strategy. As the first approach, most research laboratories preferentially use the competitive enzyme-linked immunosorbent assay (cELISA) because of the high analytical capacity and simplicity for standardization. Depending on the immobilized reagent and on the enzymatic detection step, two main formats are usually configured; that is, the antibody-coated direct cELISA and the conjugate-coated indirect cELISA (d-cELISA and i-cELISA, respectively). Immunochemical methods enabling the sensitive determination of some strobilurin fungicides in foodstuff

have been recently reported by our group (Esteve-Turrillas, Mercader, Agullo, Abad-Somovilla, & Abad-Fuentes, 2011; Esteve-Turrillas, Parra, Abad-Fuentes, Agullo, Abad-Somovilla, & Mercader, 2010; Josep V. Mercader, Parra, Esteve-Turrillas, Agulló, Abad-Somovilla, & Abad-Fuentes, 2012). In this paper, the development of novel monoclonal antibody-based immunoassays with different selectivity and in different cELISA formats is reported. Furthermore, a validation study was undertaken using blind and in-field treated samples, and statistical agreement with GC–MS was assessed. To the best of our knowledge, the determination of trifloxystrobin in food samples by immunoassay is herein reported for the first time.

2. Experimental

2.1. Reagents and instruments

Trifloxystrobin (CAS Registry No. 141517-21-7, Mw 408.13) analytical standard was kindly provided by Bayer CropScience (Leverkusen, Germany). Other pesticide standards were obtained from BASF (Limburgerhof, Germany), Fluka/Riedel-de-Haën (Seelze, Germany), Dr. Ehrenstorfer (Augsburg, Germany), or Syngenta (Basel, Switzerland). Triphenylphosphate was from Sigma/Aldrich (Madrid, Spain). Primary/secondary amine and solvents were from Scharlab (Barcelona, Spain). Enzyme immunoassays were performed using regular reagents, buffers, and plastic ware as reported in the preceding paper (López-Moreno, Mercader, Agulló, Abad-Somovilla, & Abad-Fuentes, 2013). The employed monoclonal antibodies (mAbs) and protein–hapten conjugates were those described in previous articles (López-Moreno, Mercader, Agulló, Abad-Somovilla, & Abad-Fuentes, 2013; J. V. Mercader, Suarez-Pantaleon, Agullo, Abad-Somovilla, & Abad-Fuentes,

2008). Peroxidase conjugate to rabbit anti-mouse immunoglobulin polyclonal antibody was from Dako (Glostrup, Denmark). Microplate wells were washed with an ELx405 microplate washer and absorbances were read with a PowerWave HT, both from BioTek Instruments (Winooski, VT, USA).

2.2. Competitive ELISAs

Eight-point standard curves, including a blank, were prepared in borosilicate glass tubes by 10-fold serial dilution in PBS (10 mM phosphate, pH 7.4 containing 140 mM NaCl), starting from a 100 µg L⁻¹ trifloxystrobin solution in PBS. Experimental values were fitted using the SigmaPlot software (Systat Software Inc., Chicago, IL, USA) to a four-parameter logistic equation:

$$y = (A_{\max} - A_{\min}) / [1 + (x/C)^B] + A_{\min}$$

where $A_{\max}$ is the absorbance reached in the absence of analyte, $A_{\min}$ is the background signal, C is the analyte concentration at the inflexion point of the sigmoidal curve, and B is the slope at the inflexion point.

For assay characterization, the concentration of trifloxystrobin that was necessary to induce a 50% inhibition of the antibody–conjugate reaction (IC_{50}) was taken as a reference. The analyte concentration that inhibited 10% (IC_{10}) the immunochemical reaction was considered as the limit of detection (LOD) of the assay. Curves were normalized and average values were calculated from independent experiments. Cross-reactivity (CR) was calculated as percentage value from the quotient between the IC_{50} for trifloxystrobin and the IC_{50} for the studied compound, both in molar concentration units.

2.2.1. Antibody-coated direct cELISA

Microwells were coated with 100 μ L per well of antibody solution in coating buffer (50 mM carbonate, pH 9.6) by overnight incubation at room temperature. Next day, plates were washed four times with washing solution (150 mM NaCl containing 0.05% (v/v) Tween 20). The competitive step was run with 50 μ L per well of analyte or sample solution in PBS and 50 μ L per well of enzyme tracer solution in PBST (PBS containing 0.05% (v/v) Tween 20). After 1 h incubation at room temperature, plates were washed again as described before. Finally, signal was generated with 100 μ L per well of a 0.012% (v/v) H₂O₂ and 2 mg mL⁻¹ of *o*-phenylenediamine solution in 25 mM sodium citrate and 62 mM sodium phosphate buffer, pH 5.4. The enzymatic activity was stopped after 10 min with 2.5 M sulfuric acid. Absorbance was spectrophotometrically measured at 492 nm with 650 nm as a reference wavelength.

2.2.2. Conjugate-coated indirect cELISA

Coating was performed overnight at room temperature with 100 μ L per well of OVA conjugate solution in coating buffer. Then, coated microwells were washed as previously described. Competition was done with 50 μ L per well of analyte or sample solution in PBS and 50 μ L per well of antibody solution in PBST by incubation at room temperature during 1 h. After another washing step, 100 μ L per well of enzyme-labeled secondary antibody (diluted 1/2000) in PBST was added and incubated during 1 h at room temperature. Finally, plates were washed and signal was generated and read as indicated above.

2.3. *Buffer evaluation and solvent tolerance*

A biparametric approach with a central composite design was followed using the Minitab software (Minitab Inc., State College, PA, USA). Trifloxystrobin was prepared in Milli-Q water, and antibody or tracer solutions were diluted in a series of nine buffers with different pH and ionic strength values (from 5.5 to 9.5 and from 50 to 300 mM, respectively). Those buffers were prepared from a 40 mM citrate, 40 mM phosphate, and 40 mM Tris solution as described elsewhere ([Abad-Fuentes, Esteve-Turrillas, Agulló, Abad-Somovilla, & Mercader, 2012](#)). Ionic strength and pH values of each buffer were adjusted using 2 M NaCl and 5 M HCl, respectively. All buffers contained 0.05% (v/v) Tween 20. For solvent studies, trifloxystrobin standard curves were prepared in PBS containing diverse amounts of methanol, ethanol, or acetonitrile (from 0.5 to 10% (v/v) solvent contents) whereas antibody or tracer solutions were set up in PBST.

2.4. *Sample treatment and analysis*

Strawberries were acquired from the market, and absence of trifloxystrobin was verified by GC–MS. Trifloxystrobin-sprayed and non-sprayed tomato and cucumber samples were obtained from nearby farms. Plant treatments were performed with a nebulizer using a commercial formulation from Bayer CropScience (Flint) prepared at 125 mg L⁻¹ trifloxystrobin in water containing 20% (v/v) alkyl polyglycol ether. Vegetables at the beginning of ripening (according to the BBCH scale, code 82 for tomatoes and cucumbers and code 85 for strawberries) were collected before spraying for control and at days 1, 3, 6, 10, and 15 after pesticide application. Immediately, they were homogenized with a blender and stored frozen at –20 °C until analysis. Samples were extracted using

the QuEChERS method (AOAC International official method) (Lehotay, 2007). Briefly, 20 g of homogenate was mixed with 2 g of sodium acetate and 8 g of anhydrous magnesium sulfate in 20 mL of acetonitrile containing 1% (v/v) acetic acid under vigorous stirring. The organic phase was separated by centrifugation at 2200×g during 5 min. Primary/secondary amine (500 mg) was used to clean, by vortexing, 10 mL of extract in the presence of 1.5 g of anhydrous magnesium sulfate. After a second centrifugation step, clean extracts were filtrated through a Teflon filter (0.2 µm) and stored at –20 °C.

QuEChERS extracts containing 500 µg L⁻¹ of triphenylphosphate as internal standard were employed for GC–MS. Chromatographic determinations were carried out with a 6890N GC apparatus furnished with a 7683 Series autosampler, a HP-5MS (30 m x 0.25 mm x 0.25 µm) capillary column, and a quadrupole 5973N mass detector, all from Agilent Technologies (Santa Clara, CA, USA). One microliter of clean extract was injected in splitless mode at 280 °C by employing helium as carrier with a constant flow of 1 mL min⁻¹. The temperature of the oven (110 °C) was held during 1 min; then, it was increased at a rate of 15 °C min⁻¹ until 280 °C and kept constant at the final temperature during 15 min. Electron impact ionization at 70 eV was used with the ion source at 225 °C. The employed quantification ions were *m/z* 116 and 131 for trifloxystrobin and *m/z* 325 and 326 for triphenylphosphate. For cELISA determinations, extracts were diluted in PB (100 mM sodium phosphate, pH 7.4) and antibody or tracer solutions were prepared in PBT (PB containing 0.05% (v/v) Tween 20). A trifloxystrobin standard curve in PB was run in each plate. Deming regression and Bland–Altman analysis were applied for method validation using the SigmaPlot software (version 12.0).

3. Results and discussion

3.2. Antibody specificity

In addition to the bioconjugates and mAbs to trifloxystrobin that were described in the preceding article, an antibody (mAb TF0#17) previously produced in our lab (J. V. Mercader, Suarez-Pantaleon, Agullo, Abad-Somovilla, & Abad-Fuentes, 2008), was also incorporated to this study. The term “type of antibody” refers, in the present paper, to the immunizing hapten (Fig. S1 of the Supplementary Data File) from which the antibody was derived.

Antibody selectivity was determined in homologous assays using the i-cELISA format. As competitors, the main strobilurin fungicides (kresoxim-methyl, azoxystrobin, picoxystrobin, metominostrobin, and pyraclostrobin) were evaluated. It was observed that antibodies showed varied CR patterns, and that recognition profiles were determined by the immunizing hapten. All of the mAbs that had been generated from haptens with the spacer arm at the methoxyiminoacetate toxophore or at the bridging aromatic ring (TF α -, TF ϵ -, TF ϕ -type antibodies), *i.e.* holding the linker at a distal position from the characteristic aromatic ring of trifloxystrobin (Fig. 1), were highly specific – CR values were lower than 1% for all of the assayed strobilurins. Also, no CR was observed with antibodies derived from the hapten with the linker at a central position of the molecule (TF ζ -type antibodies). Contrarily, most of the antibodies originated from haptens with the spacer arm at the trifluoromethylphenyl characteristic ring (TF ζ - and TF η -type antibodies), *i.e.* at a position distal from the common moieties of strobilurins (Fig. 1), showed significant CR with one or more of these agrochemicals. Particularly, kresoxim-methyl – a strobilurin containing the

same methoxyiminoacetate moiety as trifloxystrobin – was recognized by most of the TFf- and TFt-type antibodies (CR values were between 0.5% and 10%; see Fig. S2).

Thereafter, the ability of the mAb collection to recognize the main metabolite of trifloxystrobin (Fig. 1) was studied. As expected, no inhibition was observed with TF α -type antibodies, whereas the signal was reduced to some extent with TFe- and TFO-type mAbs – CR values were between 2% and 43% (data not shown). Remarkably, TFt#316 displayed equivalent affinity for trifloxystrobin and its main metabolite (Fig. S2), which was an unexpected finding taking into account the linker attachment site in hapten TFt. Finally, recognition towards other relevant fungicides that are commonly formulated together with trifloxystrobin, such as pyrimethanil, propiconazole, and tebuconazole, was also checked, and predictably no inhibition was observed with any of the studied mAbs.

3.3. Immunoreagent selection

The antibody characterization data with homologous and heterologous conjugates described in the preceding paper were capitalized in order to select several immunoreagent combinations for further reevaluation, with the aim of making a solid choice of the best immunoassays. In particular, three immunoreagent combinations were selected for the d-cELISA format and three combinations for the i-cELISA format, and all of these assays were repeated along ten days in independent experiments. Table 1 lists the employed concentrations and the average values of the main analytical parameters from the obtained inhibition curves. From this study, one immunoassay was selected in each cELISA format for further development and validation. The chosen direct assay employed mAb TF0#17 and a heterologous enzyme tracer (HRP–TFc), whereas the selected i-cELISA

engaged mAb TFt#316 along with the homologous conjugate (OVA–TFf). Both assays showed IC₅₀ values in the low nanomolar range, the slopes of the standard curves were moderate, and consumption of immunoreagents was low. Moreover, the two selected antibodies exhibited appealing alternative selectivity; thus, whereas mAb TF0#17 was highly specific, mAb TFt#316 displayed equivalent affinity for trifloxystrobin and its main metabolite.

3.4. Solvent and buffer studies

Tolerance of the selected immunoassays to three organic solvents (methanol, ethanol, and acetonitrile) that are commonly used for pesticide extraction was evaluated. Standard curve parameters using PBS containing different solvent concentrations were compared to the control assay in PBS with no solvent. Both assays tolerated the three studied solvents at a similar extent (5-10%), even though the d-cELISA based on mAb TF0#17 was somewhat more robust in this aspect than the indirect assay with antibody TFt#316 ([Fig. S3 of the Supplementary Data File](#)).

The influence of surfactant Tween 20 over the immunochemical reaction of the selected cELISAs was evaluated using PBST with different additive concentrations (from 0.01 to 0.2%, v/v). In both assays, a slight amount of this detergent (0.01%) increased the A_{max} value. However, higher concentrations of Tween 20 had a negative effect over the A_{max} of the TFt#316-based i-cELISA, whereas it did not significantly change the corresponding value of the direct TF0#17-based assay ([Fig. S4](#)). On the other hand, it was observed that the IC₅₀ of both immunoassays increased with the concentration of Tween 20.

Finally, a biparametric approach was applied for buffer optimization following a central composite design, consisting of a two-level full factorial design ($\alpha = 1.414$) with two factors (pH and ionic strength) and three replicates that included 12 cube, 12 axial, and 15 center points; that is, 39 randomized assays under 9 different buffer conditions (Table S1). $A_{\max}$ and IC_{50} values of the 39 resulting curves were fitted as functions of pH and ionic strength. Fig. S5 shows the contour plots for the direct assay using mAb TF0#17 and the indirect assay with antibody TFt#316. For the latter assay, lower ionic strength increased the $A_{\max}$ and IC_{50} values whereas a moderate influence on assay parameters was observed upon pH changes. On the contrary, a higher dependency on pH than on ionic strength was found with the d-cELISA. Accordingly, a buffer with high buffering capacity would be desirable in order to better stabilize assay conditions, so 100 mM phosphate was selected instead of PBS for future studies.

Final standard curves and the main analytical parameters of the two optimized cELISAs are presented in Table 2. Both immunoassays had high $A_{\max}$ values at low concentrations of immunoreagents, and showed quite similar sensitivities. Additional efforts to improve IC_{50} values through decreasing immunoreagent concentrations were unsuccessful because this simple and common approach, in this particular case, just resulted in lower maximum signals without a concomitant effect on sensitivity.

3.5. Sample analysis

Cucumber, tomato, and strawberry samples were analyzed with the two competitive ELISAs. QuEChERS extracts were fortified, adequately diluted in PB, and analyzed in triplicate wells under the optimized conditions listed in Table 2.

Trifloxystrobin concentrations were determined using a standard curve run in triplicate wells in the same plate. Initial determinations showed unexpectedly low recovery values even from control samples in buffer, so food matrix effects were not responsible for the observed flawed result. After several tests, we reached the conclusion that trifloxystrobin adsorbed to the glass tubes that were used in the preparation of standards and samples. Adsorption of organic compounds of different hydrophobicity to untreated borosilicate glass surfaces in aqueous systems has been reported by other authors, and the degree of the sorption phenomenon has been related to the proportion of cosolvent and to the ratio of solution volume to contact surface area (Muhamad, Ismail, Sameni, & Mat, 2011; Qian, Posch, & Schmidt, 2011; Song, Hsu, & Au, 1996; Wheelock, Miller, Miller, Phillips, Gee, Tjeerdema, et al., 2005). A simple, yet illustrative, experiment of trifloxystrobin adsorption to borosilicate glass tubes is shown in Fig. S6 of the Supplementary Data File, which showed the dissimilar inhibition curves obtained from two sets of standards (1 mL or 10 mL each) prepared in tubes of the same capacity (12 mL). Accordingly, trifloxystrobin standards were further prepared in acetonitrile to decrease adsorption to the test tubes, and then diluted 50-fold in PBS, paying attention to employ glass tubes of adequate capacity for the sample volume. Following these simple precautions, excellent recoveries were obtained with the three samples using the direct immunoassay, at levels below or around European MRLs (Table 3). Concerning the indirect assay, slight deviations were observed, particularly with cucumber samples. Thus, the limit of quantification for the three foodstuffs could be established at $10 \mu\text{g kg}^{-1}$ with the direct

assay, whereas with the i-cELISA the limit of quantification could be set at $30 \mu\text{g kg}^{-1}$ for tomatoes, $50 \mu\text{g kg}^{-1}$ for cucumbers, and $10 \mu\text{g kg}^{-1}$ for strawberries.

The developed cELISAs were further applied to the analysis of blind fortified extracts of trifloxystrobin-free strawberries, tomatoes, and cucumbers. Samples were properly diluted as before, and the immunoanalytical results were compared with those obtained by GC–MS ([Table S2 of the Supplementary Data File](#)). In order to evaluate the equivalency between methods, the Deming regression model, which takes into account the standard deviations of both measurements, was applied ([Fig. 2](#)). The obtained regression line for the direct assay ($r = 0.990$) had a slope of 1.07 (95% confidence interval was from 0.99 to 1.16) and an intercept of -0.018 (95% confidence interval was from -0.033 to -0.003), indicating a minor underestimation. Regarding the indirect assay, the regression line ($r = 0.994$) had a slope of 1.07 (95% confidence interval was from 0.99 to 1.16) and an intercept of 0.009 (95% confidence interval was from -0.009 to 0.026), showing that results were statistically comparable. Finally, good correlation between immunochemical and chromatography methods exists as observed by the Bland–Altman plots ([Fig. 2](#)) – values were randomly distributed around the average difference, and they were mainly inside the limits of agreement (average difference ± 1.96), meaning that only random deviations exist.

Finally, the suitability of the immunoassays for trifloxystrobin analysis was evaluated under a real situation. Tomato and cucumber plants were treated with a commercial formulation of trifloxystrobin following good agricultural practices. Samples were collected at days 1, 3, 6, 10, and 15 after pesticide application and they were QuEChERS-

extracted and analyzed by GC–MS and cELISAs (Table 4). Trifloxystrobin concentrations found both in tomato and cucumber samples were lower than their respective MRLs by the three analytical methods, which confirm that if the fungicide is applied at the recommended dose and preharvest safety intervals are respected (1 and 3 days for tomatoes and cucumbers, respectively), trifloxystrobin residues compliant with the legislation should be expected.

4. Conclusions

Immunoreagents were produced and immunoassays were developed and validated for trifloxystrobin analysis as part of a general study. In the present article, specificity and sensitivity of two cELISAs have been reported. One direct immunoassay specific to trifloxystrobin and one i-cELISA also recognizing the main trifloxystrobin metabolite were optimized and characterized in terms of solvent tolerance, and pH and ionic strength dependency. Phosphate 100 mM, pH 7.4 was chosen for optimum assay performance. The developed immunoassays showed satisfactory analytical parameters for the intended application – IC₅₀ values were between 0.7 and 1.0 µg L⁻¹ with curve slopes close to 1.0. Adsorption of trifloxystrobin to glass ware was observed and minimized by using a cosolvent and fitting tube capacity to sample volume. Both immunoassays were validated with fruit and vegetable samples. Recoveries between 85 and 116% were found using strawberry, tomato, and cucumber samples spiked at levels from 10 to 500 µg kg⁻¹. Also, comparison of the cELISAs and GC–MS results by Deming regression and Bland–Altman data analysis showed good agreement between both methods. Finally, analysis of

vegetables treated in the greenhouse with trifloxystrobin commercial formulations demonstrated the applicability of the developed immunoassays.

Acknowledgements

This work was supported by the Spanish *Ministerio de Ciencia e Innovación* (MICINN) (AGL2006-12750-C02-01/02/ALI and AGL2009-12940-C02-01/02/ALI) and cofinanced by FEDER funds. R.L.-M. was hired by MICINN under a predoctoral FPI grant associated to the above project. F.A.E.-T. and J.V.M. were hired by CSIC with postdoctoral contracts, the former under the *JAEdoc* program and the latter under the *Ramón y Cajal* program, both cofinanced by MICINN and by the European Social Fund (ESF). We thank Ana Izquierdo-Gil and Laura López-Sánchez for excellent technical assistance.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:

References

- Abad-Fuentes, A., Esteve-Turrillas, F. A., Agulló, C., Abad-Somovilla, A., & Mercader, J. V. (2012). Development of competitive enzyme-linked immunosorbent assays for boscalid determination in fruit juices. *Food Chemistry*, 135(1), 276-284.
- Abreu, S. D., Correia, M., Herbert, P., Santos, L., & Alves, A. (2005). Screening of grapes and wine for azoxystrobin, kresoxim-methyl and trifloxystrobin fungicides by HPLC with diode array detection. *Food Additives and Contaminants Part a-Chemistry Analysis Control Exposure & Risk Assessment*, 22(6), 549-556.
- Bayer. (2010). Bayer Annual Report 2010. <http://www.annualreport2010.bayer.com/>. Accessed on 2 August 2013
- Campillo, N., Vinas, P., Aguinaga, N., Ferez, G., & Hernandez-Cordoba, M. (2010). Stir bar sorptive extraction coupled to liquid chromatography for the analysis of strobilurin fungicides in fruit samples. *Journal of Chromatography A*, 1217(27), 4529-4534.
- Dubos, T., Pasquali, M., Pogoda, F., Hoffmann, L., & Beyer, M. (2011). Evidence for natural resistance towards trifloxystrobin in *Fusarium graminearum*. *European Journal of Plant Pathology*, 130(2), 239-248.
- Health Canada (2012). Pest Management Regulatory Agency. Proposed Maximum Residue Limit Trifloxystrobin. http://www.hc-sc.gc.ca/cps-spc/pest/part/consultations/_pmrl2012-27/pmrl2012-27-eng.php. Accessed on 2 August 2013
- EFSA. European Food Safety Authority. (2013). The 2010 European Union Report on Pesticide Residues in Food. *EFSA Journal*, 11(3), 3130. <http://www.efsa.europa.eu/de/efsajournal/pub/3130.htm>. Accessed on 2 August 2013.
- Esteve-Turrillas, F. A., Mercader, J. V., Agulló, C., Abad-Somovilla, A., & Abad-Fuentes, A. (2011). Development of immunoaffinity columns for pyraclostrobin extraction from fruit juices and analysis by liquid chromatography with UV detection. *Journal of Chromatography A*, 1218(30), 4902-4909.
- Esteve-Turrillas, F. A., Parra, J., Abad-Fuentes, A., Agulló, C., Abad-Somovilla, A., & Mercader, J. V. (2010). Hapten synthesis, monoclonal antibody generation, and development of competitive immunoassays for the analysis of picoxystrobin in beer. *Analytica Chimica Acta*, 682(1-2), 93-103.
- Frederick, Z., & Cox, K. D. (2012). Practical and qualitative resistance to trifloxystrobin in populations of *Venturia inaequalis* in the northeastern United States. *Phytopathology*, 102(7), 42-42.
- Lehotay, S. J. (2007). Determination of pesticide residues in foods by acetonitrile extraction and partitioning with magnesium sulfate: Collaborative study. *Journal of AOAC International*, 90(2), 485-520.
- Likas, D. T., Tsiropoulos, N. G., & Miliadis, G. E. (2007). Rapid gas chromatographic method for the determination of famoxadone, trifloxystrobin and fenhexamid residues in tomato, grape and wine samples. *Journal of Chromatography A*, 1150(1-2), 208-214.
- López-Moreno, R., Mercader, J. V., Agulló, C., Abad-Somovilla, A., & Abad-Fuentes, A. (2013). Immunoassays for trifloxystrobin analysis. Part I: Synthesis of regioisomeric haptens and production of monoclonal antibodies. *Food Chemistry*, 230, 230-236.
- Mastovska, K., Dorweiler, K. J., Lehotay, S. J., Wegscheid, J. S., & Szpylka, K. A. (2010). Pesticide Multiresidue Analysis in Cereal Grains Using Modified QuEChERS Method Combined with Automated Direct Sample Introduction GC-TOFMS and UPLC-MS/MS Techniques. *Journal of Agricultural and Food Chemistry*, 58(10), 5959-5972.

- Mercader, J. V., Parra, J., Esteve-Turrillas, F. A., Agulló, C., Abad-Somovilla, A., & Abad-Fuentes, A. (2012). Development of monoclonal antibody-based competitive immunoassays for the detection of picoxystrobin in cereal and oilseed flours. *Food Control*, 26(1), 162-168.
- Mercader, J. V., Suárez-Pantaleón, C., Agulló, C., Abad-Somovilla, A., & Abad-Fuentes, A. (2008). Hapten synthesis and monoclonal antibody-based immunoassay development for detection of the fungicide trifloxystrobin. *Journal of Agricultural and Food Chemistry*, 56(8), 2581-2588.
- Muhamad, H., Ismail, B. S., Sameni, M., & Mat, N. (2011). Adsorption study of C-14-paraquat in two Malaysian agricultural soils. *Environmental Monitoring and Assessment*, 176(1-4), 43-50.
- Qian, Y. A., Posch, T., & Schmidt, T. C. (2011). Sorption of polycyclic aromatic hydrocarbons (PAHs) on glass surfaces. *Chemosphere*, 82(6), 859-865.
- Sannino, A., Bolzoni, L., & Bandini, M. (2004). Application of liquid chromatography with electrospray tandem mass spectrometry to the determination of a new generation of pesticides in processed fruits and vegetables. *Journal of Chromatography A*, 1036(2), 161-169.
- Schurek, J., Vaclavik, L., Hooijerink, H., Lacina, O., Poustka, J., Sharman, M., Caldow, M., Nielen, M. W. F., & Hajslova, J. (2008). Control of Strobilurin Fungicides in Wheat Using Direct Analysis in Real Time Accurate Time-of-Flight and Desorption Electrospray Ionization Linear Ion Trap Mass Spectrometry. *Analytical Chemistry*, 80(24), 9567-9575.
- Song, D., Hsu, L. F., & Au, J. L. S. (1996). Binding of taxol to plastic and glass containers and protein under in vitro conditions. *Journal of Pharmaceutical Sciences*, 85(1), 29-31.
- Taylor, M. J., Hunter, K., Hunter, K. B., Lindsay, D., & Le Bouhellec, S. (2002). Multi-residue method for rapid screening and confirmation of pesticides in crude extracts of fruits and vegetables using isocratic liquid chromatography with electrospray tandem mass spectrometry. *Journal of Chromatography A*, 982(2), 225-236.
- Wheelock, C. E., Miller, J. L., Miller, M. J., Phillips, B. M., Gee, S. J., Tjeerdema, R. S., & Hammock, B. D. (2005). Influence of container adsorption upon observed pyrethroid toxicity to *Ceriodaphnia dubia* and *Hyalella azteca*. *Aquatic Toxicology*, 74(1), 47-52.

Figure legends

Fig. 1. Top: trifloxystrobin; the arrows point out the attachment sites of the C-6 hydrocarbon spacer arm of haptens. Bottom: main trifloxystrobin metabolite.

Fig. 2. Comparison of ELISA and GC–MS results from blind-spiked samples using Deming regression (left) and Bland–Altman plot (right).

Table 1
Curve parameters of the selected immunoassays before optimization.^a

Antibody	[Antibody] ($\mu\text{g L}^{-1}$)	Conjugate	[Conjugate] ($\mu\text{g L}^{-1}$)	A_{max}	Slope	IC_{50} ($\mu\text{g L}^{-1}$)	Cross-reactivity
TF0#17	300	HRP-TFc	30	0.92 ± 0.14	-1.19 ± 0.15	2.08 ± 0.49	Not found
TFe#38	1000	HRP-TFe	100	1.17 ± 0.18	-0.82 ± 0.14	4.95 ± 0.99	Metabolite (10%)
TF#316	300	HRP-TFe	100	1.70 ± 0.35	-1.04 ± 0.16	5.57 ± 0.82	Metabolite (96%)
TFe#38	30	OVA-TFo	1000	1.64 ± 0.17	-1.27 ± 0.14	5.87 ± 2.01	Metabolite (10%)
TF#211	30	OVA-TFb	1000	1.56 ± 0.12	-1.17 ± 0.14	3.70 ± 0.82	Kresoxim-methyl (9%)
TF#316	30	OVA-TFt	100	1.37 ± 0.14	-1.16 ± 0.28	4.33 ± 0.83	Metabolite (96%)

^a n=10.

Table 2
Assay conditions and analytical parameters of the optimized immunoassays.^a

cELISA		
Format	Direct (●)	Indirect (▲)
mAb	TF0#17 (300 $\mu\text{g L}^{-1}$)	TF#316 (30 $\mu\text{g L}^{-1}$)
Conjugate	HRP-TFc (50 $\mu\text{g L}^{-1}$)	OVA-TFt (100 $\mu\text{g L}^{-1}$)
A_{max}	1.760 ± 0.228	2.690 ± 0.196
A_{min}	0.027 ± 0.008	0.031 ± 0.008
Slope	-0.878 ± 0.034	-1.075 ± 0.028
IC_{50} ($\mu\text{g L}^{-1}$)	0.986 ± 0.048	0.668 ± 0.018
LOD ($\mu\text{g L}^{-1}$)	0.085	0.084
$\text{IC}_{20}\text{--}\text{IC}_{80}$ ($\mu\text{g L}^{-1}$)	0.211–4.860	0.178–2.460
Phosphate buffer	100 mM, pH 7.4	100 mM, pH 7.4
Tween 20 (%)	0.025	0.010
Time (h)	1.5	2.5
CR (%)	Negligible	Metabolite (100)

^a Values are the mean of 14 independent experiments.

Table 3 Recoveries (%) from trifloxystrobin fortified extracts.^a

	[TF] ^b (mg kg ⁻¹)	d-cELISA	i-cELISA
Tomato ^c	0.01	106 ± 19	167 ± 38
	0.03	104 ± 13	109 ± 21
	0.05	109 ± 14	109 ± 15
	0.10	96 ± 7	101 ± 10
	0.30	97 ± 9	94 ± 19
	0.50	94 ± 12	100 ± 15
Cucumber ^d	0.01	108 ± 14	245 ± 38
	0.03	98 ± 17	153 ± 18
	0.05	109 ± 19	124 ± 14
	0.10	112 ± 11	112 ± 10
	0.30	98 ± 5	109 ± 9
	0.50	92 ± 17	100 ± 8
Strawberry ^e	0.01	86 ± 15	93 ± 4
	0.03	109 ± 19	104 ± 6
	0.05	95 ± 3	96 ± 3
	0.10	92 ± 12	102 ± 5
	0.30	91 ± 6	98 ± 12
	0.50	86 ± 10	101 ± 19

^a Values are the mean of four independent experiments. Extracts were diluted 1/50. Standard curves were prepared in acetonitrile and diluted like the extracts. ^b Spiked trifloxystrobin concentration. ^c EU MRL = 0.5 mg kg⁻¹.

^d EU MRL = 0.2 mg kg⁻¹. ^e EU MRL = 0.5 mg kg⁻¹.

Table 4 Trifloxystrobin concentration from in-field treated samples.^a

Sample	Harvest day	GC-MS (mg kg ⁻¹)	d-cELISA (mg kg ⁻¹)	i-cELISA (mg kg ⁻¹)
Tomato	D1	0.116 ± 0.017	0.156 ± 0.016	0.181 ± 0.015
	D3	0.072 ± 0.011	0.091 ± 0.011	0.116 ± 0.009
	D6	0.081 ± 0.012	0.098 ± 0.009	0.149 ± 0.027
	D10	0.119 ± 0.017	0.119 ± 0.013	0.155 ± 0.025
	D15	0.062 ± 0.009	0.039 ± 0.004	0.073 ± 0.011
Cucumber	D1	0.136 ± 0.020	0.107 ± 0.009	0.176 ± 0.011
	D3	0.115 ± 0.017	0.091 ± 0.002	0.126 ± 0.008
	D6	< LOQ ^b	0.031 ± 0.002	0.058 ± 0.008
	D10	< LOQ	< LOQ	< LOQ
	D15	< LOQ	< LOQ	< LOQ

^a Values are the average of two and three independent determinations for GC-MS and cELISA, respectively. ^b The limit of quantitation of GC-MS was 0.05 mg kg⁻¹.

Figure(s)

Figure 1

Figure 2

SUPPLEMENTARY DATA

Immunoassays for trifloxystrobin analysis.

Part II. Assay development and application to residue determination in food

Josep V. Mercader, Rosario López-Moreno, Francesc A. Esteve-Turrillas, Antonio Abad-Somovilla, Antonio Abad-Fuentes

	<u>Page</u>
Fig. S1. Trifloxystrobin functionalized haptens	2
Fig. S2. Specificity of TFt-type mAbs	3
Fig. S3. Solvent tolerance	4
Fig. S4. Influence of Tween 20 concentration	5
Table S1. Studied pH and ionic strength conditions	6
Fig. S5. Influence of pH and ionic strength	6
Fig. S6. Adsorption of trifloxystrobin to glass ware	7
Table S2. ELISA and GC-MS results from blind samples	8

Fig. S1. Structure of the trifloxystrobin functionalized haptens

Fig. S2. Specificity of TFt-type mAbs towards the main strobilurins and the metabolite of trifloxystrobin. Analytes: TF, trifloxystrobin; KM, kresoxim-methyl; AZ, azoxystrobin; PC, picoxystrobin; MT, metominostrobin. Data are shown only for analytes that were recognized by at least one antibody.

Fig. S3. Solvent tolerance. Influence of different solvent concentrations on the $A_{\max}$ (left-hand axis) and IC_{50} (right-hand axis) values of the two proposed immunoassays.

Fig. S4. Influence of Tween 20 concentration on the $A_{\max}$ (left-hand axis) and IC_{50} (right-hand axis) values of the two proposed immunoassays.

Table S1

Composition of the buffers used in the immunoassay optimization studies.

Buffer	I (mM)	pH	Tween 20 (%)
#1	175.0	5.5	0.025
#2	86.6	6.1	0.025
#3	263.4	6.1	0.025
#4	50.0	7.5	0.025
#5 ^a	175.0	7.5	0.025
#6	300.0	7.5	0.025
#7	86.6	8.9	0.025
#8	263.4	8.9	0.025
#9	175.0	9.5	0.025

^a Central point of the composite design.

Fig. S5. Influence of pH and ionic strength on the $A_{\max}$ (left-hand graphs) and IC_{50} (right-hand graphs) values of the two proposed immunoassays.

d-cELISA: TF0#17 vs. HRP-TFc

i-cELISA: TFt#316 vs. OVA-TFt

Fig. S6. Adsorption of trifloxystrobin to glass ware. Identical trifloxystrobin standard curves were prepared in 12-mL borosilicate tubes, but in one case 1-mL individual standard solutions were prepared (circles), whereas in the other case 10-mL individual standard solutions were prepared (triangles). Displacement of the red curve to the right is indicative of trifloxystrobin adsorption to glass tubes because of the higher surface-to-volume ratio in 1-mL standards than in 10-mL standards.

Table S2
Comparison between chromatographic and
immunochemical methods using trifloxystrobin blind-spiked
samples.^a

Sample		GC-MS (mg kg ⁻¹)	d-cELISA (mg kg ⁻¹)	i-cELISA (mg kg ⁻¹)
Tomato	#1	0.62 ± 0.09	0.67 ± 0.06	0.61 ± 0.01
	#2	0.27 ± 0.04	0.26 ± 0.03	0.23 ± 0.05
	#3	1.89 ± 0.28	2.24 ± 0.30	2.17 ± 0.14
	#4	0.83 ± 0.12	1.07 ± 0.13	0.97 ± 0.10
	#5	1.10 ± 0.16	1.29 ± 0.11	1.35 ± 0.10
	#6	0.40 ± 0.06	0.42 ± 0.05	0.40 ± 0.05
	#7	1.34 ± 0.20	1.56 ± 0.03	1.55 ± 0.31
	#8	0.88 ± 0.13	0.98 ± 0.08	0.93 ± 0.15
Cucumber	#1	0.54 ± 0.08	0.53 ± 0.06	0.68 ± 0.02
	#2	1.92 ± 0.29	2.07 ± 0.16	2.15 ± 0.34
	#3	0.31 ± 0.05	0.31 ± 0.04	0.34 ± 0.03
	#4	1.63 ± 0.24	1.68 ± 0.28	1.77 ± 0.39
	#5	1.04 ± 0.13	1.06 ± 0.25	1.05 ± 0.15
	#6	1.18 ± 0.18	1.09 ± 0.05	1.16 ± 0.20
	#7	1.41 ± 0.21	1.25 ± 0.08	1.40 ± 0.25
	#8	1.24 ± 0.18	1.33 ± 0.12	1.26 ± 0.20
Strawberry	#1	0.38 ± 0.05	0.32 ± 0.03	0.41 ± 0.03
	#2	0.91 ± 0.13	0.86 ± 0.15	0.93 ± 0.16
	#3	0.56 ± 0.08	0.51 ± 0.02	0.46 ± 0.03
	#4	1.08 ± 0.16	1.26 ± 0.21	1.19 ± 0.10
	#5	0.66 ± 0.09	0.69 ± 0.13	0.70 ± 0.05
	#6	0.14 ± 0.02	0.06 ± 0.01	0.10 ± 0.02
	#7	0.94 ± 0.14	1.11 ± 0.19	1.05 ± 0.11
	#8	1.43 ± 0.21	1.53 ± 0.11	1.72 ± 0.19
	#9	0.16 ± 0.02	0.10 ± 0.02	0.14 ± 0.03
	#10	1.71 ± 0.25	1.77 ± 0.03	1.81 ± 0.18
	#11	1.12 ± 0.16	1.18 ± 0.16	1.09 ± 0.02
	#12	0.12 ± 0.02	0.12 ± 0.01	0.06 ± 0.01
	#13	0.29 ± 0.04	0.25 ± 0.05	0.27 ± 0.06
	#14	0.30 ± 0.04	0.25 ± 0.07	0.28 ± 0.07
	#15	1.98 ± 0.30	1.96 ± 0.05	2.13 ± 0.01
	#16	1.65 ± 0.25	1.65 ± 0.13	1.72 ± 0.09

^a Values are the average of two and three independent determinations for GC-MS and cELISA, respectively.