1
2
3
4 Effect of the antioxidants composition in diet on the
5 sensory and physical properties of frozen farmed Coho
6 salmon (*Oncorhynchus kisutch*)
7
8
9
10
11

12 Alicia Rodríguez¹, Mónica Latorre¹, Mónica Gajardo¹, Andrea Bunger¹,
13 Alejandro Munizaga², Luis López¹, and Santiago P. Aubourg^{3,*}
14
15
16
17
18
19
20
21
22
23

24 ¹ Department of Food Science and Chemical Technology, Faculty of Chemical and
25 Pharmaceutical Science, Universidad de Chile, Santiago (Chile).

26 ² Faculty of Biological Sciences. Pontificia Universidad Católica de Chile, Santiago
27 (Chile).

28 ³ Department of Food Technology, Marine Research Institute (CSIC), Vigo (Spain).

29 * Correspondent: saubourg@iim.csic.es. Fax: +34 986292762; Phone: +34 986231930.
30
31
32
33

ABSTRACT

BACKGROUND: Great attention has been addressed to the antioxidants presence in farmed fish feeds, being the replacement of synthetic antioxidants by natural ones a main objective. In the present study, Coho salmon (*Oncorhynchus kisutch*) was fed a conventional diet that was enriched with different kinds of antioxidants: synthetic antioxidants (butylated-hydroxy toluene and ethoxyquin; diet I), a tocopherols-rich mixture (diet II) and a tocopherols-rosemary extract mixture (diet III). A comparative study of the sensory and physical changes observed in the corresponding frozen products was undertaken.

RESULTS: After 18 months at -18°C , fish previously fed on diet I showed higher putrid and rancid odours and rancid taste scores, while lower mean typical odour and taste values were attained. Dripping and expressible moisture values obtained for diet II-fish were lower when compared with their counterparts belonging to the diet I; additionally, microstructure analysis revealed that Z-lines integration was better preserved in fish corresponding to diets II and III.

CONCLUSION: Diet II has been recognised as being the most profitable to be employed to maintain the sensory and physical properties of the frozen product when a long-term storage is considered. Further research is to be continued to optimise the natural antioxidants profile.

Key Words: *Oncorhynchus kisutch*; diet antioxidants; frozen storage; sensory acceptance; physical properties; quality.

Running Head: Diet antioxidants and quality retention of frozen farmed Coho salmon.

INTRODUCTION

Freezing and frozen storage have largely been employed to retain fish sensory and nutritional properties before consumption or further technological process. However, both processes may cause lysis of mitochondria and lysosomes and alter the distribution of enzymes and factors affecting the rate of enzyme reactions in tissues, so that deteriorative damage in frozen fish could be accelerated.¹⁻³ Additionally, the presence of both a highly unsaturated lipid composition and relevant prooxidant compounds can facilitate an important enzymatic and non-enzymatic rancidity development.^{4,5} As a result, important detrimental effects on fish muscle properties such as colour, odour, water leaching and texture take place.

To extend the shelf life of frozen fish species, many efforts have been based on the employment of antioxidants. Since synthetic antioxidants have been reported to behave as carcinogen and mutating agents, more attention has been addressed to natural antioxidants addition.⁶ Accordingly, recent research has focused on the employment of endogenous-type antioxidants (namely, tocopherols and organic acids)⁷ or antioxidants present in plant extracts⁸ to enhance quality of frozen fish products.

Great attention has also been addressed to the antioxidants presence in farmed fish feeds. Fish farmers have included a wide range of allowed synthetic⁹ and natural¹⁰ antioxidants in order to retain the quality of the corresponding processed food. In this sense, an increasing interest is being accorded to the employment of natural ones during the growing process of fish species;^{11,12} thus, in addition to food quality enhancement, the presence of such natural antioxidants in the diet would increase the oxidative stability of the living fish, as well as increase the functional properties in the corresponding food product.

Among farmed fish, Coho salmon (*Oncorhynchus kisutch*), also called silver salmon, has received great attention because of its increasing production in countries like Chile, Japan and Canada in parallel to important capture production in countries such as USA, Russian Federation, Canada and Japan.¹³⁻¹⁴ The present work focuses on the quality loss of this fatty species during the frozen storage. For this reason, Coho salmon (*Oncorhynchus kisutch*) was fed a conventional diet that was enriched with different kinds of antioxidants: synthetic antioxidants (butylated-hydroxy toluene and ethoxyquin; diet I), a tocopherols-rich mixture (diet II) and a tocopherols-rosemary extract mixture (diet III). The basic aim of the study was to investigate the effect of the antioxidants profile in diet on the sensory and physical properties of the corresponding frozen product.

MATERIALS AND METHODS

Fish and diets

Coho salmon fish used in this study were farmed in three different tanks by EWOS Innovation Research (Colaco, Puerto Montt, Chile), according to the experimental conditions reported in a parallel study.¹⁵ Thus, sandbed-filtered seawater (salinity range: 31.3-33.1 g kg⁻¹) was supplied to each tank over a temperature range of 11.2-12.8 °C. Feeding to satiety was carried out during the lighted period (photoperiod: 5.8-7.8 h) by supplying a diet with the following general composition (g kg⁻¹): 430 (protein), 290 (fat), 70 (moisture), 65 (ash), 13 (crude fibre) and 132 (carbohydrates). The distribution of fat composition into saturated, monounsaturated and polyunsaturated fatty acid groups was 325.4, 269.6 and 401.6 g kg⁻¹, respectively.

According to the objectives of the work, each of the three tanks was fed on a different antioxidant composition.¹⁵ Thus, the first tank was fed on an antioxidant

mixture including 19.3, 3.0 and 22.4 mg kg⁻¹ of EQ, BHT and total tocopherols, respectively (diet I); the antioxidant mixture supplied to the second tank included 2.9, 0.0 and 101 mg kg⁻¹ of EQ, BHT and total tocopherols, respectively (diet II); and the third tank was fed on an antioxidant mixture including 2.3, 0.0, 45, 18 and 13 mg kg⁻¹ of EQ, BHT, total tocopherols, carnosic acid and carnosol, respectively (diet III).

Fish processing and sampling

Once Coho salmon individuals attained ca. 2.500g weight, 30 fish per tank were withdrawn, sacrificed by a sharp blow to the head, the gills cut and bled in a water-ice mixture, beheaded, gutted and kept in ice for 24 hours until they arrived at our laboratory. The fish were then frozen at -40°C in individual polyethylene bags, this including vacuum sealing at 400 mbar. After 3 days, the fish were stored in a freezing room at -18°C. Frozen fish were taken for analysis on months 0, 3, 6, 9, 12 and 18 of storage at -18°C. From each tank under study, five different individuals were analysed independently (n = 5) at each sampling time after fish thawing; thawing was carried out by overnight storage (12 h) in a cool room (4°C).

In order to achieve the sensory and physical analyses, sampling was carried out according to the extraction methodology previously proposed¹⁶ and adapted to the actual species. Thus, the different muscle zones employed for each kind of quality analysis are indicated in Figure 1. All chemicals used were reagent grade (E. Merck; Darmstadt, Germany).

Sensory analysis

The Quantitative Descriptive Analysis (QDA) method was applied in order to assess changes in sensory properties of Coho salmon.^{14,16} Ten panellists (five females and five

males) with experience in sensory testing were selected and trained according to international standards.^{17,18} During the training sessions, the sensory descriptors for thawed- and thawed-cooked salmon were discussed and analysed by the panellists on samples of varying quality conditions.

At each sampling time of the present study, 4 cm x 4 cm x 2 cm steaks (N = 20) were removed from each individual salmon according to Figure 1 (zone 4). Individual steaks were then placed in polyethylene bags, coded with 3-digit random numbers and employed for thawed (N = 10) and thawed-cooked (N = 10) fish analysis; each panellist was given a single thawed steak and a single thawed-cooked steak of each individual fish. Cooking was accomplished by suspending the bags in a circulating water bath heated to $100^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 15 min to an internal temperature of $67^{\circ}\text{C} \pm 2^{\circ}\text{C}$ and then the samples were submitted independently to the panel.

At each sampling time, sensory evaluation began by the analysis of the salmon muscle in the raw state and was followed by the muscle analysis in the cooked state. Both kinds of samples were analysed at a single sitting, according to the previous training received by the panellists. At each sitting, samples were submitted independently and in a randomised order to the panellists; once each score sheet was filled in, the following sample to be analysed was received. Diluted tea without sugar was served to panellists to clean the mouth.

The different sensory descriptors (typical, rancid and putrid odour and taste; muscle dehydration; elasticity) were evaluated on a non-structured linear scale with numerical scores from 0 to 10. A score of 0 corresponds to an undetected value for the descriptor, while 10 score corresponds to the highest detectable value for the descriptor. Odour, muscle dehydration and elasticity were analysed on thawed fish; taste was analysed on thawed-cooked fish. Scores among panellists were averaged. For odour and

taste descriptors, score 5.0 was considered the borderline of acceptability; scores from the remaining descriptors were comparatively discussed without considering a borderline of acceptability.

In addition, the red colour appearance was evaluated by the Roche SalmoFan™ Lineal card. For it, panellists matched the salmon muscle colour with a 20-34-score card system previously established for salmonids pigmented with astaxanthin.¹⁸ In this card, 20-score and 34-score represent the lightest and the most intense salmon colours, respectively.

Physical analyses

Water loss assessment

The expressible moisture content was measured as the amount of fluid extracted from fillets subjected to compression.¹⁹ An Universal Testing Machine (LR-5K; Lloyd Instruments Limited, Hampshire, England, UK) was used with a load cell of 100 N connected to a computer and a Dapmat 40-0465 software for data analysis (version 3.05, Lloyd Instruments Limited, Hampshire, England, UK). Cylindrical samples (diameter = 20 mm, length = 20 mm) were extracted according to Figure 1 (zone 2) and tempered at 4°C for 4 h prior to analysis. Samples were compressed into 14 mm at a constant speed of 12.7 mm min⁻¹. The fluid exudate was received in a Whatman No. 4 filter paper. The expressible moisture was calculated as the weight difference of the filter paper before and after compression. Results were expressed as g exudate kg⁻¹ sample. Each analysis was conducted in quadruplicate for each fish sample.

Dripping loss was measured according to Rørå and Einen.²⁰ For it, whole fish were packed in sealed bags with an absorbent pad placed between the skin and the vacuum pack. The absorbent pad absorbed all liquid leakage from salmon during

freezing, frozen storage and thawing. At each sampling time, vacuum packs were opened and the wet absorbent pad was weighted. Water loss (dripping) was calculated as g kg⁻¹ fish.

Texture analysis

Shear test was used to evaluate the firmness and deformation in thawed salmon muscle.^{19,21} The force-deformation curve was obtained from an Universal Testing Machine (LR-5K; Lloyd Instruments Limited, Hampshire, England, UK) with a load cell of 100 N connected to a computer, this including a Dapmat 40-0465 software data analysis (version 3.05, Lloyd Instruments Limited). Samples of 4 cm x 4 cm x 2 cm from salmon muscle (zone 3; Figure 1) were extracted in quadruplicate and kept at 4°C ± 1°C for 4 h prior to analysis. The samples were sheared using a Warner-Bratzler steel blade (knife-edge 60°; 1.2 mm thick, 155 mm high, and 150 mm width) at a constant speed of 60 mm min⁻¹. In a force versus deformation curve, the peak of maximum force (N) required for shearing the sample was recorded as the shear force and represents the maximum resistance of the sample to shear (N). Deformation (mm) was measured during the downward movement of the blade and was calculated as the deformation (mm) at maximum peak force. Each analysis was conducted in quadruplicate for each salmon individual.

Microstructure analysis

Small pieces (0.5 mm thickness) (zone 1; Figure 1) of Coho salmon muscle were fixed for 18 h in a 30 g l⁻¹ solution of glutaraldehyde in 0.1 M Na-cacodylate buffer (pH 7.2) at room temperature and subsequently rinsed (3 × 20 min) in 0.1 M Na-cacodylate buffer. Samples were then post-fixed on ice for 90 min in an aqueous solution of 10 g l⁻¹

osmium tetroxide in 0.2 M Na-cacodylate, and rinsed for 30 min in distilled water. The samples were then stained for 60 min in an aqueous solution of 10 g l⁻¹ uranyl acetate, dehydrated for 20 min in acetone-increasing concentrations (500, 700, 950 and 1000 g l⁻¹) and embedded in 1/1 epon/acetone overnight. The muscle tissues were then immersed in epoxy resin, being the polymerisation of the resin carried out at 60°C for 24 h. Samples were cut to 1 micron in Sorvall MT-IIB ultramicrotome (Dupont Company, Newtown, CT, USA) and were stained with toluidine blue. Ultrathin sections obtained from the selected areas were stained with 40 g l⁻¹ uranyl acetate in methanol and then with lead citrate according to Reynolds.²² Finally, the sections were analysed by Transmission Electron Microscopy (TEM) (Philips Tecnai 12, Eindhoven, The Netherlands) to 80 kV. Analyses were carried out on fish samples corresponding to months 0, 9 and 18 of frozen storage.

Statistical analyses

Three variance checking methods (Cochran's C test, Bartlett's test, Hartley's test) were performed to verify that each of the quality parameters applied was consistent with a normal distribution. In negative cases, the nonparametric Kruskal-Wallis test was employed ($p < 0.05$); otherwise, data were subjected to variance analysis (ANOVA) to explore differences by two different ways ($p < 0.05$) in order to assess the effect of diet antioxidants supplied and the effect of the frozen storage time. In the event of any significant difference, a multiple range test (HSD, Tukey method) was applied. Correlation analyses between sensory and physical quality parameters and frozen storage time were studied according to the Pearson method; correlations of sensory and physical parameters with chemical quality indices corresponding to parallel research¹⁵ were also analysed. For it, Statgraphics Plus[®] version 5.1 software (Manugistics Inc.,

Statistical Graphics Corporation, 2001, Rockville, MA, USA) was employed, being in all cases a confidence interval at the 95% level ($p < 0.05$) considered.

RESULTS AND DISCUSSION

Odour and taste assessment

Progressive decreasing values with the frozen storage time could be observed for both typical odour ($r = 0.94-0.76$, linear fitting, $p < 0.05$; Figure 2) and taste ($r = 0.90-0.80$, logarithmic fitting, $p < 0.05$; Figure 3) in all kinds of samples; this decrease agrees with a previous work on the same species stored under similar frozen conditions (-18°C up to 15 months).¹³ Meantime, a general low rancid odour ($r = 0.97-0.87$; quadratic fitting, $p < 0.05$; Figure 2) and taste ($r = 0.98-0.92$; quadratic fitting, $p < 0.05$; Figure 3) development with storage time could be observed in all cases; for both descriptors, a negligible development in the 0-9-month period was followed by an increase in the 12-18-month interval, being all scores under 2.7. Finally, a very low putrid odour and taste development could be observed; thus, all values were included in the 0-1-score range (Table 1).

Concerning the comparative study of diets, fish belonging to diet I showed a higher quality decrease at the end of the experiment when compared with their counterparts from diets enriched with natural antioxidants; at that time, higher ($p < 0.05$) putrid and rancid odour and rancid taste scores were obtained in diet-I fish, while lower mean typical odour and taste values were attained in such samples. Between diets enriched with natural antioxidants, a lower ($p < 0.05$) putrid and rancid development in odour and taste could be detected at the end of the experiment in salmon fed on diet including high levels of tocopherols (diet II).

Rancidity development has been shown to be a limiting factor of acceptability for most fatty fish species under frozen storage conditions.^{4,7} Lipid oxidation is recognised as a complex process where different kinds of molecules are produced, most of them unstable and thus breaking down into smaller molecular weight ones or reacting with other compounds.^{23,24} Among them, secondary lipid oxidation compounds (namely, carbonyls) are known to be the most closely related to the rancid odour and taste development.^{25,26}

In a related study,¹⁵ individuals fed on diet II showed a greater rancidity stability (conjugated diene content, peroxide and anisidine values, thiobarbituric acid index, fluorescence and browning development) and polyunsaturated fatty acid retention (polyene index assessment) when compared with their counterparts from the diet I. Comparison of chemical and sensory indices related to lipid oxidation development showed that rancid odour and taste obtained their best correlation values when compared with the thiobarbituric acid index ($r = 0.87-0.75$ and $0.95-0.71$, respectively, $p < 0.05$).

The antioxidant content in salmon muscle was also analysed in a parallel study.²⁷ In it, BHT and EQ contents in fish muscle were found in all cases below 0.01 and 0.03 mg kg^{-1} concentration, respectively. Additionally, feeding on diet II led to frozen samples showing higher contents of γ - and δ -tocopherol when compared to their counterparts previously fed on diet I; however, no effect on α -tocopherol presence was implied. Further, a marked loss of α -tocopherol content was observed with frozen time for all kinds of samples, in agreement with the general rancidity development observed throughout the frozen storage in sensory and chemical parameters.

Determination of colour changes

The Roche card employment (Table 1) showed a decreasing tendency for pink colour intensity with storage time in all kinds of individuals ($r = 0.91-0.80$; linear fitting, $p < 0.05$); this result agrees with previous research carried out on Atlantic salmon (*Salmo salar*) frozen at -10 and -20°C for a 34-week period.²⁶ However, no differences could be assessed in the present study among individuals from the different diets, so that an effect of the antioxidant composition in diet on the pink colour intensity in thawed fish could not be proved.

Colour plays an important role in the appearance presentation and acceptability of seafood, especially in those related to salmonid species. Astaxanthin (AX) is well known as the main pigment responsible for the pink colour of salmonid fish species and could deteriorate either due to non-enzymatic or to enzymatic pathways.²⁸ Thus, previous studies have shown that AX content decreases during the frozen storage of Atlantic salmon (*Salmo salar*)²⁶ and rainbow trout (*Oncorhynchus mykiss*).²⁹ However, and in agreement with the present results, related research concerning colour analysis (CIE 1976 L^* , a^* , b^* ; lightness, greenish/reddish and bluish/yellowish values, respectively) in Coho salmon²⁷ showed no effect on AX content and a^* and b^* colour parameters during the frozen storage of Coho salmon corresponding to the three diets under study, while a general L^* value increase was implied. Additionally, no effect of the antioxidant profile could be depicted on AX content and colour (L^* , a^* and b^*) parameters²⁷. This AX content retention and accordingly, colour properties retention, may be explained on the basis of the protection exerted by the natural and synthetic antioxidants present in the different diets.¹⁵

Texture parameters assessment

Firmness assessment (Table 2) has shown a general increase with storage time throughout the 0-9-month period, followed by a score decrease at month 12 and a firmness increase at the end of the experiment. Concerning comparison among the different kinds of samples, some significant differences could be observed; however, a definite pattern related to the effect of the antioxidants profile in diet on firmness value could not be assumed.

Cohesivity analysis (Table 2) showed in most cases the highest mean values in month-9 samples. After that time, a general cohesivity decrease could be observed. Throughout the frozen storage, some differences were obtained among individuals from the different diets; thus, a lower mean value was obtained at the end of the experiment for samples previously fed on the diet enriched with synthetic antioxidants (diet I). However, a definite trend concerning the effect on the cohesivity score of the antioxidants profile included in the diet could not be proved.

Elasticity was evaluated by sensory procedure (Table 2), according to the muscle recovery after exerting a short pressure. A marked elasticity loss could be observed in all kinds of samples with frozen storage time ($r = 0.90-0.79$; quadratic fitting, $p < 0.05$), this leading to values included in the 4-5 range for individuals fed on diet III and diet I at month 9 and at months 12 and 18, respectively. Comparison among diets did not lead to a definite effect on elasticity values during the frozen storage period.

Texture is considered to be one of the most important quality attributes of seafood, which often determines consumer acceptance and hence, the marketability of such products. Protein denaturation has widely been described to be produced during the frozen storage of fish.^{1,2} Different hypotheses have been proposed to explain the denaturation development that finally leads to a drier and firmer muscle texture. The

connective tissue holding the cells together is reported to be degraded as muscle damage increases and blocks of cells become readily separated from each other, so that lower cohesivity and water holding capacity properties would be observed in muscle; additionally, elasticity decrease has been explained as a result of myofibrillar protein denaturation and aggregation.^{3,30}

In the case of fatty fish, the role of fat in the texture changes has been found to be important due to crosslinking of peptide chains by reaction with lipid oxidation compounds produced during processing and storage^{1,24} and leading to an increased firmness and decreased water holding capacity.³¹ In agreement with present rancid odour and taste values, a parallel study¹⁵ showed that frozen Coho salmon muscle acquired a higher rancidity stability if previously fed on diet II when compared with its counterpart fed on diet I. Present results have shown that, in spite of this lipid stability advantage, no definite effect on texture properties could be outlined in fish corresponding to diet II.

Water loss determination

In all three quality parameters tested (Table 3), increasing water loss values with frozen storage time could be observed for all kinds of samples. This general behaviour can be explained on the basis of an increasing protein denaturation, being this explained as a result of different factors such as partial protein dehydration, interaction with oxidised lipids, free fatty acids and formaldehyde and alteration of the protein micro-environment¹⁻².

Measurement of muscle dehydration showed some lower ($p < 0.05$) scores for individuals previously fed on the diet enriched with synthetic antioxidants (months 3 and 9); however, no differences ($p > 0.05$) could be detected at the end of the experiment.

Concerning dripping and expressible moisture assessments, a definite trend about the effect of the antioxidant composition in the diet could not be concluded; however, lower mean values ($p<0.05$) for both parameters were observed at month 18 in fish individuals previously fed on diet II when compared with samples corresponding to diet I.

Microstructure analysis

Concerning frozen samples at time 0 (Figure 4), striated muscle of diet I-fish consisted of an orderly structure of actin and myosin filaments. Thus, light bands (I-bands and dark A-bands) can be observed alternately arranged, while a thin dark line (Z-line or membrane of Krause) is found to bisect each I-band. A bright area (H-area or Hansen membrane) is located in the centre of the A-band, while a narrow dark line (M-line) runs through the centre of the H-band. Finally, the endomysium is observed to remain intact around each of the different fibres.

At that frozen time (month 0), the addition of a tocopherols-rich mixture in the diet of salmon (diet II) led to slight changes in the myofibrils morphology. Thus, sarcomeres showed well-marked areas for the I bands, A bands and Z line, while the H area showed a diffuse line of the Hansen membrane. The muscle fibres showed a slight shift and the Z line was observed somewhat misaligned. Concerning fish corresponding to diet III, no loss of alignment in the myofibrils morphology could be observed; thus, the sarcomeres network was composed of well-aligned and defined Z lines.

After month 9 of frozen storage (Figure 4), fish belonging to diets I and III showed some displacement and misalignment of myofibrils, this including the Z-line, M-line, and I-band; however, no loss of the sarcomeres basic unit in the striated muscle could be depicted. Misaligned sarcomeres were observed especially for fish corresponding to diet II; thus, discontinuities in the Z-line, I-band and A-band were

produced as a result of myofibrils displacement, although no loss of the basic unit of the striated muscle was revealed.

At the end of the experiment (Figure 4), marked changes related to the microstructure of the sarcomeres were observed in diet I-fish samples. Thus, the basic unit of the striated muscle did not present well-demarcated areas of the Z-line, A-band, and H-area. Moreover, I-band was observed misaligned. The disintegration of the Z-line has been consistently reported as a post-mortem phenomenon, being this fragmentation generally associated with calpain (Ca^{+2} -dependent protease) activity.³² Meantime, the microstructure of the sarcomeres remained unchanged for Coho salmon muscle previously fed on diets II and III, this indicating that the addition of natural antioxidants does not affect this parameter until month 18 of storage. It is concluded that the typical characteristics of striated muscle and the sarcomeres morphology were influenced throughout the frozen storage by the antioxidants profile included in the previous diet so that a better preservation of frozen samples previously fed on diets II and III was proved at the end of the frozen storage.

CONCLUSIONS

After 18 months at -18°C , fish previously fed on a conventional diet enriched with natural antioxidants (diets II and III) showed lower putrid and rancid odours and rancid taste scores, while higher mean typical odour and taste values were attained. Concerning physical parameters, dripping and expressible moisture values obtained at month 18 for diet II-fish were lower when compared with their counterparts belonging to the diet enriched with synthetic antioxidants (diet I). Additionally, microstructure analysis showed that the Z-lines integration was better preserved in fish corresponding to diets II and III after frozen storage for 18 months when compared with their corresponding

404 samples belonging to diet I. Between diets enriched with natural antioxidants, a lower
405 putrid and rancid development in odour and taste could be detected at the end of the
406 experiment in salmon fed on diet including an excess of tocopherols (diet II).

407 According to the present results, the employment of diet II has been recognised
408 as being the most profitable to be employed to maintain the sensory and physical
409 properties of the frozen product when a long-term storage is considered. In agreement
410 with the wide range of benefits attributed to natural antioxidants, further research is
411 currently under development to optimise the natural antioxidants profile to be employed
412 on each farmed species.

415 **Acknowledgements**

416 The authors thank EWOS Innovation Research (Colaco, Puerto Montt, Chile) for
417 kindly providing the Coho salmon fish and the “Departamento de Investigación,
418 Vicerrectoría de Investigación y Desarrollo de la Universidad de Chile”. The study was
419 supported by the Chilean-Spanish Cooperation Program “Universidad de Chile-Consejo
420 Superior de Investigaciones Científicas (CSIC)” (Project 2004 CL 0038) and the
421 FONDECYT program (Chile; project number 1120627). Prof. Julia Vinagre Leiro is
422 gratefully acknowledged for her valuable advice, expertise and encouragement that have
423 made possible the actual collaborative study.

FIGURE LEGENDS

Figure 1: Location of the different fish muscle zones employed for the different sensory and physical analyses: (1) microstructure, (2) water loss (expressible moisture); (3) texture (shear test); (4) sensory analysis (odour, taste, elasticity, colour and muscle dehydration). Dripping loss was obtained in whole fish.

Figure 2: Typical (TO) and rancid (RO) odour development* in frozen Coho salmon muscle which was previously fed on different diets**.

* Mean values of five (n = 5) replicates. Standard deviations are denoted by bars.

** Diets I, II and III include an antioxidant mixture enriched in ethoxyquin/butylated-hydroxytoluene, tocopherols and tocopherols/polyphenols (carnosic acid and carnosol), respectively, as expressed in the Material and Methods section.

Figure 3: Typical (TT) and rancid (RT) taste development in frozen Coho salmon muscle which was previously fed on different diets*.

* Mean values of five (n = 5) replicates. Standard deviations are denoted by bars. Diets employed as expressed in Figure 2.

Figure 4: Evolution of the muscle structure of frozen (a: 0 months; b: 9 months; c: 18 months) Coho salmon muscle which was previously fed on different diets*.

* Diets employed as expressed in Figure 2. The bar indicates a 1- μ m length.

REFERENCES

1. Sikorski Z and Kolakowska A, Changes in protein in frozen stored fish, in *Seafood proteins*, ed. by Sikorski Z, Sun Pan B and Shahidi F. Chapman and Hall, New York, USA, pp. 99-112 (1994).
2. Sotelo M^aC, Piñeiro C and Pérez-Martín R, Denaturation of fish proteins during frozen storage: Role of formaldehyde. *Z Lebensm Unters Forsch* **200**: 14-23 (1995).
3. Barroso M, Careche M and Borderías AJ, Quality control of frozen fish using rheological techniques. *Trends Food Sci Technol* **9**: 223-229 (1998).
4. Erickson M, Lipid oxidation: flavour and nutritional quality deterioration in frozen foods, in *Quality in frozen food*, ed. by Erickson M and Hung YC. Chapman and Hall, New York, USA, pp. 141-173 (1997).
5. Aubourg S, Stodolnik L, Stawicka A and Szczepanik G, Effect of a flax seed (*Linum ussitatissimum*) soaking treatment on the frozen storage stability of mackerel (*Scomber scombrus*) fillets. *J Sci Food Agric* **86**: 2638-2644 (2006).
6. Frankel E, Natural and biological antioxidants in foods and biological systems: their mechanism of action, applications and implications. *Lipid Technol* **July**: 77-80 (1995).
7. Medina I, Gallardo J and Aubourg S, Quality preservation in chilled and frozen fish products by employment of slurry ice and natural antioxidants. *Int J Food Sci Technol* **44**: 1467-1479 (2009).
8. Tironi V, Tomás M and Añón M^aC, Quality loss during the frozen storage of sea salmon (*Pseudoperca semifasciata*). Effect of rosemary (*Rosmarinus officinalis*) extract. *Food Sci Technol (LWT)* **43**: 263-272 (2010).

- 475 9. He P and Ackman R, HPLC determination of ethoxyquin and its major oxidation
476 products in fresh and stored fish meals and fish feeds. *J Sci Food Agric* **80**: 10-
477 16 (2000).
- 478 10. Hamre K, Berge R and Lie O, Oxidative stability of Atlantic salmon (*Salmo salar*,
479 L.) fillet enriched in α -, γ - and δ - tocopherol through dietary supplementation.
480 *Food Chem* **62**: 173-178 (1998).
- 481 11. Jittinanadana S, Brett Kenney P, Slider S, Kamireddy N and Hankins J, High dietary
482 vitamin E affects storage stability of frozen-refrigerated trout fillets. *J Food Sci*
483 **71**: C91-C95 (2006).
- 484 12. Choubert G and Baccaunaud M, Colour changes of fillets of rainbow trout
485 (*Oncorhynchus mykiss* W.) fed astaxanthin or cantaxanthin during storage under
486 controlled or modified atmosphere. *Food Sci Technol (LWT)* **39**: 1203-1213
487 (2006).
- 488 13. Rodríguez A, Losada V, Larraín M^aA, Quiral V, Vinagre J and Aubourg S,
489 Development of lipid changes related to quality loss during the frozen storage of
490 farmed Coho salmon (*Oncorhynchus kisutch*). *J Amer Oil Chem Soc* **84**: 727-
491 734 (2007).
- 492 14. Rodríguez A, Carriles N and Aubourg S, Effect of chill storage under different icing
493 conditions on sensory and physical properties of canned farmed salmon
494 (*Oncorhynchus kisutch*). *Int J Food Sci Technol* **45**: 295-304 (2010).
- 495 15. Ortiz J, Larraín M^aA, Vivanco J and Aubourg S, Rancidity development during the
496 frozen storage of farmed coho salmon (*Oncorhynchus kisutch*): Effect of
497 antioxidant composition supplied in the diet. *Food Chem* **115**: 143-148 (2009).
- 498 16. Sveinsdóttir K, Martinsdóttir E, Hyldig G, Jorgensen B and Kristbergsson K,
499 Application of quality index method (QIM) scheme in shelf-life study of farmed

500 Atlantic salmon (*Salmo salar*). *J Food Sci* **67**: 1570-1579 (2002).

501 17. ISO 3972. *Sensory analysis-methodology-method of investigating sensitivity of*
502 *taste*. International Standardization Organization, Geneva, Switzerland (1991).

503 18. Codex Alimentarius. *Directrices para la Evaluación Sensorial del Pescado y los*
504 *Mariscos en Laboratorio*. Norma Técnica CAC/GL 31-1999 (1999).
505 <http://www.pes.fvet.edu.uy/cursos/mail2.pdf>.

506 19. Jonsson A, Sigurgisladóttir S, Hafsteinsson H and Kristbergsson K, Textural
507 properties of raw Atlantic salmon (*Salmo salar*) fillets measured by different
508 methods in comparison to expressible moisture. *Aquac Nutr* **7**: 81-89 (2001).

509 20. Rørå A and Einen O, Effects of freezing on quality of cold-smoked salmon based on
510 the measurements of physiochemical characteristics. *J Food Sci* **68**: 2123-2128
511 (2003).

512 21. Sigurgisladóttir S, Hafsteinsson H, Jonsson A, Nortvedt R, Thomasses M and
513 Torrisen O, Textural properties of raw salmon fillets as related to sampling
514 method. *J Food Sci* **64**: 99-104 (1999).

515 22. Reynolds E, The use of lead citrate at high pH as an electron-opaque stain in
516 electron microscopy. *J Cell Biol* **17**: 208-212 (1963).

517 23. Howell N, Interaction of proteins with small molecules, in *Ingredient Interactions-*
518 *Effects on Food Quality*, ed. by Gaonkar A. Marcel Dekker, New York, USA,
519 pp. 269-289 (1995).

520 24. Tironi V, Tomás M and Añón M^aC, Structural and functional changes in
521 myofibrillar proteins of sea salmon (*Pseudoperca semifasciata*) by interaction
522 with malondialdehyde (RI). *J Food Sci* **67**: 930-935 (2002).

- 523 25. White P, Conjugated diene, anisidine value and carbonyl value analyses, in *Methods*
524 *to assess quality and stability of oils and fat-containing foods*, ed. by Warner K
525 and Eskin M. AOCS Press, Champaign, Illinois, USA, pp. 159-178 (1994).
- 526 26. Refsgaard H, Brockhoff P and Jensen B, Sensory and chemical changes in farmed
527 Atlantic salmon (*Salmo salar*) during frozen storage. *J Agric Food Chem* **46**:
528 3473-3479 (1998).
- 529 27. Ortiz J, Larraín M^aA, Pacheco N, Vivanco J and Aubourg S, Effect of the
530 antioxidant profile in the diet of farmed salmon (*Oncorhynchus kisutch*) on the
531 nutritional value retention during frozen storage. *Grasas y Aceites* **64**: 311-319
532 (2013).
- 533 28. Krinsky N, Antioxidant functions of carotenoids. *Free Rad Biol Med* **7**: 617-635
534 (1989).
- 535 29. Jensen C, Birk E, Jokumsen A, Skibsted L and Bertelsen G, Effect of dietary levels
536 of fat, alpha-tocopherol and astaxanthin on colour and lipid oxidation during
537 storage of frozen rainbow trout (*Oncorhynchus mykiss*) and during chill storage
538 of smoked trout. *Z Lebensm Unters Forsch* **207**: 189-196 (1998).
- 539 30. Bjerkeng B and Johnsen G, Frozen storage quality of rainbow trout (*Oncorhynchus*
540 *mykiss*) as affected by oxygen, illumination, and fillet pigment. *J Food Sci* **60**:
541 284-288 (1995).
- 542 31. Lou X, Wang C, Xiong Y, Liu B and Mims S, Physicochemical stability of
543 paddlefish (*Polyodon spathula*) meat under refrigerated and frozen storage. *J*
544 *Aquat Food Prod Technol* **9**: 27-39 (2000).
- 545 32. Morzel M, Heapes M, Reville W and Arendt E, Textural and ultrastructural changes
546 during processing and storage of lightly preserved salmon (*Salmo salar*) products.
547 *J Sci Food Agric* **80**: 1692-1697 (2000).