Phenological growth stages of longan (*Dimocarpus longan*) according to the BBCH scale

V.T. Pham^{a, c}, M. Herrero^b, J.I. Hormaza^{c, *}

^a Department of Plant Resources, Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, Ha Noi, Vietnam. Email: phamvthe@gmail.com

^b Pomology Department, Estación Experimental de Aula Dei - CSIC, Avda Montañana 1005, 50059 Zaragoza, Spain. Email: mherrero@eead.csic.es

^c Subtropical Fruit Department, Instituto de Hortofruticultura Subtropical y Mediterránea la Mayora (IHSM la Mayora-CSIC-UMA), 29750 Algarrobo-Costa, Malaga, Spain.

* Corresponding author. Tel.: +34 952548990; fax: +34 952552677. E-mail: ihormaza@eelm.csic.es (J.I. Hormaza).

ABSTRACT

Longan (*Dimocarpus longan*) is an important evergreen fruit crop grown in tropical and subtropical climates, with a clear expanding potential, but with a poorly described phenology. In this work, the different phenological growth stages of longan are characterized according to the BBCH (Biologische Bundesanstalt, Bundessortenamt und Chemische industrie) scale. From vegetative bud dormancy to fruit harvest, seven main growth stages are described for bud, leaf and shoot development, inflorescence emergence, flowering, fruit development and fruit maturity. Within them 41 secondary growth stages are described. The BBCH code will be an important tool to assist the development and implementation of longan management protocols and to standardize observations made in different edaphoclimatic and experimental conditions.

Keywords: BBCH code; *Dimocarpus longan*; Sapindaceae; Phenology

1. Introduction

The longan (*Dimocarpus longan* Lour.) is a subtropical fruit tree species native from Southern Asia. The species is a member of the Sapindaceae, a family that includes other fruit crops such as litchi (*Litchi chinensis* Sonn.), rambutan (*Nephelium lappaceum* L.), guarana (*Paullinia cupana* Kunth), Spanish lime (*Melicoccus bijugatus* Jacq.), ackee (*Blighia sapida* KD Koenig), or pulasan (*Nephelium mutabile* Blume).

The longan was first described in 1790 by the Jesuit missionary and botanist Joao de Loureiro (Loureiro, 1790) although it has a documented history of more than 2200 years in China (Huang et al., 2005), based in its excellent nutritive and pharmaceutical potential (Wong, 2000). The genus *Dimocarpus* contains six species of trees and shrubs and *D. longan* has two subspecies and five varieties distinguished mainly by the leaflet structure (Subhadrabandhu and Stern, 2005): *D. longan* ssp. *longan* var. *longan* (the most important for fruit consumption) originally distributed from Myanmar to southern China, south-west India and Sri Lanka; *D. longan* ssp. *longan* var. *longepetiolulatus* originally distributed in southern Vietnam; *D. longan* ssp. *longan* var. *obtusus* originally distributed in Indochina; *D. longan* ssp. *malesianus* var. *malesianus* and *D. longan* ssp. *malesianus* var. *echinatus* originally distributed in northern Borneo and southern Philippines (Loureiro, 1790; Leenhouts, 1971; Wong and Ketsa, 1991; Subhadrabandhu and Stern, 2005). Nowadays, longan is also cultivated in other regions with tropical to temperate climates in China and South-East Asia, Queensland in Australia, and Florida in the United States, but has a further potential in other countries. The main world longan producing countries are China, with about 1.300 million tons in 2010 (Qiu, 2014), Vietnam, with more than 550 million tons in 2012 (Ministry of Agriculture and Rural Development of Vietnam, 2012), and Thailand with more than 500 million tons in 2014 (Buara and Kumcha, 2014).

Longan is an evergreen tree that can grow up to 20 m high forming alternate paripinnate leaves with 6 to 9 opposite leaflets (Subhadrabandhu and Stern, 2005; Wong and Ketsa, 1991). As other evergreen tropical and subtropical trees, longan produces one or more flushes of shoot

growth per year (Nakasone and Paull, 1998; Diczbalis, 2002; Paull and Duarte, 2011). Usually the species has one flowering flush per year but some cultivars can produce three in a two year period (Wong, 2000). The longan inflorescences are compound dichasia, terminal, usually leafless, erect, with widely branched panicles 8-60 cm long (Subhadrabandhu and Stern, 2005; Davenport and Stern, 2005). The flowers are unisexual, small, yellow-brown with five petals and appear generally in three waves in the inflorescence with different degrees of overlap among them depending on the cultivars and the environmental conditions: a first wave of male flowers, followed by a wave of female flowers and, finally, a third wave of male flowers. The female flowers have a bicarpellate ovary although usually only one locule will develop into a fruit (Davenport and Stern, 2005). Each inflorescence contains a large number of flowers, approximate 200-1000 functionally female flowers and 1000-4000 functionally male flowers. Usually, the flowers open at night and pollination takes place in the morning by different insects, mainly honeybees (Davenport and Stern, 2005; Pham et al., 2013). The time from flowering to harvest lasts from 4 to 6 months depending on the environmental conditions and the cultivars. After fertilization, usually two fruit abscission waves take place and finally each panicle can carry about 60-80 fruits (Davenport and Stern, 2005). The edible portion of the longan fruit is a fleshy aril (Wong, 2000).

While several studies on production, floral induction, fruit quality and aril composition, have been performed in longan, little information is available on longan phenology. Previous research has studied fruit development and some phenological stages (Nakasone and Paull, 1998; Diczbalis, 2002; Paull and Duarte, 2011), but to have a complete picture additional work and a standardization of the observations are needed. Thus, in this work, we propose to apply the extended BBCH (Biologische Bundesanstalt, Bundessortenamt und Chemische industrie) scale system to describe the phenological growth stages of longan contributing to the standardization of phenology studies in this crop.

2. Materials and Methods

Data were collected from 15 years old adult trees of *Dimocarpus longan* ssp. *longan* cultivars ‘Choompoo’, ‘Fuk How’, ‘Biew Kiew’, and ‘DuanYu’, propagated by air layering, located at IHSM la Mayora in Malaga (Spain), at latitude 36°45N, longitude 4°4W and elevation 35m above sea level. The climate data of the last 50 years show a mean annual temperature for this location of 18.5°C, average maximum temperatures of 28.9°C in the hottest month (August) and average minimum temperature of 9.8°C in the coolest month (January), with an average annual rainfall of 536 mm.

Measurements and observations of vegetative and reproductive development were carried out during two annual growing seasons (2012-2014). A total of 82 buds located in 25 branches of 5 trees (5 branches of each) were marked and measurements were made twice or once per week, or twice per month depending on the developmental stage. During the experimental period the average maximum and minimum temperature were 26.7°C and 12.8°C, respectively.

The BBCH scale described in this work for longan uses 7 of the 10 principal growth stages starting with stage 0 (vegetative bud development), followed by stages 1 (leaf development), 3 (shoot development), 5 (inflorescence emergence), 6 (flowering), 7 (development of fruit), and ending with stage 8 (maturity of fruit). Secondary stages are described in detail. As in most fruit trees, principal growth stage 2 (formation of side shoots/tillering), 4 (development of harvestable vegetative plant parts or vegetative propagated organs/booting), and 9 (senescence, beginning of dormancy) are not considered for longan.

3. Results and Discussion

In this work, the phenological growth stages of longan according to the extended BBCH scale are described (Fig. 1), under the environmental conditions of Southern Spain, separating the vegetative and reproductive flushes (Fig. 2). The description is divided in seven main growth stages, three for vegetative growth (including bud, leaf, and shoot development) and four for reproductive development (including inflorescence emergence, flowering, fruit development, and fruit maturity). Within them a total of 41 secondary growth stages were described (Table 1).

3.1. Principal growth stage 0: vegetative bud development

The first vegetative buds were apparent usually two to four weeks after harvesting the fruits. This occurred in the winter from December to January and depended on the harvesting time (Fig. 2).

010. Vegetative buds dormant: foliar buds are completely closed and covered by brownish scales (Fig. 1).

011 Beginning of bud swell: bud scales begin to open.

013 End of bud swell: brownish scales are slightly separated (Fig. 1).

017 Beginning of bud break: compound leaf tips start to become visible (Fig. 1).

019 End of bud break: shoot tip is clearly visible and first compound leaves slightly separate (Fig. 1).

Additional flushes: Under climatic conditions where additional vegetative flushes are observed, additional mesostages can be added: 020, 021...

3.2. Principal growth stage 1: leaf development

Most leaves on the first vegetative growth flush emerged in winter, over January and February (Fig. 2). Growth phase 110–119 was completed in approximately 50-60 days.

110. First leaves separated: leaflets of the first compound leaf separating and showing reddish color.

111. First leaves unfolded: All leaflets of the first compound leaf unfolded, first leaflets at 10% of their full size or nearly 10% compound leaves unfolded.

113. More leaves unfolded: First leaflets at 30% of their full size or nearly 30% compound leaves unfolded (Fig. 1).

116. More leaves unfolded: First leaflets at 60% of their full size and changing color from red to red-green or nearly 60% compound leaves unfolded.

119. All leaves unfolded: all leaflets fully expanded, first leaflets nearly at 100% of their full size and changing from red to red-yellow-green (Fig. 1).

Additional flushes: Under climatic conditions where additional vegetative flushes are observed, additional mesostages can be added.

3.3. Principal growth stage 3: shoot development

The vegetative flush normally took place in spring (February/March) before inflorescence emergence (stage 5) and flowering (stage 6) (Fig. 2).

310. Beginning of shoot extension: axes of developing shoots visible. This growth stage proceeds in parallel with stage 110.

311. 10% of final shoot length. This growth stage proceeds in parallel with stage 113 (Fig. 1).

313. 30% of final shoot length. This growth stage proceeds in parallel with stage 116.

315. 50% of final shoot length. This growth stage proceeds in parallel with stage 119 (Fig. 1).

317. 70% of final shoot length: first leaflets changed to light green (Fig. 1).

319. 90% or more of final shoot length: all leaflets changed to light green, shoots completely developed (Fig. 1).

Additional flushes: Under climatic conditions where additional shoot development flushes are observed, additional mesostages can be added.

3.4. Principal growth stage 5: inflorescence emergence

Reproductive bud development occurred during the spring from April to early May (Fig. 2). At the end of this stage, flower bud developed completely. Short ovary, stigma and anthers can be observed under the binocular microscope after removing the petals and sepals (Fig. 2). Detailed anatomical images of different stages during reproductive bud development are provided in Fig. 3).

510. Reproductive buds dormant: completely closed and covered with brownish scales (Fig. 1).

511. Beginning of reproductive bud swell: brownish scales separating and flower buds visible (Fig. 1, Fig. 3).

512. Panicle axes begin to elongate: folded primary leaves are visible in the principal axes (Fig. 3).

513. Beginning of panicle development: primary branches visible, folded leaves clearly visible, usually the leaves will drop but sometimes they can unfold and grow as the compound leaves (Fig. 1, Fig. 3).

515. Inflorescences 50% of final length: secondary branches visible (Fig. 1, Fig. 3)).

517. Inflorescences 70% of final length: secondary branches elongated, tertiary branches visible (Fig. 1, Fig. 3).

519. End of inflorescence extension: tertiary branches almost fully developed; individual flower pedicels elongated, flowers separated and closed, corolla change from brownish to green (Fig. 1, Fig. 3)).

Additional flowering periods: Under climatic conditions where there is more than one flowering period, additional mesostages can be added.

3.5. Principal growth stage 6: flowering

The first flowers opened in mid spring in May and flowering continues for about 35-45 days in the same individual tree (Pham et al., 2013) (Fig. 2).

Similar to litchi, longan has three types of flowers (Davenport and Stern, 2005): staminate flowers (M1), functionally female hermaphrodite flowers (F), and functionally male hermaphrodite flowers (M2). It has been put forward that three waves of longan flowers are present in the inflorescences. The first wave consists of M1 flowers, the second of F flowers, and the third of M2 flowers (Davenport and Stern, 2005; Paull and Duarte, 2011; Jonathan, 2013), although climatic conditions or cultural management practices may modify this pattern (Paull and Duarte, 2011). Also there is an overlap of the different flower types in the same tree since not all the panicles are in the same developmental stage.

610. First flowers open: female (F) or male (M1) flowers open depending on tree and climate conditions (Fig. 1).

611. Beginning of flowering: 10% flowers open.

613. Early flowering: 30% flowers open (Fig. 1).

615. Mid bloom: 50% flowers open, both female (F) and male (M2) flowers can be found at this time (Fig. 1).

617. Full flowering: 70% flowers open, some initial fruit set can be found (Fig. 1).

619. End of flowering: 90% flowers open, most petals dried out and dropped, numerous fruits have already set (Fig. 1).

Additional flowering periods: Under climatic conditions where there is more than one period of flowering, additional mesostages can be added.

218

219 *3.6. Principal growth stage 7: fruit development*

220

221 Longan fruit growth took place during 4 months from June/July to October/November (Fig. 2).
222 Longan fruit development could be divided into two main growth stages (Ke et al., 1992; Zheng
223 et al., 1994; Chen et al., 1995; Xu et al., 1997). In the first stage, pericarp and aril development
224 started and the seed coat, embryo and endosperm can be observed with the naked eye. This
225 stage occurred 1-50 days after anthesis. The second stage is characterized by the embryo filling
226 the seed cavity, hardening of the seed coat, thinning of the pericarp, development of the fleshy
227 aril and the maturation process (Fig. 2). Detailed images of some of those steps are provided in
228 Fig. 4).

229 Zee et al. (1998) reported two waves of fruit abscission in longan in China. The first cycle
230 occurred 3–20 days after fertilization, and the second 30–60 days before harvest. In southern
231 Spain, the first fruit abscission takes place 1-6 weeks after anthesis. At harvest time, each
232 panicle can carry about 60-80 fruits (Davenport and Stern, 2005) although about 1000 female
233 flowers can be produced per panicle.

234

235 710. No ovary growth still visible

236 711. Initial ovary growth, following fertilization. First physiological fruit abscission (Fig.
237 1).

238 712. 20% of final fruit size. Beginning of ovary growth: one (two) ovary carpel(s) dried,
239 one fertilized ovary carpel growing: (Fig. 1, Fig. 4).

240 713. 30% of final fruit size, fruits show muricate skin (Fig. 1).

241 715. 50% of final fruit size: skin is still muricate, and the seed is covered by the aril (Fig. 1,
242 Fig. 4).

243 717. 70% of final fruit size: skin changes from muricate to rough and the aril becomes
244 fleshy (Fig. 1).

718. 80% of final fruit size (Fig. 1, Fig. 4).

719. 90% or more of final fruit size (Fig. 1).

Additional fruiting periods: Under climatic conditions where there is more than one period of flowering and fruiting, additional mesostages can be added.

3.7. Principal growth stage 8: fruit maturity

Fruit maturation took place at the end of the autumn, in November/December, depending on the year and the cultivar (Fig. 2). During this stage, the fruit is still growing. The fruits are ready for harvest when the pericarp is thin, smooth, tough and leathery, and its color changes from green-yellow to yellow-brown (Subhadrabandhu and Stern, 2005) (Fig. 2).

810. Physiological maturity: skin is more or less rough, skin color changes from green to light yellow-green; this growth stage proceeded parallel with stage 718 (Fig. 1).

811. Beginning of fruit ripening: fruit skin color changes from light yellow-green to bright yellow, surface skin changes from rough to smooth (Fig. 1); this growth stage proceeded parallel with stage 719.

815. Advanced ripening: fruit becomes smoother with yellow skin color (Fig. 4).

819. Fruit fully ripened, ready for consumption: fruit skin color fully changes to yellow-brown, the fruit is completely smooth, with 100% of final size and ready for harvest (Fig. 1).

While this is the general pattern, in southern Spain two additional vegetative flushes can take place depending on the reproductive flush and the flowering stages. The first additional flush appears if no inflorescences are produced in the branches. This flush occurs at the same time of the normal reproductive growth stage, in mid-spring to early summer, from April to June. A similar situation has been reported in Australia (Diczbalis, 2002). A second additional

flush can occur if all flowers, inflorescences or developing fruitlets fall down due to adverse weather conditions or lack of pollination. This flush occurs in mid-summer to early autumn during July and September. These additional flushes can be added as, additional mesostages: 020, 021, 023, 027, 029; 030, 031, 033, 037, 039. Likewise under climatic conditions where there is more than one period of flowering, additional mesostages can be added.

Considerable variation occurs when comparing times of flowering and fruiting in different parts of the world. Flowering in this work spanned from May-June, but in China, flowering takes place from late February to early April (Wong, 2000), in Thailand from late December to late February (Wong, 2000; Stern, 2005), in Queensland from August to October and October to November, depending on the location (Wong, 2000), in Florida in February/March through April and the beginning of May (Jonathan et al., 2013), and in northern Vietnam from the end of March to the beginning of April (Pham et al., 2013). The same applies for fruit maturity that in our conditions occurred in November-December, in China in July-September (Groff, 1921; Wong, 2000), in Thailand in June-August (Stern, 2005; Wong, 2000), in Queensland from January/February to March/April (Stern, 2005; Wong, 2000), in Florida in August-September (Jonathan et al., 2013), and in northern Vietnam from July to August (FAO, 2004). This variability provides a wide fresh fruit provision around the world, but makes difficult to envisage the phenology calendar for a new growing area. To provide a base line, the phenology recorded here is related to temperature (Fig. 2) showing the different temperature requirements for the different major phenology events. Thus, while vegetative growth proceeded between 12-15°C, flowering occurred between 15-20°C and fruit development was concomitant with the higher temperatures of 20-27°C.

In this work, the phenological stages in longan have been described for the first time according to an extended BBCH scale separating the different vegetative and reproductive flushes (Fig. 2). The description here presented will be useful not only to provide basic information on crop requirements, disease and pest management and control but also to

facilitate exchange of scientific information among longan experiments performed under different edaphoclimatic conditions.

Acknowledgements

This work was supported by the Spanish Ministerio de Economía y Competitividad—European Regional Development Fund, European Union (AGL2013-43732-R, and AGL2012-40239), Gobierno de Aragón A-43, and Junta de Andalucía (FEDERP08-AGR-3694). VTP was supported by a PhD scholarship in the frame of an agreement between the Consejo Superior de Investigaciones Científicas (CSIC), Spain and the Vietnamese Academy of Science and Technology (VAST).

References

- Buara, P., Kumcha, K., 2014. Fruit and Vegetables production in Thailand. http://www.fao.org/fileadmin/templates/agphome/documents/horticulture/WHO/seoul/F_V_Thailand.pdf. Accessed March 23, 2015.
- Chen, Q.X., Liao, J.S., Hu Y.L., 1995. The growth curve of longan fruit and the correlative analysis of its tissues. *Journal of Fujian Academy of Agricultural Sciences* 24, 19–22.
- Davenport, T.L., Stern R.A., 2005. Flowering. In: Menzel, C.M., Waite, G.K. (Eds.), *Litchi and Longan: Botany, Production and Uses*. CABI International, Wallingford, UK, pp. 87-113.
- Diczbalis, Y., 2002. *Longan: Improving Yield and Quality*. Rural Industries Research and Development Corporation, Canprint, Australia, 59 p.
- FAO, 2004. *Fruits of Vietnam*. Food and Agricultural Organization of the United Nations, Regional Office for Asia and the Pacific, Bangkok, Thailand. <http://www.fao.org/docrep/008/ad523e/ad523e00.htm>. Accessed March 23, 2015.
- Groff, G.W., 1921. *The Lychee and Lungan*. Orange Judd Company, New York, US, pp. 103-

327 Huang, X., Subhadrabandhu, S., Mitra, S.K., Ben-Arie, R., Stern R.A., 2005. Origin, history,
 328 production and processing. In: Menzel, C.M., Waite, G.K. (Eds.), *Litchi and Longan:*
 329 *Botany, Production and Uses*. CABI International, Wallingford, UK, pp. 1-23.
 330 Jonathan, H.C., Carlos, F.B., Steven, A.S., Ian M., 2013. Longan growing in the Florida home
 331 landscape, Fact Sheet HS-49. Inst. of Food and Agricultural Sciences, University of
 332 Florida, 11 pp.
 333 Ke, G.W., Wang, C.C., Huang, J.H., 1992. The aril initiation and ontogenesis of longan fruit.
 334 *Journal of Fujian Academy of Agricultural Sciences* 7, 22–26.
 335 Leenhouts, P.W., 1971. A revision of *Dimocarpus* (Sapindaceae). *Blumea* 19, 113-131.
 336 Loureiro, J.D., 1790. *Flora Cochinchinensis*, Ulyssipone, Typis, et expensis
 337 Academicis. Portugal, T1, pp. 233-234.
 338 Ministry of Agriculture and Rural Development of Vietnam, 2012.
 339 <http://mard.gov.vn/Pages/home.aspx>. Accessed March 23, 2015.
 340 Nakasone, H.Y., Paull, R.E., 1998. *Tropical Fruits*, CAB International, Wallingford, UK, 445 p.
 341 Paull, R.E., Duarte, O., 2011. *Tropical Fruits*, 2nd Edition Vol. 1. CAB International,
 342 Wallingford, UK, pp. 221-251.
 343 Pham, V.T., Tran, M.H., Herrero, H., Hormaza, J.I., 2013. The reproductive biology of the
 344 Longan. In: *Proceedings of the 5th National Scientific Conference on Ecology and*
 345 *Biological Resources*. Hanoi, 18 October 2013, pp. 1242-1246.
 346 Qiu, D.L., 2014, Longan production and research in China. *Acta Horticulturae* 1029, 39-46.
 347 Stern, R.A., 2005. Fruit Set, Development and Maturation: Longan. In: Menzel, C.M., Waite,
 348 G.K. (Eds.), *Litchi and Longan: Botany, Production and Uses*. CABI International,
 349 Wallingford, UK, pp. 138-140.
 350 Subhadrabandhu, S., Stern, R.A., 2005. Taxonomy, Botany and Plant Development. In: Menzel,
 351 C.M., Waite, G.K. (Eds.), *Litchi and Longan: Botany, Production and Uses*. CABI
 352 International, Wallingford, UK, pp. 25-34.
 353 Wong, K.C., 2000. *Longan Production in Asia*. Food and Agricultural Organization of the
 354 United Nations, Bangkok, Thailand, 44 pp.

355 Wong, K.C., Ketsa, S., 1991. *Dimocarpus longan* Lour. In: Verheij, E.W.M., Coronel, R.E.
 356 (Eds.), Plant Resources of South-East Asia: Edible fruits and nuts 2. Pudoc, Wageningen,
 357 the Netherlands, pp. 146-151.

358 Xu, X.D., Zheng, S.Q., Huang, J.S., Xu, J.H., Lin, Y.Q., Liu, H.Y., 1997. Studies on fruit
 359 development of extremely late maturing species of longan. II. Dynamic changes of fresh
 360 and dry weight, the requirement and distribution of moisture and solute content in fruits.
 361 Journal of Fujian Academy of Agricultural Sciences 12, 19–23.

362 Zee, F.T.P., Chan, H.T.Jr., Yen, C.R., 1998. Lychee, longan, rambutan and pulasan. In: Shaw,
 363 P.E., Chan, H.T.Jr., Nagy, S. (Eds.), Tropical and Subtropical Fruits, Agscience,
 364 Auburndale, Florida , pp. 290–335.

365 Zheng, S.Q., Huang, J.S., Xu, X.D., 1994. Studies on fruit development of seed-wilted longan:
 366 correlative analysis on fruit growing type and its characters. Journal of Fujian Academy of
 367 Agricultural Sciences 9 (4), 22–25.

368

370 **Table 1.** Description of the phenological growth stages of longan (*Dimocarpus longan* Lour.)
 371 according to the BBCH scale
 372

BBCH code	Description
<i>Principal growth stage 0: vegetative bud development</i>	
010	Vegetative buds dormant
011	Beginning of bud swell
013	End of bud swell
017	Beginning of bud break
019	End of bud break
<i>Principal growth stage 1: leaf development</i>	
110	First leaves separated
111	First leaves unfolded
113	More leaves unfolded: First leaflets at 30% of its full size
116	More leaves unfolded: First leaflets at 60% of its full size
119	All leaves unfolded: all leaflets fully expanded
<i>Principal growth stage 3: shoot development</i>	
310	Beginning of shoot extension
311	10% of final shoot length
313	30% of final shoot length
315	50% of final shoot length
317	70% of final shoot length
319	90% or more of final shoot length
<i>Principal growth stage 5: inflorescence emergence</i>	
510	Reproductive buds dormant
511	Beginning of reproductive bud swell
512	Panicle axes begin to elongate
513	Beginning of panicle development
515	Inflorescences 50% of final length
517	Inflorescences 70% of final length
519	End of inflorescence extension
<i>Principal growth stage 6: flowering</i>	
610	First flowers open
611	10% flowers open.
613	30% flowers open
615	50% flowers open
617	70% flowers open
619	90% flowers open
<i>Principal growth stage 7: fruit development</i>	
710	No ovary growth still visible
711	Initial ovary growth. First physiological fruit abscission
712	20% of final fruit size
713	30% of final fruit size
715	50% of final fruit size
717	70% of final fruit size
718	80% of final fruit size
719	90% or more of final fruit size
<i>Principal growth stage 8: maturity of fruit</i>	
810	Physiological maturity
811	Beginning of fruit ripening
815	Advanced ripening
819	Fruit fully ripened

373 Figures

374 Fig. 1. Main phenological growth stages of longan according to the extended BBCH scale

375

376 Fig. 2. Progression of longan principal growth stages (PGS) and average monthly temperatures
377 following the BBCH scale in southern Spain.

378

379 Fig. 3. Longan flower bud development according to the extended BBCH scale

380

381 Fig. 4. Longan fruit development according to the extended BBCH scale

382

383

Fig. 1

Fig. 2

Fig. 3

Fig. 4