Manuscript Number: JFOODENG-D-14-00374R2

Title: Effect of water activity in tortilla and its relationship on the acrylamide content after frying

Article Type: Research Article

Keywords: Acrylamide; Tortilla chips; Minimum integral entropy

Corresponding Author: Dr. Ricardo Salazar, Ph.D

Corresponding Author's Institution: Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional

First Author: Rosa M Delgado, Ph.D

Order of Authors: Rosa M Delgado, Ph.D; Gabriel Luna-Bárcenas, Ph.D; Gerónimo Arámbula-Villa, Ph.D; Ebner Azuara, Ph.D; Patricia López-Pérea, Ph.D; Ricardo Salazar, Ph.D

Abstract: The objective of this study was to relate the tortilla minimum integral desorption entropy with acrylamide content during processing of tortilla chips. Tortilla pieces were stored at 30 °C at a_w of 0.11-0.84 for 4 days and fried later in soybean oil at 180 °C for 25 s. The lowest acrylamide content was observed in tortilla chips made of non-stored tortilla ($a_w=0.98$) as well as in those prepared from tortilla stored in the minimum integral entropy ($a_w=0.53$). In addition, the color and texture values were similar in both cases. These results suggest that the reduction of the acrylamide content during processing of tortilla chips and other tortilla based foods thermally processed might be modified by factors such as moisture content, a_w , and the physical state of water in the tortilla. Thus, the minimum integral entropy showed to be a reliable indicator to establish the most appropriate moisture conditions to obtain tortilla chips with reduced level of acrylamide when tortilla is dehydrated.

Highlights

- Water activity affects the acrylamide content in tortilla chips.
- Minimum integral entropy is a reliable indicator to establish the most appropriate moisture conditions to get a reduction of acrylamide in tortillas chips.
- Tortilla dehydration to the minimum integral entropy reduces the acrylamide content.
- Products obtained from stored tortilla showed an acceptable texture and appearance.

Effect of water activity in tortilla and its relationship on the acrylamide content after frying

Rosa M. Delgado^a, Gabriel Luna-Bárcenas^b, Gerónimo Arámbula-Villa^b, Ebner Azuara^c, Patricia López-Peréa^d and Ricardo Salazar^{b*}.

^a *Instituto de la Grasa, Consejo Superior de Investigaciones Científicas, Avenida Padre García Tejero 4, 41012, Seville, Spain*

^b *Centro de Investigación y de Estudios Avanzados del I.P.N., Unidad Querétaro, Libramiento Norponiente # 2000, Fraccionamiento Real de Juriquilla, 76230, Querétaro, Querétaro, México*

^c *Instituto de Ciencias Básicas, Universidad Veracruzana. Av. Dr. Rafael Sánchez Altamirano s/n, Col. Industrial Animas, Carretera Xalapa-Las Trancas Km. 3.5, 91192, Xalapa, Veracruz, México*

^d *Facultad de Ciencias Agrícolas, Campus El Cerrillo, Universidad Autónoma del Estado de México. Piedras Blancas, Carretera Toluca - Ixtlahuaca Km. 15.5, 5020, Toluca, Estado de México, México.*

* Corresponding author. Tel. +52 442 211 9900; fax: +52 442 211 9938. E-mail address: rsalazar@qro.cinvestav.mx (Ricardo Salazar)

Abstract

The objective of this study was to relate the tortilla minimum integral desorption entropy with acrylamide content during processing of tortilla chips. Tortilla pieces were stored at 30 °C at a_w of 0.11-0.84 for 4 days and fried later in soybean oil at 180 °C for 25 s. The lowest acrylamide content was observed in tortilla chips made of non-stored tortilla ($a_w=0.98$) as well as in those prepared from tortilla stored in the minimum integral entropy ($a_w=0.53$). In addition, the color and texture values were similar in both cases. These results suggest that the reduction of the acrylamide content during processing of tortilla chips and other tortilla based foods thermally processed might be modified by factors such as moisture content, a_w , and the physical state of water in the tortilla. Thus, the minimum integral entropy showed to be a reliable indicator to establish the most appropriate moisture conditions to obtain tortilla chips with reduced level of acrylamide when tortilla is dehydrated.

Keyword: Acrylamide, Tortilla chips, Minimum integral entropy

Chemical compounds studied in this article

Acrylamide (PubChem CID: 6579)

1. Introduction

The Maillard reaction is the main responsible of the generation of flavor and color in thermally processed foods (Ames, 1990). However, this reaction has been also linked to the acrylamide formation, specifically with the presence of carbonyl compounds with groups capable of forming a Schiff Base with the amino acid asparagine (Hidalgo et al. 2009; Mottram et al. 2002; Stadler et al. 2002). Acrylamide, a neurotoxic compound (Spencer & Schaumburg, 1974) and probable human carcinogen (IARC, 1994), has attracted the attention of the scientific community in recent years due to it has been found in high concentrations in thermally processed foods (Tareke et al. 2002).

The effect of water activity (a_w) on acrylamide formation has been widely studied. Depending on the water activity range studied, positive or negative correlations between water activity and acrylamide content have been observed. De Vleeschouwer et al. (2007) showed very slight promoting effect of water activity ($0.34 \leq a_w \leq 0.92$) on acrylamide amount in an asparagine–glucose model system. Similar results were reported by De Vleeschouwer et al. (2008) ($0.88 \leq a_w \leq 0.99$). The same tendency was observed in a potato model system ($0.11 \leq a_w \leq 0.97$) where the acrylamide content was rather dependent upon the moisture content than upon the water activity (Mestadgh et al. 2006). However, an opposite trend was found in aqueous glycerol asparagine model system ($0.33 \leq a_w \leq 0.71$) (Hedegaard et al. 2007) and plantain paste ($0.43 \leq a_w \leq 0.97$) (Bassama et al. 2011), revealing a decrease in acrylamide formation with increasing water activity.

The above-mentioned studies have in common that the food model showed a sorption isotherm typical of sugar-rich products (Mathlouthi, 2001) where a small variation in

moisture content corresponds to a large variation in a_w for water activities lower than 0.9. For water activities higher than 0.9, a steep increase in the corresponding moisture content can be observed (De Vleeschouwer et al. 2007). In contrast, most foods are a mixture of different compounds (for example, carbohydrate, protein, fat, minerals, vitamins, salts and others). Hence, real foods tend to have a sigmoidal moisture sorption isotherm due to the presence of solutes with very strong affinity for water that could maintain a larger amount of bound moisture at lower humidity than solutes with less affinity.

Water activity influences on the stability and chemical reaction rates in foods (Labuza et al. 1972). Thus the mobility of acrylamide precursors and as a consequence, the acrylamide formation is affected by the physical state of the water sorbed during thermal processing. In this context, it is expected that the final acrylamide content will be mainly affected by the moisture content (monolayer value) where strong bonds between the water (adsorbate) and the food (adsorbent) occur.

In an attempt to understand the influence of water activity of real foods on acrylamide content, the level of acrylamide was compared in tortilla chips prepared from tortilla equilibrated at different water activities, including that corresponding to the monolayer value. The monolayer was calculated from tortilla desorption isotherms by using both, GAB equation and the minimum integral entropy criteria.

2. Materials and methods

2.1 Materials

Labeled [2,2,3- $^2\text{H}_3$]acrylamide was purchased from Sigma–Aldrich (St. Louis, MO). All other chemicals were analytical grade and purchased from Sigma (St. Louis, MO) or Merck

(Darmstadt, Germany). Soybean oil was obtained from local supermarkets in Querétaro, México.

2.2. Elaboration of nixtamalized corn flour and tortilla chips.

Nixtamalized corn flour was prepared with commercial corn variety named Pioneer 30P16 and commercial lime ($\text{Ca}(\text{OH})_2$) (El Topo, Monterrey, N.L. Mexico), commonly used in the tortilla industry. This flour was prepared by cooking (8 kg) of whole corn kernels in a solution of 16 liters of water with 80 g of $\text{Ca}(\text{OH})_2$, corresponding to 1.0 g/100 g of lime relative to the corn weight used. The corn was boiled in an aluminum pan for 25 min and steeped for 16 h at room temperature ($22 \pm 1^\circ\text{C}$). The steep liquor was removed. The cooked corn was washed with 16 liters of water, then ground into corn dough (FUMASA, M100, Querétaro, México), and finally dehydrated using a flash type dryer (Cinvestav-AV, M2000, Querétaro, México). The drying conditions were adjusted to have 250°C inlet air temperature and 90°C to the exhaust air to avoid burning the material. Before storage, the nixtamalized corn flour was milled using a hammer mill (PULVEX 200, México DF) equipped with a 0.5 mm screen.

For tortilla chips elaboration, nixtamalized corn flour (100 g) was rehydrated with enough water (118 mL) to provide fresh dough with proper consistency to make tortillas. The dough was shaped into thin disks (11cm diameter and 1.0 mm thickness) using a commercial tortilla roll machine (Casa Herrera, México, D.F.). The dough shaped into tortillas were cooked on both sides for around 1.0 min by using an iron hot plate ($270 \pm 10^\circ\text{C}$) named “comal”. The resulting tortillas were cut into circular pieces with an average area of 10 cm^2 and its a_w and pH value (AACC International, 2010) determined. Two set of

tortilla pieces were obtained. The first set was fried immediately in soybean oil at 180°C for 25 s (control). The second set was used in the construction of desorption isotherms. After frying, tortilla chips were cooled on a paper towel to remove superficial oil and the color, fracture force and acrylamide content determined.

2.3. Water desorption isotherms

Water desorption isotherms were determined using a static equilibrium method. Tortilla samples (~4.0 g) ($a_w=0.98$) were weighed in triplicate into standard weighing dishes with a circular section on the bottom. Samples were placed in separate desiccators containing saturated salt slurries in the range of water activity from 0.11 to 0.85 using the a_w values reported by Labuza et al. (1985). The samples were held at 25, 30, and 35 °C until equilibrium was reached. Values of water activity were generated using equations reported in the same paper. A small amount of toluene was placed in each desiccator to prevent the growth of fungi. The equilibrium condition was attained within 7-10 days, when the differences among two consecutive weights were within 0.001 g. The dry matter content of the tortilla was determined by the drying in a conventional oven at 105 °C for 24 h. The Guggenheim-Anderson-De Boer (GAB) equation was used in modeling water sorption (Quirijns et al. 2005):

$$M = \frac{M_0 C k a_w}{(1 - k a_w)(1 - k a_w + C k a_w)} \quad (1)$$

Where a_w is water activity; M is water content of the sample on dry basis; M_0 is the monolayer water content; C is the Guggenheim constant, given by $C = c' \exp (h_m - h_n)/RT$; where c' is the equation constant; h_m is the heat of sorption of the first layer; h_n is the heat of sorption of the multilayer; R is the gas constant; T is the absolute temperature; and k is

the constant correcting properties of multilayer molecules with respect to bulk liquid, and given by $k = k' \exp (h_l - h_n)/RT$; where k' is the equation constant; h_l is the heat of condensation of pure water. The parameters values of GAB equation (M_0 , C and k) were estimated by fitting the mathematical model to the experimental data, using non-linear regression with the Kaleidagraph 4.0 package (Synergy Software, Perkiomen, USA). Goodness of fit was evaluated using the average of the relative percentage difference between the experimental and predicted values of the moisture content or mean relative deviation modulus (P) defined by the following equation:

$$P(\%) = \frac{100}{N} \sum_{i=1}^N \frac{|Me_i - Mc_i|}{Me_i} \quad (2)$$

Where Me_i is the moisture content at observation i ; Mc_i is the predicted moisture content at that observations; and N is the number of observations. It is generally assumed that a good fit is obtained when $P < 10\%$ (Lomauro et al. 1985).

2.4 Determination of minimum desorption integral entropy

The determination of the integral (enthalpy and entropy) thermodynamic properties, and the water activity-temperature conditions where the tortilla minimum integral entropy occurred, considered as the point of maximum storage stability (monolayer value), was established as indicated by Pascual- Pineda et al. (2013) and Vigano et al. (2012). These authors have provided a thorough description of the procedure followed and equations used for this purpose. Briefly, the integral enthalpy changes $(\Delta H_{int})_T$ (J/mol) at the water-tortilla interface and, at different stages of the adsorption process, were determined using the equation of Othmer (Othmer, 1940).

$$\frac{d \ln P_v}{d \ln P_v^0} = \frac{H_v(T)}{H_v^0(T)} \quad (3)$$

144 Where: the desorbed substance is water; P_v (Pa) is the vapor pressure of water over the
 145 adsorbent; P_v^0 (Pa) is the vapor pressure of pure water at the temperature of sorption; $H_v(T)$
 146 (J/mol) is the integral molar heat of sorption, and $H_v^0(T)$ (J/mol) is the heat of
 147 condensation of pure water. Since all these terms are temperature-dependent, the equation
 148 can be integrated:

$$\ln P_v = \left[\frac{H_v(T)}{H_v^0(T)} \right]_{\phi} \ln P_v^0 + A \quad (4)$$

149 Where: A is the adsorption constant, and ϕ (J/mol) is the pressure of diffusion or surface
 150 potential. A plot of $\ln P_v$ versus $\ln P_v^0$ gives a straight line if the ratio $H_v(T)/H_v^0(T)$ is
 151 constant within the range of temperatures used.

152 The molar integral enthalpy $(\Delta H_{int})_T$ can be calculated using equation (5), at a constant
 153 pressure of diffusion(Nunes & Ronstein, 1991):

$$(\Delta H_{int})_T = \left[\frac{H_v(T)}{H_v^0(T)} - 1 \right]_{\phi} H_v^0(T) \quad (5)$$

$$\phi = \mu_{ap} - \mu_a = RT \frac{W_{ap}}{W_v} \int_0^{a_w} M d \ln a_w \quad (6)$$

154 Where: μ_{ap} (J/mol) is the chemical potential of the pure adsorbent; μ_a (J/mol) is the
 155 chemical potential of the adsorbent participating on the condensed phase; W_{ap} (g/mol) is
 156 the molecular weight of the adsorbent, and W_v (g/mol) is the molecular weight of the
 157 water.

By calculating $H_v(T)/H_v^0(T)$ from equation (4) and substituting it into equation (5) it becomes possible to calculate the integral enthalpy at different temperatures, provided that a good means of estimating $H_v^0(T)$ is available, such as that proposed by Wexler, (1976):

$$H_v^0(T) \text{ J/mol-K} = 6.15 \times 10^4 - 94.14 T + 17.74 \times 10^{-2} T^2 - 2.03 \times 10^{-4} T^3 \quad (7)$$

Using the values obtained for $(\Delta H_{int})_T$ changes, the molar integral entropy $(\Delta S_{int})_T$ can be estimated using the following equation :

$$(\Delta S_{int})_T = S_1 - S_L = - \frac{(\Delta H_{int})_T}{T} - R \ln a_w \quad (8)$$

where: $S_1 = S/N_1$ (J/mol K) is the integral entropy of water desorbed in the foodstuff; S (J/mol K) is the total entropy of water desorbed in the foodstuff; N_1 is the moles of water desorbed in the foodstuff, and S_L (J/mol K) is the molar entropy of pure liquid water in equilibrium with vapor.

2.5. Storage of tortilla before frying

Since water retention capacity in tortillas is poor, the moisture loss in tortilla after processing is often caused by starch retrogradation. Taking advantage of this phenomenon, tortilla samples were left at room temperature (25 °C) during 4 h to reduce their a_w from 0.98 to 0.85 and therefore, reducing the moisture content and preventing spoilage reactions during the time that the samples reached the desired a_w . The tortilla pH was monitored in order to confirm that acrylamide levels were not affected by pH changes during storage. Duplicate samples containing ca. 5 g of tortilla were placed in desiccators containing the same saturated salt slurries referred in section 2.3 at 30°C. Tortilla samples were withdrawn at day 5 to prepare tortilla chips.

2.6. Color determination in tortilla chip samples

Color changes were determined using a colorimeter MiniScan XE, model 45/0-L (Hunter Associates Laboratory, 11491 Sunset Hill Rd., Reston, Va., U.S.A.). Total color differences (ΔE) at the different periods of time were calculated from the determined CIELAB L^* a^* b^* values according to Hunter (1973): $\Delta E = [(\Delta L^*)^2 + (\Delta a^*)^2 + (\Delta b^*)^2]^{1/2}$; where L^* = brightness or lightness (100 = perfect white, to 0 = black); a^* = greenness/redness [negative (green) to positive (red)]; b^* = yellowness/ blueness [negative (blue) to positive (yellow)]; ΔL^* , Δa^* , and Δb^* = absolute differences of the values between the reference tile (white porcelain) and sample values; ΔE = total difference between reference and sample color. The reference values (calibration) were: $L^* = 92.22$, $a^* = -0.82$ and $b^* = 0.62$.

2.7. Texture determination in tortilla chips

The fracture force of the tortilla chips was evaluated using the Texture Analyzer TA-XT2 (Texture Technologies Corp., N. Y.). Fracture force was evaluated in freshly prepared samples. The test was carried out using a 2.03 mm diameter stainless-steel probe and a platform accessory with a hollow cylindrical base with 33.5 and 10 mm external and internal diameters, respectively. The probe traveled at a velocity of 10 mm/s until it cracked the sample (distance 6 mm).

2.8. Acrylamide determination in tortilla chips

Acrylamide was analyzed as the stable 2-bromopropenamide derivative by gas chromatography–mass spectrometry (GC–MS) using a combination of the methods of Andrawes et al. (1987) and Castle et al. (1991) as described previously (Salazar et al. 2012). Tortilla chips were ground in a mortar lab and powdered samples (~0.8 g) were

successively weighed in centrifugal tubes, spiked with 20 μ L of internal standard solution (0.5 mg/mL of deuterium-labeled [2,2,3- $^2\text{H}_3$]acrylamide in acetonitrile), and stirred with 8 mL of distilled water and 10 mL of *n*-hexane at room temperature for 5 min. After centrifugation at 2000 $\times g$ for 10 min, organic phases were removed. Co-extractives from supernatants were precipitated with 30 μ L of carrez I and 30 μ L of carrez II solutions. Later, supernatants were centrifuged at 2000 $\times g$ for 5 min and filtered. These extracts (4 mL) were treated with 0.45 g of potassium bromide, 200 μ L of sulfuric acid (10 mL/100 mL), and 300 μ L of potassium bromate solution (0.1 mol/L). After 1 h in the dark at 4°C, the bromination reaction was terminated by adding of 1 mol/L sodium thiosulfate until solutions became colorless, and solutions were extracted with 5 mL of ethyl acetate/hexane (4:1). Organic layers were recovered after centrifugation at 2000 $\times g$ for 10 min, and were dried with sodium sulfate and evaporated to dryness under nitrogen. Each sample was dissolved in 50 μ L of ethyl acetate, treated with 25 μ L of triethylamine, and analyzed by GC–MS. The ions monitored for the identification of the analyte, 2- bromopropenamide, were $[\text{C}_3\text{H}_4\text{NO}]^+ = 70$, $[\text{C}_3\text{H}_4^{79}\text{BrNO}]^+ = 149$, and $[\text{C}_3\text{H}_4^{81}\text{BrNO}]^+ = 151$, using m/z 149 for quantification. The ions monitored for the identification of the corresponding derivative 2-bromo[$^2\text{H}_2$]propenamide were $[\text{C}_2^2\text{H}_2\text{H}^{81}\text{Br}]^+ = 110$ and $[\text{C}_3^2\text{H}_2\text{H}_2^{81}\text{BrNO}]^+ = 153$, using m/z 153 for quantification.

GC–MS analyses were conducted with a Perkin Elmer GC Clarus 500 coupled with a Perkin Elmer Clarus 560 MSD (Mass Selective Detector-Quadrupole type). In most experiments, a 30 m $\times$ 0.32 mm i.d. $\times$ 0.25 μ m Elite-5MS capillary column was used. Working conditions were as follows: carrier gas helium (1 mL/min at constant flow);

injector, 250°C; oven temperature: from 50 (10 min) to 240°C at 5°C/min and then to 300°C at 10°C/min; transfer line to MSD, 280°C; ionization EI, 70 eV.

Quantification of acrylamide was carried out by preparing standard curves of this compound. Acrylamide content was directly proportional to the acrylamide/internal standard area ratio ($r = 0.999$, $p < 0.0001$). The coefficients of variation at the different concentrations were lower than 10%.

2.9. Statistical analysis

All results were expressed as mean $\pm$ SD values ($n=3$). When significant F values were obtained, group differences were evaluated by the Tukey test. All statistical procedures were carried out using the JMP 9.0 package (SAS Institute Inc., Cary, NC). The significance level was $p < 0.05$ unless otherwise indicated.

3. Results and discussion

Figure 1 shows the desorption isotherms of the tortilla used to make tortilla chips at 25, 30 and 35 °C. Sigmoidal isotherms described as type by Brunauer et al. (1940) were determined. Since desorption is an endothermic process, the increment of the temperature at constant a_w enhances the moisture desorbed from the tortilla. Similar desorption isotherms have been reported for starchy products such as potato (McLaughlin & Magee, 1998), maize (Samapundo et al. 2006) and banana (Yan et al. 2008), chestnut flour (Chenlo et al. 2011) and Japanese noodle (Inazu et al. 2001). **Table 1** gives the estimated parameters obtained by fitting the GAB equation to experimental data. As can be seen from the table, the GAB equation describes adequately the experimental data over the whole measured range of a_w . Thus the values of r^2 were very close to unit and the P value was ≤ 10 % under

the studied conditions, which it indicates a good fit (Lomauro et al. 1985). As expected, maximum water desorption by strongly binding sites (monolayer) predicted for GAB equation decreased with increasing temperature. The moisture content in the monolayer (Mo) was 10.12, 9.46 and 8.96 g water/100 g dry solid (corresponding to a_w of 0.32, 0.31 and 0.30) for 25, 30 and 35 °C, respectively. It is assumed that the value of Mo is a parameter for estimating the amount of water bound to specific polar sites in dehydrated food and in this water content; a storage product should be stable against degradation reactions (Rahman & Labuza, 1999). The constant C is a measure of the attraction force between the water and the sorption sites. The values obtained are higher than those reported by Polou et al. (1997) in commercial tortilla chips. In this study, the k values increased with increasing temperature. The value of k provides a measure of the interaction of the molecules in the multilayers with the adsorbent, and tends to fall between the energy value of the molecules in the monolayer and the liquid water. When $k = 1$, the properties of the water present in the multilayer are similar to those of the free water (Perez-Alonso et al. 2006).

Variations in the differential and integral entropy with respect to a_w of the tortilla used to make tortilla chips are represented in **Figure 2**. The intersection of the curves is found in the water content and a_w corresponding to the minimum integral entropy. It could be observed that as tortilla desorbed moisture, their integral entropy fell to a minimum. Integral entropy can be directly related to the order-disorder of water molecules sorbed on food, and therefore it is a useful function to study the effect of drying method on the stability of the product. The minimum integral entropy is expected where strong bonds between the adsorbate and the adsorbent are (Nunes & Rotstein, 1991), and hence water

molecules are less available to participate in spoilage reactions. This point has been proposed as the most suitable for storage and it is considered as the water content corresponding to the monolayer (Dominguez et al. 2007; Hill et al. 1951).

Although the value of minimum entropy may be unique, there are food products with zones in which this minimum does not vary appreciably in a defined range of moisture. As it can be observed in **Figure 2**, the minimum integral entropy at 30 °C for tortilla corresponds to a_w value of 0.53 (13.20 g water/100g dry solids). It should be noted that although the minimum integral entropy occurs in a given a_w , this minimum does not vary appreciably in a defined range of water activities (0.45 to 0.62). Differential entropy had a minimum at 7.3 g water/100 g dry solids ($a_w=0.18$). This parameter does not mean order or disorder of the total system. The differential entropy represents the algebraic sum of the integral entropy at a particular hydration level, plus the change of order or disorder after new water molecules were desorbed by the system at the same hydration level. If the values of moisture content corresponding to minimum integral entropy and minimum differential entropy are different, this particular hydration level at the minimum differential entropy cannot be considered as the maximum stability point, because not all available active sites have been occupied at that particular water content, and therefore it is possible to obtain after this point lower differential changes that provide a better ordering of the water molecules desorbed on food (Beristain et al. 2002).

The minimum integral entropy approach has been verified in several studies (see for example: Beristain et al. 1994; Bonilla et al. 2010; Carrillo -Navas et al. 2011; Dominguez et al. 2007; Rascon et al. 2011).

On the other hand, frequently the thermodynamic analysis is not in accordance with the monolayer obtained with the GAB model (Beristain & Azuara, 1990; Beristain et al. 2002; Nunes & Rotstein, 1991). In this study, the minimum integral entropy point is higher than the calculated Mo value with GAB equation. Since the BET theory, which assumes that the surface of the adsorbent is energetically uniform and does not take into account horizontal interactions between the molecules in the adsorbed layer (Dollimore et al. 1976), is also present in the GAB equation, the GAB monolayer concept has a limited applicability.

In this context, the Maillard reaction (main pathway acrylamide formation pathway) is a reaction that requires the interaction of two reagents. Therefore, those factors which affect the concentration and mobility of the reactants will have a significant impact on the reaction rate (Bell, 2007). Thus, the plasticizing effect of water significantly influences on the rate of the Maillard reaction (Bell et al. 1998; White & Bell, 1999). In the minimum integral entropy point, water is less reactive to carry out spoilage reactions in the food, so the mobility of acrylamide precursors is reduced, affecting its formation during thermal processing.

Figure 3 shows acrylamide content in tortilla chips prepared from tortillas stored at different water activities. All water activities induced the formation of acrylamide in the tortilla chips, but tortilla chips made of no-stored tortilla ($a_w=0.98$) always contained less acrylamide (688.55 ± 65 $\mu\text{g/kg}$) than those made of tortilla stored at different water activities ($p < 0.05$). This suggests a dilution of acrylamide precursors caused by high moisture content. This effect is similar to that reported for the Maillard reaction in which a reduction in reaction rate is observed in a_w close to 1 because of the reagent dilution as well as the inhibitory effect that water exerts in this reaction (Ames, 1990; Acevedo et al. 2008). Thus,

a reduction in the acrylamide content was observed in the minimum integral entropy point. The acrylamide content of tortilla chips prepared from tortilla stored at $a_w=0.53$ (741.85 ± 88 $\mu\text{g/kg}$) was similar to those prepared from no-stored tortilla ($p < 0.05$). This point is characterized by strong water-food interactions causing a reduction in the diffusion and mobility of the system and affecting to the development of chemical reactions during frying such as the acrylamide formation. On the other hand, an increase on acrylamide content on tortilla chips proportional to reduction of a_w (0.43-0.11) on tortilla was observed. This increase may be linked to the concentration of the precursors as well as the evaporation of water which takes place during the start of frying. At this stage, the temperature of the tortilla does not exceed 100°C until a certain amount of water is evaporated. Since acrylamide is not formed at temperatures below 100°C , an increase on acrylamide formation is observed when frying is carried out at low a_w where a rapid increase of the internal temperature of the food occurs (Mestdagh et al. 2006).

Foods are heterogeneous and dynamic mixtures of macromolecules, solvent and solutes, therefore, the positive or negative correlation between water activity and acrylamide content are probably due to differences in composition and structure of used food systems. According to Viveros- Contreras et al. (2013) and Vigano et al. (2011), food with the same chemical composition and different microstructure could show different moisture sorption behavior, and therefore the a_w where spoilage reactions occurs, could be modified . This fact suggests that the microstructure and physical structure of the water in the food is able to restrict the mobility of the precursors responsible for the formation of acrylamide to a specific humidity level during thermal processing and it explains the decrease of the acrylamide content of tortilla elaborated from tortillas equilibrated at the a_w corresponding

to minimum integral entropy zone. Thus, the influence of a_w on the acrylamide formation will be different depending on each particular food system.

Texture and color are considered the most important parameters of quality and acceptability of fried products. **Figure 4** shows the effect of tortilla water activity on the parameters of tortilla chips above mentioned. The color has been correlated with the acrylamide generation in thermally processed foods (Gökmen & Şenyuva, 2006; Lukac et al. 2007; Majcher & Jelen, 2007; Pedreschi et al. 2007). In this study, the appearance of the tortilla chips (**Figure 4a**) remained constant in the range of a_w from 0.43 to 0.84 and increased proportionally to a_w reduction (0.43-0.11). It should be emphasized that this change was observed in the water activity zone corresponding to the minimum integral entropy. The fracture force (**Figure 4b**) showed similar values over the whole range of water activities studied. The obtained fracture force values are close to those found (8.73–9.63 N) by Tseng et al. (1996), and higher than those (5.33–6.17 N) described in commercial tortilla chips (Lujan-Acosta & Moreira, 1997).

4. Conclusions

The effect of a_w on the acrylamide content of tortilla chips could be related to the moisture sorption behavior. The minimum integral entropy in conjunction with the corresponding a_w obtained from the desorption isotherm provided a reliable indicator to establish the most appropriate moisture conditions to obtain tortilla chips with reduced level of acrylamide when tortilla was dehydrated. Furthermore, the acrylamide content reduction was similar when tortilla chips were prepared from non-stored tortilla with a moisture content corresponding to water activities close to 1. It should be noted that a partial tortilla

dehydration to a water activity different to the minimum integral entropy results in tortilla chips with higher acrylamide content. Although a detailed sensory evaluation of tortilla chips prepared is needed, preliminary results showed that the products obtained from tortilla stored in the minimum integral entropy a_w did not show any significant change in the appearance of the food product. Therefore, the results suggest that the control of a_w , moisture content and the physical state of water in the tortillas are factors to reduce the formation of acrylamide during processing of tortilla chips. This is of the upmost importance, because it provides the food manufacturer with information of which a_w is more likely to get a low acrylamide content product over a wider range of processing conditions.

Acknowledgments

We are indebted to Juan Veles, Edmundo Gutierrez, Carlos Alberto Ávila, Araceli Mauricio and Veronica Flores from CINVESTAV Querétaro for their technical assistance.

References

- AACC International (2010). Approved Method 02-52.01. *Approved Methods of the American Association of Cereal Chemists*. (11 th Ed.). St Paul, MN, USA.
- Acevedo, N. C., Briones, V., Buera, P., & Aguilera, J. M. (2008). Microstructure affects the rate of chemical, physical and color changes during storage of dried apple discs. *Journal of Food Engineering*, 85(2), 222-231.
- Ames, J. M. (1990). Control of the Maillard reaction in food systems. *Trends in Food Science & Technology*, 1, 150-154.

378 Andrawes, F., Greenhouse, S., & Draney, D. (1987). Chemistry of acrylamide bromination
 379 for trace analysis by gas chromatography and gas chromatography—mass spectrometry.
 380 *Journal of Chromatography A*, 399, 269-275.

381 Azuara, E., Vernon-Carter, E. J., & Beristain, C. I. (2007). Thermodynamic analysis of the
 382 effect of water activity on the stability of macadamia nut. *Journal of Food Engineering*,
 383 81(3), 566-571.

384 Bassama, J., Brat, P., Bohuon, P., Hocine, B., Boulanger, R., & Günata, Z. (2011).
 385 Acrylamide kinetic in plantain during heating process: Precursors and effect of water
 386 activity. *Food Research International*, 44(5), 1452-1458.

387 Bell, L. N. (2007). Moisture effects on food's chemical stability. In G. V. Barbosa-
 388 Canovas, A. J. J. Fontana, S. J. Schmidt, & T. P. Labuza (Eds.), *Water activity in foods:
 389 Fundamentals and applications* (pp. 177–198). Ames: Blackwell Publishing Ltd.

390 Bell, L. N., Touma, E., White, K. L., & Chen, Y. (1998). Glycine loss and maillard
 391 browning as related to the glass transition in a model food system. *Journal of Food Science*,
 392 63(4), 625–628.

393 Beristain, C., & Azuara, E. (1990). Estabilidad máxima en productos deshidratados.
 394 *Ciencia*, 41(3), 229–236.

395 Beristain, C. I., Diaz, R., Garcia, H. S., & Azuara, E. (1994). Thermodynamic behavior of
 396 green whole and decaffeinated coffee beans during adsorption. *Drying Technology*, 12(5),
 397 1221-1233.

398 Beristain, C. I., Azuara, E., & Vernon-Carter, E. J. (2002). Effect of Water Activity on the
 399 Stability to Oxidation of Spray-Dried Encapsulated Orange Peel Oil Using Mesquite Gum
 400 (Prosopis Juliflora) as Wall Material. *Journal of Food Science*, 67(1), 206-211.

401 Bonilla, E., Azuara, E., Beristain, C. I., & Vernon-Carter, E. J. (2010). Predicting suitable
 402 storage conditions for spray-dried microcapsules formed with different biopolymer
 403 matrices. *Food Hydrocolloids*, 24(6), 633-640.

404 Brunauer, S., Deming, L. S., Deming, W. E., & Teller, E. (1940). On a theory of the van
 405 der Waals adsorption of gases. *Journal of the American Chemical Society*, 62(7), 1723-
 406 1732.

407 Carrillo-Navas, H., González-Rodea, D. A., Cruz-Olivares, J., Barrera-Pichardo, J. F.,
 408 Román-Guerrero, A., & Pérez-Alonso, C. (2011). Storage stability and physicochemical
 409 properties of passion fruit juice microcapsules by spray-drying. *Revista Mexicana de*
 410 *Ingeniería Química*, 10(3), 421-430.

411 Castle, L., Campos, M. J., & Gilbert, J. (1991). Determination of acrylamide monomer in
 412 hydroponically grown tomato fruits by capillary gas chromatography—mass spectrometry.
 413 *Journal of the Science of Food and Agriculture*, 54(4), 549-555.

414 Chenlo, F., Moreira, R., Prieto, D. M., & Torres, M. D. (2011). Desorption isotherms and
 415 net isosteric heat of chestnut flour and starch. *Food and Bioprocess Technology*, 4(8),
 416 1497-1504.

417 De Vleeschouwer, K., Plancken, I. V. D., Van Loey, A., & Hendrickx, M. E. (2007).
 418 Kinetics of acrylamide formation/elimination reactions as affected by water activity.
 419 *Biotechnology Progress*, 23(3), 722-728.

420 De Vleeschouwer, K., Van der Plancken, I., Van Loey, A., & Hendrickx, M. E. (2008).
 421 Investigation of the influence of different moisture levels on acrylamide
 422 formation/elimination reactions using multiresponse analysis. *Journal of Agricultural and*
 423 *Food Chemistry*, 56(15), 6460-6470.

424 Dollimore, D., Spooner, P., & Turner, A. (1976). The BET method of analysis of gas
 425 adsorption data and its relevance to the calculation of surface areas. *Surface Technology*,
 426 4(2), 121-160.

427 Gökmen, V., & Şenyuva, H. Z. (2006). Study of colour and acrylamide formation in coffee,
 428 wheat flour and potato chips during heating. *Food Chemistry*, 99(2), 238-243.

429 Hedegaard, R. V., Frandsen, H., Granby, K., Apostolopoulou, A., & Skibsted, L. H. (2007).
 430 Model studies on acrylamide generation from glucose/asparagine in aqueous glycerol.
 431 *Journal of Agricultural and Food Chemistry*, 55(2), 486-492.

432 Hidalgo, F. J., Delgado, R. M., & Zamora, R. (2009). Degradation of asparagine to
 433 acrylamide by carbonyl-amine reactions initiated by alkadienals. *Food Chemistry*, 116(3),
 434 779–784.

435 Hill, T. L., Emmett, P. H., & Joyner, L. G. (1951). Calculation of Thermodynamic
 436 Functions of Adsorbed Molecules from Adsorption Isotherm Measurements: Nitrogen on
 437 Graphon1, 2. *Journal of the American Chemical Society*, 73(11), 5102-5107.

438 Hunter, R. S. (1973). *The measurement of appearance*. Fairfax, VA: 383 Hunter Associates
 439 Laboratory.

440 IARC. (1994). IARC monographs on the evaluation of carcinogenic risks to humans (Vol.
 441 60). *International Agency for Research on Cancer*. Lyon, France

442 Inazu, T., Iwasaki, K. I., & Furuta, T. (2001). Desorption isotherms for Japanese noodle
 443 (udon). *Drying Technology*, 19(7), 1375-1384.

444 Labuza, T. P., Kaanane, A., & Chen, J. Y. (1985). Effect of temperature on the moisture
 445 sorption isotherms and water activity shift of two dehydrated foods. *Journal of Food*
 446 *Science*, 50(2), 385-392.

447 Labuza, T. P., McNally, L., Gallagher, D., Hawkes, J., & Hurtado, F. (1972). Stability of
 448 intermediate moisture foods. 1. Lipid oxidation. *Journal of Food Science*, 37(1), 154-159.

449 Lomauro, C. J., Bakshi, A. S., & Labuza, T. P. (1985). Evaluation of food moisture
 450 sorption isotherm equations. Part I: Fruit, vegetable and meat products. *Lebensmittel-*
 451 *Wissenschaft und-Technologie*, 18(2), 111-117.

452 Lujan-Acosta, J., & Moreira, R. G. (1997). Effects of different drying processes on oil
 453 absorption and microstructure of tortilla chips. *Cereal Chemistry*, 74(3), 216-223.

454 Majcher, M. A., & Jeleń, H. H. (2007). Acrylamide formation in low-fat potato snacks and
 455 its correlation with colour development. *Food Additives and Contaminants*, 24(4), 337-342.

456 Mathlouthi, M. (2001). Water content, water activity, water structure and the stability of
 457 foodstuffs. *Food Control*, 12(7), 409-417.

458 McLaughlin, C. P., & Magee, T. R. A. (1998). The determination of sorption isotherm and
 459 the isosteric heats of sorption for potatoes. *Journal of Food Engineering*, 35(3), 267-280.

460 Mestdagh, F., De Meulenaer, B., Cucu, T., & Van Peteghem, C. (2006). Role of water upon
 461 the formation of acrylamide in a potato model system. *Journal of Agricultural and Food*
 462 *Chemistry*, 54(24), 9092-9098.

463 Mottram, D. S., Wedzicha, B. L., & Dodson, A. T. (2002). Acrylamide is formed in the
 464 Maillard reaction. *Nature*, 419(6906), 448-449.

465 Nunes, R. V., & Rotstein, E. (1991). Thermodynamics of the water-foodstuff equilibrium.
 466 *Drying Technology*, 9(1), 113-137.

467 Othmer, D. F. (1940). Correlating vapor pressure and latent heat data. A new plot.
 468 *Industrial and Engineering Chemistry*, 32(6), 841-856.

469 Palou, E., Lopez-Malo, A., & Argai, A. (1997). Effect of temperature on the moisture
 470 sorption isotherms of some cookies and corn snacks. *Journal of Food Engineering*, 31(1),
 471 85-93.

472 Pascual-Pineda, L. A., Flores-Andrade, E., Alamilla-Beltrán, L., Chanona-Pérez, J. J.,
 473 Beristain, C. I., Gutiérrez-López, G. F., & Azuara, E. (2013). Micropores and Their
 474 Relationship with Carotenoids Stability: A New Tool to Study Preservation of Solid Foods.
 475 *Food and Bioprocess Technology*, 7(4), 1160-1170.

476 Pedreschi, F., Kaack, K., & Granby, K. (2004). Reduction of acrylamide formation in
 477 potato slices during frying. *LWT-Food Science and Technology*, 37(6), 679-685.

478 Pérez-Alonso, C., Beristain, C. I., Lobato-Calleros, C., Rodriguez-Huezo, M. E., &
 479 Vernon-Carter, E. J. (2006). Thermodynamic analysis of the sorption isotherms of pure and
 480 blended carbohydrate polymers. *Journal of Food Engineering*, 77(4), 753-760.

481 Quirijns, E. J., Van Boxtel, A. J., van Loon, W. K., & Van Straten, G. (2005). Sorption
 482 isotherms, GAB parameters and isosteric heat of sorption. *Journal of the Science of Food
 483 and Agriculture*, 85(11), 1805-1814.

484 Rahman, M. S., & Labuza, T. P. (1999). Water activity and food preservation. In M. S.
 485 Rahman (Ed.), *Handbook of food preservation* (pp. 339-382). New York: Marcel Dekker.

486 Rascón, M. P., Beristain, C. I., García, H. S., & Salgado, M. A. (2011). Carotenoid
 487 retention and storage stability of spray-dried encapsulated paprika oleoresin using gum
 488 Arabic and soy protein isolate as wall materials. *LWT-Food Science and Technology*, 44
 489 (2), 549-557.

490 Rizvi, S. S. H., & Benado, A. L. (1984). Thermodynamics properties of dehydrated food.
 491 *Food Technology*, 38(3), 83-92.

492 Salazar, R., Arámbula-Villa, G., Hidalgo, F. J., & Zamora, R. (2012). Mitigating effect of
 493 piquin pepper (*Capsicum annuum* L. var. *Aviculare*) oleoresin on acrylamide formation in
 494 potato and tortilla chips. *LWT-Food Science and Technology*, 48(2), 261-267.

495 Samapundo, S., Devlieghere, F., Meulenaer, B. D., Atukwase, A., Lamboni, Y., &
 496 Debevere, J. M. (2007). Sorption isotherms and isosteric heats of sorption of whole yellow
 497 dent corn. *Journal of Food Engineering*, 79(1), 168-175.

498 Spencer, P. S., & Schaumburg, H. H. (1974). A review on acrylamide neurotoxicity. Part I.
 499 Properties, uses and human exposure. *Canadian Journal of Neurological Science*, 1(2),
 500 151–169.

501 Stadler, R. H., Blank, I., Varga, N., Robert, F., Hau, J., Guy, P. A., Robert, M. C., &
 502 Riediker, S. (2002). Acrylamide from Maillard reaction products. *Nature*, 41 (6906), 449–
 503 454.

504 Tareke, E., Rydberg, P., Karlsson, P., Eriksson, S., & Tornqvist, M. (2002). Analysis of
 505 acrylamide, a carcinogen formed in heated foodstuffs. *Journal of Agricultural and Food*
 506 *Chemistry*, 50 (17), 4998–5006.

507 Tseng, Y. C., Moreira, R., & Sun, X. (1996). Total frying-use time effects on soybean-oil
 508 deterioration and on tortilla chip quality. *International Journal of Food Science &*
 509 *Technology*, 31(3), 287-294.

510 Viganó, J., Azuara, E., Telis, V., Beristain, C. I., Jiménez, M., & Telis-Romero, J. (2012).
 511 Role of enthalpy and entropy in moisture sorption behavior of pineapple pulp powder
 512 produced by different drying methods. *Thermochimica Acta*, 528, 63-71.

513 Viveros-Contreras, R., Téllez-Medina, D.I., Perea-Flores, M.J., Alamilla-Beltrán, L.,
 514 Cornejo-Mazón, M., Beristain-Guevara, C. I., Azuara-Nieto E., & Gutiérrez-López, G. F.
 515 (2013). Encapsulation of ascorbic acid into calcium alginate matrices through coacervation
 516 coupled to freeze-drying. *Revista Mexicana de Ingeniería Química*, 12(1), 29-39.

517 Wexler, A. (1976). Vapor pressure formulation for water in range 0 to 100 °C. A review.
518 *Journal of Research of the National Bureau of Standards A Physics and Chemistry*, 80(5),
519 775–785

520 White, K. L., & Bell, L. N. (1999). Glucose loss and Maillard browning in solids as
521 affected by porosity and collapse. *Journal of Food Science*, 6 (6), 1010-1014.

522 Yan, Z., Sousa-Gallagher, M. J., & Oliveira, F. A. (2008). Sorption isotherms and moisture
523 sorption hysteresis of intermediate moisture content banana. *Journal of Food Engineering*,
524 86 (3), 342-348.

Table 1

Estimated GAB parameters for tortilla desorption isotherms

	25°C	30°C	35°C
<i>Mo</i> (g H ₂ O/100 g d.s.)	10.11	9.46	8.96
<i>C</i>	14.67	15.72	14.26
<i>k</i>	0.66	0.67	0.70
<i>r</i> ²	0.99	0.99	0.99
<i>P</i> (%)	0.96	1.36	1.58

Figure captions

Figure 1. Moisture desorption isotherms of tortilla used to make tortilla chips at 25 (●), 30 (□) and 35 (Δ) °C. The continuous lines are the fitted points using the GAB model.

Figure 2. Differential (■) and integral (○) entropy changes as a function of water activity for tortilla at 30 °C.

Figure 3. Effect of tortilla water activity on acrylamide content in tortilla chips fried at 180°C for 25 s.

Figure 4. Effect of tortilla water activity on: color (**a**) and fracture force (**b**) in tortilla chips fried at 180°C for 25 s.

Figure 1

Fig 1. Moisture desorption isotherms of tortilla used to make tortilla chips at 25 (●), 30 (□) and 35 (Δ) °C. The continuous lines are the fitted points using the GAB model.

Figure 2

Fig 2. Differential (■) and integral (○) entropy changes as a function of water activity for tortilla at 30 °C.

Figure 3

Fig 3. Effect of tortilla water activity on acrylamide content in tortilla chips fried at 180°C for 25 s.

Figure 4

Fig 4. Effect of tortilla water activity on: color (a) and fracture force (b) in tortilla chips fried at 180°C for 25 s.