

# Spatial patterns in mesic savannas: the local facilitation limit and the role of demographic stochasticity.

Ricardo Martínez-García,<sup>1</sup> Justin M. Calabrese,<sup>2</sup> and Cristóbal López<sup>1</sup>

<sup>1</sup>*IFISC, Instituto de Física Interdisciplinar y Sistemas Complejos (CSIC-UIB), E-07122 Palma de Mallorca, Spain*

<sup>2</sup>*Conservation Ecology Center, Smithsonian Conservation Biology Institute, National Zoological Park, 1500 Remount Rd., Front Royal, VA 22630, USA.*

We propose a model equation for the dynamics of tree density in mesic savannas. It considers long-range competition among trees and the effect of fire acting as a local facilitation mechanism. Despite short-range facilitation is taken to the local-range limit, the standard full spectrum of spatial structures obtained in general vegetation models is recovered. Long-range competition is thus the key ingredient for the development of patterns. The long time coexistence between trees and grass, and how fires affect the survival of trees as well as the maintenance of the patterns is studied. The influence of demographic noise is analyzed. The stochastic system, under the parameter constraints typical of mesic savannas, shows irregular patterns characteristics of realistic situations. The coexistence of trees and grass still remains at reasonable noise intensities.

PACS numbers:

## I. INTRODUCTION

Savannas ecosystems are characterized by the long-term coexistence between a continuous grass layer and scattered or clustered trees [1]. Occurring in many regions of the world, in areas with very different climatic and ecological conditions, the spatial structure, persistence, and resilience of savannas have long intrigued ecologists [2–5]. However, despite substantial research, the origin and nature of savannas have not yet been fully resolved and much remains to be learned.

Savanna tree populations often exhibit pronounced, non-random spatial structures [6–10]. Much research has therefore focused on explaining how spatial patterning in savannas arises [8–13]. In most natural plant systems both facilitative and competitive processes are simultaneously present [2, 14] and hard to disentangle [15, 16]. Some savanna studies have pointed toward the existence of short-distance facilitation [7, 9], while others have demonstrated evidence of competition [6, 8, 10], with conflicting reports sometimes arriving from the same regions.

Different classes of savannas, which can be characterized by how much rainfall they typically receive, should be affected by different sets of processes. For example, in semiarid savannas water is extremely limited (low mean annual precipitation) and competition among trees is expected to be strong, but fire plays little role because there is typically not enough grass biomass to serve as fuel. In contrast, humid savannas should be characterized by weaker competition among trees, but also by frequent and intense fires. In-between these extremes, in mesic savannas, trees likely have to contend with intermediate levels of both competition for water and fire [3, 11, 17–20].

Competition among trees is mediated by roots that typically extend well beyond the crown [4, 16]. Additionally, fire can lead to local facilitation due to a protec-

tion effect, whereby vulnerable juvenile trees placed near adults are protected from fire by them [21]. We are particularly interested in how the interplay between these mechanisms governs the spatial arrangement of trees in mesic savannas, where both mechanisms may operate. On the other side, it has frequently been claimed that pattern formation in arid systems can be explained by a combination of long-distance competition and short-distance facilitation [22–28]. This combination of mechanisms is also known to produce spatial structures in many other natural systems [29]. Although mesic savannas do not display the same range of highly regular spatial patterns that arise in arid systems (e.g., tigerbush), similar mechanisms might be at work. Specifically, the interaction between long-range competition and short-range facilitation might still play a role in pattern formation in savanna tree populations, but only for a limited range of parameter values and possibly modified by demographic stochasticity.

Although the facilitation component has often been thought to be a key component in previous vegetation models [9, 26, 28, 30], Rietkerk and Van de Koppel [31], speculated, but did not show, that pattern formation could occur without short-range facilitation in the particular example of tidal freshwater marsh. In the case of savannas, as stated before, the presence of adult trees favor the establishment of new trees in the area, protecting the juveniles against fires. Considering this effect, we take the facilitation component to its infinitesimally short spatial limit, and study its effect in the emergence of spatially periodic structures of trees. To our knowledge, this explanation, and the interrelation between long-range competition and local facilitation, has not been explored for a vegetation system. One of our main results is that when considering the limit of local facilitation and nonlocal competition, clustering of trees appears.

Here we develop a minimalistic model of savannas that considers two of the factors, as already mentioned,

thought to be crucial to structure mesic savannas: tree-tree competition and fire, with a primary focus on spatially nonlocal competition. Employing standard tools used in the study of pattern formation phenomena in physics (stability analysis and the structure function) [29], we explore the conditions under which the model can produce non-homogeneous spatial distributions. A key strength of our approach is that we are able to provide a complete and rigorous analysis of the patterns the model is capable of producing, and we identify which among these correspond to situations that are relevant for mesic savannas. We further examine the role of demographic stochasticity in modifying both spatial patterns and the conditions under which trees persist in the system in the presence of fire, and discuss the implications of these results for the debate on whether the balance of processes affecting savanna trees is positive, negative, or is variable among systems. This is the framework of our study: the role of long-range competition, facilitation and demographic fluctuations (in the second part of the paper) in the spatial structures of mesic savannas. To complete our work we include an appendix where we study the effect of external fluctuations (mimicking for e.g. rainfall) on savanna dynamics.

Our model is inspired by the one presented by Calabrese et al. in [11]. It complements theirs by providing further analytical results that clearly demonstrate that this simple system, where we focus on the local limit of facilitation, can produce the full spectrum of spatial patterns reported from models employing both short-range facilitation and long-range inhibition (competition).

## II. THE DETERMINISTIC MODEL

In this section we derive the deterministic equation for the local density of trees, such that dynamics is of the logistic type and we only consider tree-tree competition and fire. We study the formation of patterns via stability analysis and provide numerical simulations of our model, showing the emergence of spatial structures.

### A. The nonlocal savanna model

Calabrese et al. [11] introduced a simple discrete-particle lattice savanna model that considers the birth-death dynamics of trees, and where tree-tree competition and fire are the principal ingredients. These mechanisms act on the probability of establishment of a tree once a seed lands at a particular point on the lattice. In the discrete model, seeds land in the neighborhood of a parent tree with a rate  $b$ , and establish as adult trees if they are able to survive both competition neighboring trees and fire. As these two phenomena are independent, the probability of establishment is  $P_E = P_C P_F$ , where  $P_C$  is the probability of surviving the competition, and  $P_F$  is the probability of surviving a fire event. From this dynamics,

we write a deterministic differential equation describing the time evolution of the global density of trees (mean field),  $\rho(t)$ , where the population has logistic growth at rate  $b$ , and an exponential death term at rate  $\alpha$ . It reads:

$$\frac{d\rho}{dt} = bP_E(\rho)\rho(t)(1 - \rho(t)) - \alpha\rho(t). \quad (1)$$

Generalizing Eq. (1), we propose an evolution equation for the space-dependent (local) density of trees,  $\rho(\mathbf{x}, t)$ :

$$\frac{\partial\rho(\mathbf{x}, t)}{\partial t} = bP_E\rho(\mathbf{x}, t)(1 - \rho(\mathbf{x}, t)) - \alpha\rho(\mathbf{x}, t). \quad (2)$$

We allow the probability of overcoming competition to depend on tree crowding in a local neighborhood, decaying exponentially with the density of surrounding trees as

$$P_C = \exp\left(-\delta \int G(\mathbf{x} - \mathbf{r})\rho(\mathbf{r}, t)d\mathbf{r}\right), \quad (3)$$

where  $\delta$  is a parameter that modulates the strength of the competition, and  $G(\mathbf{x})$  is a positive kernel function that introduces a finite range of influence. This model is related to earlier models of pattern formation in arid systems [23], and subsequent works [24, 25], but it differs from standard kernel-based models in that the kernel function accounts for the interaction neighborhood, and not for the type of interaction with the distance. Note also that the nonlocal term enters nonlinearly in the equation.

Following [11],  $P_F$  is assumed to be a saturating function of grass biomass,  $1 - \rho(\mathbf{x}, t)$ , similar to the implementation of fire of Jeltsch *et al.* in [12]

$$P_F = \frac{\sigma}{\sigma + 1 - \rho(\mathbf{x}, t)}, \quad (4)$$

where  $\sigma$  governs the resistance to fire, so  $\sigma = 0$  means no resistance to fires. Notice how our model is close to the one in [11] through the definitions of  $P_C$  and  $P_F$ , although we consider the probability of surviving a fire depending on the local density of trees, and in [11] it depends on the global density. The deterministic differential equation that considers tree-tree competition and fire for the spatial tree density is

$$\frac{\partial\rho(\mathbf{x}, t)}{\partial t} = b_{eff}(\rho)\rho(\mathbf{x}, t)(1 - \rho(\mathbf{x}, t)) - \alpha\rho(\mathbf{x}, t), \quad (5)$$

where

$$b_{eff}(\rho) = \frac{be^{-\delta \int G(\mathbf{x}-\mathbf{r})\rho(\mathbf{r}, t)d\mathbf{r}}\sigma}{\sigma + 1 - \rho(\mathbf{x}, t)}. \quad (6)$$

Thus, we have a logistic-type equation with an effective growth rate that depends nonlocally on the density itself, and which is a combination of long-range competition and local facilitation mechanisms (fire). The probability of surviving a fire is higher when the local density of trees

is higher, as can be seen from the definition in equation (4).

In figure 1 we show numerical solutions for the mean field equation (1) (lines) and the spatially explicit model (equation 5) (dots) in the stationary state ( $t \rightarrow \infty$ ) using different values of the competition. We have used a top-hat function as the competition kernel,  $G(\mathbf{x})$  (See section II B for more details on the kernel choice). We observe a very good agreement of both descriptions which becomes worse when we get closer to the critical point  $\sigma^*$ , where the model presents a phase transition from a tree-grass coexistence to a grassland state. This disagreement appears because while the mean field equation describes an infinite system, the Eq. (5) description forces us to choose a size for the system.

The model reproduces the long-term coexistence between grass and trees that is characteristic of savannas. To explore this coexistence, we study the long-time behavior of the system and analyze the homogeneous stationary solutions of Eq. (5), which has two fixed points. The first one is the absorbing state representing the absence of trees,  $\rho_0 = 0$ , and the other can be obtained, in the general case, by numerically solving

$$b_{eff}(\rho_0)(1 - \rho_0) - \alpha = 0. \quad (7)$$

In the regime where  $\rho_0$  is small (near the critical point), if competition intensity,  $\delta$ , is also small, it is possible to obtain an analytical expression for the critical value of the probability of surviving a fire,  $\sigma^*$ ,

$$\sigma^* = \frac{\alpha}{b - \alpha}. \quad (8)$$

Outside of the limit where  $\delta \ll 1$ , we can solve Eq. (7) numerically in  $\rho_0$  to show that the critical value of the fire resistance parameter,  $\sigma^*$ , does not depend on competition. A steady state with trees is stable for higher fire survival probability (Fig. 1).

The model, then, shows a transition from a state where grass is the only form of vegetation to another state where trees and grass coexist at  $\sigma^*$ . In what follows, we fix  $\alpha = 1$ , so we choose our temporal scale in such a way that time is measured in units of  $\alpha$ . This choice does not qualitatively affect our results.

## B. Linear stability analysis

The spatial patterns appearing in the nonlocal savanna model can be studied by performing a linear stability analysis [29] of the stationary homogeneous solutions of equation (5),  $\rho_0 = \rho_0(\sigma, \delta)$ . The stability analysis is performed by considering small harmonic perturbations around  $\rho_0$ ,  $\rho(\mathbf{x}, t) = \rho_0 + \epsilon e^{\lambda t - i\mathbf{k}\cdot\mathbf{x}}$ ,  $\epsilon \ll 1$ . After some calculations (A), one arrives at the dispersion relation

$$\lambda(k; \sigma, \delta) = b_{eff}(\rho_0) \frac{1 + \sigma(1 - 2\rho_0)}{\sigma - \rho_0 + 1} - b_{eff}(\rho_0) \frac{\rho_0 [2 - \rho_0 + \delta \hat{G}(k)(\rho_0 - 1)(\rho_0 - 1 - \sigma)]}{(\sigma - \rho_0 + 1)} - 1, \quad (9)$$




FIG. 1: Grass-coexistence phase transition. Stationary tree density,  $\rho_0$ , as a function of the resistance to fires parameter,  $\sigma$ . The lines come from the mean field solution, Eq. (7), and the dots from the numerical integration of Eq. (5) over a square region of 1 ha. We have chosen  $\alpha = 1$ , and  $b = 5$ . In the case of the spatial model,  $\rho_0$  involves an average of the density of trees over the studied patch of savanna.

where  $\hat{G}(k)$ ,  $k = |\mathbf{k}|$ , is the Fourier transform of the kernel,

$$\hat{G}(\mathbf{k}) = \int G(\mathbf{x}) e^{-i\mathbf{k}\cdot\mathbf{x}} d\mathbf{x}. \quad (10)$$

The critical values of the parameters of the transition to pattern,  $\delta_c$  and  $\sigma_c$ , and the fastest growing wavenumber  $k_c$  are obtained from the simultaneous solution of

$$\lambda(k_c; \sigma_c, \delta_c) = 0, \quad (11)$$

$$\left( \frac{\partial \lambda}{\partial k} \right)_{k_c; \sigma_c, \delta_c} = 0. \quad (12)$$

Note that  $k_c$  represents the most unstable mode of the system, which means that it grows faster than the others and eventually dominates the state of the system. Therefore, it determines the length scale of the spatial pattern. These two equations yield the values of the parameters  $\delta$  and  $\sigma$  at which the maximum of the curve  $\lambda(k)$ , right at  $k_c$ , starts becoming positive. This signals the formation of patterns in the solutions of Eq. (5). As Eq. (12) is explicitly written as

$$\lambda'(k_c) = b_{eff}(\rho_0) \delta \rho_0 \hat{G}'(k_c) (\rho_0 - 1), \quad (13)$$

the most unstable wavenumber  $k_c$  can be obtained by evaluating the zeros of the derivative of the Fourier transform of the kernel.

Equation (9) shows that competition, through the kernel function, fully determines the formation of patterns in the system. The local facilitation appears in  $b_{eff}(\rho_0)$  and it is not relevant in the formation of spatial structures. If the Fourier transform of  $G$  never takes positive values, then  $\lambda(k; \sigma, \delta)$  is always negative and only the homogeneous solution is stable. However, when  $\hat{G}$  can take

negative solutions then patterns may appear in the system. What does this mean in biological terms? Imagine that we have a family of kernels described by a parameter  $p$ :  $G(\mathbf{x}) = \exp(-|\mathbf{x}|/R)^p$  ( $R$  gives the range of competition). The kernels are more peaked around  $\mathbf{x} = 0$  for  $p < 2$  and more box-like when  $p > 2$ . It turns out that this family of functions has non-negative Fourier transform for  $0 \leq p < 2$ , so that no patterns appear in this case. A lengthy discussion of this property in the context of competition of species can be found in [32]. Thus, the shape of the competition kernel dictates whether or not patterns will appear in the system. If pattern formation is possible, then the values of the fire and competition parameters govern the type of solution (see below).

Our central result for nonlocal competition is that, contrary to conventional wisdom, it can, in the limit of infinitesimally short (purely local) facilitation, promote the clustering of trees. Whether or not this occurs depends entirely on the shape of the competition kernel. For large  $p$  we have a box-like shape, and in these cases trees compete strongly with other trees, roughly within a distance  $R$  from their position. The mechanism behind this counterintuitive result is that trees farther than  $R$  away from a resident tree area are not able to *invade* the zone defined by the radius  $R$  around the established tree (their seeds do not establish there), so that an exclusion zone develops around it. For smaller  $p$  there is less competition and the exclusion zones disappear.

For a more detailed analysis, one must choose an explicit form for the kernel function. Our choice is determined by the original  $P_C$  taken in [11], so that it decays exponentially with the number of trees in a neighborhood of radius  $R$  around a given tree. Thus, for  $G$  we take the step function (limit  $p \rightarrow \infty$ )

$$G(|\mathbf{r}|) = \begin{cases} 1 & \text{if } |\mathbf{r}| \leq R \\ 0 & \text{if } |\mathbf{r}| > R. \end{cases} \quad (14)$$

As noticed before, the idea behind the nonlocal competition is to capture the effect of the long roots of a tree. The kernel function defines the area of influence of the roots, and it can be modeled at first order with the constant function of equation (14). Thus the parameter  $R$ , which fixes the *nonlocal* interaction scale, must be of the order of the length of the roots [4]. Since the roots are the responsible for the adsorption of resources (water and soil nutrients), a strong long-range competition term implies strong resource depletion. For this kernel the Fourier transform is [33]  $\hat{G}(k) = 2\pi R^2 J_1(kR)/kR$  and its derivative is  $\hat{G}'(k) = -2\pi R^2 J_2(kR)/k$ , where  $k \equiv |\mathbf{k}|$ , and  $J_i$  is the  $i^{\text{th}}$ -order Bessel function. Since  $\hat{G}(k)$  can take positive and negative values, pattern solutions may arise in the system, that will in turn depend on the values of  $\delta$  and  $\sigma$ . The most unstable mode is numerically obtained as the first zero of  $\lambda'(k)$ , Eq. (13), which means the first zero of the Bessel function  $J_2(kR)$ . This value only depends on  $R$ , being independent of the resistance to fires and competition, and it is  $k_c = 5.136/R$ . Because a pattern

of  $n$  cells is characterized by a wavenumber  $k_c = 2\pi n/L$ , where  $L$  is the system size, the typical distance between clusters,  $d_t = L/n$ , using the definition of the critical wavenumber is given by  $d_t \approx 1.22R$ . In other words, it is approximately the range of interaction  $R$ . This result is also independent of the other parameters of the system.

Since we are interested in the effect of competition and fire on the distribution of savanna trees, we will try to fix all the parameters but  $\sigma$  and  $\delta$ . We will explore the effect of different values of these parameters on the results. First, we have chosen (as in [11]) the death rate  $\alpha = 1$ , and solving Eq. (7) we will roughly estimate the birth rate,  $b$ . We will work in the limit of intermediate to high mean annual precipitation, so water is non-limiting and thus we can neglect the effects of competition ( $\delta = 0$ ). At this intermediate to high mean annual precipitation the empirically observed upper limit of savanna tree cover is approximately  $\rho_0 = 0.8$  [3, 20]. To reach this upper limit in the tree cover, disturbances must also be absent, implying no fire ( $\sigma \rightarrow \infty$ ). In this limit, the mean field equation (1) is quantitatively accurate, as it is shown in Figure 1, and the stationary mean field solution of the model depends only on the birth rate

$$\rho_0(\sigma \rightarrow \infty) = \frac{b-1}{b}. \quad (15)$$

It can be solved for  $b$  for a fixed  $\rho_0 = 0.8$ , and it yields  $b = 5$  [11]. In the following we just consider the dependence of our results on  $\delta$  and  $\sigma$ . In particular,  $\rho_0 = \rho_0(\sigma, \delta)$ .

The phase diagram of the model, computed numerically, is shown in Fig. 2, where we plot the spatial character of the steady solution (homogeneous or inhomogeneous) as a function of  $\delta$  and  $\sigma$ . Note that increasing competition enhances the inhomogeneous or pattern solution. This is because, as we are now in the case of a kernel giving rise to clusters, increasing  $\delta$  makes it more difficult to enter the exclusion zones in-between the clusters. For very strong competition (high, unrealistic,  $\delta$ ), fire has no influence on the pattern.

The critical line separating these two solutions (pattern and homogeneous) can be analytically computed as a function of the parameters  $\delta$ ,  $\sigma$ ,  $\rho_0$  and  $\hat{G}(k_c)$  (see Eq. (B1) in B). In Figure 2 we have plotted (with crosses) this critical line separating homogeneous and pattern solutions for the step kernel. Note that the stationary density of trees,  $\rho_0$ , must be computed numerically from Eq. (7).

With  $b = 5$ , in the absence of fire ( $\sigma \rightarrow \infty$ ), and for weak competition, we can take the limits  $\delta \rightarrow 0$  and  $\sigma \rightarrow \infty$  of the dispersion relation Eq. (9), leading to

$$\lambda(k; \delta \rightarrow 0, \sigma \rightarrow \infty) = 4 - 10\rho_0. \quad (16)$$

In Fig. 1, for large  $\sigma$ , it can be seen that typically  $\rho_0 > 0.4$ , so Eq. (16) becomes negative. This result means that in this limit, trees are uniformly distributed in the system as there is no competition, and space does not play a relevant role in the establishment of new trees.




FIG. 2: Phase diagram of the mean field equation (5) for  $b = 5.0$ ,  $\alpha = 1.0$ , and a step kernel. The absorbing-active transition is shown at  $\sigma^*$  with circles (o). The homogeneous-pattern transition (Eq. (B1)) is indicated with crosses (x). The diamond, the square, and the up-triangle show the value of the parameters  $\sigma$  and  $\delta$  taken in Figures 3(a)-(c) respectively. The stars point out the transition to inhomogeneous solutions in the stochastic model as described in Section III, with  $\Gamma = 0.2$ .

Such situation could be interpreted as favorable to forest leading to a fairly homogeneous density of trees. This result agrees with the phase plane plotted in Figure 2. In biological terms, there are no exclusion zones in the system because there is no competition.

### C. Numerical simulations

The previous analysis provides information, depending on the competition and fire parameters, about when the solution is spatially homogenous and when trees arrange in clusters. However, the different shapes of the patterns have to be studied via numerical simulations [34] of the whole equation of the model. We have taken a finite square region of savanna with an area of 1 ha., allowed competition to occur in a circular area of radius  $R = 8 m$ , and employed periodic boundary conditions and a finite differences algorithm to obtain the numerical solution. Similarly to what has been observed in studies of semi-arid water limited systems [26, 28], different structures, including gaps, stripes, and tree spots, are obtained in the stationary state as we increase the strength of competition for a fixed value of the fire parameter or, on the other hand, as we decrease the resistance to fires for a given competition intensity. In both equivalent cases, we observe this spectrum of patterns as far as we go to a more dry state of the system, where resources (mainly water) are more limited (see Figs. 3(a)-3(c)) and competition is consequently stronger. This same sequence of appearance of patterns has been already observed in the presence of different short-range facilitation mechanisms [26, 35]. It indicates that, when  $\delta$  is increased (i.e. the probability of surviving competition is decreased), new trees cannot establish in the exclusion areas so clustering




FIG. 3: (a) Grass spots ( $\delta = 7.0$ ), (b) striped grass vs. tree ( $\delta = 8.0$ ), and (c) tree spots ( $\delta = 11.0$ ) patterns in the deterministic model in a square patch of savanna of 1 ha.  $\sigma = 2.9$ ,  $R = 8.0 m$ ,  $b = 5.0$  and  $\alpha = 1.0$  in all the plots.

is enhanced.

On the other hand, in the case of fire-prone savannas, previous works had only shown either tree spot [36] or grass spots [37]. Therefore, at some values of the parameter space (see Fig. 3b), the patterns in our deterministic approach are not observed in mesic savannas, and should correspond to semiarid systems. However, we will show in the following sections that under the parameter constraints of a mesic savanna, and considering the stochastic nature of the tree growth dynamics in the system (i.e. demographic noise), our model shows realistic spatial structures.

A much more quantitative analysis of the periodicity in the patterns can be performed via the structure function. This will be helpful to check the previous results and, especially, for the analysis of the data of the stochastic model of the next section, for which we will not present analytical results. The structure function is defined as the modulus of the spatial Fourier transform of the density of trees in the stationary state,

$$S(k) = \left\langle \left| \int d\mathbf{x} e^{i\mathbf{k}\cdot\mathbf{x}} \rho(\mathbf{x}, t \rightarrow \infty) \right|^2 \right\rangle, \quad (17)$$

where the average is a spherical average over the wavevectors with modulus  $k$ . The structure function is helpful to study spatial periodicities in the system, similar to the power spectrum of a temporal signal. Its maximum identifies dominant periodicities, which in our case are the distances between tree clusters. Note that the geometry of the different patterns cannot be uncovered with the structure function, since it involves a spherical average. In Fig. 4, we show the transition to patterns using the maximum of the structure function as a function of the competition parameter. A peak appears when there are




FIG. 4: Maximum of the structure function for different values of the competition parameter  $\delta$  at long times. The fire parameter is fixed at  $\sigma = 2.9$ . Black circles refers to the deterministic model and red squares to the stochastic model,  $\Gamma = 0.20$ .




FIG. 5: Numerical computation of the structure function defined in equation (17) for different values of the demographic noise intensity.  $\delta = 9.8$ ,  $\sigma = 2.9$ ,  $R = 8 m$ ,  $\alpha = 1.0$ ,  $b = 5.0$ .

spatial structures in the system, so  $Max[S(k)] \neq 0$ . However, we do not have information about the values where the shapes of the patterns change. Taking  $R = 8 m$ , the peak is always at  $\lambda_c = 10m$  for our deterministic savanna model, independently of the competition and fire resistance parameters, provided that they take values that ensure the emergence of patterns in the system (see the line labeled by  $\Gamma = 0$  in Fig. 5; for the definition of  $\Gamma$  see next section). This result is in good agreement with the theoretical result provided for the wavelength by the linear stability analysis  $\lambda = 2\pi/k_{max} = 9.78 m$ , which is also independent of competition and resistance to fires.

### III. STOCHASTIC MODEL

The perfectly periodic patterns emerging in Fig. 3 from the deterministic model seem to be far from the disordered ones usually observed in aerial photographs of mesic savannas and shown by individual based models [6–8, 11]. We have so far described a savanna system in terms of the density of trees with a deterministic dynam-

ics. The interpretation of the field  $\rho(\mathbf{x}, t)$  is the density of tree (active) sites in a small volume,  $V$ . If we think of trees as reacting particles which are born and die probabilistically, then to provide a reasonable description of the underlying individual-based birth and death dynamics, we have to add a noise term to the standard deterministic equation. It will take into account the *intrinsic* stochasticity present at the individual level in the system.

If we take a small volume,  $V$ , the number of reactions taking place is proportional to the number of particles therein,  $N$ , with small deviations. If  $N$  is large enough, the central limit theorem applies to the sum of  $N$  independent random variables and predicts that the amplitude of the deviation is of the order of  $\sqrt{N} \propto \sqrt{\rho(\mathbf{x}, t)}$  [38]. This stochasticity referred to as demographic noise. The macroscopic equation is now stochastic,

$$\begin{aligned} \frac{\partial \rho(\mathbf{x}, t)}{\partial t} = & b_{eff}(\rho)[\rho(\mathbf{x}, t) - \rho^2(\mathbf{x}, t)] - \\ & - \alpha \rho(\mathbf{x}, t) + \Gamma \sqrt{\rho(\mathbf{x}, t)} \eta(\mathbf{x}, t), \end{aligned} \quad (18)$$

where  $\Gamma \propto \sqrt{b_{eff}}$  (but we take it as a constant, [39]) modulates the intensity of  $\eta(\mathbf{x}, t)$ , a Gaussian white noise term with zero mean and correlations given by Dirac delta distributions

$$\langle \eta(\mathbf{x}, t) \eta(\mathbf{x}', t') \rangle = \delta(\mathbf{x} - \mathbf{x}') \delta(t - t'). \quad (19)$$

The complete description of the dynamics in Eq.(18) should have the potential to describe more realistic patterns.

We first investigate the effect of demographic noise on the persistence of trees in the system. We show in (Fig. 6) that the critical point,  $\sigma^*$ , depends on the value of the competition parameter  $\delta$ . This effect is rather small, so that when  $\delta$  increases the transition to the grassland state appears only for a slightly larger  $\sigma$  (i.e, less frequent fire). The reason seems to be that fire frequency and intensity depend on grass biomass. Seasonally wet savannas support much more grass biomass that serves as fuel for fires during the dry season [40, 41]. Dry savannas have much lower grass biomass, so they do not burn as often or as intensely. The shift of the critical value of  $\sigma$  when competition is stronger is consistent with the one showed in [11], as can be seen comparing Figure 2 in [11] with Figure 6 here. Besides, the values obtained for  $\sigma^*$  are larger when we consider the demographic stochasticity [42] neglected in the deterministic field approach.

We explore numerically the stochastic savanna model using an algorithm developed in [39] (See C). Note that the noise makes the transition to pattern smoother so the change from homogeneous to inhomogeneous spatial patterns is not as clear as it is in the limit where the demographic noise vanishes (See Fig. 4). The presence of demographic noise in the model, as shown in Fig. 2 (red stars), also decreases the value of the competition strength at which patterns appear in the system, as has been observed in other systems. Mathematically, these new patterns appear since demographic noise maintains




FIG. 6: Active-absorbing phase transition in the deterministic (Circles) and the stochastic model (Squares). In the later case, we integrate the Eq. (18) with  $\Gamma = 0.2$  and average the density of trees in the steady state.

Fourier modes of the solution which, due to the value of the parameters, would decay in a deterministic approach [43]. Biologically, exclusion zones are promoted by demographic noise, since it does not affect regions where there are not trees. On the other hand in vegetated areas fluctuations may enhance tree density, leading to stronger competition. The presence of demographic noise in the model allows the existence of patterns under more humid conditions. This result is highly relevant for mesic savannas, as we expect competition to be of low to intermediate strength in such systems. We show two examples of these irregular patterns in Fig. 7(a) and Fig. 7(b). Unrealistic stripe-like patterns no longer appear in the stochastic model.

We have studied the dynamics of the system for some values of the fire and competition parameters. Demographic noise influences the spatial structures shown by the model. The deterministic approach shows a full spectrum of patterns which are not visually realistic for mesic savannas (but for arid systems). The role of the noise is to transform this spectrum of regular, unrealistic patterns into more irregular ones (Figures 7(a)-7(d)) that remind the observed in aerial photographs of real mesic savannas. On the other hand, these patterns are statistically equivalent to the deterministic ones, as it is shown with the structure function in Fig. 5. The dominant scale in the solution is given by the interaction radius,  $R$ , and it is independent of the amplitude of the noise (see the structure function in Figure 5, peaked around  $\lambda = 10$  m independently of the noise). Besides, over a certain threshold in the amplitude, demographic noise destroys the population of trees. Therefore, the model presents an active-absorbing transition with the noise strength,  $\Gamma$ , being the control parameter.

#### IV. DISCUSSION

Understanding the mechanisms that produce spatial patterns in savanna tree populations has long been an area of interest among savanna ecologists [6–10]. A key step in such an analysis is defining the most parsimo-




FIG. 7: Patterns of the stochastic model in a square patch of savanna of 1 ha.  $\sigma = 2.9$ ,  $R = 8.0$  m,  $b = 5.0$  and  $\alpha = 1.0$  in all the plots. (a)  $\Gamma = 0.2$ ,  $\delta = 3.0$ . (b)  $\Gamma = 0.2$ ,  $\delta = 5.0$ . (c)  $\Gamma = 0.1$ ,  $\delta = 10.0$ . (d)  $\Gamma = 0.2$ ,  $\delta = 10.0$ .

nous combination of mechanisms that will produce the pattern in question. In this paper the combination of long-range competition for resources and the facilitation induced by fire are considered the responsible of the spatial structures, in the line of studies of vegetation pattern formation in arid systems, where also a combination of long-range inhibition and short-range facilitation is introduced [22, 23, 26, 27]. The main difference is that the facilitation provided by the protection effect of adult trees against fires in our savanna model takes the short-range facilitation to its infinitesimally short limit (i.e, local limit). Under this assumption we have studied the conditions under which our model could account for patterns. We have shown that nonlocal competition combined with local facilitation induces the full range of observed spatial patterns, provided the competition term enters nonlinearly in the equation for the density of trees, and that competition is strong enough.

The key technical requirement for this effect to occur is that the competition kernel must be an almost constant function in a given competition region, and decay abruptly out of the region. We verify this condition working with supergaussian kernel functions. In practice, this means that competition kernels whose Fourier transform takes negative values for some wavenumber values, will lead to competition driven clustering.

The other mechanism we have considered for a minimalistic but realistic savanna model, fire, has been shown to be relevant for the coexistence of trees and grass and for the shape of the patterns. However, competition is the main ingredient allowing pattern solutions to exist in the model. If the shape of the kernel allows these types of solutions, then the specific values of fire and competition parameters determine the kind of spatial structure

that develops. It is also worth mentioning that one can observe the full spectrum of patterns in the limit where fires vanish ( $\sigma \rightarrow \infty$ ), so there is no facilitation at all, provided competition is strong enough. However, when there is no competition,  $\delta = 0$ , no patterns develop regardless of the value of the fire term. Therefore, we conclude that the nonlocal competition term is responsible for the emergence of clustered distributions of trees in the model, with the fire term playing a relevant role only to fix the value of the competition parameter at which patterns appear. In other words, for a given competition strength, patterns appear more readily when fire is combined with competition. A similar mechanism of competitive interactions between species has been shown to give rise to clusters of species in the context of classical ecological niche theory. Scheffer and van Nes [44] showed that species distribution in niche space was clustered, and Pigolotti et al. [32] showed that this arises as an instability of the nonlocal nonlinear equation describing the competition of species.

Long-distance competition for resources in combination with the local facilitation due to the protection effect of adult trees in the establishment of juvenile ones can explain the emergence of realistic structures of trees in mesic savannas. In these environmental conditions, competition is limited, so we should restrict to small to intermediate values of the parameter  $\delta$ , and the effect of fires is also worth to be taken into account. However, these two ingredients give a full range of patterns observed in vegetated systems, but not in the particular case of savannas. It is necessary to consider the role of demographic noise, which is present in the system through the stochastic nature of the birth and death processes of individual trees. In this complete framework our model shows irregular patterns of trees similar to the observed in real savannas.

The other important feature of savannas, the characteristic long-time coexistence of trees and grass is well captured with our model (Figures 1 and 6). Besides, the presence of demographic noise, as it is shown in Figure 6, makes our approach much more realistic, since the persistence of trees in the face of fires is related to the water in the system. On the other hand, demographic stochasticity causes tree extinction at lower fire frequencies (larger  $\sigma$ ) than in the deterministic case. This is because random fluctuations in tree density are of sufficient magnitude that this can hit zero even if the deterministic stationary tree density (for a given fire frequency) is greater than zero. This effect vanishes if we increase the system size. The demographic noise is proportional to the density of trees (proportional to  $(L_x \times L_y)^{-1}$ ), so fluctuations are smaller if we study bigger patches of savannas. As usually happens in the study of critical phenomena in Statistical Mechanics, the extinction times due to demographic noise increase exponentially with the size of the system for those intensities of competition and fire that allow the presence of trees in the stationary state. Over the critical line, this time will follow a power law scaling,

and a logarithmic one when the stationary state of the deterministic model is already absorbing (without trees) [45].

## V. SUMMARY

We have shown the formation of patterns in a minimal savanna model, that considers the combination of long-range competition and local facilitation mechanisms as well as the transition from trees-grass coexistence to a grass only state.

The salient feature of the model is that it only considers nonlocal (and nonlinear) competition through a kernel function which defines the length of the interaction, while the facilitation is considered to have an infinitesimally short influence range. Our model thus differs from standard kernel-based savanna models that feature both short-range facilitation and long-range competition. The same sequence of spatial patterns appears in both approaches, confirming Rietkerk and van de Koppel's [31] suggestion that short-range facilitation does not induce spatial pattern formation by itself, and long-distance competition is also needed. It also suggests that long-range competition could be not only a necessary, but also a sufficient condition to the appearance of spatial structures of trees.

Inspired by [11], we have proposed a nonlocal deterministic macroscopic equation for the evolution of the local density of trees where fire and tree-tree competition are the dominant mechanisms. If the kernel function falls off with distance very quickly (the Fourier transform is always positive) the system only has homogenous solutions. In the opposite case, patterns may appear depending on the value of the parameters ( $\delta$  and  $\sigma$ ), and in a sequence similar to the spatial structures appearing in standard kernel-based models. Under less favorable environmental conditions, trees tend to arrange in more robust structures to survive (Fig. 3(d)). Biologically, trees are lumped in dense groups, separated by empty regions. Entrance of new trees in these *exclusion zones* is impossible due to the intense competition they experience there.

A great strength of our approach is that our deterministic analysis is formal, and we have shown the different spatial distributions of the trees that occur as competition becomes more intense, concluding that self organization of trees is a good mechanism to promote tree survival under adverse conditions [26]. Trees tend to cluster in the high competition (low resources) limit (Fig. 3(d)), due to the formation of exclusion zones caused by non-local competition, and not as a result of facilitation. However, because we are dealing with a deterministic model, the patterns are too regular and the transition between the grass-only and a tree-populated states is independent of tree competition. We therefore considered stochasticity coming from the stochastic nature of individual birth and death events, to provide a more realistic description of savanna dynamics. Calabrese et al. [11] also

noted that savanna-to-grassland transition was independent of competition intensity in the mean field approach, but not when demographic noise was included. In the present model, both the grassland to savanna transition and the spatial structures that develop are influenced by demographic stochasticity. In the case of spatial structures, demographic noise is specially relevant, since it turns much of the unrealistic patterns of the deterministic model into more realistic ones, that remind the observed in real savannas. It also allows the existence of periodic arrangements of trees in more humid systems, which means environmental conditions closer to mesic savannas.

We have quantified the characteristic spacing of spatial patterns through the structure function. The irregular patterns produced by the stochastic model still have a dominant wavelength whose value is the same as in the deterministic model and depends only on the value of the range of the interaction,  $R$ , in the kernel function. The match between the typical spatial scale of the patterns and the characteristic distance over which nonlocal competition acts indicates that competition is responsible for the presence of clustered spatial structures.

## VI. ACKNOWLEDGMENTS

R.M-G is supported by the JAEPredoc program of CSIC. R.M-G. and C.L. acknowledge support from MICINN (Spain) and FEDER (EU) through Grant No. FIS2007- 60327 FISICOS. We acknowledge Federico Vázquez and Emilio Hernández-García for their comments and discussion. We also acknowledge the detailed reading and insightful comments of three anonymous referees which greatly helped to improve this manuscript.

### Appendix A: Linear stability analysis

This appendix shows the details of the linear stability analysis, in particular how it is obtained the dispersion relation in Eq. (9). We consider the stationary solution  $\rho_0$  plus a small harmonic perturbation,

$$\rho(\mathbf{x}, t) = \rho_0 + \epsilon e^{\lambda t - i\mathbf{k} \cdot \mathbf{x}}, \quad (\text{A1})$$

where  $\epsilon \ll 1$ . Substituting Eq. (A1) into the original equation (5), and retaining only linear terms in  $\epsilon$ , we arrive to the relation dispersion

$$\lambda(k) = bC\sigma(\rho_0 - \rho_0^2) \left[ \frac{1}{(\sigma+1-\rho_0)^2} - \frac{\hat{G}(k)\delta}{\sigma+1-\rho_0} \right] + bC\sigma \frac{1-2\rho_0}{\sigma+1-\rho_0} - 1, \quad (\text{A2})$$

where  $\hat{G}(k)$  is the Fourier transform of the kernel,  $\hat{G}(k) = \int G(x) \exp(\lambda t - i\mathbf{k} \cdot \mathbf{x})$ , and  $C \equiv \exp(-\delta\rho_0)$ , provided that we deal with normalized kernels. Equation (A2) can be written as Eq. (9) using the definition of  $b_{eff}(\rho_0)$ .

### Appendix B: Expression of the transition to pattern critical line.

We show here the analytical expression for the critical line in the transition from homogeneous to inhomogeneous solutions. Starting from Eq. (11) it is possible to write an expression for the value of the resistance to fires parameter,  $\sigma$ , at which the macroscopic equation (5) starts showing pattern solutions, as a function of the competition parameter,  $\delta$ , and the most unstable mode  $k_c$ . Considering the value of the parameters taken in our study,  $b = 5$  and  $\alpha = 1$ , it is

$$\sigma_c = \frac{(\rho_0-1)[5(\rho_0-1)(\delta\hat{G}(k_c)\rho_0-1)-2e^{\delta\pi R^2\rho_0}]}{10[1-2\rho_0+\delta\hat{G}(k_c)\rho_0(1+\rho_0)-e^{\delta\pi R^2\rho_0/5}]} + \frac{(\rho_0-1)\sqrt{5[5(\rho_0-1)^2(\delta\hat{G}(k_c)\rho_0-1)^2-4e^{\delta\pi R^2\rho_0}\rho_0]}}{10[1-2\rho_0+\delta\hat{G}(k_c)\rho_0(1+\rho_0)-e^{\delta\pi R^2\rho_0/5}]} \quad (\text{B1})$$

This complicated expression must be evaluated numerically together with the solution of Eq. (7) for the stationary density of trees, which is also a function of the competition and fire parameters. We show the results in Figure 2, where the curve, represented with the black crosses, fits perfectly with the numerical results from the linear stability analysis.

### Appendix C: Numerical algorithm for the integration of the equation (18).

The integration of stochastic equations where the noise amplitude depends on the square root of the variable,  $\rho$ , and there are absorbing states (i.e, states where the system stays indefinitely), has awoken a great interest, specially in the study of critical phenomena (i.e, properties of the system that appear when it is close to the critical point, often the absorbing state). The amplitude of the fluctuations tends to zero there, and thus numerical instabilities may appear. Recently [46, 47] a very efficient method has been developed, but we have used in this work an older one, presented in [39], since its implementation is easier and it gives precise results working far from the transition point. It consists on discretizing the Langevin equation, taking a step size  $\Delta\rho$  in the variable.

To apply the method to equation (18), first of all we discretize the space. Particularly, we compute the integral in the exponential term approximating it by a sum of the field evaluated in the nodes of the discrete space

$$\int \rho(\mathbf{x}, t) G(\mathbf{x} - \mathbf{x}') dx \approx \sum_{i=1}^{N_x} \sum_{j=1}^{N_y} \rho_{i,j} G_{i,j;i',j'} \Delta x \Delta y. \quad (\text{C1})$$

Then, we integrate the temporal dependence. The key of the algorithm is to prevent  $\rho + \Delta\rho$  to take negative values. From a general equation

$$\frac{d\rho}{dt} = f(\rho) + \sqrt{\rho}\psi(t), \quad (\text{C2})$$

where  $\psi(t)$  is a gaussian white noise with zero mean and delta correlated, it is

$$\Delta\rho = f(\rho)\Delta t + \sqrt{\rho}\Delta W, \quad (\text{C3})$$

where  $\Delta W = \sqrt{\Delta t}Y$ .  $Y$  is a Gaussian number with zero mean and unit variance. At this point, to prevent  $\rho + \Delta\rho$  to take negative values, the author in [39] proposes to discretize the density setting  $\rho = n\rho_{min}$  and to truncate the gaussian distribution from where  $Y$  is obtained symmetrically so that  $|Y| \leq Y_{max}$ . The negatives values are avoided requiring  $Y_{max}\sqrt{\Delta t} \leq \rho_{min}$ . It can be done in many ways but following [39] we use

$$\begin{aligned} Y_{max} &= \frac{|\ln \Delta t|}{3}, \\ \rho_{min} &= \frac{(\ln \Delta t)^2 \Delta t}{9}. \end{aligned} \quad (\text{C4})$$

Finally, rescaling the equation, we can achieve a discretized version in which positive and zero-mean noise are ensured at the cost of a ‘‘quantized’’ density.

#### Appendix D: The effect of rainfall: Random switching between death and birth

One of the key ingredients for the long coexistence between grass and trees is the largely inhomogeneous temporal distribution of precipitations over time [3, 13, 34]. We have studied this environmental variability following the idea in [30], considering the switching between unstressed vegetation growth, given by the first term in (5), and drought-induced vegetation decay, represented with the second term in Eq. (5). These processes take place each time step with probability  $P$  and  $1 - P$ , respectively. From now on, we call

$$\begin{aligned} f_b[\rho(\mathbf{x}, t)] &= b_{eff}(\rho) [\rho(\mathbf{x}, t) - \rho^2(\mathbf{x}, t)], \\ f_d[\rho(\mathbf{x}, t)] &= -\alpha\rho(\mathbf{x}, t), \end{aligned} \quad (\text{D1})$$

and

$$f_{\pm}[\rho(\mathbf{x}, t)] = \frac{1}{2} [f_b[\rho(\mathbf{x}, t)] \pm f_d[\rho(\mathbf{x}, t)]]. \quad (\text{D2})$$

The random dynamics of the system is written in terms of a stochastic partial differential equation,

$$\frac{\partial\rho(\mathbf{x}, t)}{\partial t} = f_+[\rho(\mathbf{x}, t)] + f_-[\rho(\mathbf{x}, t)]\xi_{dn}(t), \quad (\text{D3})$$

where  $\xi_{dn}(t)$  is a dichotomous noise (DMN), assuming values  $+1$  (wet season) and  $-1$  (dry season) with probability  $P$  and  $1 - P$ , respectively.

If the rate of random switching, taken as the inverse of the integration time step, is relatively fast respect to the rate of convergence to equilibrium in each of the two states, we can replace the noise term in Eq. (D3) with its average value,  $\langle \xi_{dn}(t) \rangle = 1 - 2P$ . It is meaningful since the rainfall seasons are much shorter than the time needed to reach one of the equilibrium stationary states of death and birth processes,  $\rho(\mathbf{x}, t) = 0, 1$ , respectively. This substitution leads to a deterministic equation

$$\frac{\partial\rho(\mathbf{x}, t)}{\partial t} = f_+[\rho(\mathbf{x}, t)] + f_-[\rho(\mathbf{x}, t)](1 - 2P), \quad (\text{D4})$$

where we will be able to perform linear stability analysis as usual. The new dispersion relation is easily obtained,

$$\begin{aligned} \lambda(k; \sigma, \delta) &= b_{eff}(\rho_0)P \frac{1+\sigma(1-2\rho_0)}{\sigma-\rho_0+1} - (1 - P) \\ &- b_{eff}(\rho_0)P \frac{\rho_0[2-\rho_0+\delta\hat{G}(k)(\rho_0-1)(\rho_0-1-\sigma)+2\sigma]}{(\sigma-\rho_0+1)}, \end{aligned} \quad (\text{D5})$$

which means that the main effect of the dichotomous noise is to renormalize the rates  $\alpha$  and  $b$ . The patterns observed now are the same as the ones in the deterministic case, though the regions where they emerge change in accordance with this renormalization. Thus, the effect of stochastic precipitation, as modeled with this random switching mechanism, is a change of the parameter values for the different transitions observed in the deterministic continuum model Eq. (5).

According to the value of  $P$ , an absorbing-active phase transition is observed,  $P_c \approx 0.20$ . Small values of  $P$ , meaning long dry season, lead to an absorbing state while increasing the probability of raining implies the appearance of trees in the system. In this latter case, the solution can be either homogeneous or showing spatial patterns, depending on fire and competition.

This attempt to model rainfall has not been very successful and does not give a lot of new information. Much effort of future research should be put on this point, trying to get much more realistic modelling of external environmental variability, according to empirical observations, with long runs of dry years and rare wet years.

---

[1] G. Sarmiento, The ecology of Neotropical savannas. Harvard Univ. Press, Cambridge, MA, 1984.  
[2] R. J. Scholes, and S. R. Archer. Tree-grass interactions in savannas. Annual Review of Ecology and Systematics, 28, (1997), 517-544.

[3] M. Sankaran, N.P. Hanan, R.J. Scholes, J. Ratnam, D.J. Augustine, B.S. Cade, J. Gignoux, et al. Determinants of woody cover in African savannas. Nature, 438, (2005), 846-849.  
[4] F. Borgogno, P. D’Odorico, F. Laio, and L. Ridolfi. Math-

- ematical models of vegetation pattern formation in ecohydrology. *Reviews of Geophysics*, 47, RG1005, (2009).
- [5] A.J. Belsky. Influences of trees on savanna productivity: tests of shade, nutrients, and tree-grass competition. *Ecology*, 75, (1994), 922-932.
- [6] S. Barot, J. Gignoux, and J.C. Menaut. Demography of a savanna palm tree: predictions from comprehensive spatial pattern analyses. *Ecology*, 80, 6, (1999), 1987-2005.
- [7] K.K. Caylor, H.H. Shugart, P.R. Dowty, and T.M. Smith. Tree spacing along the Kalahari transect in southern Africa. *Journal of Arid Environments*, 54, 2, (2003), 281-296.
- [8] F. Jeltsch, K. Moloney, and S.J. Milton. Detecting process from snapshot pattern: lessons from tree spacing in the southern Kalahari. *Oikos* 85(3), (1999), 451-466.
- [9] T.M. Scanlon, K.K. Caylor, S.A. Levin, and I. Rodriguez-Iturbe. Positive feedbacks promote power-law clustering of Kalahari vegetation. *Nature*, 449, 7159, (2007), 209-212.
- [10] C. Skarpe. Spatial patterns and dynamics of woody vegetation in an arid savanna. *Journal of Vegetation Science*, 2, 4, (1991), 565-572.
- [11] J.M. Calabrese, F. Vázquez, C. López, M. San Miguel, and V. Grimm. The individual and interactive effects of tree-tree establishment competition and fire on savanna structure and dynamics. *The American Naturalist*, 175, 3, (2010).
- [12] F. Jeltsch, S.J. Milton, W.R.J. Dean, and N. van Rooyen. Tree spacing and coexistence in semiarid savannas. *Journal of Ecology*, 84, (1996), 583-595.
- [13] F. Vázquez, C. López, J.M. Calabrese, and M.A. Muñoz. Dynamical phase coexistence: a simple solution to the "savanna problem". *Journal of Theoretical Biology*, 264, 2, (2010), 360366.
- [14] O.R. Vetaas. Micro-site effects of trees and shrubs in dry savannas. *Journal of Vegetation Science*, 3, 3, (1992), 337-344.
- [15] K. E. Veblen. Season and herbivore dependent competition and facilitation in a semiarid savanna. *Ecology*, 89, (2008), 1532-1540.
- [16] N. Barbier, P. Couteron, R. Lefever, V. Deblauwe, and O. Lejeune. Spatial decoupling of facilitation and competition at the origin of gapped vegetation patterns. *Ecology*, 89, (2008), 1521-1531.
- [17] M. Sankaran, J. Ratnam, and N. Hanan. Woody cover in African savannas: the role of resources, fire and herbivory. *Global Ecology and Biogeography*, 17, (2008), 236-245.
- [18] W.J. Bond, G.F. Midgley, and F.I. Woodward. What controls South African vegetation: climate or fire? *South African Journal of Botany*, 69, (2003), 79-91.
- [19] W.J. Bond. What limits trees in C4 grasslands and savannas? *Annual Review of Ecology, Evolution, and Systematics*, 39, (2008), 641-659.
- [20] G. Bucini, and N.P. Hanan. A continental-scale analysis of tree cover in African Savannas. *Global Ecology and Biogeography*, 16, (2007), 593-605.
- [21] R.M. Holdo. Stem mortality following fire in Kalahari sand vegetation: effects of frost, prior damage, and tree neighbourhoods. *Plant Ecology*, 180, (2005), 77-86.
- [22] A. Klausmeier. Regular and irregular patterns in semiarid vegetation. *Science*, 284, (1999), 1826-1828.
- [23] R. Lefever, and O. Lejeune. On the origin of the tiger brush. *Bulletin Mathematical Biology*, 59, (1997), 263-294.
- [24] R. Lefever, N. Barbier, P. Couteron, and O. Lejeune. Deeply gapped vegetation patterns: On crown/root allometry, criticality and desertification. *Journal of Theoretical Biology*, 2, 261, (2009), 194-209.
- [25] R. Lefever, and J. W. Turner. A quantitative theory of vegetation patterns based on plant structure and the non-local FKPP equation. *Comptes Rendus Mcanique*, 340, 1112, (2012), 818828.
- [26] M. Rietkerk, M.C. Boerlijst, and F. van Langevelde. Self-Organization of Vegetation in Arid Ecosystems. *The American Naturalist*, 160, (2002), 524-530.
- [27] J. von Hardenberg, E. Meron, M. Shachak, and Y. Zarmi. Diversity of vegetation patterns and desertification. *Physical Review Letters*, 97, (2001), 197101.
- [28] P. D'Odorico, F. Laio, and L. Ridolfi. Patterns as indicators of productivity enhancement by facilitation and competition in dryland vegetation. *Journal Geophysical Research*, 111, (2006), G03010.
- [29] M. C. Cross, and P. C. Hohenberg. Pattern formation out of equilibrium. *Reviews of Modern Physics*, 65, (1993), 851-1112.
- [30] P. D'Odorico, F. Laio, and L. Ridolfi. Vegetation patterns induced by random climate fluctuations. *Geophysical Research Letters*, 33, (2006), L19404.
- [31] M. Rietkerk, and J. van de Koppel. Regular Pattern Formation in Real Ecosystems. *Trends in Ecology and Evolution*, 23, 3, (2008).
- [32] S. Pigolotti, C. López, and E. Hernández-García. Species clustering in competitive Lotka-Volterra models. *Physical Review Letters*, 98, (2007), 258101.
- [33] C. López, and E. Hernández-García. Fluctuations impact on a pattern-forming model of population dynamics with non-local interactions. *Physica D: Nonlinear Phenomena*, 199, (2004), 223-234.
- [34] L. Ridolfi, P. D'Odorico, and F. Laio. *Noise-induced Phenomena in the Environmental Sciences*. Cambridge Univ. Press, Cambridge, 2011.
- [35] O. Lejeune, and M. Tlidi. A model for the explanation of vegetation stripes (tiger bush). *Journal of Vegetation Science*, 10, 2, (1999), 201-208.
- [36] O. Lejeune, M. Tlidi, and P. Couteron. Localized vegetation patches: A self-organized response to resource scarcity. *Physical Review E*, 66, (2002), 010901(R).
- [37] P. D'Odorico, F. Laio, A. Porporato, L. Ridolfi, and N. Barbier. Noise-induced vegetation patterns in fire-prone savannas. *Journal of Geophysical Research*, 112, G02021, (2007).
- [38] C.W. Gardiner. *Handbook of Stochastic Methods for Physics, Chemistry and Natural Sciences*. Springer-Verlag, Berlin-Heidelberg, 1985.
- [39] R. Dickman. Numerical study of a field theory for directed percolation. *Physical Review E*, 50, (1994), 4404.
- [40] P. D'Odorico, F. Laio, and L. Ridolfi. A probabilistic analysis of fire-induced tree-grass coexistence in savannas. *The American Naturalist*, 167, (2006), E79-E87.
- [41] N.P. Hanan, W.B. Sea, G. Dangelmayr, and N. Govenader. Do fires in savannas consume woody biomass? A comment on approaches to modeling savanna dynamics. *The American Naturalist*, 171, (2008), 851-856.
- [42] H.E. Stanley. *Introduction to phase transitions and critical phenomena*. Oxford University Press, Oxford, (1971).
- [43] T. Butler, and N. Goldenfeld. Robust ecological pattern formation induced by demographic noise. *Physical Re-*

- view E, 80, (2009), 030902(R).
- [44] M. Scheffer, and E.H. van Nes. Self-organized similarity, the evolutionary emergence of groups of similar species. *Proceedings of the National Academy of Sciences USA*, 103, (2006), 62306235.
- [45] J. Marro and R. Dickman. *Nonequilibrium Phase Transitions in Lattice Models*. Cambridge University Press, Cambridge, 1999.
- [46] I. Dornic, H. Chaté, and M.A. Muñoz, Integration of Langevin Equations with Multiplicative Noise and the Viability of Field Theories for Absorbing Phase Transitions. *Physical Review Letters*, 94, (2005), 100601.
- [47] L. Pechenik, and H. Levine. Interfacial velocity corrections due to multiplicative noise. *Physical Review E*, 59, (1999), 3893.