Synthesis

The lithic microbial ecosystems of Antarctica's McMurdo Dry Valleys

ASUNCIÓN DE LOS RÍOS, JACEK WIERZCHOS and CARMEN ASCASO

Museo Nacional de Ciencias Naturales (CSIC), Serrano 115 dpdo, Madrid-28006, Spain

arios@mncn.csic.es

Abstract: We review the lithic microbial ecosystems of the McMurdo Dry Valleys as the main form of terrestrial colonization in this region, and assess the role of environmental controls such as temperature, solar radiation, water availability, wind, nutrient availability, salinity and the physicochemical properties of the colonized rock. Epilithic communities, especially those dominated by lichens, are able to withstand extreme environmental conditions but subsurface endolithic microhabitats provide more tolerant conditions. Endolithic microbial communities can be grouped into two main classes: eukaryotic communities (dominated by lichenized fungi and algae) and prokaryotic communities (dominated by cyanobacteria). Heterotrophic bacteria and non-lichenized algae and fungi (mainly black fungi) are also components of these communities. These lithobiontic microorganisms generally have effective mechanisms against freezing temperatures and desiccation. Extracellular polymeric substances play an important role not only in protecting microbial cells but also in community organization and in mitigating microenvironmental conditions. Antarctic lithobiontic communities are comprised of microbial consortia within which multiple interactions between the different biological and abiotic components are essential for microbial survival, whilst fossils and biomarkers provide evidence of earlier successful microbial life in Antarctic deserts. Finally, the uniqueness of the present lithobiont assemblages suggests they are the outcome of geographical isolation during the evolution of the continent and not merely the descendants of a subset of globally distributed taxa that have adapted to the extreme environmental conditions.

Received 18 April 2013, accepted 7 March 2014

Key words: endoliths, lichens, lithic microhabitat, lithobiontic microorganisms

Introduction

The geological evolutionary history and geographical isolation of the Antarctic continent have produced a unique environment inhabited by species adapted to extreme conditions (Hodgson *et al.* 2010). Terrestrial ecosystems in Antarctica are restricted to a few ice-free areas (0.34% of the continent), most of which are small and isolated from one another (Convey 2011). These areas span from the deserts, nunataks, cliffs and boulder/scree fields of the continental interior to the moist cryptogam-dominated fellfields of Maritime Antarctic (Convey 2011). The largest ice-free area in Antarctica, the McMurdo Dry Valleys (Fig. 1), is among the most remote and harsh environments for terrestrial life (Doran *et al.* 2002a).

Antarctica is the only continent dominated by microorganisms that exclusively drive the biogeochemical transformations of carbon (Hopkins *et al.* 2009, Namsaraev *et al.* 2010). The presence and abundance of microbial communities in the rocks of Antarctica's terrestrial areas indicate their ability to colonize the lithic substrate. Microbial life in Antarctica retreats to niches that formed as a result of biotic–abiotic relationships on, under or within rocks (Pointing & Belnap 2012). Green algal-fungal symbiosis gives rise to diverse lichen species as the most visible components of lithic microbial ecosystems, especially when they grow on rock surfaces (Fig. 2a). These epilithic forms may scarcely cover rocks or, in areas where conditions are less extreme such as Maritime Antarctic, rocks may be heavily colonized. Under harsher environmental conditions, such as those of the Dry Valleys, crustose and foliose lichens preferentially occupy cavities or protected areas (Fig. 2b) and chasmoendolithic forms can be observed in weathered rocks (Fig. 2c). Some lichen symbiotic associations hide within the rock interior (Friedmann 1982) only to

be seen after the rock has been fractured (Fig. 3a & b). Cyanobacteria dominated communities also inhabit these rocks (Fig. 3c & d) (Friedmann *et al.* 1988, De los Ríos *et al.* 2007, Büdel *et al.* 2008). Here we review current knowledge of the lithic microbial ecosystems of Antarctica's Dry Valleys with special attention paid to their microhabitats and specific adaptations to these habitats.

The lithobiontic ecological niche

Microorganisms and lichen associations in Antarctica appear both on rock surfaces (epilithic forms) and hidden inside rocks (endolithic forms). Antarctic endolithic microorganisms adapt to different ecological niches within the lithic substrate. They can be found living beneath rock surfaces as interstitial habitats of cracks and fissures of weathered siliceous rocks (chasmoendolithic, arrow in Fig. 4a) but also in pore spaces between mineral grains of porous rocks (cryptoendolithic, arrows in Fig. 4b) (Golubic *et al.* 1981). The type of colonization is determined mainly by the micromorphological, structural, physical and mineralogical properties of the rock substrate.

The development of the “(SEM-BSE) technique” (Wierzchos & Ascaso 1994), which enabled the study of the lithobiontic community within its rock microhabitat, has permitted the characterization of the different ecological niches and the spatial organization of lithic microbial ecosystems (Wierzchos & Ascaso 2001, Ascaso & Wierzchos 2003, De los Ríos *et al.* 2003, 2005a, 2005b, Wierzchos *et al.* 2003, 2005). Chasmoendolithic and cryptoendolithic communities share some features. In both, two main classes of community are defined: eukaryotic communities dominated by lichens and algal components, and prokaryotic cyanobacteria-dominated communities (Friedmann *et al.* 1988, De la Torre *et al.* 2003). Both classes can be found in close proximity but they occupy different zones within the rock: fissures or cracks for chasmoendolithic communities (Broady 1981, De los Ríos *et al.* 2005b, 2007) and colonization of pore spaces between mineral grains in the case of cryptoendolithic communities (Friedmann 1982). Chasmoendolithic growth forms have been reported in granite, marble, silicified sandstone, gypsum crusts and anorthosite (Broady 1981, Friedmann 1982, Ascaso & Wierzchos 2003, De los Ríos *et al.* 2005a, 2007, Guglielmin *et al.* 2005, Büdel *et al.* 2008), while cryptoendolithic microorganisms have been detected mainly in porous sandstones (Friedmann 1977, 1982, Friedmann *et al.* 1988, Siebert *et al.* 1996, Wierzchos & Ascaso 2001, De la Torre *et al.* 2003, Blackhurst *et al.* 2005, Selbmann *et al.* 2005, Wierzchos *et al.* 2005, Pointing *et al.* 2009).

The first endolithic Antarctic communities described were cryptoendolithic microorganisms inhabiting sandstones (Friedmann & Ocampo 1976, Friedmann 1982, Friedmann *et al.* 1988). Shortly afterwards, Ascaso & Wierzchos (2002) and Wierzchos & Ascaso (2002) reported the presence of microbial fossils within these rocks. Cryptoendolithic microbial colonization normally occurs as different coloured bands running parallel to the rock surface (Friedmann 1982). These bands are composed of different groups of microorganisms at varying depths. Layers of photosynthetic microorganisms (Fig. 5) alternate with layers dominated by fungi (Fig. 6a & b). Typically from the surface to greater rock depths, the pattern consists of a black zone under the crust followed by a white zone, a green and sometimes a blue-green zone (Fig. 3b) (Friedmann 1982, Ruisi *et al.* 2007). Although the sequence of black, white and green is generally constant, any one or two of these zones can be absent (Friedmann 1982). The first black layer exclusively contains melanized rock fungi (Fig. 6a), the white layer harbours cryptoendolithic lichen associations (Figs 5b & 6b); and the green layer (Fig. 6c) contains several species of non-lichenized chlorophycean algae, such as *Hemichloris antarctica* Tschermak-Woess & Friedmann, coexisting with cyanobacteria of the genera *Choroococcidiopsis* and *Gloeocapsa* (Friedmann *et al.* 1988). Finally, a layer of cyanobacterial colonization may be observed in some rock samples (Figs 5d & 6d). In addition, melanized rock fungi can appear in the different layers (arrows in Fig. 6b, c & d).

The fissures and cracks appearing in weathered siliceous rocks can also feature extensive chasmoendolithic microbial growth in the Dry Valleys (Broady 1981, Ascaso & Wierzchos 2003, De los Ríos *et al.* 2005a, 2005b, 2007). Rock freeze-fracturing could favour the formation of chasmoendolithic microhabitats (Cowan & Tow 2004). As with the

cryptoendolithic communities, some zonation is observed for chasmoendoliths. Certain fissures are occupied by cyanobacterial colonies frequently associated with heterotrophic bacteria (Fig. 7a), while others harbour algal-fungal lichen associations (Fig. 7b). Free-living fungi may also occur (Fig. 7c), generally close to areas colonized by lichen associations. Fissures only occupied by non-lichenized algae have also been detected in some areas (Fig. 7d). Lichen-dominated chasmoendolithic growth to some extent follows the lichen thallus structure. In fissure zones close to the surface, algal and fungal symbionts intermix to form something similar to an algal layer (Fig. 7b), while only mycobiont cells inhabit deeper areas, as in the medulla of a lichen thallus (Fig. 7e). Chasmoendolithic microbial communities in the Dry Valleys are generally observed in areas lacking epilithic growth but may also be detected under epilithic lichen thallus (arrow in Fig. 7f).

The zonation shown by cryptoendolithic microbial communities has been correlated with the steep light gradient that occurs in the top few millimetres of the rock (Nienow *et al.* 1988). The occurrence of cyanobacteria in the lowermost zone may reflect their ability to grow at lower light levels and in microaerophilic conditions, whereas green algae in the outer zones may require higher light and oxygen levels (Broady 1981). Chasmoendolithic colonization by algae and cyanobacteria is also determined by the existence of sufficient light for a community to become established. Most extensive colonization by cyanobacteria has been described in close-to-surface areas of a fissure (De los Ríos *et al.* 2007). The algae of chasmoendolithic lichens also show a preference for zones that are closer to the surface and are frequently observed under the more translucent minerals (De los Ríos *et al.* 2005a).

Besides epilithic and endolithic colonization of Antarctica's terrestrial environments, hypolithic colonization also exists as microbial assemblages present on the undersides of translucent rocks (Cowan 2009, Pointing *et al.* 2009, Cowan *et al.* 2011). While this zone acts as a refuge for microorganisms from extreme environmental conditions (Cowan *et al.* 2010) and a high diversity of microbial communities have been reported (Chan *et al.* 2012, 2013, Khan *et al.* 2011), their description is outside the scope of this review.

Features of the Antarctic Dry Valley microhabitat

Microhabitat conditions are an important determining factor for the establishment of lithic microbial ecosystems (Friedmann 1980) and are generally less extreme than external environmental conditions. We review the main features of the Antarctic lithic microhabitat below.

Water availability

Water is essential for biological processes and is, hence, a requirement for microbial life. In Antarctic lithic microbial ecosystems, water availability varies according to geographical and local topographical factors. In Maritime Antarctic, although conditions are milder (Green *et al.* 2007) and liquid water is available for microbial and lichen growth. However, liquid water availability is one of the most limiting factors for the distribution and abundance of terrestrial organisms in the Dry Valleys (Doran *et al.* 2002a, Ruisi *et al.* 2007). These differences in water availability are reflected in lichen growth rates, which are close to two orders of magnitude lower in the Dry Valleys than on the Antarctic Peninsula (Sancho *et al.* 2007). The presence of liquid water in the Dry Valleys is limited to transient ice or snow melt driven by solar heating when rock temperatures are near or above 0°C (summer) (Friedmann 1978, Broady 1981, Doran *et al.* 2002b). Humidity within the rock can be two-fold that detected in the atmospheric air (Green *et al.* 2007). Water can enter the endolithic habitat via percolation into subsurface pore spaces or through fissures or weathered areas (Friedmann 1978, McKay & Friedmann 1985) where evaporation is limited due to reduced gas exchange between the air and subsurface, especially when rock varnish is present (Friedmann & Ocampo-Friedmann 1984). Hence, the existence of points of snow or ice accumulation has a positive effect on the development of lithobiontic communities (Broady 1981, Winkler *et al.* 2000) (Fig. 2a). In addition, the physical structure and water-adsorbing capacity of the lithic substrate helps to retain moisture following precipitation events (Kappen *et al.* 1981,

Friedmann *et al.* 1987, 1988). These properties confer the endolithic microhabitat advantages over other habitats in the Dry Valleys, where water is inherently scarce (Omelson 2008). For example, Büdel *et al.* (2008) observed that chasmoendolithic cyanobacteria could use available moisture from dew/rime deposition on rocks.

Temperature

Low temperatures and temperature fluctuations are among the most decisive factors for the existence of lithobiontic microbial communities in Antarctica because they can arrest metabolic activity (Friedmann & Weed 1987). Specific climatic conditions, i.e. sunny days with a gentle breeze, generate frequent temperature oscillations (within few minutes), while low-frequency oscillations occur (McKay & Friedmann 1985). Rapid temperature fluctuations due to gusty winds are common in the Dry Valleys (Friedmann 1982). These conditions give rise to freeze-thaw cycles that can severely impact microorganisms that do not have specific anti-freeze mechanisms (Nienow *et al.* 1988). Depending on topographical factors, such as aspect and altitude, mean annual air temperatures in the McMurdo Dry Valleys range from -15–30°C, with differences between the absolute maximum and minimum of *c.* 50°C (Doran *et al.* 2002a, 2002b). Field microclimate measurements have revealed that the rock surface and shallow subsurface are considerably warmer, up to 20°C higher, than the ambient air during sunny periods at high altitudes. These conditions are attributed to solar radiative heating of the rock surface, and the heat capacity and transparency of the rock (Friedmann 1977, Kappen *et al.* 1981, McKay & Friedmann 1985, Green *et al.* 2007). Furthermore, the heat absorption capacity of the rock dampens rapid temperature oscillations, especially important when temperatures are *c.* 0°C (McKay & Friedmann 1985).

Light

Light is a vital factor for the development and functioning of lithic microbial ecosystems because it drives autotrophic processes. In the Dry Valleys, mean annual solar flux ranges from 73–117 W m⁻² and photosynthetic active radiation (PAR) ranges from 170–238 $\mu\text{M s}^{-1}$ in relation to topographical effects (Doran *et al.* 2002b). Rock surfaces and narrow or deep valleys that face north receive more sunlight and are, therefore, more frequently colonized by endolithic microbial communities than south facing surfaces (Friedmann 1977, McKay & Friedmann 1985).

Light attenuation patterns have been linked to the limited distribution depth of algae and cyanobacteria within colonized rocks (Matthes *et al.* 2001). Given the reduced light that reaches microbial ecosystems in the endolithic habitat, the amount of insolation on rock surfaces becomes a more relevant factor for the distribution of endolithic communities. Using mathematical methods, Nienow *et al.* (1988) showed that within the endolithic sandstone habitat light was diminished by up to 90% per millimetre of depth and that, in addition to the physicochemical features of the rock, light attenuation depended on the presence of water in the rock and/or its microbial colonization. Greater light penetration was observed through a hydrated rock but light penetration was markedly reduced if the rock was colonized by epilithic and endolithic microorganisms.

Microbial photosynthetic activity is dependent on a sufficient amount of PAR reaching the microbial community. However, an excess of radiation can also be detrimental; this is especially true in Antarctica where ultra violet (UV-A and UV-B) transmission to the lower atmosphere and surface is substantially higher than elsewhere in the world as a result of a depleted ozone layer (Madronich *et al.* 1998). Incoming solar UV radiation is harmful to microorganisms and is, therefore, an important factor controlling the surface development of biological colonizers (Caldwell *et al.* 1998, Hughes *et al.* 2003, Tosi *et al.* 2005, Hughes 2006). To protect themselves against excessive radiation and UV-B rays, Antarctic microbiota synthesize UV-photoprotective pigments (such as scytonemin or mycosporine-like amino acids) and photoprotective-quenching pigments (e.g. carotenoids), and also have special DNA repair mechanisms (Ruisi *et al.* 2007, Selbmann *et al.* 2011, Singh *et al.* 2011). A number of Antarctic endolithic communities take advantage of geoprotective screens, such as iron oxide (haematite), goethite or calcium sulphate (gypsum), that protect them from potential lethal radiation (Villar *et al.* 2005).

Other factors

Although temperature, radiation and water availability critically influence the composition and survival of microbial communities, other factors such as wind, nutrient availability, salinity and the physicochemical properties of local colonized rock, which can vary at the microscale level, are also important.

Wind has both direct and indirect effects on the distribution of lithobiontic communities. While summer temperatures are strongly controlled by coastal winds, warming as they move inland and cooling with inland elevation, sporadic katabatic windstorms have a pronounced effect on non-summer temperatures (Doran *et al.* 2002b). Broady (1981) noted that snowdrift from the continental ice plateau is a source of moisture upon melting, promoting the survival of microbial colonizers. However, high winds enhance evaporation, making lithobiontic colonization especially difficult at or close to the rock surface (Ruisi *et al.* 2007). In addition, erosion due to a sandblasting effect induced by wind can prevent microbial colonization, as shown by Broady (1981) in the Vestfold Hills area. Finally, interactions between topography and air masses can greatly influence climate in microhabitats (Doran *et al.* 2002b) and subsequently the possibility of lithobiontic colonization.

Nutrients are scarce in Antarctic lithic habitats. Birds and other animals are a source of nutrients, especially nitrogen and phosphorous, in coastal areas (Green *et al.* 2007) but this supply is minimal in other regions. In the endolithic habitat, nutrients may be obtained from *in situ*-weathered bedrock or allochthonous sediments, such as dust (Johnston & Vestal 1989, Omelon *et al.* 2007, Walker & Pace 2007), and thereby depend on the local physicochemical properties of the colonized rocks. However, the nutrient requirements of slow-growing communities cannot be too great and there seem to be no signs of nutrient limitation for cryptoendolithic photosynthetic metabolism (Johnston & Vestal 1991).

In Antarctica, high evaporation rates often lead to high concentrations of salts on certain rocks and soil surfaces (Nishiyama 1977). Furthermore, strong winds can pick up salt spray from the sea and saline lakes and deposit this salt on downwind land areas (Broady 1981). Salinity may be a stress factor for microbial colonization and determine community structure (Broady 1981, Ruisi *et al.* 2007).

Diversity of lithic microbiota

Epilithic communities

The colonization of rock surface in the Dry Valleys is dominated by lichens. Although lichens are among the most diverse organisms inhabiting Antarctica, the magnitude of this diversity is still unclear mainly due to difficulties in identification related to the presence of numerous sterile forms and extensively modified thalli, including those of endolithic growth forms (Kappen 2004, Pérez-Ortega *et al.* 2012). Øvstedal & Lewis-Smith (2011) have estimated the number of lichen species in Antarctica and South Georgia at *c.* 500. Several taxonomic (Hertel 1987, 1988, Ruprecht *et al.* 2010) and floristic (Castello *et al.* 1995, Øvstedal & Lewis-Smith 2001, 2009, 2011, Castello 2010, Seppelt *et al.* 2010, Green *et al.* 2011a) studies exist, but DNA-based diversity estimates including modified forms have only been undertaken recently for three McMurdo ice-free valleys: Garwood, Marshall and Miers (Pérez-Ortega *et al.* 2012). Higher lichen diversity estimates are obtained when morphological and molecular characters are used for identification. In Continental Antarctica, a much lower diversity of lichens has been reported than in the Antarctic Peninsula region (Green *et al.* 2007, Armstrong 2010). Lichen communities, composed of foliose and crustaceous lichens, are practically the only types of epilithic community in Continental Antarctica. A clear diversity gradient exists in the Antarctic Peninsula, with a strong decline in species richness, albeit in steps, from 62°S to around 70°S (Peat *et al.* 2007). However, despite a lack of evidence for a major drop in lichen diversity with increasing latitude in Victoria Land and the Transantarctic Mountains, lichen diversity hotspots have been discovered in these areas (Green *et al.* 2007, 2011a, Peat *et al.* 2007, Seppelt *et al.* 2010). These observations suggest that the distribution of lichens in Continental Antarctica could be more highly influenced by microclimatic features, such as the concurrent presence of warmth,

water, light and shelter in a microhabitat, and historical events than by climate effects (Schroeter *et al.* 2010, Green *et al.* 2011b).

To examine the diversity of lithic microbial ecosystems dominated by lichens, as well as knowledge of mycobiont diversity (determining lichen nomenclature), the diversity of the photobionts forming lichen associations needs to be known. A good understanding of the diversity of photobionts and specificity of their associations acquires special relevance in extreme environments, such as Antarctica, where the availability of algae could be a limiting factor for lichen-dominated microbial communities (De los Ríos *et al.* 2005a). In a recent study, Pérez-Ortega *et al.* (2012) found that photobiont diversity in three of the McMurdo Dry Valleys was not much lower than in non-Antarctic regions. Five putative species of lichen photobionts were associated with 27 species of lichen-forming fungi in the three valleys examined. There was a general trend of low selectivity but a number of species showed high selectivity. Other studies have also shown low selectivity of the photobiont for the mycobiont in Maritime Antarctica (Romeike *et al.* 2002, Wirzt *et al.* 2003) and in lecideoid lichens in Continental and Maritime Antarctic (Ruprecht *et al.* 2012).

Endolithic communities

Before the introduction of molecular biology methods, the diversity of lithobiontic communities was described based on cultures and microscopic observations of morphological and ultrastructural features. Using these methods, Friedman *et al.* (1988) distinguished and characterized the diversity of the different classes of cryptoendolithic community inhabiting the Ross Desert: cyanobacteria-dominated communities and two types of eukaryotic communities, lichen-dominated and *Hemichloris* communities.

Cyanobacteria of the genera *Gloeocapsa* found under quartz-like rocks in southern Victoria Land were the first primary producers described in Antarctica's cold desert ecosystems (Friedmann & Ocampo-Friedmann 1976). Since then, different cyanobacteria-dominated communities have been described within cryptoendolithic and chasmoendolithic ecological niches. In total, 17 cryptoendolithic cyanobacteria from the McMurdo Dry Valleys were described in cyanobacteria-dominated communities from Beacon Supergroup sandstone (Friedman *et al.* 1988). Depending on the dominating cyanobacterium, three communities were defined: *Gloeocapsa*, *Hormathonema-Gloeocapsa* and *Chroococcidiopsis*. In Taylor Valley, cyanobacteria-dominated communities were composed of the cyanobacterium *Chroococcidiopsis* sp., with the occasional presence of *Cyanothece* cf. *aeruginosa* (Nägeli) Komárek and *Nostoc* sp. (Büdel *et al.* 2008). In the Granite Harbour area, two different chasmoendolithic cyanobacteria-dominated communities have been described (De los Ríos *et al.* 2007). One of them, loosely adhered to the substrate, formed by a cyanobacteria closely related to *Acaryochloris marina* Miyashita et Chihara (the only organism known to contain chlorophyll *d*). The second community, more closely associated with the rock minerals, harbours *Gloeocapsa*-like cells.

The non-lichenized algae most commonly detected in the cryptoendolithic niches of the McMurdo Dry Valleys belong to the genus *Hemichloris* (Friedmann & Ocampo-Friedmann 1984, Tschermak-Woess & Friedmann 1984). In addition, several black fungi have been isolated from cryptoendolithic lichen-dominated communities living within sandstones in the McMurdo Dry Valleys (Friedmann & Ocampo-Friedmann 1976, Friedmann 1982, Nienow & Friedmann 1993, Onofri *et al.* 2007a, Selbmann *et al.* 2008). These fungi form a heterogeneous, taxonomic, phylogenetic group characterized by thick, heavily-pigmented cell walls, apparent lack of reproductive structures and slow growth. Members of this group show different growth styles, such as the formation of daughter cells by yeast-like multilateral or polar budding (black yeasts) or the formation of aggregates of thick-walled, melanized cells, which enlarge and reproduce by isodiametric division (meristematic black fungi) (Ruibal *et al.* 2009). The biodiversity of Antarctic black fungi has only recently been addressed and although sampling is still limited, several new genera have been described: *Friedmanniomyces*, *Cryomyces*, *Acidomyces*, *Recurvomyces* and *Elasticomyces* within the class Dothideomycetes (Selbmann *et al.* 2005, 2008). With the exception of the genus *Cryomyces* considered to be Dothideomycetes incertae

sedis (Ruibal *et al.* 2009), all these genera have been ascribed to the order Capnodiales (Selbmann *et al.* 2008, Ruibal *et al.* 2009). Among the filamentous hyphomycetes, the other ecological group of rock-inhabiting fungi represented in Antarctica, a strain of *Verticillium* sp. has been identified in a microbial endolithic community inhabiting gypsum crusts at Two Step Cliffs on Alexander Island (Hughes & Lawley 2003).

De la Torre *et al.* (2003) conducted the first rRNA-based molecular survey of the two main classes of endolithic communities (lichen-dominated and cyanobacteria-dominated) described by Friedmann *et al.* (1998) in Beacon sandstone. In both classes, a dominant fungal and algal phylotype was detected. The fungal phylotype was closely related to the epilithic *Buellia* spp. and the algal phylotype was *Trebouxia jamesii* (Hildreth & Ahmadjian) Gärtner. The most commonly detected cyanobacterium was closely related to *Plectonema* spp. In cyanobacteria-dominated communities, many sequences of the Thermus-Deinococcus bacterial phylogenetic group and a representative of the α -proteobacteria have also been detected. Pointing *et al.* (2009) reported that endolithic communities in the McKelvey Valley were dominated by different cyanobacteria and showed a great complexity including 16 phyla. *Chroococcidiopsis* and Nostocales phylotypes were the most common endoliths. An additional source of bacterial phototrophy was indicated by the presence of Chloroflexi. A lower phylogenetic diversity was detected for chlorophytes with two closely related groups within the Trebouxiophyceae and a single *Brateococcus* phylotype.

Physiological peculiarities of microbiota in Antarctica

Antarctic microorganisms generally have specific physiological adaptations in order to cope with the harsh environmental conditions, such as metabolic activity at subzero temperatures, the ability to persist in metabolically inactive states under unfavourable conditions and the presence of cell protection strategies (e.g. photoprotective mechanisms and extracellular polymeric substances (EPS) production). These mechanisms are probably not driven by the extreme climate conditions per se, but by the extreme range of conditions encountered (Friedmann 1993, Yergeau *et al.* 2007, Omelon 2008). In endolithic communities, physiological adaptations act in combination with the inherent protective properties of the host rock microenvironment, leading to successful colonization.

Microbial growth and metabolism are clearly limited at low temperature. Cold-adapted microorganisms include both psychrophilic and psychrotrophic organisms. In the lithic habitat, psychrotrophic microorganisms are probably more common than psychrophilic forms. Psychrophiles include organisms whose optimum growth temperatures are < 15°C and upper limits are < 20°C, while the optimum and maximum growth temperatures of psychrotrophs are > 15°C (although they are still capable of growing at or close to zero) (Morita 1975). Maximum photosynthetic rates of 10°C suggest a psychrophilic temperature response for photosynthesis in cryptoendolithic lichen communities, which contrasts with psychrotrophic cryptoendolithic cyanobacterial communities that achieve maximum photosynthetic rates at 20–30°C (Johnston & Vestal 1991). Only a few Antarctic cyanobacteria can be considered true psychrophiles and these have been mostly associated with the polar freshwater ecosystem (Pandey 2004). Siebert *et al.* (1996) observed psychrophilic and psychrotrophic cryptoendolithic bacteria in cultures of sandstone samples obtained from Linneus Terrace. Antarctic black fungi have also been described as psychrotolerant. These fungi are able to grow at *c.* 0°C but optimal growth is observed at much higher temperatures than those to which they are generally exposed. This enables them to tolerate daily and seasonal temperature fluctuations (Selbmann *et al.* 2005).

Effective strategies against freezing temperatures and desiccation (Selbmann *et al.* 2005, 2011, Kranner *et al.* 2008, Billi 2009) enable microorganisms to persist in Antarctic terrestrial environments but the *in situ* primary production is very limited, including in the abundant cryptoendolithic microbial communities (Johnston & Vestal 1991, Hopkins *et al.* 2009). Most microorganisms exist in a metabolic dry and/or frozen state for most of the year and are only active during the short summer period, when temperatures rise and water from snow or ice melt is available (Friedmann 1982, Selbmann *et al.* 2005, Schroeter *et al.* 2011, Sterflinger *et al.* 2012).

Lichens are able to survive under extreme environmental conditions that are adverse to most other organisms owing to their anhydrobiotic behaviour. Lichens are even able to recover their metabolic activity after exposure to space conditions (Sancho *et al.* 2007, Raggio *et al.* 2011). Some lichens show positive net photosynthesis at subzero temperatures of -5–20°C (Lange 1965, Kappen & Redon 1987, Schroeter *et al.* 1994, Bártak *et al.* 2007). However, Antarctic lichens seem to feature lower net rates of photosynthesis than lichens in other parts of the world (Sancho *et al.* 2000, Green *et al.* 2007). Low molecular weight antioxidants, such as glutathione, and reactive oxygen species (ROS)-processing enzymes and peroxidases have been described as key components in the desiccated state and during subsequent rehydration to avoid the effects of oxidative stress (Kranner *et al.* 2008). In response to high light intensities, photoprotective mechanisms via non-photochemical energy dissipation, synthesis of antioxidants and production of light-screening compounds by mycobiont hyphae forming the upper cortex have also been described in Antarctic lichens (Kappen *et al.* 1998, Edwards *et al.* 2004, Bártak *et al.* 2012). Photobionts and mycobionts feature different mechanisms and those facilitating tolerance are more effective in the whole lichen than in the separate symbionts (Kranner *et al.* 2005). Additionally, Friedmann & Sun (2005) observed that the contribution of community adaptation to heat was greater than the mechanisms functioning at the organism level.

Meristematic black fungi that make up some cryptoendolithic communities are not only able to tolerate high levels of stress, but are even specialized to do so. The simple morphology, scarcely differentiated structure and reduced life cycle make these fungi particularly suited to live in the harsh environment of the McMurdo Dry Valleys. These fungi can complete their life cycle during the short period of suitable conditions (Selbmann *et al.* 2005). Thick-walled and intensely melanized cells produce UV-B absorbing substances, such as mycosporine-like amino acids (Kogej *et al.* 2006), conferring these fungi drought- and UV-resistance and thus protecting the cell DNA (Selbmann *et al.* 2011).

The extremotolerance exhibited by cyanobacteria found in Antarctic lithobiontic communities, such as *Chroococcidiopsis* and *Nostoc*, has also been attributed to protection mechanisms and efficient repair systems (Shirkey *et al.* 2003, Billi 2009, 2012). It has been proposed that EPS production is a crucial factor for the stress tolerance of cyanobacteria to photosynthesis during desiccation and freezing-thawing cycles (Grilli Caiola *et al.* 1996, De los Ríos *et al.* 2003, Tamaru *et al.* 2005). In these anhydrobiotic cyanobacteria genera, EPS could act in synergy with trehalose accumulation in the cytoplasm (Billi 2012). The mechanisms causing the death of a few *Chroococcidiopsis* cells within a dried aggregate are thought to contribute to the survival of the remaining cells, by providing physical protection and/or nutrient supply upon rehydration (Billi 2009). Cyanobacterial mechanisms including light-harvesting pigments, light-screening pigments, ROS-quenching compounds (such as carotenoids), membrane fluidity at low temperatures and cold-stable proteins allow for slow yet steady growth in Antarctica (Vincent 2007, Banerjee & Verma 2009). Cryptoendolithic cyanobacteria within Antarctic rocks obtain nitrogen through overnight nitrogen fixation using energy generated by daytime photosynthesis or heterotrophic fixation using lichen osmolytes (Banerjee & Verma 2009).

The presence of EPS is a common feature of lithobiontic communities (De los Ríos *et al.* 2005b). In lithic microbial ecosystems, microorganisms and mineral components are embedded in an organic matrix resulting from the excretion of polymers by resident microorganisms, leading to formation of a protective biofilm (De los Ríos *et al.* 2003). The low-temperature scanning electron microscopy (LTSEM) image in Fig. 8a shows endolithic cyanobacterial cells covered by an EPS matrix. These EPS often confers a compact structure to the community (Fig. 8b) and can play an important role in adapting to Antarctic conditions (De los Ríos *et al.* 2005b). As with other types of biofilm, EPS could have structural and protective functions in these lithic microbial communities, such as helping to maintain structure, acting as a protective barrier, and facilitating water retention and the sorption of organic compounds and ions (Davey & O'Toole 2000, De los Ríos *et al.* 2003, Omelon 2008). At the physiological level, the EPS matrix could facilitate the photosynthesis of endolithic cyanobacteria. Decho *et al.* (2003) determined that high EPS concentrations enable

sediment-associated, densely-packed photosynthetic cells to trap photons more efficiently. The EPS matrix has also been considered to be important for long-term accumulation of carbon and energy (Wolfaardt *et al.* 1999), and may act as a cryoprotectant under cold temperatures (De los Ríos *et al.* 2004).

Functioning of lithic microbial ecosystems

Microorganisms in rock habitats are integrated within a complex ecological entity, or ecosystem. The dynamics of this integration are driven by multiple relationships involving not only the biological components but also the immediate environment, or lithic substrate, in which they are found. Ecosystem dynamics are driven by independent controls, such as climate, host rock, topography, potential biota and time, and interactive controls which directly affect the activity of the ecosystem and are controlled by feedback (Chapin *et al.* 1997). Dynamics include microclimate, nutrient supply, activity of the lithobiontic community and human activity. In Antarctic environments, the latter is minimal.

Lithobiontic communities are made up of microbial consortia in which several microorganism species make physical, and probably biochemical, contact with one another. In Antarctica, the lithic microbial ecosystem is simple. There are no higher predators, only an autochthonous microflora comprised of cyanobacteria and lichenized algae, playing a small but crucial role in primary production, with mycobionts and non-lichenized fungi as consumers and non-photosynthetic bacteria as decomposers (Friedmann 1982, De la Torre *et al.* 2003, Selbmann *et al.* 2008, Pointing *et al.* 2009). The presence of heterotrophic microorganisms points to the existence of functional trophic structures. Given that the exogenous input of organic carbon is almost negligible, the nutrient supply for consumers and decomposers is limited to products released by the photosynthetic primary producers, or to the organic matter available after cell lysis or death (Siebert & Hirsch 1988, De los Ríos *et al.* 2007, Billi 2009). Microscopy has shown close physical associations between the different types of microorganisms (Fig. 8c) indicating possible nutritional relationships (De los Ríos *et al.* 2004, 2007, Pointing *et al.* 2009). Heterotrophic bacteria in Antarctic endolithic communities are frequently detected in close association with photosynthetic microorganisms, both cyanobacteria (Fig 8d) and algae (Fig. 8c). Fungal and algal cells forming endolithic lichen associations show close interactions, resulting from symbiont behaviour (Fig. 7b) (De los Ríos *et al.* 2005a), but mycobiont and photobiont cells also establish close relationships with black pigmented free-living fungi in cryptoendolithic communities (Fig. 6b) (Friedmann 1982, Onofri *et al.* 2007b). Microbial interactions have also been observed at the biochemical level. For example, *Micrococcus* and *Deinococcus* isolated from Antarctic rocks of the Dry Valleys showed a preference for sugar alcohols and organic acids, compounds commonly released by lichens, moulds and algae, as the remaining components of cryptoendolithic ecosystems (Siebert & Hirsch 1988). The beneficial use of spent algal medium, or the co-culture with algae, when culturing heterotrophic endolithic bacteria also point to possible nutritional interactions such as the exchange of vitamins and other resources; these interactions could be important for all community populations (Siebert *et al.* 1996).

Not all the biotic components of lithic microbial ecosystems contribute equally to ecosystem processes. In Antarctic lithobiontic communities, some microorganisms probably only play a potential role within the community (active yet non-productive), others have a functional role and contribute to the production of biomass (active and productive), and the remaining microorganisms only represent particulate organic matter (moribund and dead) (Vestal 1988, De los Ríos *et al.* 2004, Wierzbos *et al.* 2004). Fluorescence microscopy (Fig. 8e) shows the coexistence of cells in different physiological states in Antarctic endolithic communities. Transmission electron microscopy (TEM) reveals living cells with a healthy ultrastructural appearance (arrow in Fig. 8f) associated with cells that have lost their integrity (arrowhead in Fig. 8f) and cellular remains (asterisk in Fig. 8f). The abundant presence of moribund cells is notorious in Antarctic lithobiontic communities; these cells retain their morphological integrity longer than in other environments (Vestal 1988, De los Ríos *et al.* 2004). This physiological heterogeneity and the high proportion of moribund or decayed cells may confer these isolated communities an ecological advantage. Leakage and lysis products from decaying

microorganisms, which break down all their major polymer constituents (except lipids and DNA), support the activity of the survivors (Roszak & Colwell 1987, Billi 2009). The close relationship observed among microorganisms in different physiological states in lithobiontic communities (Fig. 8f) also their contribution to recycling processes. Collectively, these findings suggest that microbial viability in Antarctic lithic ecosystems works more at the community than individual level.

The long-term survival of Antarctic lithobiontic communities has been linked to microbial interactions on the lithic substrate and their effects (Johnston & Vestal 1993). An intimate relationship between the rock substrate and microorganisms gives rise to complex mineral–microorganism interactions, which result in physical and chemical alterations to the substrate. Epilithic colonizers and their actions can promote the formation of cracks in the substrate (Fig. 7f); in turn, these cracks provide a microhabitat for chasmoendolithic microorganisms (Ascaso *et al.* 1990, Sun & Friedmann 1999, Ascaso & Wierzchos 2002, Wierzchos *et al.* 2003, De los Ríos *et al.* 2005b). Endolithic microorganisms, especially fungi, also interact with subsurface zones altering the rock (Fig. 9a). Mechanical actions are the main force behind these microhabitat modifications, but in most cases chemical actions also play a role. Antarctic endolithic microorganisms show evidence of calcium and potassium biomobilization processes (Ascaso & Wierzchos 2003, De los Ríos *et al.* 2003, 2005a). In epilithic and endolithic lichens, calcium oxalate accumulation as a result of a biomineralization is also a common phenomenon (Fig. 9b) (Ascaso *et al.* 1990, Johnston & Vestal 1993, Ascaso & Wierzchos 2003, De los Ríos *et al.* 2005a, 2005b). The build-up of calcium oxalate may lead to enhanced mineral dissolution, providing nutrients and increasing rock porosity, and ultimately may affect the functioning of the microbial community (Johnston & Vestal 1993, Blackhurst 2005). Calcium oxalate could also protect against solar radiation (Modenesi *et al.* 2000), especially in the presence of EPS. Most microorganism–mineral interactions in a lithic ecosystem occur within the EPS matrix; EPS chemically react with minerals and accelerate dissolution by up to several orders of magnitude. Thereby, EPS participates in the biomobilization of elements (Wierzchos & Ascaso 1996), biomineralization phenomena (Banfield *et al.* 1999, Barker *et al.* 1998) and clay mineral transformation processes (Wierzchos & Ascaso 1998). This effect is amplified by increases in the amount of reactive surface area produced through the physical action of microbial cells (Barker & Banfield 1996), such as the exfoliation of biotite and other clays layers (De los Ríos *et al.* 2003, 2007).

Past life and fossilization processes

The 1990s and early 2000s saw the description of indirect biomarkers of the past activity of Antarctic lithobionts (Kappen 1993, Sun & Friedmann 1999, Wierzchos & Ascaso 2001). Iron leaching bioweathering patterns observed on the surface layers of rocks were first interpreted as signs of previous microbial activity (Friedmann & Weed 1987). The biogeophysical weathering of rock surfaces forming characteristic mosaic exfoliation patterns indicating the previous presence and activity of microbial communities was also considered a biomarker (Kappen 1993, Sun & Friedmann 1999). The introduction of the SEM-BSE visualization strategy (Wierzchos & Ascaso 1994) made it possible to better characterize the mechanical and chemical changes to minerals induced by lithobiontic microorganisms that may give rise to physical or chemical traces of their previous presence along with microbial fossil formation processes (Wierzchos & Ascaso 2002, Ascaso & Wierzchos 2003, Wierzchos *et al.* 2005). Nanocrystals of diagenic iron hydroxide and biogenic clays transformed by the actions of chasmoendolithic microorganisms have been reported as biomarkers of prior chasmoendolithic life in the Granite Harbour area (Wierzchos *et al.* 2003). In other regions, physical bioweathering patterns have been proposed. These take the form of exfoliated, undulated biotite sheets and microdivided clay minerals, along with biogeochemical microbial action effects resulting in inorganic deposits such as calcium oxalate and silica (Ascaso & Wierzchos 2003).

Furthermore, there is direct evidence of past microbial life in Antarctic deserts in the form of mineralized structures recognizable as microorganism fossils. In extreme regions of Antarctica, microbial fossil traces formed in the past and are currently forming (Friedmann 1986). The survival of microbial cells in this setting depends upon a precarious balance among biological, geological and climate factors, and any change in the external conditions upsetting this balance will lead to their death. This may then be followed by trace microfossil formation (Friedmann 1993, Wierzchos & Ascaso 2001, Wierzchos *et al.* 2003). Progressive cell damage stages can be observed in Antarctic endolithic colonizers (De los Ríos *et al.* 2004, Wierzchos *et al.* 2005). These stages are the steps that precede the start of the fossilization process. Depending on the environmental conditions prevailing at the time of death, fossilization may preserve the remains of this organic material (Schopf 1993, Westall & Folk 2003) or lead to complete cell mineralization (Wierzchos & Ascaso 2002). The relicts of cell structures in these mineralized cells can be observed through the combined use of SEM-BSE and energy dispersive x-ray spectroscopy (EDS) (Wierzchos & Ascaso 2001). Careful interpretation of morphological features, such as preserved cell walls, cytoplasm elements and occasionally chloroplast membranes, will lead to the identification of mineralized cells as microfossils (Wierzchos & Ascaso 2002, Ascaso & Wierzchos 2003, Wierzchos *et al.* 2005). Presumably, in the presence of liquid water, organic matter is silicified generating coloured areas on the surface of the rock and colonized area (Friedmann & Weed 1987). Although a limited quantity of liquid water is sufficient, its presence is a prerequisite for the fossilization of microorganisms. Wierzchos & Ascaso (2002) proposed a scenario for microbial fossil formation in rocks of the Mount Fleming area. The first stage of microbial cell biomineralization occurs when microorganisms are biologically active and/or after their decay; pores containing microbial fossils become infiltrated with jarosite and gypsum. Wierzchos *et al.* (2005) later proposed that an organic matter template may guide the diffusion of chemical elements and give rise to the characteristic distribution pattern observed inside the fossilized cells. Mineralization leading to good preservation could be the result of the extensive biomobilization of elements occurring when the microorganisms are still biologically active or after their decay, and seems to be favoured by the extraordinary slow growth rate and the long lifespan of Antarctic cryptoendoliths (Wierzchos & Ascaso 2002). Apart from silicon, which mineralizes the core of algal cells, the presence of elements such as aluminium and iron has been detected in mineralized algae; aluminium usually occupies a more peripheral location in the cytoplasm whereas iron appears on the wall or close to the wall inside cells (Ascaso & Wierzchos 2002). It has not been clearly established if it is the microorganisms that induce the mineral precipitation associated with microbial fossilization processes. Although Wierzchos *et al.* (2005) consider that the organic template is the main factor provoking the initial diffusion of mineral elements into decaying cells, they propose that inorganic processes occurring after cell death are also key factors for microbial fossil formation.

Biogeography

Owing to the geographical and environmental features of Antarctica, its lithic microbial ecosystems are probably not consistent with the 'global ubiquity hypothesis' (Finlay 2002). This hypothesis proposes that microorganisms are ubiquitous and that it is the environment that selects out resistant and/or tolerant organisms. The existence of highly-specialized communities characterized by a low biodiversity of taxa tolerant to extreme cold and dry conditions, along with some extent of endemism in areas of the Dry Valleys, points to the uniqueness in Continental Antarctica of its microbial assemblages (de la Torre *et al.* 2003, Selbmann *et al.* 2005, Convey & Stevens 2007, Peat *et al.* 2007, Ruisi *et al.* 2007, Convey *et al.* 2008, De Weber *et al.* 2009, Pointing *et al.* 2009, Hodgson *et al.* 2010).

Isolation could be a structuring factor for the biogeography of lithobiontic microorganisms in Antarctica. In addition to the harsh environmental conditions, the geographical, oceanic and atmospheric isolation of Antarctica's terrestrial ecosystems will have hindered the dispersal, survival and establishment of lithobiontic microbial communities and promote dependance on stochastic processes (Convey 2011). The presence of good colonizers, defined as those adapted

for efficient dispersal with phenotypic flexibility and an ability to reproduce freely under suitable conditions (Block 1994), is an insufficient explanation for the current distribution of lithobiontic microorganisms, and their arrival would have had to coincide with the appropriate conditions. The dispersal of local species may be more efficient than long-distance dispersal determining the considerable influence of intense environmental selection and microbial adaptive strategies, especially for subsequent development and maintenance (Franzmann 1996, De Wever *et al.* 2009, Hodgson *et al.* 2010). In lithic microbial ecosystems, the physicochemical features of the lithic substrate, along with the contribution of lithobiont activity in the formation of proper microenvironments for settling, growth and development, may condition colonization success (De los Rios *et al.* 2002, Pearce *et al.* 2009). Hence, the establishment and development of the lithobiontic community could depend on the success of processes that operate over short time scales, such as dispersal, colonization and extinction, but this success will be, in turn, strongly affected by regional changes in geophysical and climatic conditions (Selbmann *et al.* 2005, Vyverman *et al.* 2010, Green *et al.* 2011a, 2011c).

Final remarks

Rocks as habitats across the planet have been scarcely explored, including those in extreme environments such as Antarctica's ice-free areas where lithic microbial ecosystems constitute most of the terrestrial life. Although our knowledge of these microbial ecosystems has expanded since early work on the lithobiontic communities of Antarctica's Dry Valleys led by Professor E.I. Friedmann, there is still much to be learnt. If we are to fill this void, novel ice-free areas of Antarctica will need to be explored in surveys using both conventional and molecular biology techniques.

Acknowledgements

The authors thank all those who kindly provided the Antarctic rock samples used in our investigations, especially Prof Sancho and Dr I.E. Friedmann, and A. Burton for reviewing the English. We also thank Drs Alan Green and Chris McKay for facilitating the collection of samples from the McMurdo Dry Valleys, and the investigators of the Antarctic New Zealand Program for their excellent scientific and logistic organization of the Dry Valley expeditions. We are very grateful to D.W. Walton for the invitation to write this review and two anonymous reviewers for their constructive comments. This work was supported by grant CTM2012-38222-CO2-02 and CGL2010-16004 from the Spanish Ministry of Economy and Competitiveness.

References

- ARMSTRONG, R. 2010. Growth of crustose lichens: a review. *Geografiska Annaler*, **92A**, 3–17.
- ASCASO, C., SANCHO, L. & RODRIGUEZ-PASCUAL, C. 1990. The weathering action of saxicolous lichens in maritime Antarctica. *Polar Biology*, **11**, 33–39.
- ASCASO, C. & WIERZCHOS, J. 2002. New approaches to the study of Antarctic lithobiontic microorganisms and their inorganic traces, and their application in the detection of life in Martian rocks. *International Microbiology*, **5**, 215–222.
- ASCASO, C. & WIERZCHOS, J. 2003. The search for biomarkers and microbial fossils in Antarctic rock microhabitats. *Geomicrobiology Journal*, **20**, 439–450.
- BANERJEE, M. & VERMA, V. 2009. Nitrogen fixation in endolithic cyanobacterial communities of the McMurdo Dry Valley, Antarctica. *Science Asia*, **35**, 215–219.
- BANFIELD, J.F., BARKER, W.W., WELCH, S.A. & TAUNTON, A. 1999. Biological impact on mineral dissolution: application of the lichen model to understand mineral weathering in the rhizosphere. *Proceedings of the National Academy of Science*, **96**, 3404–3411.
- BARKER, W.W. & BANFIELD, J.F. 1996. Biologically versus inorganically mediated weathering reactions: relationship

- between minerals and extracellular microbial polymers in lithobiontic communities. *Chemical Geology*, **132**, 55–69.
- BARKER, W.W., WELCH, S.A., CHU, S. & BANFIELD, J.F. 1998. Experimental observations of the effects of bacteria on aluminosilicate weathering. *American Mineralogist*, **83**, 1551–1563.
- BARTÁK, M., HÁJEK, J., & OČENÁŠOVÁ, P. 2012. Photoinhibition of photosynthesis in Antarctic lichen *Usnea antarctica*. I. Light intensity- and light duration-dependent changes in functioning of photosystem II. *Czech Polar Reports*, **2**, 42–51.
- BARTÁK, M., VACZI, P., HAJEK, J. & SMYKLA, J. 2007. Low temperature limitation of primary photosynthetic processes in Antarctic lichens *Umbilicaria antarctica* and *Xanthoria elegans*. *Polar Biology*, **31**, 47–51.
- BILLI, D. 2009. Subcellular integrities in *Chroococcidiopsis* sp. CCMEE 029 survivors after prolonged desiccation revealed by molecular probes and genome stability assays. *Extremophiles*, **13**, 49–57.
- BILLI, D. 2012. Anhydrobiotic rock-inhabiting cyanobacteria: potential for astrobiology and biotechnology. In STAN-LOTTER H. & SERGIU F., eds. *Adaptation of microbial life to environmental extremes*. Wien, NY: Springer, 119–232.
- BLACKHURST, R. 2005. Cryptoendolithic alteration of Antarctic sandstones: pioneers or opportunists? *Journal of Geophysical Research*, **110**, 10.1029/2005JE002463.
- BLOCK, W. 1994. Terrestrial ecosystems: Antarctica. *Polar Biology*, **14**, 293–300.
- BROADY, P.A. 1981. The ecology of chasmolithic algae at coastal locations of Antarctica. *Phycologia*, **20**, 259–272.
- BÜDEL, B., BENDIX, J., BICKER, F.R., & GREEN, T.G.A. 2008. Dewfall as a water source frequently activates the endolithic cyanobacterial communities in the granites of Taylor Valley, Antarctica. *Journal of Phycology*, **44**, 1415–1424.
- CALDWELL, M., BJÖRN, L., BORNMAN, J., FLINT, S., KULANDAIVELU, G., TERAMURA, A. & TEVINI, M. 1998. Effects of increased solar ultraviolet radiation on terrestrial ecosystems. *Journal of Photochemistry and Photobiology*, **46**, 40–52.
- CASTELLO, M. & NIMIS, P. 1995. A critical revision of Antarctic lichens described by C. W. Dodge. *Bibliotheca Lichenologica*, **57**, 71–92.
- CASTELLO, N. 2010. Notes on the lichen genus *Rhizoplaca* from Continental Antarctica and on some other species from Northern Victoria Land. *Lichenologist* **42**, 429–32.
- CHAN, Y., LACAP, D.C., LAU, M.C., HA, K.Y., WARREN-RHODES, K.A., COCKELL, C.S., MCKAY, C.P. & POINTING, S.B. 2012. Hypolithic microbial communities: between a rock and a hard place. *Environmental Microbiology*, **14**, 2272–2282.
- CHAN, Y., VAN NOSTRAND, J.D., ZHOU, J., POINTING, S.B., AND FARRELL, R.L. 2013. Functional ecology of an Antarctic Dry Valley. *Proceedings of the National Academy of Sciences*, **110**, 8990–8995.
- CHAPIN, F.S.I., WALKER, B.H., HOBBS, R.J., HOOPER, D.U., LAWTON, J.H., SALA, O.E. & TILMAN, D. 1997. Biotic control over the functioning of ecosystems. *Science*, **277**, 500–504.
- CONVEY, P. 2011. Antarctic terrestrial biodiversity in a changing world. *Polar Biology*, **34**, 1629–1641.
- CONVEY, P., GIBSON, J.A.E., HILLENBRAND, C.D., HODGSON, D.A., PUGH, P.J.A., SMELLIE, J. & STEVENS, M.I. 2008. Antarctic terrestrial life challenging the history of the frozen continent?. *Biological Reviews*, **83**, 103–117.
- CONVEY, P. & STEVENS, M.I. 2007. Ecology. Antarctic biodiversity. *Science*, **317**, 1877–1878.
- COWAN, D.A. 2009. Cryptic microbial communities in Antarctic deserts. *Proceedings of the National Academy of Sciences*, **106**, 19 749–19 750.
- COWAN, D.A., KHAN, N., POINTING, S.B. & CARY, C. 2010. Diverse hypolithic refuge communities in the McMurdo

- Dry Valleys. *Antarctic Science*, **22**, 714–720.
- COWAN, D.A. & TOW, L.A. 2004. Endangered Antarctic environments. *Annual Review of Microbiology*, **58**, 649–690.
- COWAN, DA, POINTING, S.B., STEVENS M.I., CRAIG CARY, S., STOMEIO, F. & TUFFIN, I.M. 2011a. Distribution and abiotic influences on hypolithic microbial communities in an Antarctic Dry Valley. *Polar Biology*, **34**: 307–311.
- DAVEY, M.E. & O'TOOLE, G.A. 2000. Microbial Biofilms: from ecology to molecular genetics. *Microbiology and Molecular Biology Reviews*, **64**, 847–867.
- DE LA TORRE, J., GOEBEL, B.M., FRIEDMANN, E.I. & PACE, N.R. 2003. Microbial diversity of cryptoendolithic communities from the McMurdo Dry Valleys, Antarctica. *Applied and Environmental Microbiology*, **69**, 3858–3867.
- DE LOS RÍOS, A., GRUBE, M., SANCHO, L.G. & ASCASO, C. 2007. Ultrastructural and genetic characteristics of endolithic cyanobacterial biofilms colonizing Antarctic granite rocks. *FEMS Microbiology Ecology*, **59**, 386–395.
- DE LOS RÍOS, A., SANCHO, L.G., GRUBE, M., WIERZCHOS, J. & ASCASO, C. 2005b. Endolithic growth of two *Lecidea* lichens in granite from continental Antarctica detected by molecular and microscopy techniques. *New Phytologist*, **165**, 181–190.
- DE LOS RÍOS, A., WIERZCHOS, J. & ASCASO, C. 2002. Microhabitats and chemical microenvironments under saxicolous lichens growing on granite. *Microbial Ecology*, **43**, 181–188.
- DE LOS RÍOS, A., WIERZCHOS, J., SANCHO, L.G. & ASCASO, C. 2003. Acid microenvironments in microbial biofilms of Antarctic endolithic microecosystems. *Environmental Microbiology*, **5**, 231–237.
- DE LOS RÍOS, A., WIERZCHOS, J., SANCHO, L.G., GREEN, A. & ASCASO, C. 2005a. Ecology of endolithic lichens colonizing granite in continental Antarctica. *The lichenologist*, **37**, 383–395.
- DE LOS RÍOS, A., WIERZCHOS, J., SANCHO, L.G. & ASCASO, C. 2004. Exploring the physiological state of continental Antarctic endolithic microorganisms by microscopy. *FEMS Microbiology Ecology*, **50**, 143–152.
- DE WEVER, A., LELIAERT, F., VERLEYEN, E., VANORMELINGEN, P., VAN DER GUCHT, K., HODGSON, D.A., SABBE, K. & VYVERMAN, W. 2009. Hidden levels of phylodiversity in Antarctic green algae: further evidence for the existence of glacial refugia. *Proceedings: Biological Sciences*, **276**, 3591–3599.
- DECHO, A.W., KAWAGUCHI, T., ALLISON, M.A., LOUCHARD, E.M., REID, P., STEPHENS, F.C., VOSS, K.J., WHEATCROFT, R. A. & TAYLOR, B.B. 2003. Sediment properties influencing up-welling spectral reflectance signatures: the biofilm gel effect. *Limnology Oceanography*, **48**, 431–443.
- DORAN, P.T., PRISCU, J.C., LYONS, W.B., WALSH, J.E., FOUNTAIN, A.G., MCKNIGHT, D.M., MOORHEAD, D.L., VIRGINIA, R.A., WALL, D.H., CLOW, G.D., FRITSEN, C.H., MCKAY, C.P. & PARSONS, A.N. 2002a. Antarctic climatic cooling terrestrial ecosystem response. *Nature*, **415**, 517–520.
- DORAN, P., MCKAY, C., CLOW, G., DANA, G., FOUNTAIN, A., NYLEN, T. & LYONS, W. 2002b. Valley floor climate observations from the McMurdo dry valleys, Antarctica, 1986–2000. *Journal of Geophysical Research*, **107**, 10.1029/2001JD002045.
- EDWARDS, H.G.M., WYNN-WILLIAMS, D.D. & VILLAR, S.E.J. 2004. Biological modification of haematite in Antarctic cryptoendolithic communities. *Journal of Raman Spectroscopy*, **35**, 470–474.
- FINLAY, B. 2002. Global dispersal of free-living microbial eukaryote species. *Science*, **296**, 1061–1063.
- FRANZMANN, P.D. 1996. Examination of Antarctic prokaryotic diversity through molecular comparisons. *Biodiversity and Conservation*, **5**, 1295–1305.
- FRIEDMANN, E.I. 1977. Microorganisms in Antarctic desert rocks from dry valleys and Dufek Massif. *Antarctic Journal of the United States*, **12**, 26–29.
- FRIEDMANN, E.I. 1978. Melting snow in the dry valleys is a source of water for endolithic microorganisms. *Antarctic*

Journal of the United States, **9**, 162–163.

- FRIEDMANN, E. I. 1980. Endolithic microbial life in hot and cold deserts. *Origins of life*, **10**, 223–235.
- FRIEDMANN, E.I. 1982. Endolithic microorganisms in the Antarctic cold desert. *Science*, **215**, 1045–1053.
- FRIEDMANN, E.I. 1986. The Antarctic cold desert and the search for traces of life on Mars. *Advances in Space Research*, **6**, 265–268.
- FRIEDMANN, E.I. & OCAMPO-FRIEDMANN, R. 1976. Endolithic blue-green algae in the dry valleys: primary producers in the Antarctic desert ecosystem. *Science*, **193**, 1247–1249.
- FRIEDMANN, E.I. & OCAMPO-FRIEDMANN R. 1984. The Antarctic cryptoendolithic ecosystem: relevance to exobiology. *Origins of Life and Evolution of the Biosphere* **14**, 771–776.
- FRIEDMANN, E.I. 1993. Extreme environments limits of adaptation and extinction. In GUERRERO R. & PEDRÓS-ALIÓ C., eds *Trends in microbial ecology*. Barcelona: Spanish Microbiology Society, 9–12.
- FRIEDMANN, E.I., HUA, M. & OCAMPO-FRIEDMANN, R. 1988. Cryptoendolithic lichen and cyanobacterial communities of the Ross Desert, Antarctica. *Polarforschung*, **58**, 251–259.
- FRIEDMANN, E.I., MCKAY, C.P. & NIENOW, J.A. 1987. The cryptoendolithic microbial environment in the Ross Desert, Antarctica: satellite-transmitted continuous nanoclimate data, 1984 to 1986. *Polar Biology*, **7**, 273–287.
- FRIEDMANN, E.I. & WEED, R. 1987. Microbial trace-fossil formation, biogenous and abiotic weathering in the Antarctic cold desert. *Science*, **236**, 703–705.
- FRIEDMANN, E.I. & SUN, H. J. 2005. Communities Adjust their Temperature Optima by Shifting Producer-to-Consumer Ratio, Shown in Lichens as Models: I. Hypothesis. *Microbial Ecology*, **49**, 523–527.
- GOLUBIC, S., FRIEDMANN, I. & SCHNEIDER, J. 1981. The lithobiontic ecological niche, with special reference to microorganisms. *Journal of Sedimentary Petrology*, **51**, 475–478.
- GREEN, T., SANCHO, L., TÜRK, R., SEPPELT, R. & HOGG, I.D. 2011a. High diversity of lichens at 84° S, Queen Maud Mountains, suggests preglacial survival of species in the Ross Sea region, Antarctica. *Polar Biology*, **34**, 1211–1220.
- GREEN, T.G.A., BRABYN, L., BEARD, C. & SANCHO, L.G. 2011b. Extremely low lichen growth rates in Taylor Valley, Dry Valleys, continental Antarctica. *Polar Biology*, **35**, 535–541.
- GREEN, T.G.A., SANCHO, L.G., PINTADO, A. & SCHROETER, B. 2011c. Functional and spatial pressures on terrestrial vegetation in Antarctica forced by global warming. *Polar Biology*, **34**, 1643–1656.
- GREEN, T.G.A., SCHROETER, B., SANCHO, L.G. 2007. Plant life in Antarctica. In PUGNAIRE, F.I. & VALLADARES F., eds. *Functional Plant Ecology*. Boca Ratón: CRC Press, 389–434.
- GRILLI CAIOLA, M., BILLI, D. & FRIEDMANN, E.I. 1996. Effect of desiccation on envelope of the cyanobacterium *Chroococcidiopsis* (Chroococcales). *European Journal of Phycology*, **31**, 97–105.
- GUGLIELMIN, M., CANNONE, N., STRINI, A. & LEWKOWICZ, A.G. 2005. Biotic and abiotic processes on granite weathering landforms in a cryotic environment, Northern Victoria Land, Antarctica. *Permafrost and Periglacial Processes*, **16**, 69–85.
- HERTEL, H. 1987. Progress and problems in taxonomy of Antarctic saxicolous lecideoid lichens. In PEVELING E., ed. *Progress and problems in lichenology in the eighties*. Berlin: J. Cramer, 219–242.
- HERTEL, H. 1988. Problems in monographing Antarctic crustose lichens. *Polarforschung*, **58**, 65–76.
- HODGSON, D. A., CONVEY, P., VERLEYEN, E., VYVERMAN, W., MCINNES, S. J., SANDS, C. J., FERNÁNDEZ-CARAZO, R., WILMOTTE, A., DE WEVER, A. & PEETERS, K. 2010. The limnology and biology of the Dufek Massif, Transantarctic Mountains 82° South. *Polar Science*, **4**, 197–214.
- HOPKINS, D.W., SPARROW, A.D., GREGORICH, E.G., ELBERLING, B., NOVIS, P., FRASER, F., SCRIMGEOUR, C., DENNIS,

- P.G., MEIER-AUGENSTEIN, W. & GREENFIELD, L.G. 2009. Isotopic evidence for the provenance and turnover of organic carbon by soil microorganisms in the Antarctic dry valleys. *Environmental Microbiology*, **11**, 597–608.
- HUGHES, K. 2006. Solar UV-B radiation, associated with ozone depletion, inhibits the Antarctic terrestrial microalga, *Stichococcus bacillaris*. *Polar Biology*, **29**, 327–336.
- HUGHES, K. A., LAWLEY, B. & NEWSHAM, K.K. 2003. Solar UV-B Radiation Inhibits the Growth of Antarctic Terrestrial Fungi. *Applied and Environmental Microbiology*, **69**, 1488–1491.
- JOHNSTON, C.G. & VESTAL, J.R. 1991. Photosynthetic carbon incorporation and turnover in Antarctic cryptoendolithic microbial communities: are they the slowest-growing communities on earth? *Applied and Environmental Microbiology*, **57**, 2308–2311.
- JOHNSTON, C.G. & VESTAL, J.R. 1989. Distribution of inorganic species in two Antarctic cryptoendolithic microbial communities. *Geomicrobiology Journal*, **7**, 137–153.
- JOHNSTON, C.G. & VESTAL, J.R. 1993. Biogeochemistry of oxalate in the Antarctic cryptoendolithic lichen-dominated community. *Microbial Ecology*, **25**, 305–319.
- KAPPEN, L. 1993. Lichens in the Antarctic region. In FRIEDMANN, E.I. *Antarctic Microbiology*. New York: Wiley-Liss, 433–490.
- KAPPEN, L. 2004. The diversity of lichens in Antarctica a review and comments. *Bibliotheca Lichenologica*, **88**, 331–343.
- KAPPEN, L., FRIEDMANN, E.I. & GARTY, J. 1981. Ecophysiology of lichens in the dry valleys of southern Victoria Land, I. Microclimate of the cryptoendolithic lichen habitat. *Flora*, **171**, 216–235.
- KAPPEN, L. & REDON, L. 1987. Photosynthesis and water relations of three antarctic lichen species. *Flora*, **179**, 215–229.
- KAPPEN, L., SCHROETER, B., GREEN, T.G.A. & SEPPELT, R.D. 1998. Chlorophyll a fluorescence and CO₂ exchange of *Umbilicaria aprina* under extreme light stress in the cold. *Oecologia*, **113**, 325–331.
- KHAN, N., TUFFIN, M.I., STAFFORD, W., CARY, C.S., LACAP, D., POINTING, S.B. & COWAN, D. 2011. Hypolithic microbial communities of quartz rocks from Miers Valley, McMurdo Dry Valleys, Antarctica. *Polar Biology* **34**, 1657–1668.
- KOGEJ, T., GORBUSHINA, A.A. & GUNDE-CIMERMAN, N. 2006. Hypersaline conditions induce changes in cell-wall melanization and colony structure in a halophilic and a xerophilic black yeast species of the genus *Trimmatostroma*. *Mycological Research*, **110**, 713–724.
- KRANNER, I., BECKETT, R.P., HOCHMAN, A. & NASH, T. 2008. Desiccation-tolerance in lichens: a review. *Bryologist*, **111**, 576–593.
- KRANNER, I., CRAM, W.J., ZORN, M., WORNIK, S., YOSHIMURA, I., STABENTHEINER, E. & PFEIFHOFFER, H.W. 2005. Antioxidants and photoprotection in a lichen as compared with its isolated symbiotic partners. *Proceedings of the National Academy of Sciences*, **102**, 3141–3146.
- LANGE, O.L. 1965. Der CO₂-gaswechsel von Flechten bei tiefen temperaturen. *Planta*, **64**, 1–19.
- MADRONICH, S., MCKENZIE, R., BJÖRN, L. & CALDWELL, M. 1998. Changes in biologically active ultraviolet radiation reaching the earth's surface. *Journal of Photochemistry and Photobiology*, **46**, 5–19.
- MATTHES, U., TURNER, S.J. & LARSON, D.W. 2001. Light attenuation by limestone rock and its constraint on the depth distribution of endolithic algae and cyanobacteria. *International Journal of Plant Sciences*, **162**, 263–270.
- MCKAY, C.P. & FRIEDMANN, E.I. 1985. The cryptoendolithic microbial environment in the Antarctic cold desert: temperature variations in nature. *Polar Biology*, **4**, 19–25.
- MODENESI, P., PIANA, M., GIORDANI, P., TAFANELLI, A. & BARTOLI, A. 2000. Calcium oxalate and medullary architecture in *Xanthomaculina convoluta*. *Lichenologist*, **32**, 505–512.

- MORITA, R.Y. (1975). Psychrophilic bacteria. *Bacteriological Reviews* **39**, 144.
- NAMSARAEV, Z., MANO, M.J., FERNÁNDEZ, R. & WILMOTTE, A. 2010. Biogeography of terrestrial cyanobacteria from Antarctic ice-free areas. *Annals of Glaciology*, **51**, 171-177.
- NIENOW, J.A., MCKAY, C.P. & FRIEDMANN, I.E. 1988. The cryptoendolithic microbial environment in the Ross Desert of Antarctica: Light in the photosynthetically active region. *Microbial Ecology*, **16**, 271-289.
- NIENOW, J.A. & FRIEDMANN, I.E. 1993. In FRIEDMANN, I.E., ed. *Terrestrial lithophytic (rock) communities*. New York: Wiley-Liss, 343-412.
- NISHIYAMA, T. 1977. Studies on evaporite minerals from Dry Valley, Victoria Land, Antarctica. *Antarctic Record* **58**, 171-185.
- OMELON, C. 2008. Endolithic microbial communities in polar desert habitats. *Geomicrobiology Journal*, **25**, 404-414.
- OMELON, C.R., POLLARD, W. & FERRIS, F.G. 2007. Inorganic species distribution and microbial diversity within high arctic cryptoendolithic habitats. *Microbial Ecology*, **54**, 740-752.
- ONOFRI, S., SELBMANN, L., DE HOOG, G.S., GRUBE, M., BARRECA, D., RUISI, S. & ZUCCONI, L. 2007a. Evolution and adaptation of fungi at boundaries of life. *Advances in Space Research*, **40**, 1657-1664.
- ONOFRI, S., ZUCCONI, D., SELBMANN, L., DE HOOG, G.S., DE LOS RÍOS, A., RUISI, S. & GRUBE, M. 2007b. Fungal associations at the cold edge of life. In SECKBACH, J., ed. *Algae and cyanobacteria in extreme environments*. Dordrecht: Springer, 739-757.
- ØVSTEDAL, D. & LEWIS-SMITH, R. 2001. *Lichens of Antarctica and South Georgia: a guide to their identification and ecology*. Cambridge: Cambridge University Press, 411 pp.
- ØVSTEDAL, D.O. & LEWIS SMITH, R. 2009. Further additions to the lichen flora of Antarctica and South Georgia. *Nova Hedwigia*, **88**, 157-168.
- ØVSTEDAL, D. & LEWIS-SMITH, R. 2011. Four additional lichens from the Antarctic and South Georgia, including a new *Leciophysma* species. *Folia Cryptogamica Estonica*, **48**, 65-68.
- PANDEY, K. 2004. Cyanobacteria in Antarctica: ecology, physiology and cold adaptation. *Cellular and Molecular Biology*, **50**, 575-584.
- PEARCE, D.A., BRIDGE, P. D., HUGHES, K. A., SATTLER, B., PSENNER, R. & RUSSELL, N.J. 2009. Microorganisms in the atmosphere over Antarctica. *FEMS Microbiology Ecology*, **69**, 143-157.
- PEAT, H., CLARKE, A. & CONVEY, P. 2007. Diversity and biogeography of the Antarctic flora. *Journal of Biogeography*, **34**, 132-146.
- PÉREZ-ORTEGA, S., ORTIZ, R., GREEN, T.G.A. & DE LOS RÍOS, A. 2012. Lichen myco- and photobiont diversity and their relationships at the edge of life (McMurdo Dry Valleys, Antarctica). *FEMS Microbial Ecology*, **82**, 429-448.
- POINTING, S.B., CHAN, Y., LACAP, D.C., LAU, M.C.Y., JURGENS, J. A. & FARRELL, R.L. 2009. Highly specialized microbial diversity in hyper-arid polar desert. *Proceedings of the National Academy of Sciences*, **106**, 19 964-19 969.
- POINTING, S.B. & BELNAP, J. 2012. Microbial colonization and controls in dryland systems. *Nature Reviews in Microbiology*, **10**, 551-562.
- RAGGIO, J., PINTADO, A., ASCASO, C., DE LA TORRE, R., DE LOS RÍOS, A., WIERZCHOS, J., HORNECK, G. & SANCHO, L.G. 2011. Whole lichen thalli survive exposure to space conditions: results of lithopanspermia experiment with *Aspicilia fruticulosa*. *Astrobiology*, **11**, 281-92.
- ROMEIKE, J., FRIEDL, T., HELMS, G. & OTT, S. 2002. genetic diversity of algal and fungal partners in four species of *Umbilicaria* (Lichenized Ascomycetes) along a transect of the Antarctic Peninsula. *Molecular Biology and Evolution*, **19**, 1209-1217.

- ROSZAK, D.B. & COLWELL, R.R. 1987. Survival strategies of bacteria in natural environment. *Microbiological Reviews*, **51**, 365–379.
- RUIBAL, C., GUEIDAN, C., SELBMANN, L., GORBUSHINA, A., CROUS, P., GROENEWALD, J.Z., MUGGIA, L., GRUBE, M., ISOLA, D., SCHOCH, C., STALEY, J., LUTZONI, F. & DE HOOG, G.S. 2009. Phylogeny of rock-inhabiting fungi related to *Dothideomycetes*. *Studies in Mycology*, **64**, 123–133.
- RUISI, S., BARRECA, D., SELBMANN, L., ZUCCONI, L. & ONOFRI, S. 2007. Fungi in Antarctica. *Reviews in Environ Science and Biotechnology*, **6**, 127–141.
- RUPRECHT, U., BRUNAUER, G. & PRINTZEN, C. 2012. Genetic diversity of photobionts in Antarctic lecideoid lichens from an ecological viewpoint. *The Lichenologist*, **44**, 661–678.
- RUPRECHT, U., LUMSCH, H., BRUNAUER, G., GREEN, T.G.A. & TÜRK, R. 2010. Molecular data reveal unexpected diversity of Lecidea (Lecideaceae, Ascomycota) species in continental Antarctica (Ross Sea Region) corroborated by previously overlooked morphological characters. *Antarctic Science*, **22**, 727–741.
- SANCHO, L.G., DE LA TORRE, R., HORNECK, G., ASCASO, C., DE LOS RÍOS, A., PINTADO, A., WIERZCHOS, J. & SCHUSTER, M. 2007. Lichens survive in space: results from the 2005 LICHENS experiment. *Astrobiology*, **7**, 443–454.
- SANCHO, L.G., VALLADARES, F., SCHROETER, B. & KAPPEN, L. 2000. Ecophysiology of Antarctic versus temperate populations of a bipolar lichen: the key role of the photosynthetic partner. In DAVISON W., HOWARD-WILLIAMS C. & BROADY, P. *Antarctic ecosystems: models for wider ecological understanding*. Christchurch: The Caxton Press.
- SCHOPF, J.W. 1993. Microfossils of the early archean apex chert: new evidence of the antiquity of life. *Science*, **260**, 640–646.
- SCHROETER, B., GREEN, T.G.A., KAPPEN, L. & SEPPELT, R.D. 1994. Carbon dioxide exchange at subzero temperatures. Field measurements on *Umbilicaria aprina* in Antarctica. *Cryptogamic Botany*, **4**, 233–241.
- SCHROETER, B., GREEN, T.G.A., PANNEWITZ, S., SCHLENSOG, M. & SANCHO, L.G. 2010. Fourteen degrees of latitude and a continent apart: comparison of lichen activity over two years at continental and maritime Antarctic sites. *Antarctic Science*, **22**, 681–690.
- SCHROETER, B., GREEN, T.G.A., PANNEWITZ, S., SCHLENSOG, M. & SANCHO, L.G. 2011. Summer variability, winter dormancy: lichen activity over 3 years at Botany Bay, 77°S latitude, continental Antarctica. *Polar Biology*, **34**, 13–22.
- SELBMANN, L., DE HOOG, G. S., MAZZAGLIA, A., FRIEDMANN, E.I. & ONOFRI, S. 2005. Fungi at the edge of life: cryptoendolithic black fungi from Antarctic desert. *Studies in Mycology*, **51**, 1–32.
- SELBMANN, L., DE HOOG, G. S., ZUCCONI, D., RUISI, S., GERRITS VAN DEN ENDE, A.H.G., RUIBAL, C., DE LEO, F., URZI, C. & ONOFRI, S. 2008. Drought meets acid: three new genera in a dothidealean clade of extremotolerant fungi. *Studies in Mycology*, **61**, 1–20.
- SELBMANN, L., ISOLA, D., ZUCCONI, L. & ONOFRI, S. 2011. Resistance to UV-B induced DNA damage in extreme-tolerant cryptoendolithic Antarctic fungi: detection by PCR assays. *Fungal Biology*, **115**, 937–944.
- SEPPELT, R.D., TÜRK, R., GREEN, T.G.A., MOSER, G., PANNEWITZ, S., SANCHO, L.G. & SCHROETER, B. 2010. Lichen and moss communities of Botany Bay, Granite Harbour, Ross Sea, Antarctica. *Antarctic Science*, **22**, 691–702.
- SHIRKEY, B., MCMASTER, N.J., SMITH, S.C., WRIGHT, D. J., RODRIGUEZ, H., JARUGA, P., BIRINCIOGLU, M., HELM, R.F. & POTTS, M. 2003. Genomic DNA of *Nostoc commune* (Cyanobacteria) becomes covalently modified during long-term (decades) desiccation but is protected from oxidative damage and degradation. *Nucleic Acids Research*, **31**, 295–300.
- SIEBERT, J. & HIRSCH, P. 1988. Characterization of 15 Selected Coccal Bacteria Isolated from Antarctic Rock and Soil

- Samples from the McMurdo-Dry Valleys (South-Victoria Land). *Polar Biology*, **9**, 37–44.
- SIEBERT, J., HIRSCH, P., HOFFMANN, B. & GLIESCHE, C.G. 1996. Cryptoendolithic microorganisms from Antartic sandstone of Linnaeus Terrace (Asgard Range): diversity, properties and interactions. *Biodiversity and Conservation*, **5**, 1337–1363.
- SINGH, J., DUBEY, A. & SINGH, R. 2011. Antarctic terrestrial ecosystem and role of pigments in enhanced UV-B radiations. *Reviews in Environmental Science and Biotechnology*, **10**, 63–77.
- STERFLINGER, K., TSEI, D. & ZAKHAROVA, K. 2012. Fungi in hot and cold deserts with particular reference to microcolonial fungi. *Fungal Ecology*, **5**, 453–462.
- SUN, H.J. & FRIEDMANN, E.I. 1999. Growth on geological time scales in the Antarctic cryptoendolithic microbial community. *Geomicrobiology Journal*, **16**, 193–202.
- TAMARU, Y., TAKANI, Y., YOSHIDA, T. & SAKAMOTO, T. 2005. Crucial role of extracellular polysaccharides in desiccation and freezing tolerance in the terrestrial cyanobacterium. *Applied and Environmental Microbiology*, **71**, 7327–7333.
- TOSI, S., ONOFRI, S., BRUSONI, M., ZUCCONI, L. & VISHNIAC, H. 2005. Response of Antarctic soil fungal assemblages to experimental warming and reduction of UV radiation. *Polar Biology*, **28**, 470–482.
- TSCHERMAK-WOESS, E. & FRIEDMANN, E.I. 1984. *Hemichloris antarctica* gen. et sp. nov. (Chlorococcales, Chlorophyta) a cryptoendolithic alga from Antarctica. *Phycologia*, **23**, 443–454.
- VESTAL, J.R. 1988. Biomass of the cryptoendolithic microbiota from the Antarctic desert. *Applied and Environmental Microbiology*, **54**, 957–959.
- VILLAR, S.E.J., EDWARDS, H.G.M. & SEAWARD, M.R.D. 2005. Raman spectroscopy of hot desert, high altitude epilithic lichens. *The analyst*, **130**, 1–9.
- VINCENT, W.F. 2007. Cold tolerance in cyanobacteria and life in the cryosphere. In SECKBACH, J. *Algae and cyanobacteria in extreme environments*. Heidelberg: Springer, 287–301.
- VYVERMAN, W., VERLEYEN, E., WILMOTTE, A., HODGSON, D. A., WILLEMS, A., PEETERS, K., VAN DE VIJVER, B., DE WEVER, A., LELIAERT, F. & SABBE, K. 2010. Evidence for widespread endemism among Antarctic microorganisms. *Polar Science*, **4**, 103–113.
- WALKER, J.J. & PACE, N.R. 2007. Phylogenetic composition of Rocky Mountain endolithic microbial ecosystems. *Applied and Environmental Microbiology*, **73**, 3497–3504.
- WESTALL, F. & FOLK, R.L. 2003. Exogenous carbonaceous microstructures in Early Archaean cherts and BIFs from the Isua greenstone belt: Implications for the search for life in ancient rocks. *Precambrian Research*, **126**, 313–330.
- WIERZCHOS, J. & ASCASO, C. 1994. Application of back-scattered electron imaging to the study of the lichen rock interface. *Journal of Microscopy*, **175**, 54–59.
- WIERZCHOS, J. & ASCASO, C. 1996. Morphological and chemical features of bioweathered granitic biotite induced by lichen activity. *Clays and Clay Minerals*, **44**, 652–657.
- WIERZCHOS, J. & ASCASO, C. 1998. Mineralogical transformation of bioweathered granitic biotite, studied by HRTEM: evidence for a new pathway in lichen activity. *Clays and clay minerals*, **46**, 446–452.
- WIERZCHOS, J. & ASCASO, C. 2001. Life, decay and fossilisation of endolithic microorganisms from the Ross Desert, Antarctica: suggestions for *in situ* further research. *Polar Biology*, **24**, 863–868.
- WIERZCHOS, J. & ASCASO, C. 2002. Microbial fossil record of rocks from the Ross Desert Antarctica: implications in the search for past life in Mars. *International Journal of Astrobiology*, **1**, 51–60.
- WIERZCHOS, J., ASCASO, C., SANCHO, L. G. & GREEN, A. 2003. Iron-rich diagenetic minerals are biomarkers of microbial activity in Antarctic rocks. *Geomicrobiology Journal*, **20**, 15–24.

- WIERZCHOS, J., DE LOS RÍOS, A., SANCHO, L.G. & ASCASO, C. 2004. Viability of endolithic micro-organisms in rocks from the McMurdo Dry Valleys of Antarctica established by confocal and fluorescence microscopy. *Journal of Microscopy*, **216**, 57–61.
- WIERZCHOS, J., SANCHO, L.G. & ASCASO, C. 2005. Biomineralization of endolithic microbes in rocks from the McMurdo Dry Valleys of Antarctica: implications for microbial fossil formation and their detection. *Environmental Microbiology*, **7**, 566–575.
- WINKLER, J.B., KAPPEN, L. & SCHULZ, F. 2000. Snow and ice as an important ecological factor for the cryptogams in the maritime Antarctic. In HOWARD-WILLIAMS, C. & DAVISON, W., eds. *Antarctic ecosystems: models for wider ecological understanding*. Christchurch: The Caxton Press, 220–224.
- WIRTZ, N., LUMBSCH, T., GREEN, A., TÜRK, R., PINTADO, A., SANCHO, L.G. & SCHROETER, B. 2003. Lichen fungi have low cyanobiont selectivity in maritime Antarctica. *New Phytologist*, **160**, 177–183.
- WOLFAARDT, G.M., LAWRENCE, J.R. & KORBER, D.R. 1999. Function of EPS. In WINGENDER, J., NEU, T.R. & FLEMMING, H.C., eds. *Microbial extracellular polymeric substances*. Berlin: Springer, 171–195.
- YERGEAU, E., BORKHORST, S., HUISKES, A.H.L., BOSCHKER, H.T.S., AERTS, R. & KOWALCHUK, G.A. 2007. Size and structure of bacterial, fungal and nematode communities along an Antarctic environmental gradient. *FEMS Microbiol Ecology* **29**, 436–451.

Fig. 1. General landscape of **a.** Miers Valley and **b.** University Valley in the McMurdo Dry Valleys.

Fig. 2. **a.** Granite boulder showing epilithic lichen colonization and snow deposits, which could favour lichen community development. **b.** *Umbilicaria aprina* Nyl. thalli occupying protected areas of the rock surface. **c.** Lecideoid lichen colonizing weathered granite.

Fig. 3. **a.** Sandstones from University Valley showing a lack of epilithic colonization. **b.** Fractured sandstone showing bands of microbial colonization in the area indicated with a square in **a.** **c.** Granite boulder from the Miers Valley showing a lack of epilithic colonization. **d.** Fractured granite boulder showing signs of endolithic cyanobacterial colonization corresponding to the area indicated with a square in **c.**

Fig. 4. **a.** Scanning electron microscope - backscattered electron (SEM-BSE) image of chasmoendolithic colonization of granite rock. Arrow shows the fungal colonized fissure. **b.** SEM-BSE image showing cryptoendolithic colonization in pores of sandstones (arrows).

Fig. 5. **a.** Closer view of the cryptoendolithic microbial colonization of sandstone. **b.** Same area as in **a.** visualized by epifluorescence microscopy showing autofluorescence emitted by algal cells (red signal). **c.** Microscopy image of a sandstone area colonized by cyanobacteria. **d.** Same area as in **c.** visualized by epifluorescence microscopy showing autofluorescence emitted by phototrophic microorganisms (red signal).

Fig. 6. Cryptoendolithic colonization of Antarctic sandstones. Arrows indicate the presence of black fungi. **a.** Microscopy image of an area colonized by black fungi on sandstone. **b.** Scanning electron microscope - backscattered electron (SEM-BSE) image of the white band occupied by algal (A) and fungal (F) associations. **c.** SEM-BSE image of free-living green algae colonizing a sandstone pore. **d.** SEM-BSE image of a sandstone pore colonized by cyanobacteria (C).

Fig. 7. Chasmoendolithic colonization of Antarctic granites. **a.** Scanning electron microscope - backscattered electron (SEM-BSE) image of fissures colonized by cyanobacteria. **b.** SEM-BSE image of fissures colonized by lichen algal-fungal associations. **c.** SEM-BSE image of fissures colonized by free-living fungi (arrow). **d.** SEM-BSE image of fissures colonized by free-living algae. **e.** Confocal scanning light microscopy image of glutaraldehyde-fixed endolithic lichen showing colonization of a rock fissure. **f.** Epilithic lichen colonizing granite. Arrow indicates the separation of mica layers on the rock surface and endolithic colonization through these layers.

Fig. 8. **a.** Low-temperature scanning electron microscope (LTSEM) image of endolithic cyanobacteria embedded in an extracellular polymeric substance (EPS) matrix (arrows). **b.** LTSEM image of a cyanobacteria community showing fractured cells surrounded by an EPS matrix. **c.** Transmission electron microscopy (TEM) image of endolithic microbial cells showing living bacterial cells in close proximity to green alga cells (A). **d.** TEM image of endolithic *Gloeocapsa* cyanobacteria cells showing heterotrophic bacteria (arrows) in the cyanobacterial EPS matrix. **e.** *In situ* 3D reconstruction of a lithobiontic microbial community visualized by epifluorescence microscopy operated in structural illumination mode. **f.** TEM image of a group of cyanobacterial cells showing living cells (arrow), cells that have lost their cell contents (arrowhead) and remnants of cells (asterisk).

Fig. 9. **a.** Scanning electron microscope - backscattered electron (SEM-BSE) image of fungal endolithic colonization associated with damaged micas. **b.** Calcium oxalate crystals (arrows) associated with endolithic lichen growth.