

Review Article

Melanins and Melanogenesis: From Pigment Cells to Human Health and Technological Applications.

Marco d'Ischia,^{1*} Kazumasa Wakamatsu,² Fabio Cicoira,³ Eduardo Di Mauro,⁴ Josè Carlos Garcia-Borron,⁵ Stephane Commo,⁶ Ismael Galván,⁷ Ghanem Ghanem,⁸ Koike Kenzo,⁹ Paul Meredith,¹⁰ Alessandro Pezzella,¹ Clara Santato,⁴ Tadeusz Sarna,¹¹ John. D. Simon,¹² Luigi Zecca,¹³ Fabio A. Zucca,¹³ Alessandra Napolitano,¹ Shosuke Ito.²

¹Department of Chemical Sciences, University of Naples “Federico II”, Via Cintia 4, I-80126 Naples, Italy

²Department of Chemistry, Fujita Health University School of Health Sciences, Toyoake, Aichi 470-1192, Japan

³Department of Chemical Engineering, École Polytechnique de Montréal, 2900 Boulevard Edouard-Montpetit, Montréal, QC H3T 1J4, Canada.

⁴Department of Engineering Physics, École Polytechnique de Montréal, 2900 Boulevard Edouard-Montpetit, Montréal, QC H3T 1J4, Canada.

⁵School of Medicine, University of Murcia, and IMIB-Arrixaca, Murcia, Spain

⁶L'Oréal Recherche & Innovation, 1 Avenue E.Schueller - 93600 Aulnay sous Bois France

⁷Departamento de Ecología Evolutiva, Estación Biológica de Doñana - CSIC, c/ Américo Vespucio s/n, 41092 Sevilla, Spain

⁸LOCE, Institut J. Bordet, Université Libre de Bruxelles, Belgium

⁹Development Research - Hair Care Products, KAO Corporation, 2-1-3, Bunka, Sumida, Tokyo, 131-8501, JAPAN

¹⁰Centre for Organic Photonics and Electronics, School of Mathematics and Physics, University of Queensland, St Lucia campus, Brisbane, Queensland 4072 Australia

¹¹Department of Biophysics, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Gronostajowa 7, 30-387 Krakow, Poland

¹²Department of Chemistry, University of Virginia, Charlottesville, VA 22904-4319

¹³Institute of Biomedical Technologies – National Research Council of Italy, 20090 Segrate, Milan, Italy.

*Address correspondence to:

Professor Marco d'Ischia, Department of Chemical Sciences, University of Naples “Federico II”, Via Cintia 4, I-80126 Naples, Italy, Phone: +39-081-674132, Fax: +39-081-674393, e-mail: dischia@unina.it

Total word counts: abstract (177) , text and figure legends (12,475) , references (9,700).

Summary

During the past decade, melanins and melanogenesis have attracted growing interest for a broad range of biomedical and technological applications. The burst of polydopamine-based multifunctional coatings in materials science is just one example, and the list may be expanded to include melanin-based organic semiconductors, drug delivery systems, functional nanoparticles and biointerfaces, sunscreens, environmental remediation devices. Despite considerable advances, applied research on melanins and melanogenesis is still far from being mature. Closer intersectoral interactions between research centers is essential to raise the interests and increase the awareness on the part of the biomedical, biomaterials science and hi-tech sectors of the manifold opportunities offered by pigment cells and related metabolites and pathways. Starting from a survey of biological roles and functions at various levels of the evolutionary scale, the present review aims at providing an interdisciplinary perspective of melanin pigments and related pathway with a view to showing how it is possible to translate current knowledge about physical and chemical properties and control mechanisms into new biomedical, dermocosmetic and sustainable technological solutions.

Key words: Melanogenesis control, bioelectronics, antioxidants, polydopamine, dermocosmetic applications, melanosomes, extracutaneous melanins.

Running title: 50 characters Melanins for health and technological applications

Introduction

Melanin-producing cells, the melanocytes, are endowed with unique and fascinating properties that make them a most valuable source of inspiration for new exciting discoveries and for new approaches and technological solutions to important biomedical problems from a multidisciplinary and interdisciplinary perspective. Much of the uniqueness of pigment cells is due to the pigment itself, melanin, as a most distinguishing trait that has attracted the attention of scientists and clinicians since a long time.

As a matter of fact, over the past decade melanins and melanogenesis have been increasingly appreciated outside the pigment cell community, as a source of novel research opportunities in the fields of biomedicine, dermocosmetics, nanotechnology and materials science. The burst of interest in polydopamine is a paradigm example of this trend (Liu et al., 2014). Following a preceding consensus review (d'Ischia et al., 2013), this paper has been conceived as a means of bringing together for the first time researchers from broadly diverse disciplines and from both academic and industrial settings to provide an interdisciplinary and intersectoral perspective of melanins and melanogenesis. Far from being comprehensive, the paper aims mainly to offer an inspiring overview of melanin roles and functions to stimulate the interests of academic and industrial laboratories active in the fields of human health and sustainable technology (Figure 1). For a more detailed discussion of the topics of this paper, the readers are referred to the supporting information, as well as to a number of excellent reviews (for example, Prota, 1992; Riley, 1997; Galvan and Solano, 2009; Solano 2014) and a recent book (Borovansky and Riley, 2011).

Terminology

The term *melanins* denotes pigments of diverse structure and origin derived by the oxidation and polymerization of tyrosine in animals or phenolic compounds in lower organisms. *Eumelanins* are the black-brown subgroup of insoluble melanin pigments derived at least in part from the oxidative polymerization of L-dopa via 5,6-dihydroxyindole intermediates. *Pheomelanins* are yellow-to-reddish brown subgroup of melanin pigments derived from the oxidation of cysteinyl-dopa precursors via benzothiazine and benzothiazole intermediates. It is noticed that very often the terms melanin and eumelanin are used as synonymous, though this practice is not recommended. Melanocytes are responsible for the synthesis of melanin within specialized membrane-bound organelles termed *melanosomes*, and the subsequent transfer of the melanosomes to surrounding epidermal cells, the keratinocytes. Melanosomes are thus distinct from “*melanin granules*” which refer to the pigment particles produced by enzymatic or chemical reactions. “*Melanogenesis*” is the complex process of melanin synthesis in nature or, by extension, in an in vitro system. A more in-depth discussion of melanin nomenclature, definitions and classification is given in the previous consensus paper (d'Ischia et al., 2013).

Biological roles and functions of melanins and melanogenesis

Skin and hair

Photoprotection of the skin is apparently one of the most important biological functions of melanin pigments, since skins with higher levels of constitutive pigmentation appear to be less susceptible to the deleterious effects of ultraviolet radiation (UVR) (Brenner and Hearing, 2008; Coelho et al., 2009). Subjects with white skin are approximately 70 times more likely to develop skin cancer than subjects with black skin. All types of skin, including those with red hair, contain a higher proportion of eumelanin than pheomelanin. Generally speaking, black and Asian skin contain several-fold and two-fold higher levels of melanin than fair skin, respectively (Tadokoro et al., 2003). Although the epidermal melanin confers skin protection factor of only 2-3, it provides sufficient photoprotection against UVR-induced damage of nuclear DNA by shielding nuclei by supranuclear melanin caps in melanocytes and keratinocytes (Kobayashi et al., 1998). Other important properties of eumelanin include its function as a free radical scavenger reducing the production of reactive oxygen species (ROS) (Meredith and Sarna, 2006).

Hair pigmentation is one of the most conspicuous phenotypes in humans. It is highly variable in color, ranging from black, dark brown, light brown, and blond to red. The underlying genetic basis for this diversity in human hair color has been a subject of extensive studies in recent years (for a review, see Ito and Wakamatsu, 2011; Sturm and Duffy, 2012). A pioneer work in those studies was reported in 1995 by Valverde et al. (Valverde et al., 1995) showing that specific mutations at the melanocortin-1 receptor (MC1R) gene are responsible for the switch from eumelanin to pheomelanin underlying the red hair phenotype (RHP). Chemical basis underlying the diversity of hair color was studied by using chemical degradative approach (Ito et al., 2011b; Wakamatsu et al., 2003). The contents of eumelanin and pheomelanin (chemical phenotype) in human hairs of black, dark brown, brown, light brown, blond, and red color (visual phenotype) were measured. Eumelanin contents decrease in that order, with a trace but constant level of pheomelanin, except for red hair which contains about equal levels of pheomelanin and eumelanin. Thus, the chemical phenotype correlates well with the visual phenotype (Ito et al., 2011b). It also helped evaluate the precise effects of various hypopigmentary disorders.

Melanins, in particular pheomelanin, can also have toxic properties, especially upon exposure to UVR. Pheomelanin is prone to photodegradation (Chedekel, 1982; Greco et al., 2009; Wakamatsu et al., 2012a) and is believed to contribute to the damaging effects of UVR because it can generate hydrogen peroxide and superoxide anions (Chedekel et al., 1978; Felix et al., 1978; Simon and Peles, 2010) and might cause mutations in melanocytes (Harsanyi et al., 1980). The ability of pheomelanin to induce antioxidant depletion without UVR has recently been disclosed (Panzella et al., 2014). In relation to (photo)toxic properties of melanin pigments, several studies have recently addressed possible roles of eumelanin or pheomelanin in inducing DNA damage and ultimately melanomagenesis both with/without UVR. Mitra et al. (2012) showed that by using a conditional, melanocyte-targeted allele of the melanoma oncoprotein BRAF(V600E) into yellow mice carrying the $Mc1r^{e/e}$ allele of the *Mc1r* gene, a high incidence of invasive melanoma was

observed without UVR. On the other hand, selective absence of pheomelanin synthesis was proved to be protective against melanoma development.

Eyes

The mammalian eye contains melanin-producing cells, which are present in different tissues and have different embryonic origin (Sarna, 1992; Hu, 2008). While the pigment epithelial cells of the retina (RPE), the iris and the ciliary are derived from the neural ectoderm, the uveal melanocytes of the choroid, the stroma of the iris and ciliary body, are developed from the neural crest. The human pigment epithelium is typically densely pigmented in all races and in all eye colors, and it mainly contains eumelanin. Melanosomes in the human RPE are formed early in the fetal development and although pigment deposition in these organelles may continue for some time after birth, there is very little metabolic turnover of the formed pigment granules. Pigmentation in human uveal melanocytes exhibits significant variation; these cells contain both eumelanin and pheomelanin, and their relative content and total amount depends on the race and iris color (Prota et al., 1998; Wakamatsu et al., 2008; Wielgus and Sarna, 2005).

The biological role of ocular pigmentation is only partially understood. There is little doubt that melanin in the iris and the RPE acts as an optical absorber, which prevents the retina from excess of light that impinges on the eye and reduces spurious signals that could arise from light reflection of the surface of the Bruch's membrane. Interestingly, the incidence of two important eye diseases, uveal melanoma and age-related macular degeneration (AMD), appears to be correlated with the pigmentation of the iris. A population-based study revealed that the incidence of uveal melanoma was highest in non-Hispanic white followed by Hispanics, Asians and blacks (Weis et al., 2006). Although the relationship between iris color and AMD is not so conclusive as that in uveal melanoma, AMD is an order of magnitude more prevalent in the white population than in darkly pigmented races (Friedman et al., 1999).

Although the correlation between AMD and pigmentation of the iris is supported by convincing epidemiological data, whether or not such a correlation can be viewed as causal is still unclear. Nevertheless, it has been postulated by many researchers that excess of light that reaches the retina is a contributing factor to AMD. Therefore light-pigmented iris, which transmits more light, could be responsible for enhanced risk of developing AMD.

Acting as an antioxidant, uveal and RPE melanin can also protect the pigmented tissues against oxidative damage, induced by photic or oxidative stress (Sarna, 1992; Meredith and Sarna, 2006). Indeed, it was recently demonstrated that phagocytized porcine RPE melanosomes protected ARPE-19 cells, a human RPE cell line, against oxidative stress induced by hydrogen peroxide (Burke et al., 2011).

Metal accumulation, especially iron, has been reported in eye melanosomes, suggesting that melanosomes may contribute to iron homeostasis (Kaczara et al., 2012). The lack of metabolic turnover of melanin in post-mitotic cells, such as RPE, raises an intriguing question whether antioxidant and photoprotective abilities of this pigment decrease with aging and, if so, what physicochemical modifications of melanin are responsible for such changes. EPR (Sarna et al., 2003) and chemical analysis (Ito et al., 2013) of human RPE melanin

and of bovine RPE melanosomes exposed to intense blue light demonstrated an age-related loss of the content of human RPE melanin, which undergoes progressive modifications through oxidative cleavage and cross-linking. These changes are most likely induced by in situ chronic photooxidation of the melanin. Chloroquine can be detected in eye melanin one year after administration, a phenomenon possibly associated with retinopathies.

Human brain

Neuromelanin (NM) containing neurons are ubiquitous in human brain and occur with the highest number in substantia nigra (SN) and locus coeruleus (LC), regions that are the main targets of Parkinson's disease (PD) (Cowen, 1986; Zecca et al., 2008a). The NM pigments are found also in neurons of monkeys, dogs, horses, rats and frogs (Barden, 1971; DeMattei et al., 1986).

NM pigments are contained within double membrane organelles along with lipid bodies and proteins (Duffy and Tennyson 1965; Sulzer et al., 2008; Zecca et al., 2008a). Recent investigations show that these NM-containing organelles are a special type of lysosomes derived from fusion with autophagic vacuoles. NM deposition begins very early in life and the pigment concentration increases linearly with age reaching values up to 3.7 $\mu\text{g}/\text{mg}$ tissue between 50 and 90 years of age in SN and up to 3.1 $\mu\text{g}/\text{mg}$ tissue in LC (Zecca et al., 2002, 2004). In PD, a preferential loss of SN dopamine neurons containing NM occurs with respect to non-pigmented neurons and it has been long discussed if NM can increase neuronal vulnerability in PD (Hornykiewicz and Kish, 1987; Hirsch et al., 1988; Marsden, 1983). NM concentration is 50-60 % lower than in age-matched controls, due to the loss of NM-containing neurons (Zecca et al., 2002, 2004). A relationship between the vulnerability of dopamine neurons and their NM content was reported (Kastner et al., 1992).

NM pigments are comprised of spherical aggregates which are ~200-400 nm in size, with varying shapes, composed of smaller spherical substructures of ~30 nm in diameter (Bush et al., 2006, 2009). The structure proposed for the melanic portion of NMs is that of a pheomelanin core surrounded by eumelanin, forming spherical electron-dense aggregates. The oxidation potential of NM surface is not sufficiently reductive to cause a high level of oxidative stress, attributed to the presence of eumelanin on the surface (Bush et al. 2006). In NM structures there are melanic groups bound to proteins and aliphatic chains (Zecca et al. 1992, 2000). The melanic groups, composed of dihydroxyindole and benzothiazine rings are covalently bound to aliphatic chains and peptide moieties (Wakamatsu et al., 2003). An organic radical located on the melanic portion is present in all NM pigments (Aime et al., 1997; Enochs et al., 1993; Zecca et al., 1996, 2008a). The amount of lipids covalently bound to NM pigment of the SN is about 20 %, while lipid content is higher for NMs from other brain areas (Zecca et al., 2000; Engelen et al., 2012). Dolichols and dolichoic acids (containing 14-22 isoprene units) are the main component of NM lipids, and to a lesser extent cholesterol, ubiquinone-10, α -tocopherol, phospholipids, and lactones (Ward et al., 2007, 2009; Engelen et al., 2012; Zecca et al., 2000, 2008a). In NM pigments there is a soluble portion composed of dolichols bound to

proteins and melanic oligomers, and an insoluble portion with a more extended melanic component (Ward et al., 2007, 2009; Engelen et al., 2012).

The synthesis of NM pigment depends on the cytosolic concentrations of catechols that are not accumulated in synaptic vesicles (Sulzer et al., 2000). Catechols can be oxidized to quinones and semiquinones via iron catalysis, and these can react with aggregated proteins accumulated in cytosol. The polymerization process forms a melanin-protein component, which can also bind high amounts of metals, especially iron. The resulting undegradable material is taken into autophagic vacuoles and after fusion with lysosomes, it can interact with lipids and other proteins forming the NM-containing organelle (Engelen et al., 2012; Zucca et al., 2014).

The synthesis of NM is itself a protective mechanism that traps excessive cytosolic quinones and semiquinones in an inactive form, thereby blocking their toxicity (Sulzer et al., 2000). Moreover the dihydroxyindole groups of NM pigment are able to bind several potentially toxic metals forming stable complexes (Zecca and Swartz, 1993; Zecca et al., 1994, 2008a), and in particular, the redox-active ferric ions, mitigating the formation of hydroxyl radicals (Zecca et al., 2008b). NM pigment can initially play a protective role also by sequestering several toxins, such as 1-methyl-4-phenylpyridinium ion (D'Amato et al., 1986) and the herbicide paraquat (Lindquist et al., 1988).

Catecholamine neurons of SN and LC which degenerate in PD express the major histocompatibility complex class I (MHC-I) and this is highly concentrated in NM-containing organelles. MHC-I can bind antigenic peptides and expose them on neuronal membrane, so that cytotoxic T cells target these neurons inducing cell death. The finding of MHC-I in pigmented catecholamine neurons of SN and LC provides an explanation for selective cell death of neurons containing NM, suggesting the presence of a novel neurodegenerative process of autoimmune type that is specific for PD (Cebrian et al., 2014). In parkinsonian SN, NM is released from dying neurons and remains in the extracellular space, along with sustained microglial activation (Langston et al., 1999; McGeer et al., 1988). Due to its insolubility, NM pigment could remain for long time in the extracellular milieu of the SN, and may become an agent of chronic inflammation, slowly releasing the metals and toxic components encased in its structure. It was found that this NM can activate microglia causing release of tumor-necrosis factor- α , interleukin-6, nitric oxide and hydrogen peroxide. Then activated microglia and these molecules induce degeneration of dopaminergic neurons thus fueling a continuous vicious cycle of neuroinflammation and neurodegeneration causing the progression of PD (Wilms et al., 2003; Zhang et al., 2011; Zecca et al., 2008c).

Inner ear

Melanin-producing cells are present in the inner ear (Meyer zum Gottesberge, 1988), with the higher number of melanocytes being found in the intermediate layer of the stria vascularis, where they are interplaced with the secretory cells and are in contact with the capillary vessels (Breathnach, 1988). Melanin of the inner ear from the utricles and posterior ampullae of colored guinea pigs is chemically similar to the pigments from hair and eyes (Inoue et al., 1985).

The function of melanin in the inner ear remains unclear. Likely hypotheses for melanin roles include biological reservoir for divalent ions, ion exchanger, and intracellular buffering system for calcium. In relation to this latter aspect, local disturbances of calcium homeostasis in the inner ear induce a set of morphological responses of melanocytes involving intracellular movement and intraepithelial deposition of melanosomes, motility, *de novo* melanogenesis and enhanced exocytotoxic/endocytotic activity (Meyer zum Gottesberge, 1988). Besides metal ions, inner ear melanin can bind drugs (Wasterstrom et al., 1986), a property which has been implicated in the mechanism of ototoxicity of aminoglycoside-type drugs, e.g. kanamycin (Gratacap et al., 1989). A more profound hearing loss was observed in pigmented guinea pigs treated with kanamycin compared to albinos, supporting a role of kanamycin affinity to inner ear melanin in ototoxicity. In vitro studies on melanocyte cultures showed that kanamycin inhibits melanogenesis and lowers the levels of antioxidant enzymes, suggesting that the undesired effects of the drug may be the result of lack of the antioxidant defense mechanism in the cell at least in part due to the melanin pigment (Wrzesniok et al., 2013). A study of the relationship between skin pigmentation and noise-induced temporary hearing loss (temporary threshold shift, TTS) showed that the most pigmented subjects are more protected against TTS (Barrenas, 1994). Similar studies on the high-frequency threshold and latency of stapes reflex indicated that vitiligo is an effective factor in hearing loss, with men more susceptible than women, reinforcing the hypothesis of a protective function of melanin-containing cells in the inner ear (Ardic et al., 1998). A role of melanocytes in maintaining the integrity of the intrastrial fluid-blood barrier and hearing function has also been demonstrated (Zhang et al., 2012).

Birds, reptiles, amphibians and fish

Among non-mammalian vertebrates, birds are the group where the properties and functions of melanins have been so far most intensively investigated. Many of the genes encoding the main components of the mammalian melanogenesis pathway, i.e. pro-opiomelanocortin (POMC), five melanocortin receptors (MCRs) and agouti, have all been characterized in birds (Takeuchi et al., 2003). Perhaps the most striking difference is the fact that, while in mammals it is the agouti-signalling protein (ASIP) that antagonizes MC1R signalling on melanocytes, in birds this role may be fulfilled by the agouti-related protein (AGRP), which is involved in the regulation of energy balance in mammals (Boswell and Takeuchi, 2005). Thus, the melanogenesis pathway is remarkably similar in mammals and birds and at least the basic mechanisms seem to have evolved early (Schiöth et al., 2003).

Despite the similarity in the mechanisms of melanin production, the functions of melanins in birds seem to be diverse than those so far described in mammals. Thus, feathers colored by melanins have evolved as signals in many species with which individuals can honestly reveal their genotypic quality to conspecifics and thus maximize their reproductive success or performance during social interactions (reviewed in McGraw, 2008). The fact that birds, like mammals, produce the two main forms of melanin (i.e., eumelanin and pheomelanin) opens interesting perspectives about the signalling potential of melanin-based colors in birds. (Galván and Solano, 2009). Variations in melanization levels of skin and its protective role against UV radiation in birds have remained however unexplored (Bortolotti, 2006).

Non-signalling functions of melanins must also be considered. Melanins deposited in tegumentary structures such as feathers and bills increase hardness (Bonser, 1995) and resistance of these structures against abrasion caused by airborne particles (Schreiber et al., 2006), although eumelanin and pheomelanin do not seem to differ in their capacity to increase plumage strength (Pannkuk et al., 2010). Melanins also protect feathers from the damaging effects of feather-degrading bacteria (Burt and Ichida, 2004), although they may even serve as a substrate for some species of bacteria (Grande et al., 2004). Melanins in plumage may also be involved in thermoregulatory functions in birds, as the dark colours typically conferred by these pigments absorb more radiant energy than light colours, although there is no clear evidence of the actual importance of plumage melanization in covering the thermoregulatory needs of birds (reviewed in Bortolotti, 2006). The capacity of dark colours to absorb radiant energy may however represent a physiological cost, which may have led to the evolution of compensatory mechanisms (Negro et al., 2006; Bamford et al., 2010). Lastly, studies with birds have increased our understanding of the function of melanins from non-classical melanocytes that are found in internal organs. Thus, the presence of melanin in testes have evolved in species of birds with high mtDNA substitution rates, probably an adaptive response related to the protective capacity of melanins against the oxidative stress generated by mitochondrial dysfunction caused by mutations (Galván et al., 2011a). In the same context, studies with birds have suggested that melanins in internal organs such as the brain may function as protective agents against reactive quinones and toxic metals (Galván and Møller, 2011; Galván et al., 2011b).

Melanins are also present in poikilotherms. The most obvious difference in the mechanism of melanin production between homeotherms and poikilotherms is the fact that in the latter the transfer of melanosomes from epidermal melanocytes to the target structures does not occur; instead dermal melanophores have the ability to disperse and aggregate their melanosomes (Kawasaki et al., 2008). The change in melanization patterns in poikilotherms can thus be a dynamic and rapid process that occurs in response to a diversity of hormonal and environmental stimuli as opposed to homeotherms, in which changes in melanization depend on the turnover rate of the dead keratinized structures where melanosomes are transferred (Fujii, 2000). The melanogenesis pathway of poikilotherms, however, does not seem to greatly differ from that of homeotherms (Cerdá-Reverter et al., 2011). Gallone et al. (2007) described that the liver melanin of amphibians (*Rana esculenta*) is composed of DHI-rich eumelanin showing characteristics very similar to those of *Sepia* melanin, although it contained little pheomelanin. Indeed, up to recently, it was believed that pheomelanin was only synthesized by higher vertebrates (i.e., birds and mammals), but the presence of this pigment has been discovered in the shell of a reptile (the Eastern Hermann's tortoise *Eurotestudo boettgeri*; Roulin et al., 2013) and in the skin of adults and tadpoles of an amphibian (the African dwarf frog *Hymenochirus boettgeri*; Wolnicka-Glubisz et al., 2012).

Given the dependence of poikilotherm vertebrates on the temperature of the environment to conduct their vital processes and the capacity of the dark colours conferred by melanins to absorb radiant energy, it is expected that melanins play a crucial role in the thermoregulation of poikilotherms, as it actually does by facilitating the adaptation of some individuals to cooler environments (Clusella-Trullas et al., 2007). The

universal function of (eu)melanin as a protective agent against UVR has also been investigated in poikilotherm vertebrates, particularly in amphibians, where skin melanization in response to increase in UVR has been shown to be crucial for the survival of larvae (e.g., García et al., 2004). Melanization as a response to increase in UVR ('suntanning') has also been observed in both adults (Adachi et al., 2005) and larval fish (Mueller and Neuhauss, 2014).

Cephalopod ink

A black insoluble eumelanin is the major visible constituent of cephalopod ink, which is produced by the activity of a gland located in the bottom of the ink sac. The ink gland consists of an immature inner portion and a larger mature portion. Pigmentation in ink gland cells is an index of the degree of maturation, a process which culminates in the secretion of melanin and other cell constituents into the lumen of the gland (Derby, 2014). A central role in the regulation of melanogenesis in the ink gland is played by the glutamate/NO/cGMP pathway (Palumbo et al., 2000). Extensive protein tyrosine nitration has been detected in the mature melanin-producing cells of *Sepia* ink but not in the immature unpigmented cells suggesting a role for redox imbalance and control in the maturation process (Fiore et al., 2009).

Tyrosine metabolism is the central biochemical feature of the ink defence system, however the actual significance of this process, with special reference to the biological functions of the melanin in the ink, is still little understood. The occurrence of large amounts of tyrosinase in the ink (Palumbo, 2003) together with another melanogenic enzyme which catalyzes the rearrangement of dopachrome to 5,6-dihydroxyindole (DHI) (Palumbo et al., 1994) was supposed to ensure efficient conversion of catechol constituents into toxic quinones preventing the attack of predators. DOPA and dopamine are also found in the melanin-free fraction of cephalopod ink (Fiore et al., 2004) and can affect squid olfactory neurons (Di Cristo et al., 2007). It can thus be suggested that the melanin granules in the secreted ink serve as sink/carrier for bioactive substances such as dopamine, preventing dilution after ejection and ensuring efficient interactions with target organs. In addition, *sepia* tyrosinase was found to be toxic to a variety of cell lines, including PC12 cells (Russo et al., 2003), leading eventually to an irreversible apoptotic process.

Insects

Melanogenesis is a prominent expression of tyrosine metabolism in insects which depends on fine-tuned systems of enzymes (Sugumaran, 2002; Vavricka et al., 2014), including phenoloxidase and dopachrome (decarboxylating) isomerase, and is mainly related to cuticle sclerotization and innate immune response. Sclerotization is a complex process involving the rapid hardening of the soft and pale freshly produced cuticle to prevent dehydration and to afford protection to eggs and animals at the various levels of development, from larvae to adults. Sclerotization is initiated by the action of phenoloxidase on catecholamine sclerotizing precursors, such as *N*-acetyldopamine and *N*- β -alanyldopamine, to generate their corresponding quinones which in turn, rapidly form adducts not only with the reactive amino acid side chains

of structural proteins, but also with the side chain hydroxyl groups of chitin polymer, which is the major structural component of the cuticle.

Besides its role in cuticle sclerotization, melanogenesis provides an important mechanism in the defence against parasites and pathogens, due to the lack of immunoglobulins and other components of mammalian immune system (Shao et al., 2012; Nappi and Christensen, 2005). The aim of melanogenic reactions is to prevent the uncontrolled growth and the damage caused by the intruder. In response to an alarm signal insect phenoloxidase in the haemolymph is activated and adheres to the invader inducing melanization by quinone formation. Melanin formation then blocks the parasite and prevents its growth. (Clark and Strand, 2013)

Plants, fungi and bacteria

Plants usually incorporate melanins for cell wall strengthening purposes. In contrast to animal pigments, plant melanins are devoid of nitrogen and their color may range from dark brown to totally black, dependings on the nature of the main unit oxidized. Most common precursors include catechol, 1,8-dihydroxynaphthalene (DHN) and a number of acids, such as caffeic, chlorogenic, protocatechuic, and gallic acids (Figure 2), yet their structure is still little understood.

In the sclerotization reactions of plants, polyphenol oxidases are liberated from vesicles in response to injury and, in the presence of oxygen, catalyze the two-electron oxidation of tannins to produce very reactive quinones. Cross-linking by quinones occurs typically through reaction with a nucleophilic group in a protein, such as a thiol or amine, to give a melanoprotein. The result is blackening and hardening at a site of injury on a plant, a reaction known as enzymatic browning.

Increasing interest is currently being devoted to fungal melanins (Nosanchuk and Casadevall, 2003; Langfelder et al., 2003; Eisenman and Casadevall, 2012; Jacobson, 2000) due to their biotechnological importance and their central role in the virulence of plant and human pathogenic fungi. In fungi, melanogenesis occurs during specific developmental stages. Melanin may be synthesized in internal vesicles and then transported to the cell wall where they are likely cross-linked to polysaccharides. Main phenolic precursors of fungal melanins include (DHN), which is typical of Ascomycetes, and γ -glutaminy-4-hydroxybenzene and catechol (Dunn, 1986), characteristic of Basidiomycetes.

Some fungi can synthesize two types of melanin. The neuropathogenic basidiomycetous *Cryptococcus neoformans* (Casadevall et al., 2000) can attack human brain and contains a phenol oxidase capable of oxidizing dopamine and norepinephrine to NM-like polymers (Polacheck and Kwon-Chung, 1988) which can increase the virulence of the infection. The same fungus oxidizes homogentisic acid (HGA) to synthesize pyomelanin in the form of an insoluble fluorescent material that can likewise increase the virulence. *Aspergillus fumigatus*, an airborne fungal pathogen in immunosuppressed humans, is also able to produce DHN-melanin or pyomelanin starting from L-tyrosine through HGA (Schmaler-Ripcke et al., 2009), while *Aspergillus nidulans* can form DHN-melanin and DOPA-melanin (Goncalves et al., 2012).

Melanins are also found in bacteria, where they were initially identified in tyrosinase-containing *Streptomyces*, *Marinomonas mediterranea* (Lopez-Serrano et al., 2004) and *Bacillus thuringiensis* (Patel et

al., 1996). This topic has been the subject of much interest also in view of the potential applications and extensively reviewed (Riley, 1997; Solano, 2014; Dadachova and Casadevall, 2008; Plonka and Grabacka, 2006). Bacterial melanins usually contain nitrogen, although non-nitrogenous variants from catecholic precursors have been identified. *S. marcescens* (Trias et al., 1989) can produce a brown pigment by oxidation of 3,4-dihydroxyphenylacetate, as a means of counteracting accumulation of this catabolic intermediate. Some *Pseudomonas* produced a pyomelanin from 4-hydroxyphenylpyruvate. Besides pathogenesis, melanin synthesis in microorganisms is part of the defense systems against environmental stresses such as UVR and oxidizing agents, and more in general against the consequences of environmental cues. It is relevant in this connection that after the Chernobyl disaster black and highly melanized fungi responded to ionizing radiation with enhanced growth (Dadachova and Casadevall, 2008).

Properties of melanins and control of melanogenesis

Melanins and the related process of melanogenesis have been shown to serve a variety of roles and functions that depend on the hosting system and its environment. Although these roles are often unknown or at best speculative, they depend either on the specific set of physicochemical properties of melanins or on the chemical and biochemical nature of the precursors and intermediates in the underlying pathways. Such a relationship is summarized in Table 1.

Biological control of melanogenesis

Biological control of melanogenesis and melanin pigmentation in living organisms derives from the interplay of different factors and mechanisms. For example, melanin-forming enzymes are not functional unless they enter the melanosomes and become activated in the endoplasmic reticulum (Lamoreux et al., 2010). Correct control over this process is necessary for normal pigmentation and to prevent activation of toxic melanogenesis-related pathways outside the melanosome.

Melanin-type. Within melanocytes, the receptor for α MSH (melanocortin 1 receptor, Mc1r) encoded by the *extension* locus is the major determinant of the type of pigment formed in mouse coat. Activation of Mc1r stimulates eumelanogenesis, whereas loss-of-function mutations at *extension* are associated with pheomelanin coats (Robbins et al., 1993). Likewise, human MC1R regulates the amount and type of pigment formed in human skin, with the wild-type receptor promoting eumelanogenesis whereas variants with impaired signaling are associated with pheomelanogenesis and a red hair color phenotype (García-Borrón et al., 2014).

Activation of the MC1R by its endogenous agonists α MSH and ACTH stimulates tyrosinase activity by transcriptional stimulation of *Tyrosinase* gene expression as well as by post-transcriptional mechanisms. The molecular events leading to upregulation of *Tyrosinase* gene expression downstream of MC1R are relatively well understood, and rely on activation of the cAMP signaling pathway and the Microphthalmia transcription factor. MC1R signaling also triggers a cAMP-dependent increase of the melanosomal pH from acidic to near-neutral values, which enhances the catalytic efficiency of tyrosinase (Cheli et al., 2009). Surprisingly,

little is known about possible effects of MC1R signaling on the intracellular pool of low molecular weight thiols in mammalian melanocytes. However, it has been shown that cysteine depletion promotes eumelanogenesis in human melanoma cells (del Marmol et al., 1996). Conversely, high levels of cysteine in the culture media of mouse melanocytes, coupled with tyrosinase inhibition with phenylthiourea in the presence of the agouti signaling protein (ASIP) dramatically increase the pheomelanin/eumelanin ratio, up to 200-fold (Hida et al., 2009).

Treatment of normal human melanocytes with α MSH or ACTH also increases expression of the other melanogenic enzymes, the tyrosinase-related proteins TRP1 and TRP2 (Abdel Malek et al., 1995). Given that the 3 proteins of the tyrosinase family most likely do not undergo identical fold-stimulations, it is conceivable that the ratios of their enzymatic activities would not be the same in resting versus stimulated melanocytes. These variations may have a subtle and still uncharacterized impact on the final structure of the pigment, for instance through the modification of the ratio of carboxylated monomeric units incorporated to eumelanins.

MC1R signaling is regulated by at least two endogenous ligands in addition to α MSH and ACTH. In mice, ASIP blocks the stimulatory effect of α MSH on Tyrosinase, and decreases *TRP-1* and *TRP-2* gene expression and total melanin production (Hida et al., 2009, Le Pape et al., 2009). In fact, ASIP behaves as a potent inverse agonist (Siegrist et al., 1997), since it decreases agonist-dependent signaling as well as the basal constitutive activity of the receptor (Sánchez-Mas et al., 2004). The effects of ASIP on cultured normal human melanocytes are similar (Suzuki et al., 1997, Swope et al., 2012), and polymorphisms in human *ASIP* are associated with skin, hair and eye pigmentation phenotypic changes (Bonilla et al., 2005; Kanetsky et al., 2002). A third MC1R ligand, the small secreted protein β -Defensin-3 (BD3), binds MC1R with nanomolar affinity (Candille et al., 2007). Human BD3 behaves as a neutral antagonist unable to activate cAMP formation in normal human melanocytes, but efficiently blocking α MSH-induced cAMP production and activation of tyrosinase (Swope et al., 2012).

The knowledge of the structure and function of MC1R ligands, and of the signaling pathways that they engage to regulate melanin pigmentation and non-pigmentary defensive mechanisms against UVR-induced cellular damage may direct the rational design of small peptides with potential pharmacological applications (Abdel-Malek et al., 2009), and of receptor-independent strategies to achieve physiological responses such as tanning (D'Orazio et al., 2006).

Hair graying. Gray hair condition varies between individuals, worldwide, with various ages of incidence and distinct progression in magnitude (Panhard et al., 2012). The graying of hair is due to a progressive depletion of melanocyte stem cells (MSC) in hair follicles, leading to the defect of the hair pigmentation unit renewal at the telogen to anagen transition during the hair cycle, and to the growth of white hair, eventually (Commo et al., 2004a, Nishimura et al., 2005). It is noteworthy that the exhaustion of bulb melanocytes is progressive and during the graying of hair, transitional gray hairs show a reduced, albeit not null, melanocyte number in the bulb with reduced melanin content in the hair shaft, as compared to fully pigmented hair (Commo et al., 2004a).

Following the demonstration that hair graying was due to MSC depletion, Arck (Arck et al., 2006) showed a selective oxidative stress sensitivity of hair follicle melanocytes compared to other hair follicle cells, namely keratinocytes and fibroblasts, along with increased mitochondrial DNA deletion in the bulb of gray hair.

Genetic mutations in mice also pointed out the particularly high susceptibility of the MSC to oxidative stress-associated DNA damage leading to premature hair graying and furthermore highlighted determinant genes for coat pigmentation maintenance in mice, e.g. BRCA1, ATM, Bcl2 (Cao et al., 2003; Inomata et al., 2009; Yamamura et al., 1996). Although of strong interest as mechanistic approach, there is still no evidence whether imbalance in the activity of these genes is involved in natural hair graying in humans.

Of interest, human hair follicle melanocytes (Caucasians, Asians and Africans) do not express the melanogenic enzyme dopachrome tautomerase (DCT) or TRP2, as opposed to epidermal melanocytes (Commo et al., 2004b). Despite the lack of DCT, not all hairs are gray. Actually, hairs on a given scalp are different and behave differently. These differences could also impact melanocyte stem cell (MSC) maintenance and hair graying, as detailed below, in addition to the lack of DCT. Depending on the scalp areas, distinct hair follicle environments would impact differently MSC fate/maintenance, e.g. distinct stimuli or distinct stress sources. Moreover, hairs do not have hair cycles of the same duration, do not have the same number of cycles through life, and the number of MSC is various when comparing hair follicle from the same individual and between individuals (similarly, in mice the distribution of melanocytes in hair is different depending on the body area, e.g. belly vs back, due to cell migration process at during development).

Beside its function in melanin synthesis, DCT provides a beneficial effect to melanocytes exposed to oxidative stress (Michard et al., 2008a). Likewise, cyto-protective property of DCT was demonstrated in other distinct models (Michard et al., 2008b; Sendoel et al., 2010). Hence, the specific lack of DCT in hair follicle melanocytes likely contributes to their particular susceptibility to deleterious conditions, and the observation that DCT expression could decrease with age in human hair, further strengthens a possible relation between DCT expression and hair graying (Commo et al., 2012). Interestingly, in vitro comparison of hair follicle melanocytes isolated from young and old donors suggested a decrease of catalase expression in melanocytes in the elderly (Kausser et al., 2011).

Chemical control of melanogenesis

Both eumelanin and pheomelanin derive from the common precursor dopaquinone formed by oxidation of L-tyrosine by tyrosinase (Figure 3; Raper, 1927; Cooksey et al., 1997). Intramolecular cyclization of dopaquinone then produces cyclodopa which, following redox exchange with dopaquinone itself, gives dopachrome and dopa, respectively. Dopachrome then rearranges to generate mostly DHI and to a lesser extent DHICA (Edge et al., 2006) which are then further oxidized and polymerized to produce eumelanin. Sulfhydryl compounds (such as cysteine), if present at sufficient levels, may deviate the normal course of the pathway to give thiol adducts of dopa, *i.e.* 5-S-cysteinyl-dopa (5SCD) along with a minor isomer 2-S-

cysteinyl-dopa (2SCD) (Ito and Prota, 1977). Further oxidation of these thiol adducts leads to the formation of 1,4-benzothiazine intermediates which are then converted to pheomelanin (Figure 3).

Tse et al. (1976) showed that the addition of sulfhydryl compounds proceeds very quickly to generate thiol adducts. Reduction to the parent catechols through redox exchange proceeds as quickly as thiol addition (Tse et al., 1976), and therefore, these two reactions are competitive (Fig. S1). The redox reaction proceeds with reducing agents (QH₂) such as ascorbic acid, DHI, and 5SCD in biological system. Intermolecular reactions with amine compounds do not proceed as quickly. However, in cases when the amino group is present within the same molecule, the amino group rapidly undergo either an intramolecular addition reaction to give an aminochrome (such as dopachrome) or an intramolecular condensation reaction to give an *o*-quinonimine (such as a cyclization product from cysteinyl-dopaquinone, see Figure 3, Figure S1). It should be emphasized that all of these reactions are controlled by the intrinsic *chemical* reactivity of *o*-quinones. The high reactivity of *o*-quinones may result in their strong cytotoxicity (Riley, 2003). This subject is beyond the scope of this review, but it should be mentioned that cysteinyl residues of proteins are able to bind to *o*-quinones of biological interest such as dopaquinone and dopamine-quinone (Ito et al., 1988). It is thus possible that sulfhydryl enzymes may be inactivated by *o*-quinones in the course of melanogenesis.

How dopaquinone controls the course of mixed melanogenesis has been discussed in detail (see Ito and Wakamatsu, 2008). Kinetic studies suggested a three-step pathway for mixed melanogenesis (Ito, 2003): the initial stage is the production of cysteinyl-dopas, which continues as long as the cysteine concentration is above 0.13 μ M. The second stage is the oxidation of cysteinyl-dopas to produce pheomelanin, which continues as long as cysteinyl-dopas are present at concentrations above 9 μ M. The last stage is the production of eumelanin, which begins only after most cysteinyl-dopas and cysteine are depleted. Therefore, the ratio of eumelanin to pheomelanin is determined by tyrosinase activity and the availability of tyrosine and cysteine in melanosomes (Land et al., 2003). A “crossover value” for switching between eumelanogenesis and pheomelanogenesis has been determined *in vitro* when cysteine concentration reaches 0.8 μ M.

Recently, Greco et al. (2011) provided chemical evidence supporting the casing model for mixed melanogenesis, by showing that 5SCD-melanin oxidizes dopa into a black insoluble polymer (eumelanin) which encapsulates the active cysteinyl-dopa-melanin core (pheomelanin). Although this model certainly stands true for the synthesis of NM in the substantia nigra of the brain where dopamine and cysteine act as precursors (Bush et al., 2006, Wakamatsu et al., 2012b), it is less clear whether it can be applied to the mixed melanogenesis in melanosomes where tyrosine (not dopa) and cysteine act as precursors (Ozeki et al., 1997).

Physical properties of melanins

Supramolecular structure and aggregation. Until very recently no consensus has been reached on the precise nature of melanins' physical structure. The nature of the supramolecular structure of eumelanin has been addressed by several studies (Arzillo et al., 2012, Clancy and Simon 2001, Liu and Simon 2003a, Liu and Simon 2003b, Reale et al., 2012, Capozzi et al. 2006, Ghiani et al., 2008, Stark et al., 2005, Kaxiras et al.,

2006, Meng and Kaxiras, 2008). At the molecular level, cross-linking of eumelanin basic building blocks can occur at a number of sites – and this confers upon the melanin structure a degree of disorder which gives rise to many of its physical and chemical properties. This model of chemical disorder, that does not exclude the possible presence of preferential aggregation and aggregation-dependent functional properties in the eumelanin biopolymer (Panzella et al., 2013), is now generally accepted. As proposed by Meredith and colleagues (Watt et al., 2009), accepting the simple proposition that melanin structure is defined by disorder at all levels controlled by the feedstock and synthetic environment, then the question of its precise structure would become moot. However, based on monomer supply and experimental conditions, the degree of disorder at the various levels may vary, which may entail a certain level of control over melanin properties and functions.

Chemical disorder has implications for higher levels of structure within the system. Watt et al. (2009) recently produced compelling electron microscopy evidence that despite the primary-level disorder, melanins both natural and synthetic can self assemble into stacked structures defined by hetero-aromatic non-covalent interactions with characteristic inter-plane spacing of $\sim 3.7\text{\AA}$. This work confirms earlier propositions (both theoretical and partially observed with atomic force microscopy) that indolic oligomers could self assemble in this fashion. The extent of the sheets seems dependent upon the synthetic environment – in natural melanins the secondary sheets can be many hundreds of nanometers if not microns in size, whilst in synthetic melanin they are much smaller and limited by the relatively poor solvation environment, suggesting that solvophobic interactions drive sheet stacking and subsequent aggregation. Watt et al. (2009) and Bothma et al. (2008) went on to show that the optical absorption of melanin is related to the monomer composition and mode of coupling since de-stacking the sheets did not influence the shape of the optical absorption. Armed with this knowledge, they also used de-stacking methods to create device quality, solution processable melanin thin films. It is also worth noting that a number of studies point to the similarities between the structural and optical features of eumelanin and sp^2 partially disordered carbon materials, such as carbon black (Jastrzebska et al., 2010, Gallas et al., 1999).

Hydration. The historical model of hydration structure of melanin granules envisages both “strongly” and a “weakly” bound water fractions (Bridelli and Crippa, 2010). The removal of water possibly produces irreversible structural modifications, affecting eumelanin properties. The dried material exposed again to water does not easily swell and regains only part of the water originally bound (Crippa et al., 1989).

Thermogravimetric analyses (TGA) showed that Sigma eumelanin loses weakly bound water below 125°C and strongly absorbed water at about 125°C . Above 250°C , it decomposes with release of carbon dioxide. TGA on Sigma eumelanin powder samples indicate that weight loss up to 140°C , mainly attributable to water, ranges from $11.6\pm 1.0\text{ wt\%}$ at 60% RH to 16.8 ± 0.7 at 90% RH (Wünsche et al., 2015)

Paramagnetic properties. The presence of unpaired electrons in melanin is one of the key features of the pigment that determines some of its basic physicochemical properties. Based on results of a thorough study, in which an array of complementary physical methods were employed, Mostert et al. (2012a) have concluded that melanin is not an amorphous semiconductor as conventionally believed. The authors demonstrated that

upon absorption of water by dehydrated synthetic eumelanin free charge carriers were produced. The carriers – extrinsic free radicals (electrons) and hydronium ions (protons) – resulted from the comproportionation reaction of the melanin basic units (Figure 5).

The second study by Mostert et al. (2012b) has shown that the solid-state EPR signal of a synthetic eumelanin is dominated by a carbon-centered radical (g-factor 2.0032) with a significant contribution from a semiquinone radical (g-factor 2.0045) from the comproportionation reaction. Although the actual structure of melanin radicals is unknown, EPR analysis allowed to distinguish signals in melanins that derive from species in which the spin density is localized on carbon atoms, and from species which are oxygen-centered radicals, with the spin density being localized on oxygen atoms (semiquinone radicals). The spin density of these radicals can be changed by the addition of either water or base, which increases the concentration of semiquinone radicals. On the other hand the water-dependent decrease in the content of the carbon-centered radical was explained by its destruction due to the wet environment. Such a destruction of the carbon-centered radicals, which are normally protected from extrinsic reactants being located in the center of the stacked oligomeric units, could be promoted by a destacking mechanism induced by extra negative charge carried by the semiquinone radicals, which disrupts the π - π stacking arrangements. As a result, the carbon-centered radical becomes more exposed to the aqueous environment.

The spin chemistry of eumelanin is probably best illustrated by a time-resolved EPR (TR EPR) study to monitor the photochemistry of radical pairs from melanin in porcine retinal pigment epithelial cells on the sub-microsecond time scale (Wang et al., 2009). Two distinct radical pair species were seen: one of enhanced absorption/emission pattern at early times and one mostly emissive at later times. Interestingly, while the early time signal can be satisfactorily simulated by assuming a population coming exclusively from the singlet excited level, the later time component to the spectrum suggests the triplet mechanism of spin polarization. Excited triplet states play important role in photochemistry of many compounds, including variety of photosensitizers. This is because of relatively long lifetime of triplet excited states, in comparison with lifetimes of singlet excited states, which increases the probability of a chemical reaction mediated by triplet excited states to occur. Based on previous studies, in which time-resolved techniques were used such as laser flash photolysis and photoacoustic spectroscopy, no involvement of melanin excited triplet states has been detected suggesting that virtually all of the energy by melanin absorbed photons was safely dissipated into heat. The suggested involvement of a triplet species in the observed electron spin polarization phenomena in eumelanin is intriguing because it is not consistent with the widely accepted view of melanin photodynamics, but its actual significance for light-induced processes in melanins depends on the quantum yield value of such a process. Thus, although the TR EPR signals in porcine RPE cells are small and do not challenge the accepted view that the majority of light energy absorbed by melanin is efficiently dissipated into heat, the role of such minority process in melanin photochemistry and phototoxicity remains to be assessed.

Electrical properties. Eumelanin in the solid-state is a hybrid conductor (Mostert et al., 2012a), i.e. it can sustain both electronic and ionic currents. The electrical conductivity of melanin films and pelletized

powders is strongly dependent upon the state of hydration of the material. Mostert et al. (2012a) showed that for a DOPA-derived synthetic eumelanin the system shows an almost “percolation-like” transition at between 10-15% by weight water wherein the electrical conductivity increases in a super-exponential manner (Figure 6). The origin of this behaviour is a process now referred to as “chemical self-doping” where the comproportionation equilibrium reaction drives the release of protons at the catechol sites. These protons are transported through the material via the Grotthuss mechanism (“concerted transfer of protons within extended hydrogen-bonded water chains”, Kreuer et al., 2004) with the adsorbed water acting as the conducting matrix. In particular, each oxygen atom of the water molecules simultaneously passes and receives a proton through the cleavage and formation of covalent -O-H bonds.

It has been proposed that the mechanism could be generic in hydrophilic conducting bio-macromolecules, and that furthermore, the phenomenon could be utilized to create ionic circuitry and even ion-to-electron transducers for use in primary bioelectronic interfaces (Meredith et al., 2013, d’Ischia et al., 2009). A similar process has been recently observed in ionic field effect transistors utilizing chitosan channels (Meredith et al., 2013).

Melanins have also been observed to be “photo-conductive” in the solid-state, i.e. their resistance decreases under illumination with UV or visible light (Mostert et al., 2012a, Meredith et al., 2013). The effect has a time constant of order 10s of seconds with pronounced hysteresis. The magnitude of the photo-current is dependent upon the state-of-hydration of the material and there is an order of magnitude difference in the effect above and below the conductivity transition described above (Mostert et al., 2012b). Given the influence of water and the characteristic response times, Mostert et al. have suggested that the photo-conductive effect is also related to the comproportionation equilibrium with light acting to drive oxidation of the hydroxyquinone to semiquinone – in the process freeing mobile protons.

Optical properties. Melanins have characteristically broad, monotonic optical absorption (Tran et al., 2006). This was originally attributed to its semiconductivity, but more recently the “chemical disorder” model has become accepted as an appropriate description (Meredith and Sarna, 2006)-see however different views on melanin disorder. In this model, the featureless absorption is derived from the superimposition of individual inhomogeneously broadened chromophore absorptions resulting from the diversity of chemical species (Meredith et al., 2006). Melanins also demonstrate near unity conversion of absorbed photons into vibrational energy –photoluminescence quantum yields (PLQYs) of < 0.1% (Meredith and Riesz, 2005) –and this is at the core of their role as photo-protectants. Although melanins exhibit variable degrees of photostability (Sarna et al., 2003), in principle eumelanins could be suited for the absorbing component within an optically-driven transducing element for bioelectronics. The actual photostability of melanin depends very much on the exact experimental conditions, such as hydration of melanin, pH, presence of redox-active metal ions, oxygen concentration and the supermolecular structure of the pigment granules. It also depends on the type of melanin with pheomelanin being more photolabile than eumelanin. Indeed, a

recently published paper (Ito et al., 2013) reports that a distinct oxidative photodegradation of melanin occurs even in the human retinal pigment epithelium.

Chemical properties of melanins

A major chemical property of eumelanins is its redox behavior. Multiple oxidation states can be demonstrated by electrochemical reduction with Ti(III) and oxidation with Fe(III) or oxygen. Eumelanins can also catalyze the reduction of Fe (III) by NADH. Redox properties of eumelanins are usually attributed to shuttling of indole units between the catechol and quinone states through semiquinone free radicals. Moreover, eumelanin can either oxidize or reduce metals with single-electron transfer and free radical generation, including ROS, as in Fenton-type reactions (Zecca et al. ,1994, 2008a).

The redox properties of a synthetic eumelanin from 5,6-dihydroxyindole (DHI) have been characterized by means of an organic electrochemical transistor (OECT), a powerful tool for biosensing, bioelectronics and nanomedical applications (Tarabella et al., 2013). Gate current measurements on fine aqueous eumelanin suspensions suggested evolution of the polymer from a far-from-the-equilibrium redox state toward a more stable electronic arrangement promoted by redox exchange with the gate electrode.

As anticipated in previous sections (see Human brain), metal chelation is one of the properties of eumelanin that have important consequences on its biological functions (Meredith and Sarna, 2006, Hong and Simon, 2007; Liu et al., 2004; Chen et al., 2009; Sarna et al., 1980; Froncisz and Sarna, 1980; Hong and Simon, 2006; Samokhvalov et al., 2004). Eumelanin has a strong binding affinity for Fe(III) and other heavy metal ions such as Cu(II), Sr(II), Cd(II), La(III), Gd(III), Pb(II). In contrast, lighter metal ions including Ca(II), Mg(II), Zn(II), Na(I), K(I) bind less strongly to eumelanin (Hong and Simon, 2007, Liu et al., 2004, Chen et al., 2009, Sono et al., 2012). Studies based on EPR (Sarna et al., 1980, Froncisz and Sarna, 1980), IR absorption (Chen et al., 2009, Hong and Simon, 2006), and Raman spectroscopy (Samokhvalov et al., 2004) revealed that the binding site depends on the type of metal ion, its concentration, and pH. Recently, specific dimers, trimers, and tetramers of DHI have been identified as potential bi-, tri-, and tetradentate metal chelators (Meng and Kaxiras, 2008, d'Ischia et al., 2011).

Eumelanins can also serve as free radical scavengers and antioxidants via H-atom transfer. In vitro studies have shown that DHICA-melanin exhibits potent hydroxyl radical-scavenging properties in the Fenton reaction, whereas DHI melanin does not. (Pezzella et al., 1996; Jiang et al., 2010) DHICA melanin exhibits more potent free radical scavenger properties than DHI and dopa melanins in three different assays, the DPPH, ABTS and NO scavenging assays (Panzella et al., 2013). Photophysical studies on oligomers from DHI and DHICA have shown that the excited state lifetimes increase on passing from DHI monomer to a dimer, but decrease on passing from DHICA to its dimers and trimers, suggesting more efficient UV dissipation mechanisms in the latter case. (Gauden et al., 2008; Corani et al., 2014)

These and other observations can be rationalized considering the marked differences in the structural, chemical and aggregation properties of DHI and DHICA melanins (Figure 7): whereas the former consists of planar and relatively stacked components, the latter is less efficiently aggregated, is less stabilized by

electronic delocalization and is more prone to react with free radical species. Both antioxidant and energy dissipation properties of DHICA melanin offer a plausible chemical explanation as to why nature selected DHICA instead of the more pigmentogenic DHI as main eumelanin building block.

As mentioned in previous sections, eumelanins can bind a range of drugs such as methotrexate, chlorpromazine, cocaine (Bridelli et al., 2006); amphetamines; doxorubicin, quinidine, disopyramide, and metoprolol (Buzman and Rozanska, 2003). Highly specific melanin-binding sites for iodobenzamides (Moreau et al., 2005) can be exploited to diagnose and stage melanoma by using radiolabeled iodine-containing compounds.

Thioureylene derivatives such as propylthiouracil can be incorporated as false thioureylene-based melanin precursors into eumelanins during synthesis and have been proposed as melanoma seekers for both diagnosis and therapy of the tumor (Napolitano et al., 1996b; Palumbo et al., 1997; Larsson, 1993; Mars and Larsson, 1995). Oxidative breakdown of the pigment by exposure to strong alkali (bleaching) has been extensively studied because of its practical importance in cosmetics (Ghiani et al., 2008, Liu et al., 2003). The degradation process apparently involves a fast solubilization of the pigment (granules disruption), followed by a slower bleaching, eventually resulting in a pale yellow solution (Wolfram et al., 1970; Korytowski and Sarna, 1990; Borges et al., 2001; Ghiani et al., 2008).

Melanosome architecture and surface properties

An extensive and detailed coverage of melanosome biology is provided in a recent book and various reviews (Borovansky and Riley, 2011; Simon et al., 2008; Solano, 2014). The surface of the intact melanosome plays a critical role in the binding and sequestering of metal cations and determines its functional capacity to mitigate oxidative stress and to participate in photo-generation of ROS (Hong and Simon, 2007; Simon et al., 2008). Imaging studies of red and black human hair melanosomes indicate that the morphology of the two differs, suggesting that the three-dimensional structure of the melanosomes is influenced by its constituent melanin (Liu et al., 2005). More importantly, the photoionization thresholds of pheomelanin and eumelanin are markedly different (326 nm and 282 nm, respectively) (Samokhvalov et al., 2005), suggesting that pheomelanin can induce ROS formation upon exposure by UVA light while eumelanin can not (Wenczl et al., 1998; Takeuchi et al., 2004). Thus, the morphology of melanosomes with mixed pigment composition, like those commonly found in the skin and eyes may be of interest in relation to the possible presence of photosensitizing pheomelanin exposed on the organelle surface.

In this connection, the “casing” model (pheomelanin encased by eumelanin), proposed also for NM pigments of the human brain (Bush et al., 2006, 2009; Ito, 2006; Zecca et al., 2008a), has received indirect support by several other studies (delMarmol et al., 1996; Ozeki et al., 1997; Smit et al., 1997). Because different photoionization thresholds characterize the two pigments, spatially-resolved imaging of the ionization thresholds of the melanosome surface can uniquely determine whether or not both pigments comprise its surface. Such studies have been reported on melanosomes isolated from the stroma of human irides, containing eumelanin:pheomelanin ratios varying from 14.8 to 1.3 (over an order of magnitude change, dark

brown to blue-green irides, respectively) (Peles et al., 2009). The imaging analyses of these melanosomes establish that only eumelanin is present on or near their surfaces (Liu et al., 2005, Peles et al., 2009). These data provide direct evidence for a casing model (Figure 4) . The finding that pheomelanin is encapsulated by an excellent electron scavenger (eumelanin) in natural systems would suggest that the structure of the melanosome and NM pigments mitigates the adverse photochemical properties of pheomelanin. Scanning electron microscopy and atomic force microscopy studies show that the intact melanosome as well as NM pigments is made up of ~30 nm diameter spherical substructures (Bush et al., 2006), whereby changes in the relative amounts of eumelanin and pheomelanin would be manifested by changes in the diameter of the core and the thickness of the outer eumelanin coat. Damage to this coating, and or significant reduction in the amount of eumelanin present could jeopardize the protective ability of eumelanin, providing mechanism(s) for the exposure of pheomelanin and consequently contributing to oxidative stress.

Surface properties of melanosomes can also be examined using nano-indentation techniques aimed at probing changes in the “stickiness” of the surface through the tip used in AFM measurements (Guo et al., 2008). Studies of the changes in the surface properties of human RPE melanosomes as a function of age (Rozanowska et al., 2002) showed that the aerobic reactivity of RPE melanosomes increases with age: thus, photoinduced oxygen uptake is a factor of 2.4 greater at 80 years of age than at 40 years at age (Boulton, 1991). In addition, RPE melanosomes exhibit significant changes in their photophysical properties with ages as well (e.g., the fluorescence yield increases with age. Nano-indentation experiments combined with spatially-resolved imaging of ionization thresholds (Hong et al., 2006) quantitatively established age-related changes in the photophysical and photo-oxidative properties of RPE melanosomes, likely related to the adhesion of lipofuscin to the melanosome surface. Several conditions have been shown to produce alteration in melanosome shape e.g. pMel inactivation (Hellstrom et al., 2011).

Applications

Dermocosmetic applications.

Dermocosmetic applications of melanins and melanogenesis include mainly the use of the melanin pathway to control skin coloration (Solano et al., 2006; Guerrero, 2012), the use of the pigment from various sources in skin photoprotection formulations, the use of melanin precursors for hair dyeing and the development of novel strategies for hair recoloration.

Hair repigmentation. As described previously, hair follicle melanocytes appear deficient in proteins protecting melanocytes from oxidative stress, particularly in the elderly. This particular phenotype likely contributes at least in part to the specific hair follicle melanocyte exhaustion that occurs during hair graying. These and previously reported observations suggest new approaches in anti-hair graying aiming at protecting melanocyte stem cells (MSC) from oxidative stress and DNA damage. Although the use of antioxidants may appear efficient to reach the goal, the demonstration that numerous antioxidants, e.g. phenolic compounds, flavonoids, vitamins C and E, act as limiting factors for melanin synthesis through

various properties, e.g. copper chelators and tyrosinase alternative substrates both acting as tyrosinase inhibitors, reducers of melanin synthesis intermediates leading to reduced melanin polymerization (Briganti et al., 2003, Smit et al., 2009), make this approach neither easy nor obvious to get a satisfactory effect. Alternatively, the demonstration that the bulb of gray hair (by contrast to the bulb of white hair) still contains active melanocytes (actual synthesis of melanin) albeit reduced in number (Commo et al., 2004a), suggests that the use of melanin synthesis stimulating compounds may lead to the correction of gray hair. Indeed, this approach would require fighting hair graying at a time where residual bulb melanocytes remain active in gray hairs. Likewise, acting on the genes shown to be involved in melanocyte stem cell maintenance e.g. BRCA1, ATM, and Bcl2 appears hardly realistic to correct hair graying in humans. This remains a most challenging area of applied pigment cell research of considerable scientific, economic and industrial relevance.

Of interest, human hair follicle melanocytes do not express the melanogenic enzyme DCT, as opposed to epidermal melanocytes (Commo et al., 2004b). This was observed in hair of various origins (Caucasians, Asians and Africans). Interesting enough however, it was shown that DCT is expressed in melanocytes in eyelashes (Thilbaut et al., 2010). There are different points demonstrating that hairs on a given scalp are actually different to each other and behave differently. These differences could also impact MSC maintenance and hair graying, in addition to the lack of DCT.

Hair dyeing with melanin precursors. The color of human hair comes from melanin, but the ability to produce melanin in the hair follicle gradually decreases with age. Loss of melanin pigments is thus associated with aesthetical issues that have considerable impact on the market, such as hair graying. Conventional hair dyes, especially oxidative hair dyes, are very common and convenient, so have been used for several decades. They have still some inconvenience and liability, however, such as color fading, hair damage, and complicated handling. Melanin pigments, both natural and synthetic, are too large in size to penetrate into hair, so, they cannot be used as a dyestuff. A useful group of bioinspired ingredients for dyestuffs is provided by melanin precursors, because they can be converted to natural-looking pigments by air oxidation without enzymes or other oxidants, such as hydrogen peroxide, and their sizes are small enough to penetrate into hair. Since the 1960s a considerable number of patents has reported the use of eumelanin precursors for hair dyeing. Hair dyeing with DHI and related compounds, e.g. diacetoxyindole or DHICA, although commercially very appealing, is methodologically challenging because of the difficulty to obtain a natural hue in the final dye, the relatively low penetration and affinity for hair and, as a main problem, the poor oxidative chemistry with hydrogen peroxide.

A valuable biotechnological approach to cover gray hair with melanin precursors has been developed using *Aspergillus oryzae*, a fungus widely used in Japanese traditional industries, especially brewing of *Sake* (alcohol), and *Shoyu* (soy sauce), and recently shown to contain a strong tyrosinase activity (Koike and Hata, 2010; Nakamura et al., 2012). Pure DOPA (>99%) extracted and purified from a plant was conveniently used for the production of 5,6-dihydroxyindole (DHI) as the melanin precursor for hair dyeing using *Aspergillus*

tyrosinase. The basic formulation for the dyeing test on Japanese or Chinese gray hair includes ammonia or ethanolamine 0.5-3% as alkaline components and melanin precursors (0.1-0.5%) (Koike and Ebato, 2013). As indices of hair damage, formation of cysteic acid and release of a surface lipid, 18-methyleicosanoic acid (18-MEA), can be measured by gas chromatography and/or high performance liquid chromatography (Jones and Rivett, 1997; Rogers and Koike, 2009). The dyeing ability of the DHI-based formulation is higher at alkaline pH, with a concentration dependence up to approx. 0.5%. The formulation leads to lower levels of hair damage than with traditional oxidative hair dyes and the skin staining level is lower than with direct dye systems (Koike and Ebato, 2013).

Pigmentation in human cutaneous melanoma as a basis for targeted therapies

Melanins and the pathway of melanogenesis have long been regarded as a possible means of achieving targeted therapy for eradication of melanoma, and have been used as markers of melanoma progression, particularly 5-S-cysteinyl-dopa (Rorsman et al., 1973; Rorsman et al., 1983; Wakamatsu et al., 2002) and isomeric 5-methoxy-6-hydroxy and 5-hydroxy-6-methoxyindol-2-carboxylic acids (Matous et al., 1980). A most notable example is the boron neutron capture therapy based on the uptake of a DOPA analogue, 10B-p-boronophenylalanine. This approach has long been investigated for specific and effective killing of melanotic and amelanotic melanoma, including brain metastasis (Iwakura et al., 2002). In the latter case the therapy was made effective by enhancing melanogenesis via tyrosinase gene-transfer activation (Hiratsuka et al., 2000). Generation of melanoma selective cytotoxic agent by tyrosinase mediated activation of phenol compounds to reactive quinones has also been pursued as strategy for chemotherapy of melanomas (Riley et al., 1997; Yamada et al., 1991; Jimbow et al., 1993).

TRP1 (Journe et al., 2011), TRP2 (Mandrizzato et al., 2006) and pigmentation in sentinel lymph nodes (van Lanschot et al., 2014) were found to correlate with poor clinical outcome. These findings seem apparently in contradiction with the fact that these markers as well as other melanosomal proteins are considered as differentiation markers. In any case, even if pigment production can be impaired in melanoma both in vitro and in vivo at various stages of the disease, it seems that the ability to reactivate melanogenesis remains intact (Pinner et al., 2009).

After decades of dormancy, recent therapeutic advances in melanoma have been prompted by the discovery of the genetic bases of this disease, pointing to a range of potential targets for therapy, including chiefly *BRAF* mutations (Flaherty et al., 2012; Li et al., 2012; Dong et al., 2012; Dynodt et al., 2013). Altogether, these data support that the evaluation of pigmentation (e.g. pigment, markers and regulators) may also be valuable for the management of melanoma patients.

Polydopamine-based multifunctional coatings and films

Inspired by the adhesion properties of mussel byssus proteins, involving a complex balance of quinone-conjugation reactions at dopa-rich sites of mussel foot proteins (Waite and Tanzer, 1981; Waite, 2008),

Messersmith and colleagues reported in 2007 a universal eumelanin-like coating material, which was produced by the spontaneous oxidative polymerization of dopamine at pH 8.5 in the presence of oxygen (Lee et al., 2007). The resulting material, commonly referred to as polydopamine (PDA), forms highly adhesive polymeric films which can coat many types of surfaces, including superhydrophobic ones. Several studies addressed the nature and main structural features of PDA (see for example; Hong et al., 2012; Della Vecchia et al., 2013; Liebscher et al., 2013). The current view is that PDA consists of oligomeric scaffolds exhibiting a three-component structure comprising uncyclized amine-containing units and typical eumelanin units, namely DHI units as well as pyrrolecarboxylic acid moieties derived from the oxidative breakdown of indole units (d'Ischia et al., 2014) (Figure 8).

PDA adhesion properties depend on the preparation conditions, (Bernsmann et al., 2011) and it has been demonstrated that changing dopamine concentration or using Tris buffer markedly affects PDA structure and possibly its efficiency depending on applications.

Because of its robustness, universal adhesion properties, biocompatibility, reversible and pH-switchable permselectivity for both cationic and anionic redox-active probe molecules, PDA-based coating technology is expanding rapidly to include energy applications, sensing, bioengineering and nanomedicine for e.g. nanoparticle functionalization, drug delivery and interfacing with cells (Kang et al., 2012, Dreyer et al., 2012; Liu et al., 2014). As an example, multilayered polyelectrolyte films made from PDA particles and polyamines like poly-(L-lysine hydrobromide) (PLL) and pure PDA grains in suspension in the cell culture medium were found to affect melanoma cell growth in different manners (Eap et al., 2013).

Copolymerization of dopamine with aromatic amines or 5-*S*-cysteinyldopamine is a useful means of controlling and modifying PDA structure to change electrical properties, suggesting a versatile behavior for a range of applications in organic electronics (Ambrico et al., 2014a; 2014b).

Bioelectronics and biosensing

Synthetic eumelanins feature a range of chemical and morphological diversity (d'Ischia et al., 2013) which affects a number of key properties of relevance for bioelectronic applications, in particular electrical conduction (Mostert et al., 2012a, Mostert et al., 2012b, Meredith et al., 2013). Several groups have published various device architectures with applications such as memory (metal-insulator-semiconductor geometries) (Ambrico et al., 2011, Ambrico et al., 2012), batteries (Kim et al., 2013) and biomimetic interfaces (Wünsche et al., 2013a, Ambrico et al., 2014b). The device fabrication, and consequently the device performance, is affected by parameters such as melanin precursor and oxidizing system/conditions during eumelanin synthesis.

Eumelanin films are potentially amenable to device applications and readily accessible to chemical and morphological characterization techniques. Significant progress has been made in eumelanin thin film fabrication and processing using different methodologies (Pezzella et al., 2015; Ambrico et al., 2011, Sangaletti et al., 2009, da Silva et al., 2004, Bothma et al., 2008, Bettinger et al., 2009, Abbas et al., 2011, Bloisi et al., 2011a, Bloisi et al., 2011b, Kim et al., 2011, Diaz et al., 2005, Orive et al., 2009, Subianto et al.,

2005). Thin film processing of eumelanin films in $\text{NH}_3(\text{aq})$ significantly increases the N content in the films, as shown by XPS analyses (Wünsche et al., 2013a). Eumelanin has proved to gain in terms of performances and range of applications after *ad hoc* mixing with eumelanin-like macromolecules and inorganic structures. Oliveira et al. prepared an intercalated hybrid material by reacting DOPA with a $\text{V}_2\text{O}_5 \cdot n\text{H}_2\text{O}$ gel preserving the lamellar structure of V_2O_5 and demonstrating the reduction of V(V) to V(IV) ions with associated stability of the V_2O_5 electrochromic response and conductivity (Oliveira et al., 2000). Recently, a bulk heterojunction of porous silicon and eumelanin was obtained, featuring an increased photocarrier collection efficiency at longer wavelengths with respect to bare porous silicon matrices (Mula et al., 2012). Qu et al. polymerized DOPA onto Au-nanoparticles (AuNP) and demonstrated tumor cell imaging with the hybrid nanoparticles (Qu et al., 2010). AuNP were also coated with polydopamine for sensing applications (Li et al., 2011). Fei et al. prepared polydopamine-coated carbon nanotubes and further functionalized them with AuNP (Fei et al., 2008). Gonzáles Orive et al. fabricated eumelanin-iron coated AuNP on HOPG with a strong catalytic activity for hydrogen peroxide electroreduction and hydrogen evolution reaction (Gonzales-Orive et al., 2011). Melanin hybrid structures can be fabricated in very mild conditions (“chimie douce”) as it has been demonstrated by the in-situ formation of the eumelanin-coated TiO_2 nanoparticles through direct reaction of the precursors (Pezzella et al., 2013).

When using eumelanin in devices, particular attention has to be given to the choice of the metal of the electrodes. Indeed, for thin films of synthetic eumelanin included between gold electrodes, under prolonged electrical biasing in humid air, gold dissolution has been observed (Wünsche et al., 2013b). Nanoaggregates of gold and eumelanin generate dendrites, in turn forming conductive pathways between the electrodes. This phenomenon was observed with eumelanins from different sources, as long as they had active phenolic hydroxyl groups. It was suggested that the dissolution of the Au electrodes is enabled by Cl^- present in eumelanin and that the electrochemical reducing ability and metal binding ability of eumelanin was responsible for a strong enhancement of Au dissolution.

In general, eumelanin can act either as electron acceptor or as electron donor, according to a biphasic mechanism reminiscent of the electron transfer processes in redox-conducting films deposited as solid electrodes (Mostert et al., 2012a, Manimala and Horak, 1986). Irreversible oxidation peaks have been observed in measurements on eumelanin synthesized by enzymatic oxidation of L-dopa and incorporated into a carbon paste electrode (Serpentini et al., 2000) as well as, indirectly, on aqueous eumelanin suspensions, where synthetic eumelanin had been prepared by DHI oxidative polymerization (Tarabella et al., 2013).

The metal chelation properties of eumelanin offer interesting possibilities for eumelanin-based metal ion sensing. Huang et al. demonstrated a eumelanin-coated piezoelectric sensor with particularly high sensitivity for Hg(II) (Huang et al., 2007).

Nanotechnology and nanomedicine.

Melanins are increasingly exploited for nanotechnology and nanomedicine applications. PDA nanoparticles that are <100 nm have been shown to exhibit good biocompatibility to HeLa cells after appropriate surface modification and a marked ability to reduce 2,2-diphenyl-1-picrylhydrazyl (DPPH), a commonly used free radical scavenger assay (Ju et al., 2011). The potential of eumelanins for clinical MRI diagnosis has been proposed based on the development of synthetic melanin-like nanoparticles complexed with paramagnetic Fe³⁺ ions as contrast agent. Such nanoparticles show much higher relaxivity values than existing MRI T₁ contrast agents based on gadolinium (Gd) or manganese (Mn) leading to significant enhancement to MRI contrast (Ju et al., 2013). Melanin nanoparticles have also been proposed as a new nanocarrier for pH-responsive drug release. Impregnation of melanin with metronidazole using supercritical carbon dioxide led to a material with a strong pH-dependent drug release profile which combined with lack of cytotoxicity suggested potential application as nanocarrier in formulations for colon and intestine targeted drug delivery (Araujo et al., 2014).

Melanin-covered nanoparticles prepared by enzymatic polymerization on silica have been shown to provide efficient protection of bone marrow against radiotoxicity during radioimmunotherapy and in some cases external beam radiation therapy, permitting the administration of significantly higher doses to tumors during radiation therapy of cancer (Schweitzer et al., 2010).

Photoresponsive and photodegradable hydrogel networks composed of commonly used biocompatible synthetic polymers and natural melanin nanoparticles that may act as photothermal sensitizer and undergo light induced disintegration have been reported (Ninh et al., 2014). These new photoreconfigurable hydrogels provide a useful entry to photodegradable polymers that can find application in clinical studies. Food grade melanin extracted from actinobacterium *Nocardioopsis alba* and melanin-mediated synthesis of silver nanostructures opened interesting perspectives toward the development of antimicrobial nanostructures for food industrial applications (Kiran et al., 2014).

Challenges and Perspectives

Despite a growing number of applications, the way toward a full appreciation of melanins' potential in biomedicine and technology is still a long one. For substantial steps forward in this direction, it is critical that increasing interdisciplinary and intersectoral research is devoted to settle the manifold challenges that hinder exploitation of melanins. These include: a) elucidation of structure-property-function relationships, b) processability, and c) the design of highly conductive films and structures. However, recent progress in eumelanin and polydopamine chemistry and physics (d'Ischia et al., 2014) would suggest that very soon these and other challenges can be met through elucidation of structure-property relationships and the development of efficient tailoring strategies.

A look at melanin-based patents in 2012-2014 (see SI, not including the over 100 polydopamine-based patents) provides a scent of what may happen in the nearest future. Bringing melanin research to a definitive maturity may be possible however only by promoting translational research from the academy to the industry

and the clinics, while encouraging industries and companies to invest more on melanin-based sustainable technology for human health .

Acknowledgments.

This review is a consensus document prepared by members of the EuMelaNet special interest group of the European Society for Pigment Cell Research (ESPCR). More extensive and detailed contributions on specific issues dealt with in this paper with relevant references are provided as Supporting Information.

References

- Abbas, M., Ali, M., Shah, S. K., D'Amico, F., Postorino, P., Mangialardo, S., Cestelli Guidi, M., Cricenti, A., and Gunnella, R. (2011) Control of structural, electronic, and optical properties of eumelanin films by electrospray deposition. *J Phys Chem B* *115*, 11199-11207,
- Abdel-Malek ,Z., Swope, V. B., Suzuki, I., Akcali, C., Harriger, M.D., Boyce, S.T., Urabe, K., and Hearing, V.J. (1995). Mitogenic and melanogenic stimulation of normal human melanocytes by melanotropic peptides. *Proc Natl Acad Sci U S A* *92*, 1789-93.
- Abdel-Malek, Z.A., Ruwe, A., Kavanagh-Starner, R., Kadekaro, A.L., Swope, V., Haskell-Luevano, C., Koikov, L., and Knittel, J.J. (2009) alpha-MSH tripeptide analogs activate the melanocortin 1 receptor and reduce UV-induced DNA damage in human melanocytes. *Pigment Cell Melanoma Res* *22*, 635-644
- Adachi, K., Kato, K., Wakamatsu, K., Ito, S., Ishimaru, K., Hirata, T., Murata, O., and Kumai, H. (2005). The histological analysis, colorimetric evaluation, and chemical quantification of melanin content in 'suntanned' fish. *Pigment Cell Res.* *18*, 465-468.
- Aime, S., Bergamasco, B., Biglino, D., Digilio, G., Fasano, M. Giamello, E., and Lopiano, L. (1997) EPR investigations of the iron domain in neuromelanin. *Biochim Biophys Acta* *1361*, 49-58.
- Ambrico, M., Ambrico, P. F. , Cardone, A., Ligonzo, T., Cicco, S. R., Di Mundo, R., Augelli, V.,and Farinola, G. M. (2011) Melanin Layer on Silicon: an Attractive Structure for a Possible Exploitation in Bio-Polymer Based Metal-Insulator-Silicon Devices. *Adv Mater* *23*, 3332-3336.
- Ambrico, M., Ambrico, P. F. , Ligonzo, T., Cardone, A. , Cicco, S. R. , Lavizzera, A., Augelli ,V., and Farinola G. M. (2012) Memory-like behaviour as a feature of electrical signal transmission in melanin-like bio-polymers. *App.Phys. Lett.* *100*, 253702
- Ambrico, M., Ambrico, P. F., Cardone, A. , Cicco, S., Palumbo, F., Ligonzo, T., Di Mundo, R., Petta, V. , Augelli, V. , Favia, P., and Farinola, G. (2014a) Melanin-like polymer layered on a nanotextured silicon surface for a hybrid biomimetic interface. *J. Mater. Chem. C* *2*, 573-582.
- Ambrico, M., Della Vecchia, N.F., Ambrico, P.F., Cardone, A., Cicco, S.R., Ligonzo, T., Avolio, R., Napolitano, A., and d'Ischia, M. (2014b) A pphotoresponsive red hair-inspired polydopamine-based copolymer for hybrid photocapacitive sensors. *Adv. Funct. Mater.* *24*,7161-7172.
- Andersen, S.O. (2010) Insect cuticular sclerotization: A review. *Insect Biochem.Molecular Biol.* *40*, 166e178
- Araujo, M., Viveiros, R, Correia, T.R., Correia, I.J., Bonifacio, V.D.B., Casimiro, T., and Aguiar-Ricardo, A. (2014) Natural melanin: A potential pH-responsive drug release device. *Int. J. Pharm.* *469*, 140-145.
- Arck, P.C., Overall, R., Spatz, K., Liezman, C., Handjiski, B., Klapp, B.F., Birch-Machin, M.A. and Peters, E.M. (2006) Towards a "free radical theory of graying": melanocyte apoptosis in the aging human hair follicle is an indicator of oxidative stress induced tissue damage. *FASEB J.* *20*, 1567-1569.
- Ardic, F. N., Aktan, S., Kara, C. O., and Sanli, B (1998) High-frequency hearing and reflex latency in patients with pigment disorder *Am.J. Otolaryngol.* *19*,365-369.
- Arzillo, M., Mangiapia, G., Pezzella, A., Heenan, R. K., Radulescu, A., Paduano, L.,and d'Ischia, M. (2012) Eumelanin buildup at the nanoscale: aggregate growth/assembly and visible absorption development in biomimetic 5,6-dihydroxyindole polymerization. *Biomacromolecules.* *13*, 2379-90

- Bamford, A.J., Monadjem, A. and Hardy, I.C.W. (2010). Associations of avian facial flushing and skin colouration with agonistic interaction outcomes. *Ethology* *116*, 1163-1170.
- Barden, H. (1971) The histochemical distribution and localization of copper, iron, neuromelanin and lysosomal enzyme activity in the brain of aging rhesus monkey and the dog. *J Neuropathol Exp Neurol.* *30*, 650-667.
- Barrenas, M. L. (1994) The influence of a melanin -binding drug on temporary threshold shift in humans. *Scandinavian audiol.* *23*, 93-98.
- Bernsmann, F., Ball, V., Addiego, F., Ponche, A., Michel, M., de Almeida Gracio, J. J., Toniazzo, V., and Ruch, D. (2011) Dopamine-melanin film deposition depends on the used oxidant and buffer solution. *Langmuir* *27*, 2819-2825.
- Bettinger, C. J. , Bruggeman, J. P., Misrac, A., Borensteind, J. T., and Langer, R. (2009) Biocompatibility of biodegradable semiconducting melanin films for nerve tissue engineering. *Biomaterials* *30*, 3050-3057
- Bloisi, F., Pezzella, A. Barra, M. Chiarella, F. Cassinese A. and Vicari, L. (2011 a) Matrix assisted pulsed laser deposition of melanin thin films. *J Appl Phys* *110*, 026105-026108.
- Bloisi, F., Pezzella, A. Barra, M. Chiarella, F. Cassinese A. and Vicari, L. (2011 b) Effect of substrate temperature on MAPLE deposition of synthetic eumelanin films. *App. Phys.a-Mater.Sci. Processing* *105*, 619-627.
- Bonilla, C., Boxill, L.A., Donald, S.A., Williams, T., Sylvester, N., Parra, E. J., Dios, S., Norton, H. L., Shriver, M. D., and Kittles, R. A. (2005) The 8818G allele of the agouti signaling protein (ASIP) gene is ancestral and is associated with darker skin color in African Americans. *Hum Genet* *116*, 402-406.
- Bonser, R.H.C. (1995). Melanin and the abrasion resistance of feathers. *Condor* *97*, 590-591.
- Borges, C.R., Roberts, J.C., Wilkins, D.G., and Rollins, D.E.(2001) Relationship of melanin degradation products to actual melanin content: application to human hair” *J. Anal. Biochem.* *290*, 116–125.
- Borovansky, J., and Riley, P.A. (2011) Physiological, and Pathological Functions of Melanosomes, in *Melanins and Melanosomes: Biosynthesis, Biogenesis, Physiological, and Pathological Functions* (Eds: J. Borovansky & P.A. Riley) Wiley-VCH , Germany, *Ch. 12*, 343– 381.
- Bortolotti, G.R. (2006). Natural selection and coloration: protection, concealment, advertisement, or deception? In *Bird Coloration, Volume II: Function and Evolution*, G.E. Hill and K.J. McGraw, eds (Cambridge, MS: Harvard University Press), pp. 3-35.
- Boswell, T., and Takeuchi, S. (2005). Recent developments in our understanding of the avian melanocortin system: its involvement in the regulation of pigmentation and energy homeostasis. *Peptides* *26*, 1733-1743.
- Bothma, J., de Boor, J. D. , Divakar, U., Schwenn, P., and Meredith P. (2008) Device-quality electrically conducting melanin thin films. *Adv Mater* *20*, 3539-3542.
- Boulton, M. (1991) Aging of the retinal pigment epithelium. *Progr. Retinal Res.* *11*, 125-151.
- Breathnach, A.S. (1988) Extra-cutaneous melanin. *Pigment Cell Res.* *1*, 234–237.
- Brenner, M, and Hearing, V.J. (2008) The protective role of melanin against UV damage in human skin. *Photochem Photobiol.* *84*, 539-549.

Bridelli, M.G., Ciati, A., and Crippa, P.R. (2006) Binding of chemicals to melanins re-examined: adsorption of some drugs to the surface of melanin particles. *Biophys Chem.* *119*, 137-145.

Bridelli, M. G., and Crippa, R. (2010) Infrared and water sorption studies of the hydration structure and mechanism in natural and synthetic melanin. *J. Phys. Chem. B* *114*, 9381-9390.

Briganti, S., Camera, E., and Picardo, M. (2003) Chemical and instrumental approaches to treat hyperpigmentation *Pigment Cell Res.* *16*, 101-110.

Burke, J.M., Kaczara, P., Skumatz, C.M.B., Zareba, M., Raciti, M.W., and Sarna, T. (2011) Dynamic analyses reveal cytoprotection by RPE melanosomes against non-photoc stress. *Mol. Vis.* *17*, 2864–2877.

Burt, E.H., Jr., and Ichida, J.M. (2004). Gloger's Rule, feather-degrading bacteria, and color variation among Song Sparrows. *Condor* *106*, 681-686.

Bush, W.D., Garguilo, J., Zucca, F.A., Albertini, A., Zecca, L., Edwards, G.S., Nemanich, R.J., and Simon, J.D. (2006) The surface oxidation potential of human neuromelanin reveals a spherical architecture with a pheomelanin core and a eumelanin surface. *Proc Natl Acad Sci U S A.* *103*, 14785-14789

Bush, W.D., Garguilo, J., Zucca, F.A., Bellei, C., Nemanich, R.J., Edwards, G.S., Zecca, L., and Simon, J.D. (2009) Neuromelanins isolated from different regions of the human brain exhibit a common surface photoionization threshold. *Photochem Photobiol.* *85*, 387-390.

Buszman, E., and Rózańska, R. (2003) Interaction of quinidine, disopyramide and metoprolol with melanin in vitro in relation to drug-induced ocular toxicity. *Pharmazie* *58*, 507-511.

Candille, S.I., Kaelin, C.B., Cattanch, B.M., Yu, B., Thompson, D.A., Nix, M.A., Kerns, J.A., Schmutz, S.M., Millhauser, G.L., and Barsh, G.S. (2007) A -defensin mutation causes black coat color in domestic dogs. *Science* *318*, 1418-1423.

Cao, L., Li, W., Kim, S., Brodie, S.G., and Deng, C.X. (2003) Senescence, aging, and malignant transformation mediated by p53 in mice lacking the Brca1 full-length isoform. *Genes Dev* *17* 201-213.

Capozzi, V., Perna, G., Carmone, P., Gallone, A., Lastella, M., Mezzenga, E., Quartucci, G., Ambrico, M., Augelli, V., Biagi, P., Ligonzo, R., Minafra, A., Schiavulli, L., Pallara, M., and R. Cicero (2006) Optical and photoelectronic properties of melanin. *Thin Solid Films* *511*, 362-366.

Casadevall, A., Rosas, A. L., and Nosanchuk, J. D. (2000) Melanin and virulence in *Cryptococcus neoformans*. *Curr. Opin. Microbiol.* *3*, 354–358.

Cebrián, C., Zucca, F.A, Mauri, P., Steinbeck, J.A., Studer, L., Scherzer, C.R., Kanter, E., Budhu, S., Mandelbaum, J., Vonsattel, J.P., Zecca, L., Loike, J.D., and Sulzer, D. (2014) MHC-I expression renders catecholaminergic neurons susceptible to T-cell-mediated degeneration. *Nat Commun.* *5*, 3633

Cerdá-Reverter, J.M., Agulleiro, M.J., Guillot, R., Sánchez, E., Ceinos, R., and Rotllant, J. (2011). Fish melanocortin system. *Eur. J. Pharmacol.* *660*, 53-60.

Chedekel, M.R. (1982) Photochemistry and photobiology of epidermal melanins *Photochem Photobiol.* *35*, 881-885.

Chedekel, M.R., Smith, S.K., Post, P.W., Pokora, A., and Vessell, D.L. (1978) Photodestruction of pheomelanin: role of oxygen. *Proc Natl Acad Sci U S A.* *75*, 5395-5399.

Cheli, Y., Luciani, F., Khaled, M., Beuret, L., Bille, K., Gounon, P., Ortonne, J.P., Bertolotto, C., and Ballotti, R. (2009) α -MSH and Cyclic AMP elevating agents control melanosome pH through a protein kinase A-independent mechanism. *J. Biol Chem* 284, 18699-706.

Chen, S. G., Xue, C. H. , Wang, J. , Feng, H. , Wang, Y., Ma, Q., and Wang, D. (2009) Adsorption of Pb(II) and Cd(II) by squid ommastrephes bartrami melanin. *Bioinorg Chem Appl.* 2009, 901563.

Clancy, C.M., and Simon, J. D.(2001) Ultrastructural organization of eumelanin from *Sepia officinalis* measured by atomic force microscopy. *Biochemistry* 40, 13353-13360.

Clark, K. D., and Strand, M. R. (2013) Hemolymph melanization in the silkworm *bombyx mori* involves formation of a high molecular mass complex that metabolizes tyrosine. *J. Biol. Chem.* 288, 14476-14487.

Clusella-Trullas, S., van Wyk, J.H., and Spotila, J.R. (2007). Thermal melanism in ectotherms. *J. Thermal Biol.* 32, 235-245.

Coelho, S. G., Zhou, Y., Bushar, H.F., Miller, S.A., Zmudzka, B.Z., Hearing, V.J., and Beer, J.Z. (2009) Long-lasting pigmentation of human skin, a new look at an overlooked response to UV. *Pigment Cell Melanoma Res.* 22, 238-241

Commo, S., Gaillard, O. and Bernard, B.A. (2004a) Human hair greying is linked to a specific depletion of hair follicle melanocytes affecting both the bulb and the outer root sheath. *Br J Dermatol* 150, 435-443

Commo, S., Gaillard, O., Thibaut, S., and Bernard, B.A. (2004b) Absence of TRP-2 in melanogenic melanocytes of human hair. *Pigment Cell Res* 17, 488-497.

Commo, S., Wakamatsu, K., Lozano, I., Panhard, S., Loussouarn, G., Bernard, B.A., and Ito, S. (2012) Age-dependent changes in eumelanin composition in hairs of various ethnic origins. *Int. J. Cosmet.Sci.* 34 102-107.

Cooksey, C. J., Garratt, P.J., Land, E.J., Pavel, S., Ramsden, C.A., Riley, P.A., and Smit, N.P.M. (1997). Evidence of the indirect formation of the catecholic intermediate substrate responsible for the autoactivation kinetics of tyrosinase. *J. Biol. Chem.* 272, 26226-26235.

Corani, A., Huijser, A., Gustavsson, T., Markovitsi, D., Malmqvist, P.Å., Pezzella, A., d'Ischia, M., and Sundström, V. (2014) Superior photoprotective motifs and mechanisms in eumelanins uncovered. *J Am Chem Soc.* 136, 11626-11635.

Cowen, D. (1986) The melanoneurons of the human cerebellum (nucleus pigmentosus cerebellaris) and homologues in the monkey. *J Neuropathol Exp Neurol.* 45, 205-221.

Crippa, R., Horak, V., Prota, G., Svoronos, P., and Wolfram, L. (1989). *The Alkaloids*. Ed.: A. Brossi Academic Press: New York, vol. 36, pp. 255-323.

Dadachova, E., and Casadevall, A. (2008) Ionizing radiation: how fungi cope, adapt, and exploit with the help of melanin *Curr. Opin. Microbiol.* 11, 525–531.

D'Amato, R.J., Lipman, Z.P., and Snyder, S.H. (1986) Selectivity of the parkinsonian neurotoxin MPTP: toxic metabolite MPP⁺ binds to neuromelanin. *Science* 231, 987-989

daSilva, M. I. N., Deziderio, S. N., Gonzalez, J. C., Graeff, C. F. O., and Cotta, M. A. (2004) Synthetic melanin thin films: Structural and electrical properties. *J Appl Phys* 96, 5803-5807.

Della Vecchia, N. F., Avolio, R., Alfe, M., Errico, M. E., Napolitano, A., and d'Ischia, M. (2013) Building-block diversity in polydopamine underpins a multifunctional eumelanin-type platform tunable through a quinone control point. *Adv. Functional Mat.* *23*, 1331-1340.

del Marmol, V., Ito, S., Bouchard, B., Libert, A., Wakamatsu, K., Ghanem, G., and Solano, F. (1996) Cysteine deprivation promotes eumelanogenesis in human melanoma cells. *J Invest Dermatol.* *107*, 698-702.

DeMattei, M., Levi, A.C., and Fariello, R.G. (1986) Neuromelanin pigment in substantia nigra neurons of rats and dogs. *Neurosci Lett.* *72*, 37-42.

Derby, C.D. (2014) Cephalopod Ink: Production, Chemistry, Functions and Applications. *Mar. Drugs* *12*, 2700-2730

Díaz, P., Gimeno, Y., Carro, P., González, S., Schilardi, P. L., Benítez, G., Salvarezza, R. C., and Creus, A. H. (2005) Electrochemical self-assembly of melanin films on gold. *Langmuir* *21*, 5924-5930.

Di Cristo, C., Fiore, G., Scheinker, V., Enikolopov, G., d'Ischia, M., Palumbo, A., and Di Cosmo, A. (2007) Nitric oxide synthase expression in the central nervous system of *Sepia officinalis*: an in situ hybridization study. *Eur J Neurosci.* *26*, 1599-1610.

d'Ischia, M., Napolitano, A., Ball, V., Chen, C-T., and Buehler, M.J. (2014) Polydopamine and eumelanin: from structure-property relationships to a unified tailoring strategy. *Acc. Chem. Res.* *47*, 3541-3550

d'Ischia, M., Wakamatsu, K., Napolitano, A., Briganti, S., Garcia-Borron, J. C., Kovacs, D., Meredith, P., Pezzella, A., Picardo, M., Sarna, T., Simon, J. D., and Ito, S. (2013) Melanins and melanogenesis: methods, standards, protocols. *Pigment Cell Melanoma Res.* *26*, 616-633.

d'Ischia, M., Napolitano, A., and Pezzella, A. (2011) 5,6-Dihydroxyindole chemistry: unexplored opportunities beyond eumelanin. *Eur J Org Chem* *2011*, 5501-5516.

d'Ischia, M., Napolitano, A., Pezzella, A., Meredith, P., and Sarna, T. (2009) Chemical and structural diversity in eumelanins: unexplored bio-optoelectronic materials. *Angew Chem Int Edit* *48*, 3914-3921.

Dong, C., Wang, H., Xue, L., Dong, Y., Yang, L., Fan, R., Yu, X., Tian, X., Ma, S., and Smith, G.W. (2012). Coat color determination by miR-137 mediated down-regulation of microphthalmia-associated transcription factor in a mouse model. *RNA* *18*, 1679-1686.

D'Orazio, J.A., Nobuhisa, T., Cui, R., Arya, M., Spry, M., Wakamatsu, K., Igras, V., Kunisada, T., Granter, S.R., Nishimura, E.K., Ito, S., and Fisher, D.E. (2006) Topical drug rescue strategy and skin protection based on the role of Mc1r in UV-induced tanning. *Nature* *443*, 340-344

Duffy, P., and Tennyson, V.M. (1965) Phase and electron microscopic observations of Lewy bodies and melanin granules in the substantia nigra and locus coeruleus in Parkinson's Disease. *J Neuropathol Exp Neurol.* *24*, 398-414.

Dunn, P. E. (1986) Biochemical aspects of insect immunology. *Ann. Rev. Entomol.* *31*, 321-339.

Dynoodt, P., Mestdagh, P., Van Peer, G., Vandesompele, J., Goossens, K., Peelman, L.J., Geusens, B., Speeckaert, R.M., Lambert, J.L., and Van Gele, M.J. (2013) Identification of miR-145 as a Key Regulator of the Pigmentary Process. *J Invest Dermatol.* *133*, 201-209.

Eap, S. , Ferrand, A. , Machi, V., Ball, V., Huck, O., and Benkirane-Jessel, N. (2013) Polydopamine particles effect on melanoma cells proliferation and melanin secretion. *Adv Chem. Engin. Sci.* 3, 1-10.

Edge, R., d'Ischia, M., Land, E.J., Napolitano, A., Navaratnam, S., Panzella, L. Pezzella, A. , Ramsden, C.A. and Riley, P.A. (2006) Dopaquinone redox exchange with dihydroxyindole and dihydroxyindole carboxylic acid. *Pigment Cell Res.* 19, 443-450.

Eisenman, H. C., and Casadevall, A. (2012) Synthesis and assembly of fungal melanin. *Appl. Microbiol. Biotechnol.* 93, 931–940.

Engelen, M., Vanna, R., Bellei, C., Zucca, F.A., Wakamatsu, K., Monzani, E., Ito, S., Casella, L., and Zecca, L. (2012) Neuromelanins of human brain have soluble and insoluble components with dolichols attached to the melanic structure. *PLoS One.* 7, e48490.

Enochs ,W.S., Nilges , M.J., and Swartz, H.M. (1993) Purified human neuromelanin, synthetic dopamine melanin as a potential model pigment, and the normal human substantia nigra: characterization by electron paramagnetic resonance spectroscopy. *J Neurochem.* 61, 68-79.

Fei, B., Qian, B., Yang, Z., Wang, R., Liu, W., Mak C. L., and Xina, J. H. (2008) Coating carbon nanotubes by spontaneous oxidative polymerization of dopamine. *Carbon* 46, 1795–1797.

Felix, C.C., Hyde, J.S., Sarna, T., and Sealy, R.C. (1978) Melanin photoreactions in aerated media: electron spin resonance evidence for production of superoxide and hydrogen peroxide. *Biochem Biophys Res Commun.* 84, 335-341.

Fiore, G., Poli, A., Di Cosmo, A., d'Ischia, M., and Palumbo, A.(2004) Dopamine in the ink defence system of *Sepia officinalis*: biosynthesis, vesicular compartmentation in mature ink gland cells, nitric oxide (NO)/cGMP-induced depletion and fate in secreted ink. *Biochem J.* 378, 785-791.

Fiore, G., Mattiello, T., Tedeschi, G., Nonnis, S., d'Ischia, M., and Palumbo, A. (2009) Protein nitration is specifically associated with melanin production and reveals redox imbalance as a new correlate of cell maturation in the ink gland of *Sepia officinalis*. *Pigment Cell Melanoma Res.* 22, 857-859.

Flaherty, K.T., Infante, J.R., Daud, A., Gonzalez, R., Kefford, R.F., Sosman, J, Hamid, O., Schuchter, L, et al. (2012) Combined BRAF and MEK inhibition in melanoma with BRAF V600 mutations. *N. Engl J Med.* 367, 1694-1703

Friedman, D. S., Katz, J., Bressler, N. M., Rahmani B., and Tielsch, J. M. (1999) Racial differences in the prevalence of age-related macular degeneration: The Baltimore Eye Survey. *Ophthalmology* 106, 1049–1055.

Froncisz, W., and Sarna, T. (1980) Cu²⁺ probe of metal-ion binding sites in melanin using electron paramagnetic resonance spectroscopy. I. Synthetic melanins. *Arch Biochem Biophys* 202, 289-303.

Fujii, R. (2000). The regulation of motile activity in fish chromatophores. *Pigment Cell Res.* 13, 300-319.

Gallas, J. M. , Littrell, K. C., Seifert, S., Zajac, G. W., and Thiyagarajan, P. (1999) Solution structure of copper ion-induced molecular aggregates of tyrosine melanin. *Biophys. J.* 77, 1135-1142.

Gallone, A., Sagliano, A., Guida, G., Ito, S., Wakamatsu, K., Capozzi, V., Perna, G., Zanna, P., and Cicero, R. (2007). The melanogenic system of the liver pigmented macrophages of *Rana esculenta* L.- tyrosinase activity. *Histol. Histopathol.* 22, 1065-1075.

Galván, I., and Alonso-Alvarez, C. (2009). The expression of melanin-based plumage is separately modulated by exogenous oxidative stress and a melanocortin. *Proc. R. Soc. B* 276, 3089-3097.

Galván, I., and Solano, F. (2009). The evolution of eu- and pheomelanin traits may respond to an economy of pigments related to environmental oxidative stress. *Pigment Cell Melanoma Res.* 22, 339-342.

Galván, I., and Møller, A.P. (2011). Brain size and the expression of pheomelanin-based colour in birds. *J. Evol. Biol.* 24, 999-1006.

Galván, I., Møller, A.P., and Erritzøe, J. (2011a). Testicular melanization has evolved in birds with high mtDNA mutation rates. *J. Evol. Biol.* 24, 988-998.

Galván, I., Mousseau, T.A., and Møller, A.P. (2011b). Bird population declines due to radiation exposure at Chernobyl are stronger in species with pheomelanin-based coloration. *Oecologia* 165, 827-835.

García, T.S., Stacy, J., and Sih, A. (2004). Larval salamander response to UV radiation and predation risk: color change and microhabitat use. *Ecol. Appl.* 14, 1055-1064.

García-Borron, J.C., Abdel-Malek, Z., and Jimenez-Cervantes, C. (2014) MC1R, the cAMP pathway, and the response to solar UV: extending the horizon beyond pigmentation. *Pigment Cell Melanoma Res* 27, 699-720.

Gauden, M., Pezzella, A., Panzella, L., Neves-Petersen, M.T., Skovsen, E., Petersen, S.B., Mullen, K.M., Napolitano, A., d'Ischia, M., and Sundstrom, V. (2008) Role of solvent, pH, and molecular size in excited-state deactivation of key eumelanin building blocks: implications for melanin pigment photostability. *J. Am. Chem. Soc.* 130, 17038-17043

Ghiani, S., Baroni, S., Burgio, D., Digilio, G., Fukuhara, M., Martino, P., Monda, K., Nervi, C., Kiyomine, A. and Aime, S. (2008) Characterization of human hair melanin and its degradation products by means of magnetic resonance techniques. *Magn Reson Chem* 46, 471-479..

Gonçalves, R. C. R., Lisboa, H. C. F., and Pombeiro-Sponchiado, S. R. (2012) Characterization of melanin pigment produced by *Aspergillus nidulans*. *World J. Microbiol. Biotechnol.* 28, 1467–1474.

González Orive, A. , Grumelli, D. , Vericat, C., Ramallo-López, J. C. , Giovanetti, L. , Benitez, G. , Corthey, A. J. C., G., Requejo, F. M. H., Hernández Creus, A., and Salvarezza, R. C. (2011) Naked gold nanoparticles supported on HOPG: melanin functionalization and catalytic activity. *Nanoscale* 3, 1708-1716.

Grande, J.M., Negro, J.J., and Torres, M.J. (2004). The evolution of bird plumage colouration: A role for feather-degrading bacteria? *Ardeola* 51, 375-383.

Gratacap, B., Attard, A., Laurent, A., Stoebner, P., Smirou, D., and Charachon R. (1989) Melanin in the inner ear. An experimental study with control and kanamycin-intoxicated colored guinea pig. *Archives of oto-rhino-laryngology* 246, 235-237.

- Greco, G., Panzella, L., Gentile, G., Errico, M.E., Carfagna, C., Napolitano, A., and d'Ischia, M. (2011). A melanin-inspired pro-oxidant system for dopa(mine) polymerization: mimicking the natural casing process. *Chem Commun.* 47, 10308-10310.
- Greco, G., Wakamatsu, K., Panzella, L., Ito, S., Napolitano, A., and d'Ischia, M. (2009) Isomeric cysteinyl dopas provide a (photo)degradable bulk component and a robust structural element in red human hair pheomelanin. *Pigment Cell Melanoma Res.* 22, 319–327.
- Guerrero, D. (2012) Dermocosmetic management of hyperpigmentations. *Ann. Dermatol. Venereol.* 139 S166-169.
- Guo, S. Hong, L., Akhremitchev, B. B. and Simon J. D. (2008) Surface elastic properties of human retinal pigment epithelium melanosomes. *Photochem. Photobiol.* 84, 671-678.
- Harsanyi, Z.P., Post, P.W., Brinkmann, J.P., Chedekel, M.R., and Deibel, R.M. (1980) Mutagenicity of melanin from human red hair. *Experientia* 36, 291-292
- Hellström, A.R., Watt, B., Fard, S.S., Tenza, D., Mannström, P., Narfström, K., Ekesten, B., Ito, S., Wakamatsu, K., Larsson, J., et al. (2011) Inactivation of Pmel alters melanosome shape but has only a subtle effect on visible pigmentation. *PLoS Genet.* 7, e1002285.
- Hida, T., Wakamatsu, K., Sviderskaya, E.V., Donkin, A.J., Montoliu, L., Lamoreux, L.M., Yu, B., Millhauser, G.L., Ito, S., Barsh, G.S., Jimbow, K., and Bennett, D.C. (2009) Agouti protein, mahogunin, and attractin in pheomelanogenesis and melanoblast-like alteration of melanocytes: a cAMP-independent pathway. *Pigment Cell Melanoma Res* 22, 623-634.
- Hiratsuka, J., Kondoh, H., Tsuboi, T., Yoshino, K., Imajo, Y., Mishima, Y. (2000) Selective uptake of para-boronophenylalanine increases in amelanotic melanoma cells transfected by the tyrosinase gene. *Melanoma Res.* 10, 297-302.
- Hirsch, E., Graybiel, A.M., and Agid, Y.A. (1988) Melanized dopaminergic neurons are differentially susceptible to degeneration in Parkinson's disease. *Nature* 334, 345-348.
- Hong, L., and Simon, J. D. (2006) Insight into the binding of divalent cations to sepia eumelanin from IR absorption spectroscopy. *Photochem Photobiol* 82, 1265-1269.
- Hong, L., and Simon, J. D. (2007) Current understanding of the binding sites, capacity, affinity, and biological significance of metals in melanin. *J Phys Chem B* 111, 7938-7947.
- Hong, L., Garguilo, J., Anzaldi, L., Edwards, G. S. , Nemanich, R. J., and Simon, J. D. (2006) Age-dependent photoionization thresholds of melanosomes and lipofuscin isolated from human retinal pigment epithelium cells. *Photochem Photobiol.* 82, 1475-1481
- Hornykiewicz, O., and Kish, S.J. (1987) Biochemical pathophysiology of Parkinson's disease. *Adv Neurol.* 45 , 19-34.
- Hu, D-N.(2008) Role of ocular melanin in ophthalmic physiology and pathology. *Photochem. Photobiol.* 84, 639-644.
- Huang, G. S. , Wang, M. T. , Su, C. , and Chen, Y. (2007) . Picogram detection of metal ions by melanin-sensitized piezoelectric sensor. *Biosens Bioelectron* 23, 319-325.

- Inomata, K., Aoto, T., Binh, N.T., Okamoto, N., Tanimura, S., Wakayama, T., Iseki, S., Hara, E., Masunaga, T., Shimizu, H., and Nishimura, E. K. (2009) Genotoxic stress abrogates renewal of melanocyte stem cells by triggering their differentiation. *Cell* *137* 1088-1099.
- Inoue, K.; Ito, S.; Kimura, E.; Nabeshima, M.; Takayama, M., and Ishii, T. (1985) Chemical analysis of melanin in guinea pig inner ear . *Ear Research Japan* *16*, 68-71
- Ito, S. (2003). A chemist's view of melanogenesis. *Pigment Cell Res.* *16*, 230-236.
- Ito, S. (2006). Encapsulation of a reactive core in neuromelanin. *Proc. Natl. Acad. Sci. USA* *103*, 14647-14648.
- Ito, S., and Prota, G. (1977). A facile one-step synthesis of cysteinyl dopas using mushroom tyrosinase. *Experientia* *33*, 1118-1119.
- Ito, S. and Wakamatsu, K. (2008). Chemistry of mixed melanogenesis – pivotal roles of dopaquinone. *Photochem. Photobiol.* *84*, 582-592.
- Ito S, and Wakamatsu, K. (2011). Human hair melanins: what we have learned and have not learned from mouse coat color pigmentation. *Pigment Cell Melanoma Res.* *24*, 63-74
- Ito, S., Kato, T, and Fujita, K. (1988). Covalent binding of catechols to proteins through the sulphydryl group. *Biochem. Pharmacol.* *37*, 1707-1710.
- Ito, S., Wakamatsu, K., d'Ischia, M., Napolitano, A., and Pezzella, A. (2011a). in *Melanins and Melanosomes: Biosynthesis, Biogenesis, Physiological, and Pathological Functions* (Eds: P.A. Riley, J. Borovansky) Wiley-VCH , Germany, *Ch. 6*, 167 – 185.
- Ito, S., Nakanishi, Y., Valenzuela, R.K., Brilliant, M.H., Kolbe, L., and Wakamatsu K. (2011b) Usefulness of alkaline hydrogen peroxide oxidation to analyze eumelanin and pheomelanin in various tissue samples: application to chemical analysis of human hair melanins. *Pigment Cell Melanoma Res.* *24*, 605-613.
- Ito, S., Pilat, A. , Gerwat, W., Skumatz, C.M.B., Ito, M., Kiyono, A., Zadlo, A., Makanishi, Y., Kolbe, L., Burke, J.M., Sarna, T. and Wakamatsu, K. (2013) Photoaging of human retinal pigment epithelium is accompanied by oxidative degradation of its eumelanin. *Pigment Cell Melanoma Res.* *26*, 357-366.
- Iwakura, M., Kondoh, H., Hiratsuka, J., Ehara, K., Tamaki, N., Mishima, Y. (2002) Effect of boron neutron capture therapy for melanotic and amelanotic melanoma transplanted into mouse brain *Pigment Cell Res.* *15*, 67-75.
- Jacobson, E. S. (2000) Pathogenic roles for fungal melanins. *Clin. Microbiol. Rev.* *13*, 708-717.
- Jastrzebska, M., Mróz, I., Barwiński, B., Wrzalik, R., and Boryczka, S. (2010) AFM investigations of self-assembled DOPA-melanin nano-aggregates. *J. Mater. Sci.* *45*, 5302-5308.
- Jiang, S., Liu, X. M., Dai, X., Zhou, Q. , Lei, T. C. , Beermann, F. , Wakamatsu, K., and Xu, S. Z. (2010) *Free Radic. Biol. Med.* *48*, 1144-1151
- Jimbow, K., Iwashina, T., Alena, F., Yamada, K., Pankovich, J., Umemura, T. (1993) Exploitation of pigment biosynthesis pathway as a selective chemotherapeutic approach for malignant melanoma. *J. Invest Dermatol.* *100*, 231S-238S.

- Jones, L. N., Rivett, D. E. (1997) The role of 18-methyleicosanoic acid in the structure and formation of mammalian hair fibres. *Micron*. 28, 469-485.
- Journe, F., Id Boufker, H., Van Kempen, L., Galibert, M-D., Wiedig, M., Salès, F., Theunis, A., Nonclercq, D., Frau, A., Laurent, G., Awada, A., and Ghanem, G. (2011) TYRP1 mRNA expression in melanoma metastases correlates with clinical outcome. *Br J Cancer* 105, 1726–1732.
- Ju, K-Y., Lee, Y., Lee, S., Park, S. B. , and Lee, J-K. (2011) Bioinspired polymerization of dopamine to generate melanin-like nanoparticles having an excellent free-radical-scavenging property. *Biomacromolecules* 12, 625–632.
- Ju, K.-Y., Lee, J.W., Im, G.H., Lee, S., Pyo, J., Park, S.B., Lee, J.H., and Lee, J-K. (2013) Bio-inspired, melanin-like nanoparticles as a highly efficient contrast agent for T_1 -weighted magnetic resonance imaging. *Biomacromolecules* 14, 3491–3497
- Kaczara, P., Zareba, M., Herrnreiter, A., Skumatz, C.M., Zazdło, A., Sarna, T., and Burke, J.M. (2012) Melanosome-iron interactions within retinal pigment epithelium-derived cells. *Pigment Cell Melanoma Res.* 25, 804–814.
- Kanetsky, P.A., Swoyer, J., Panossian, S., Holmes, R., Guerry, D., and Rebbeck, T.R. (2002) A polymorphism in the agouti signaling protein gene is associated with human pigmentation. *Am. J. Hum. Genet.* 70, 770-775.
- Kang, S. M., Hwang, N. S., Yeom, J., Park, S. Y., Messersmith, P. B., Choi, I. S., Langer, R., Anderson, D. G., and Lee, H. (2012) One-step multipurpose surface functionalization by adhesive catecholamine. *Adv. Funct. Mater.* 22, 2949–2955.
- Kastner, A., Hirsch, E.C., Lejeune, O., Javoy-Agid, F., Rascol, O., and Agid, Y. (1992) Is the vulnerability of neurons in the substantia nigra of patients with Parkinson's disease related to their neuromelanin content? *J Neurochem.* 59, 1080-1089.
- Kauser, S., Westgate, G.E., Green, M.R., and Tobin, D.J. (2011) Human hair follicle and epidermal melanocytes exhibit striking differences in their aging profile which involves catalase. *J. Invest Dermatol.* 131, 979-982.
- Kaxiras, E. , Tsolakidis, A. , Zonios, G., and Meng, S. (2006) Structural model of eumelanin. *Phys.Rev. Lett.* 97, 218102 .
- Kawasaki, A., Kumasaka, M., Satoh, A., Suzuki, M., Tamura, K., Goto, T., Asashima, M., and Yamamoto, H. (2008). Mitf contributes to melanosome distribution and melanophore dendricity. *Pigment Cell Melanoma Res* 21, 56-62.
- Kim, I. G., Nam, H. J. , Ju Ahn, H., and Jung, D.-Y. (2011) Electrochemical growth of synthetic melanin thin films by constant potential methods. *Electrochim Acta* 56, 2954-2959.
- Kim, Y. J., Wu, W. , Chun, S. E., Whitacre, J. F., and Bettinger, C. J. (2013) Biologically derived melanin electrodes in aqueous sodium-ion energy storage devices. *Proc. Natl. Acad. Sci. USA* 110, 20912-20917.
- Kiran, G.S., Dhasayan, A., Lipton, A.N., Selvin, J., Arasu, M. V. and Al-Dhabi, N. A. (2014) Melanin-templated rapid synthesis of silver nanostructures *J. Nanobiotechnol.* 12, 18.

- Kobayashi, N., Nakagawa, A., Muramatsu, T., Yamashina, Y., Shirai, T., Hashimoto, M.W., Ishigaki, Y., Ohnishi, T., and Mori, T. (1998) Supranuclear melanin caps reduce ultraviolet induced DNA photoproducts in human epidermis. *J Invest Dermatol.* *110*, 806-810.
- Koike, K. and Ebato, A. (2013) One-pack hair dye compositions containing indole compounds. *Jpn. Tokkyo Koho* (2013), JP5363703B220131211.
- Koike, K., and Hata, Y. (2010) A new bioconversion process with *Aspergillus* tyrosinase for production of dihydroxyindole and a novel coloring technology for gray hair coverage with the product. *Fragrance J.* *38*, 16-20.
- Korkina, L.G. (2007) Phenylpropanoids as naturally occurring antioxidants: from plant defense to human health. *Cell Mol. Biol.* *53*, 15-25.
- Korytowski, W. and Sarna, T., (1990) Bleaching of melanin pigments. Role of copper ions and hydrogen peroxide in autooxidation and photooxidation of synthetic dopa-melanin. *J. Biol. Chem.* *265*, 12410-12416.
- Kreuer, K.D., Paddison, S.J., Spohr, E., and Schuster, M. (2004) Transport in proton conductors for fuel-cell applications: simulations, elementary reactions, and phenomenology. *Chem. Rev.* *104*, 4637-4678.
- Lamoreux, L.M., Delmas, V., Larue, L., and Bennett, D. (2010) *The colors of mice: a model genetic network*, Wiley-Blackwell London .
- Land, E.J., Ito, S., Wakamatsu, K., and Riley, P.A. (2003). Rate constants for the first two chemical steps of eumelanogenesis. *Pigment Cell Res.* *16*, 487-493.
- Langfelder, K. , Streibel, M., Jahn, B. , Haase, G., and Brakhage, A. A. (2003) Biosynthesis of fungal melanins and their importance for human pathogenic fungi. *Fungal Gen. Biol.* *38*, 143–158.
- Langston, J.W., Forno, L.S., Tetrud, J., Reeves, A.G., Kaplan, J.A., and Karluk, D. (1999) Evidence of active nerve cell degeneration in the substantia nigra of humans years after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine exposure. *Ann Neurol.* *46*, 598-605.
- Larsson, B.S. (1993) Interaction between chemicals and melanin. *Pigment Cell Res.* *6*, 127-33.
- Lee, H., Dellatore, S.M., Miller, W.M., and Messersmith, P.B. (2007) *Science* *318*, 426-430.
- Le Pape, E., Passeron, T., Giubellino, A., Valencia, J.C., Wolber, R., and Hearing, V.J. (2009) Microarray analysis sheds light on the dedifferentiating role of agouti signal protein in murine melanocytes via the Mc1r. *Proc Natl Acad Sci U S A* *106*, 1802-1807.
- Li, F., Yang, L., Zhao, C., and Dua, Z. (2011) Electroactive gold nanoparticles/polyaniline/polydopamine hybrid composite in neutral solution as high-performance sensing platform. *Anal. Methods* *3*, 1601-1606.
- Li, J., Liu, Y., Xin, X., Kim, T.S., Cabeza, E.A., Ren, J., Nielsen, R., Wrana, J.L., and Zhang, Z. (2012). Evidence for positive selection on a number of microRNA regulatory interactions during recent human evolution. *PLoS Genet.* *8*, e1002578.
- Liebscher, J., Mrowczynski, R., Scheidt, H. A., Filip, C., Hadade N. D., Turcu, R., Bende, A., and Beck, S. (2013) Structure of Ppolydopamine: a never-ending story? *Langmuir* *29*, 10539–10548.
- Lindquist, N.G., Larsson, B.S., and Lydén-Sokolowski, A. (1988) Autoradiography of [¹⁴C]paraquat or [¹⁴C]diquat in frogs and mice: accumulation in neuromelanin. *Neurosci Lett.* *93*, 1-6.

- Liu, Y., and Simon, J. D. (2003a) Isolation and biophysical studies of natural eumelanins: Applications of imaging technologies and ultrafast spectroscopy *Pigm Cell Res* *16*, 606-618.
- Liu, Y., and Simon, J.D. (2003b) The effect of preparation procedures on the morphology of melanin from the ink sac of *Sepia officinalis*. *Pigment Cell Res* *16*, 72-80
- Liu, Y., Hong, L., Kempf, V. R., Wakamatsu, K., Ito S. and Simon, J. D. (2004) Ion-exchange and adsorption of Fe(III) by *Sepia* melanin. *Pigment Cell Res* *17*, 262-269.
- Liu, Y., Hong, L., Bowers, C.R., Wakamatsu, K., Ito, S., and Simon, J.D. (2005). Comparison of the structural, chemical, and spectroscopic properties of human black and red hair melanosomes. *Photochem. Photobiol.* *81*, 135-144.
- Liu, Y., Ai, K., and Lu, L. (2014) Polydopamine and its derivative materials: synthesis and promising applications in energy, environmental, and biomedical fields *Chem. Rev.* *114*, 5057–5115.
- L'opez-Serrano, D., Solano, F., and Sanchez-Amat, A. (2004) Identification of an operon involved in tyrosinase activity and melanin synthesis in *Marinomonas mediterranea*. *Gene* *342*, 179–187.
- Mandrizzato, S., Callegaro, A., Turcatel, G., Francescato, S., Montesco, M.C., Chiarion-Sileni, V., Mocellin, S., Rossi, C.R., Biciato, S., Wang, E., Marincola, F.M., and Zanovello, P. (2006) A gene expression signature associated with survival in metastatic melanoma. *J Transl Med* *4*, 50.
- Manimala, M., and Horak, V. (1986) Time-separated 2-step oxidation of poly(2,6-naphthoquinone) film" *J Electrochem Soc* *133*, 1987-1988.
- Mårs, U., and Larsson, B.S. (1995) New thioureas and related substances intended for melanoma targeting. *Pigment Cell Res.* *8*, 194-201.
- Marsden, C.D. (1983) Neuromelanin and Parkinson's disease. *J Neural Transm Suppl.* *19*, 121-141.
- Matouš, B., Mechl, Z., šopková, B., Duchoň, J., Pavel, S., Buděšinská, A., Kočent, A. (1980) The excretion of Thormählen positive melanogens in melanoma patients and its clinical significance, *Eur. J. Cancer* *16*, 383-388.
- McGeer, P.L., Itagaki, S., Boyes, B.E., and McGeer, E.G. (1988) Reactive microglia are positive for HLA-DR in the substantia nigra of Parkinson's and Alzheimer's disease brains. *Neurology* *38*, 1285-1291.
- McGraw, K.J. (2003). Melanins, metals, and mate quality. *Oikos* *102*, 402-406.
- McGraw, K.J. (2008). An update on the honesty of melanin-based color signals in birds. *Pigment Cell Melanoma Res.* *21*, 133-138.
- Meng, S., and Kaxiras, E. (2008) Theoretical models of eumelanin protomolecules and their optical properties. *Biophys. J.* *94*, 2095-2105.
- Meredith, P., and Riesz, J. (2005) Radiative relaxation quantum yields for synthetic eumelanin. *Photochem. Photobiol.* *79*, 211-216.
- Meredith, P., and Sarna T. (2006). The physical and chemical properties of eumelanin. *Pigment Cell Res.* *19*, 572-594.
- Meredith, P., Powell, B. J., Nighswander-Rempel, S., Pederson, M., and Moore, E. (2006) Towards structure-property-function relationships for eumelanin. *Soft Matter* *2*, 37-44.

Meredith, P., Bettinger, C. J., Irimia-Vladu, M., Mostert, A. B., and Schwenn, P. E. (2013) Electronic and optoelectronic materials and devices inspired by nature. *Rep. Progr. Phys.* *76*, 034501.

Meyer zum Gottesberge, A.M. (1988) Physiology and pathophysiology of inner ear melanin. *Pigment Cell Res.* *1*, 238-49.

Michard, Q., Commo, S., Belaidi, J.P., Alleaume, A.M., Michelet, J.F., Daronnat, E., Eilstein, J., Duche, D., Marrot, L. and Bernard, B.A. (2008a) TRP-2 specifically decreases WM35 cell sensitivity to oxidative stress. *Free Radic. Biol. Med.* *44*, 1023-1031.

Michard, Q., Commo, S., Rocchetti, J., El Houari, F., Alleaume, A.M., Wakamatsu, K., Ito, S. and Bernard, B.A. (2008b) TRP-2 expression protects HEK cells from dopamine- and hydroquinone-induced toxicity. *Free Radic. Biol. Med.* *45*, 1002-1010.

Mitra, D., Luo, X., Morgan, A. et al. (2012). An ultraviolet-radiation-independent pathway to melanoma carcinogenesis in the red hair/fair skin background. *Nature* *491*, 449-453.

Moreau, M.F., Papon, J., Labarre, P., Moins, N., Borel, M., Bayle, M., Bouchon, B., and Madelmont, J.C. (2005) Synthesis, in vitro binding and biodistribution in B16 melanoma-bearing mice of new iodine-125 spermidine benzamide derivatives. *Nucl Med Biol.* *32*, 377-84.

Mostert, A. B., Powell, B. J., Pratt, F. L., Hanson, G. R., Sarna, T., Gentle, I. R., and P. Meredith (2012 a) Role of semiconductivity and ion transport in the electrical conduction of melanin. *Proc. Natl. Acad. Sci. USA* *109*, 8943-8947.

Mostert, A. B., Powell, B. J., Gentle, I. R., and Meredith, P. (2012 b) On the origin of electrical conductivity in the bio-electronic material melanin. *Appl. Phys. Lett.* *100*, 093701.

Mueller, K.P., and Neuhauss, S.C.F. (2014). Sunscreen for fish: co-option of UV light protection for camouflage. *PLoS ONE* *9*, e87372.

Mula, G., Manca, L., Setzu, S., and Pezzella, A. (2012) Photovoltaic properties of PSi impregnated with eumelanin. *Nanoscale Res. Lett.* *7*, 377.

Nakamura, Y., Yamanaka, H., Hata, Y., Ebato, A., and Koike, K. (2012) Development of the novel hair coloring system using enzymatically-prepared melanin precursors by *Aspergillus oryzae* tyrosinase. *Seibutsu Kogaku Kaishi* *90*, 115-121.

Napolitano, A., Palumbo, A., d'Ischia, M., and Prota, G. (1996b) Mechanism of selective incorporation of the melanoma seeker 2-thiouracil into growing melanin. *J Med Chem.* *39*, 5192-5201

Nappi, A. J., and Christensen, B. M. (2005) Melanogenesis and associated cytotoxic reactions: Applications to insect innate immunity. *Insect Biochem. Molecular Biol.* *35*, 443-459

Negro, J.J., Sarasola, J.H., Fariñas, F., and Zorrilla, I. (2006). Function and occurrence of facial flushing in birds. *Comp. Biochem. Physiol. A* *143*, 78-84.

Ninh, C., Cramer, M., and Bettinger, C.J. (2014) Photoresponsive hydrogel networks using melanin nanoparticle photothermal sensitizers. *Biomater. Sci.* *2*, 766-774.

Nishimura, E.K., Granter, S.R., and Fisher, D.E. (2005) Mechanisms of hair graying: incomplete melanocyte stem cell maintenance in the niche. *Science* *307*, 720-724.

Nosanchuk, J. D. , and Casadevall, A. (2003) The contribution of melanin to microbial pathogenesis. *Cell. Microbiol.* *5*, 203–223.

Oliveira, H. P., Graeff, C. F. O., Zanta, C. L. P. S. , Galina, A. C., and Gonçalves, P. J. (2000) Synthesis, characterization and properties of a melanin-like/vanadium pentoxide hybrid compound. *J. Mater. Chem.* *10*, 371-375.

Orive, A. G. , Gimeno, Y., Creus, A. H., Grumell, D., Vericat, C. , Benitez, G. and Salvarezza, R. C. (2009) Electrochemical preparation of metal-melanin functionalized graphite surfaces. *Electrochim Acta* *54*, 1589-1596.

Ozeki, H., Ito, S., Wakamatsu, K., and Ishiguro, I. (1997). Chemical characterization of pheomelanogenesis starting from dihydroxyphenylalanine or tyrosine and cysteine. Effects of tyrosinase and cysteine concentrations and reaction time. *Biochim. Biophys. Acta* *1336*, 539-548.

Palumbo, A.(2003) Melanogenesis in the ink gland of *Sepia officinalis*. *Pigment Cell Res.* *16*, 517-522

Palumbo, A., d'Ischia, M., Misuraca, G., De Martino, L., and Prota, G. (1994) A new dopachrome-rearranging enzyme from the ejected ink of the cuttlefish *Sepia officinalis*. *Biochem. J.* *299*, 839-844

Palumbo, A., Mårs, U., De Martino, L., d'Ischia, M., Napolitano, A., Larsson, B.S., and Prota, G. (1997) Selective incorporation of the prototype melanoma seeker thiourea into nascent melanin: a chemical insight. *Melanoma Res.* *7*, 478-485.

Palumbo, A., Poli, A., Di Cosmo, A., and d'Ischia, M. (2000) N-Methyl-D-aspartate receptor stimulation activates tyrosinase and promotes melanin synthesis in the ink gland of the cuttlefish *Sepia officinalis* through the nitric Oxide/cGMP signal transduction pathway. A novel possible role for glutamate as physiologic activator of melanogenesis. *J. Biol. Chem.* *275*, 16885-90.

Panhard, S, Lozano, I., and Loussouarn, G. (2012) Greying of the human hair. A worldwide survey, re-visiting the "50" rule of thumb. *Br. J. Dermatol.* *167* , 865-873.

Panzella, L., Gentile, G., D'Errico, G., Della Vecchia, N.F., Errico, M. E., Napolitano, A., Carfagna, C., and d'Ischia, M. (2013) Atypical structural and π -electron features of a melanin polymer that lead to superior free-radical-scavenging properties. *Angew. Chem., Int. Ed.* *52*, 12684-12687.

Panzella, L., Leone, L., Greco, G., Vitiello, G., D'Errico, G., Napolitano, A., and d'Ischia, M. (2014). Red human hair pheomelanin is a potent pro-oxidant mediating UV-independent contributory mechanisms of melanomagenesis. *Pigment Cell and Melanoma Res.* *27*, 244-252.

Pannkuk, E.L., Siefferman, L.M., and Butts, J.A. (2010). Colour phases of the eastern screech owl: a comparison of biomechanical variables of body contour feathers. *Funct. Ecol.* *24*, 347-353.

Patel, K. R., Wyman, J. A., Patel, K. A., and Burden, B. J. (1996) A mutant of *Bacillus thuringiensis* producing a dark-brown pigment with increased UV resistance and insecticidal activity. *J. Invertebrate Pathol.* *67*, 120–124.

Peles, D.N., Hong, L., Hu, D-N., Ito, S., Nemanich, R.J., and Simon, J.D. (2009) Human iridal stroma melanosomes of varying pheomelanin content possess a common eumelanin outer surface. *J. Phys.Chem. B.*, *113*, 11346-11351.

Pezzella, A., Capelli, L., Costantini, A., Luciani, G., Tescione, F., Silvestri, B., Vitiello, G., and Branda, F. (2013) Towards the development of a novel bioinspired functional material: Synthesis and characterization of hybrid TiO₂/DHICA-melanin nanoparticles. *Mat Sci Eng C-Mater* 33, 347-355.

Pezzella, A., Barra, M., Musto, A., Navarra, A., Alfe, M., Manini, P., Parisi, S., Cassinese, A., Criscuolo, V., and d'Ischia, M. (2015) Stem cell-compatible eumelanin biointerface fabricated by chemically controlled solid state polymerization. *Mater. Horizons* 2, 212-220.

Pezzella, A., Napolitano, A., d'Ischia, M., and Prota, G. (1996) Oxidative polymerisation of 5,6-dihydroxyindole-2-carboxylic acid to melanin: a new insight. *Tetrahedron* 52, 7913-7920.

Pinner, S., Jordan, P., Sharrock, K., Bazley, L., Collinson, L., Marais, R., Bonvin, E., Goding, C., and Sahai, E. (2009) Intravital imaging reveals transient changes in pigment production and Brn2 expression during metastatic melanoma dissemination. *Cancer Res.* 69, 7969–7977.

Plonka, P. M., and Grabacka, M. (2006) Melanin synthesis in microorganisms biotechnological and medical aspects, *Acta.Bochimica.Polonica* 53, 429–443.

Polacheck, I., and Kwon-Chung, K. J. (1988) Melanogenesis in *Cryptococcus neoformans*. *J. Gen. Microbiol.* 134, 1037–1041.

Prota, G. (1992). *Melanins and Melanogenesis*, Academic Press: San Diego, CA.

Prota, G., Hu, D.-N., Vincensi, M. R., McCormick, S. A., and Napolitano A. (1998) Characterization of melanins in human irides and cultured uveal melanocytes from eyes of different colors. *Exp. Eye Res.* 67, 293–299.

Qu, W.G., Wang, S.M., Hu, Z.J., Cheang, T.Y., Xing, Z.H., Zhang, X.J., and Xu, A.W. (2010) In situ synthesis of Gold@3,4-dihydroxy-l-phenylalanine core-shell nanospheres used for cell imaging. *J. Phys. Chem. C.* 114, 13010.

Raper, H.S. (1927). The tyrosinase-tyrosine reaction. VI. Production from tyrosine of 5,6-dihydroxyindole and 5,6-dihydroxyindole-2-carboxylic acid – the precursors of melanin. *Biochem. J.* 21, 89-96.

Reale, S., Crucianelli, M., Pezzella, A., d'Ischia M., and F. De Angelis, (2012) Exploring the frontiers of synthetic eumelanin polymers by high-resolution matrix-assisted laser/desorption ionization mass spectrometry. *J Mass Spectrom* 47, 49-53.

Riley, P.A. (1997) Melanin. *Int. J. Biochem Cell Biol.* 29, 1235-1239.

Riley, P.A. (2003) Melanogenesis and melanoma. *Pigment Cell Res.* 16, 548-552

Riley, P.A., Cooksey, C.J., Johnson, C.I., Land, E.J., Latter, A.M., Ramsden, C.A. (1997) Melanogenesis-targeted anti-melanoma pro-drug development: effect of side-chain variations on the cytotoxicity of tyrosinase-generated ortho-quinones in a model screening system. *Eur. J. Cancer.* 33, 135-143.

Robbins, L.S., Nadeau, J.H., Johnson, K.R., Kelly, M.A., Roselli-Rehfuss, L., Baack, E., Mountjoy, K.G., and Cone, R.D. (1993) Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function. *Cell* 72, 827-834

Rogers, G., and Koike, K. (2009) Laser capture microscopy in a study of expression of structural proteins in the cuticle cells of human hair. *Exp. Dermatol.* 18, 541–547.

Rorsman, H., Agrup, G., Hansson, C., and Rosengren, E. (1983) Biochemical recorders of malignant melanoma. In *Pigment Cell*. vol. 6. R. M. Mackie (ed.), pp. 93-115, Karger, Basel.

Rorsman, H., Rosengren, A.M., and Rosengren, E. (1973) Determination of 5-S-cysteinyl-dopa in melanoma with a fluorimetric method. *Yale J. Biol. Med.* 46, 516-522.

Roulin, A., Mafli, A., and Wakamatsu, K. (2013). Reptiles produce pheomelanin: evidence in the Eastern Hermann's Tortoise (*Eurotestudo boettgeri*). *J. Herpetol.* 47, 258-261.

Rożanowska, M., Korytowski, W., Rożanowski, B., Skumatz, C., Boulton, M. E., Burke, J. M., and Sarna, T. (2002) Photoreactivity of aged human RPE melanosomes: a comparison with lipofuscin. *Invest Ophthalmol Vis Sci*, 43, 2088-2096.

Russo, G.L., De Nisco, E., Fiore, G., Di Donato, P., d'Ischia, M., and Palumbo, A. (2003) Toxicity of melanin-free ink of *Sepia officinalis* to transformed cell lines: identification of the active factor as tyrosinase. *Biochem Biophys Res Commun.* 308, 293-299

Samokhvalov, A., Liu, Y., and Simon, J. (2004) Characterization of the Fe(III)-binding site in *Sepia* eumelanin by resonance Raman confocal microspectroscopy. *Photochem. Photobiol.* 80, 84-88.

Samokhvalov, A., Hong, L., Liu, Y., Garguilo, J., Nemanich, R.J., Edwards G.S., and Simon, J.D. (2005). Oxidation potentials of human eumelanosomes and pheomelanosomes. *Photochem. Photobiol.* 81 145-148

Sanchez-Mas, J., Hahmann, C., Gerritsen, I., Garcia-Borron, J.C., and Jimenez-Cervantes, C. (2004) Agonist-independent, high constitutive activity of the human melanocortin 1 receptor. *Pigment Cell Res* 17, 386-395.

Sangaletti, L., Borghetti, P., Ghosh, P., Pagliara, S., Vilmercati, P., Castellarin-Cudia, C., Floreano, L., Cossaro, A., Verdini, A., Gebauer, R., and A. Goldoni (2009) Polymerization effects and localized electronic states in condensed-phase eumelanin. *Phys. Rev. B* 80, 174203.

Sarna, T. (1992) Properties and function of the ocular melanin: a photobiophysical view. *J. Photochem. Photobiol. B: Biol.* 12, 215–258.

Sarna, T., Froncisz, W., and Hyde, J. (1980) Cu²⁺ probe of metal-ion binding sites in melanin using electron paramagnetic resonance spectroscopy. II. Natural melanin. *Arch. Biochem. Biophys.* 202, 304-313

Sarna, T., Burke, J.M., Korytowski, W., Rożanowska, M., Skumatz, C.M.B., Zareba, A., and Zareba, M. (2003). Loss of melanin from human RPE with aging: possible role of melanin photooxidation. *Exp. Eye Res.* 76, 89–98.

Schiöth, H.B., Raudsepp, T., Ringholm, A., Fredriksson, R., Takeuchi, S., Larhammar, D., and Chowdhary, B.P. (2003). Remarkable synteny conservation of melanocortin receptors in chicken, human, and other vertebrates. *Genomics* 81, 504-509.

Schmalzer-Ripcke, J., Sugareva, V., Gebhardt, P., Winkler, R., Kniemeyer, O., Heinekamp, T., and Brakhage, A.A. (2009) Production of pyomelanin, a second type of melanin, via the tyrosine degradation pathway in *Aspergillus fumigatus*. *Appl. Environ. Microbiol.* 75, 493–503.

Schreiber, R.W., Schreiber, E., Peele, A.M., and Burt, E.H., Jr. (2006). Pattern of damage to albino Great Frigatebird flight feathers supports hypothesis of abrasion by airborne particles. *Condor* 108, 736-741.

Schweitzer, A.D., Revskaya, E., Chu, P., Pazo, V., Friedman, M., Nosanchuk, J.D., Cahill, S., Frases, S., Casadevall, A., and Dadachova, E. (2010) Melanin-covered nanoparticles for protection of bone marrow during radiation therapy of cancer. *Int. J. Rad. Oncol. Biol. Phys.* *78*, 1494–1502.

Sendoel, A., Kohler, I., Fellmann, C., Lowe, S.W., and Hengartner, M.O. (2010) HIF-1 antagonizes p53-mediated apoptosis through a secreted neuronal tyrosinase. *Nature* *465*, 577-583.

Serpentini, C., Gauchet, C., de Montauzon, D., Comtat, M., Ginestar, J., and Paillous, N. (2000) First electrochemical investigation of the redox properties of DOPA-melanins by means of a carbon paste electrode. *Electrochim. Acta* *45*, 1663-1668.

Shao, Q., Yang, B., Xu, Q., Li, X., Lu, Z., Wang, C., Huang, Y., Soederhaell, K. and Ling, E. (2012) Hindgut innate immunity and regulation of fecal microbiota through melanization in insects. *J. Biol. Chem.* *287*, 14270-14279.

Siegrist, W., Drozd, R., Cotti, R., Willard, D.H., Wilkison, W.O., and Eberle, A.N. (1997) Interactions of alpha-melanotropin and agouti on B16 melanoma cells: evidence for inverse agonism of agouti. *J. Recept. Signal Transduct. Res.* *17*, 75-98.

Simon, J.D., Hong, L., and Peles, D.N. (2008). Insights into melanosomes and melanins from some interesting spatial and temporal properties, invited Centennial Feature Article. *J. Phys. Chem. B* *112*, 13201 – 13217

Simon, J.D., Peles, D.N. (2010) The red and the black. *Acc. Chem. Res.* *43*, 1452-1460

Smit, N., VanderMeulen, H., Koerten, H. K., Kolb, R. M., Mommaas, A. M., Lentjes, E., and Pavel, S. (1997) Melanogenesis in cultured melanocytes can be substantially influenced by L-tyrosine and L-cysteine. *J. Investig. Dermatol.* *109*, 796-800.

Smit, N., Vicanova, J., and Pavel, S. (2009) The hunt for natural skin whitening agents. *Int. J. Mol. Sci.* *10*, 5326-5349.

Solano, F. (2014). Melanins: Skin pigments and much more-types, structural models, biological functions, and formation routes. *New J. Sci.* Article ID 498276.

Solano, F., Briganti, S., Picardo, M., and Ghanem, G. (2006) Hypopigmenting agents: an updated review on biological, chemical and clinical aspects. *Pigment Cell Res.* *19*, 550-571.

Sono, K., Lye, D., Moore, C., Boyd, W., Gorlin, T., and Belitsky, J. (2012) Melanin-based coatings as lead-binding agents. *Bioinorg. Chem. Appl.* 361803

Stark, K. B., Gallas, J. M., Zajac, G. W., Golab, J. T., Gidanian, S., McIntire T., and Farmer, J. P. (2005) Effect of stacking and redox state on optical absorption spectra of melanins-Comparison of theoretical and experimental results. *J. Phys. Chem. B* *109*, 1970-1977.

Sturm, R.A., and Duffy, D.L. (2012) Human pigmentation genes under environmental selection. *Genome Biol.* *13*, 248.

Subianto, S., Will, G., and Meredith, P. (2005) Electrochemical synthesis of melanin free-standing films. *Polymer* *46*, 11505-11509.

Sugumaran, M. (2002) Comparative biochemistry of eumelanogenesis and the protective roles of phenoloxidase and melanin in insects *Pigment Cell Research* *15*, 2-9.

Sulzer, D., Bogulavsky, J., Larsen, K.E., Behr, G., Karatekin, E., Kleinman, M.H., Turro, N., Krantz, D., Edwards, R.H., Greene, L.A., and Zecca, L. (2000) Neuromelanin biosynthesis is driven by excess cytosolic catecholamines not accumulated by synaptic vesicles. *Proc Natl Acad Sci U S A.* *97*, 11869-11874.

Sulzer, D., Mosharov, E., Talloczy, Z., Zucca, F.A., Simon, J.D., Zecca, L. (2008) Neuronal pigmented autophagic vacuoles: lipofuscin, neuromelanin, and ceroid as macroautophagic responses during aging and disease. *J Neurochem.* *106*, 24-36.

Suzuki, I., Tada, A., Ollmann, M.M., Barsh, G.S., Im, S., Lamoreux, M.L., Hearing, V.J., Nordlund, J.J., and Abdel-Malek, Z.A. (1997) Agouti signaling protein inhibits melanogenesis and the response of human melanocytes to alpha-melanotropin. *J Invest Dermatol* *108*, 838-842.

Swope, V.B., Jameson, J.A., McFarland, K.L., Supp, D.M., Miller, W.E., McGraw, D.W., Patel, M.A., Nix, M.A., Millhauser, G.L., Babcock, G.F., and Abdel-Malek ZA (2012) Defining MC1R regulation in human melanocytes by its agonist alpha-melanocortin and antagonists agouti signaling protein and beta-defensin 3. *J Invest. Dermatol.* *132*, 2255-2262.

Tadokoro, T., Kobayashi, N., Zmudzka, B.Z., Ito, S., Wakamatsu, K., Yamaguchi, Y., Korossy, K.S., Miller, S.A., Beer, J.Z., and Hearing, V.J. (2003) UV-induced DNA damage and melanin content in human skin differing in racial/ethnic origin. *FASEB J.* *17*, 1177-1179.

Takeuchi, S., Takahashi, S., Okimoto, R., Schiöth, H.B., and Boswell, T. (2003). Avian melanocortin system: α -MSH may act as an autocrine/paracrine hormone. *Ann. N. Y. Acad. Sci.* *994*, 366-372.

Takeuchi, S., Zhang, W., Wakamatsu, K., Ito, S., Hearing, V. J., Kraemer, K. H., and Brash, D. E. (2004) Melanin acts as a potent UVB photosensitizer to cause an atypical mode of cell death in murine skin. *Proc. Natl. Acad. Sci.* *101*, 15076-15081.

Tarabella, G., Pezzella, A., Romeo, A., D'Angelo, P., Coppedè, N., Calicchio, M., d'Ischia, M., Mosca, R., and Iannotta, S. (2013) Irreversible evolution of eumelanin redox states detected by an organic electrochemical transistor: en route to bioelectronics and biosensing. *J. Mater. Chem. B* *1*, 3843-3849.

Thibaut, S., De Becker, E., Caisey, L., Baras, D., Karatas, S., Jammayrac, O., Pisella, P.J., and Bernard, B.A. (2010) Human eyelash characterization. *Br. J. Dermatol.* *162*, 304-10.

Tran, M. L., Powell, B. J., and Meredith, P. (2006) Chemical and structural disorder in eumelanins: a possible explanation for broadband absorbance. *Biophys. J.* *90*, 743-752.

Trias, J., Vinas, M., Guinea, J., and Loren, J. G. (1989) Brown pigmentation in *Serratia marcescens* cultures associated with tyrosine metabolism. *Can. J. Microbiol.* *35*, 1037-1042

Tse, D.C.S., McCreery, R.L., and Adams, R.N. (1976). Potential oxidative pathways of brain catecholamines. *J. Med. Chem.* *19*, 37-40.

Valverde, P., Healy, E., Jackson, I., Rees, J.L., and Thody, A.J. (1995) Variants of the melanocyte-stimulating hormone receptor gene are associated with red hair and fair skin in humans. *Nat. Genet.* *11*, 328-330.

- van Lanschot, C.G., Koljenović, S., Grunhagen, D.J., Verhoef, C., and van Akkooi, A.C. (2014) Pigmentation in the sentinel node correlates with increased sentinel node tumor burden in melanoma patients. *Melanoma Res.* 24, 261-266
- Vavricka, C. J., Han, Q., Mehre, P., Ding, H., Christensen, B. M., and Li, J. (2014) Tyrosine metabolic enzymes from insects and mammals: A comparative perspective. *Insect Science* 21, 13-19.
- Waite, J. H. (2008) Surface chemistry: Mussel power. *Nature Mat.* 7, 8-9.
- Waite, J.H., and Tanzer, M.L. (1981) Polyphenolic substance of *Mytilus edulis*: Novel adhesive containing L-Dopa and hydroxyproline. *Science* 212, 1038–1040.
- Wakamatsu, K., Kageshita, T., Furue, M., Hatta, N., Kiyohara, Y., Nakayama, J., Ono, T., Saida, T., Takata, M., Tsuchida, T., Uhara, H., Yamamoto, A., Yamazaki, N., Naito, A., and Ito, S. (2002) Evaluation of 5-S-cysteinyldopa as a marker of melanoma progression: 10 years' experience. *Melanoma Res.* 12, 245-253.
- Wakamatsu, K., Fujikawa, K., Zucca, F.A., Zecca, L., and Ito, S. (2003). The structure of neuromelanin as studied by chemical degradative methods. *J. Neurochem.* 86, 1015-1023
- Wakamatsu, K., Hu, D. N., McCormick S. A., and Ito S. (2008) Characterization of melanin in human iridal and choroidal melanocytes from eyes with various colored irides. *Pigment Cell Res.* 21, 97–105.
- Wakamatsu, K., Nakanishi, Y., Miyazaki, N., Kolbe, L., and Ito, S. (2012a). UVA-induced oxidative degradation of melanins: fission of indole moiety in eumelanin and conversion to benzothiazole moiety in pheomelanin. *Pigment Cell Melanoma Res.* 25, 434-445.
- Wakamatsu, K., Murase, Y., Zucca, F.A., Zecca, L., and Ito, S. (2012b). Biosynthetic pathway to neuromelanin and its aging process. *Pigment Cell Melanoma Res.* 25, 793-802.
- Wang, A., Marino, A.R., and Norris, J.R. Jr. (2009) Investigation of Photoexcited States in Porcine Eumelanin through their Transient Radical Products *J. Phys Chem B* 113, 10480-10482.
- Ward, W.C., Guan, Z., Zucca, F.A., Fariello, R.G., Kordestani, R., Zecca, L., Raetz, C.R., and Simon, J.D. (2007) Identification and quantification of dolichol and dolichoic acid in neuromelanin from substantia nigra of the human brain. *J Lipid Res.* 48, 1457-1462.
- Ward, W.C., Zucca, F.A., Bellei, C., Zecca, L., and Simon, J.D. (2009) Neuromelanins in various regions of human brain are associated with native and oxidized isoprenoid lipids. *Arch Biochem Biophys.* 484, 94-99.
- Wasterstrom, S. A., Bredberg, G., Lindquist, N. G., Lyttkens, L., and Rask-Anderson, H. (1986) Ototoxicity of kanamycin in albino and pigmented guinea pigs. I. A morphologic and electrophysiologic study *Am. J. Otol.* 7, 11-8.
- Watt, A. A. R, Bothma, J.P., and Meredith, P. (2009) The supramolecular structure of melanin. *Soft Matter* 5, 3754–3760.
- Weis, E., Shah, C. P., Lajous, M., Shields, J. A., and Shields C. L. (2006). The association between host susceptibility factors and uveal melanoma: A meta-analysis. *Arch. Ophthalmol.* 124, 54–60.

- Wenczl, E., Van der Schans, G. P., Roza, L., Kolb, R. M., Timmerman, A. J., Smit, N. P. M., Pavel, S., and Schothorst, A. A. (1998) (Pheo)melanin photosensitizes UVA-induced DNA damage in cultured human melanocytes. *J. Invest. Dermatol.* *111*, 678-682.
- Wielgus, A. R., and Sarna T. (2005) Melanin in human irides of different color and age of donors. *Pigment Cell Res.* *18*, 454-464.
- Wilms, H., Rosenstiel, P., Sievers, J., Deuschl, G., Zecca, L., and Lucius, R. (2003) Activation of microglia by human neuromelanin is NF-kappaB dependent and involves p38 mitogen-activated protein kinase: implications for Parkinson's disease. *FASEB J.* *17*, 500-502
- Wolfram, L.J., Hall, K., and Hui, I. J. (1970) The Mechanism of Hair Bleaching. *Soc. Cosmet. Chem.* *21*, 875-900.
- Wolnicka-Glubisz, A., Pecio, A., Podkowa, D., Kolodziejczyk, L.M., and Plonka, P.M. (2012). Pheomelanin in the skin of *Hymenochirus boettgeri* (Amphibia: Anura: Pipidae). *Exp. Dermatol.* *21*, 537-540.
- Wünsche, J., Cicoira, F., Graeff, C. F. O., and Santato, C. (2013 a) Eumelanin thin films: solution-processing, growth, and charge transport properties. *J. Mat. Chem. B* *1*, 3836-3842.
- Wünsche, J., Cardenas, L., Rosei, F., Cicoira, F., Graeff, C. F. O., Poulin, S. P. A., and Santato, C. (2013 b) In situ formation of dendrites in eumelanin thin films between gold electrodes. *Adv. Funct. Mater.* *45*, 5591-5598
- Wünsche, J., Deng, Y., Kumar, P., Di Mauro, E., Josberger, E., Sayago, J., Pezzella, A., Soavi, F., Cicoira, F., Rolandi, M., and Santato, C. (2015) Protonic and electronic transport in hydrated thin films of the pigment eumelanin. *Chem. Mater.* *27*, 436-442
- Yamada, I., Seki, S., Ito, S., Suzuki, S., Matsubara, O., Kasuga, T. (1991) The killing effect of 4-S-cysteaminyphenol, a newly synthesised melanin precursor, on B16 melanoma cell lines. *Br. J. Cancer* *63*, 187-190.
- Yamamura, K., Kamada, S, Ito, S, Nakagawa, K., Ichihashi, M., and Tsujimoto, Y. (1996) Accelerated disappearance of melanocytes in bcl-2-deficient mice. *Cancer Res.* *56* 3546-3550.
- Zecca, L, and Swartz, H. M. (1993) Total and paramagnetic metals in human substantia nigra and its neuromelanin. *J Neural Transm Park Dis Dement Sect.* *5*, 203-213.
- Zecca, L., Mecacci, C., Seraglia, R., and Parati E. (1992) The chemical characterization of melanin contained in substantia nigra of human brain. *Biochim. Biophys. Acta.* *1138*, 6-10.
- Zecca, L., Pietra, R., Goj, C., Mecacci, C., Radice, D., and Sabbioni, E. (1994) Iron and other metals in neuromelanin, substantia nigra, and putamen of human brain. *J Neurochem.* *62*, 1097-2101.
- Zecca, L., Shima, T., Stroppolo, A., Goj, C., Battiston, G.A., Gerbasi, R., Sarna, T., and Swartz, H.M. (1996) Interaction of neuromelanin and iron in substantia nigra and other areas of human brain. *Neuroscience*, *73*, 407-415.
- Zecca, L., Costi, P., Mecacci, C., Ito, S., Terreni, M., and Sonnino, S. (2000) Interaction of human substantia nigra neuromelanin with lipids and peptides. *J Neurochem.* *74*, 1758-1765.

Zecca, L., Fariello, R., Riederer, P., Sulzer, D., Gatti, A., and Tampellini, D. (2002) The absolute concentration of nigral neuromelanin, assayed by a new sensitive method, increases throughout the life and is dramatically decreased in Parkinson's disease. *FEBS Lett.* *510*, 216-220.

Zecca, L., Stroppolo, A., Gatti, A., Tampellini, D., Toscani, M., Gallorini, M., Giaveri, G., Arosio, P., Santambrogio, P., Fariello, R.G., Karatekin, E., Kleinman, M.H., Turro, N., Hornykiewicz, O., and Zucca, F.A. (2004) The role of iron and copper molecules in the neuronal vulnerability of locus coeruleus and substantia nigra during aging. *Proc Natl Acad Sci U S A.* *101*, 9843-9848.

Zecca, L., Bellei, C., Costi, P., Albertini, A., Monzani, E., Casella, L., Gallorini, M., Bergamaschi, L., Moscatelli, A., Turro, N.J., Eisner, M., Crippa, P.R., Ito, S., Wakamatsu, K., Bush, W.D., Ward, W.C., Simon, J.D., and Zucca, F.A. (2008a) New melanic pigments in the human brain that accumulate in aging and block environmental toxic metals. *Proc Natl Acad Sci U S A.* *105*, 17567-17572.

Zecca, L., Casella, L., Albertini, A., Bellei, C., Zucca, F.A., Engelen, M., Zadlo, A., Szweczyk, G., Zareba, M., and Sarna, T. (2008b) Neuromelanin can protect against iron-mediated oxidative damage in system modeling iron overload of brain aging and Parkinson's disease. *J. Neurochem.* *106*, 1866-1875.

Zecca, L., Wilms, H., Geick, S., Claasen, J.H., Brandenburg, L.O., Holzknecht, C., Panizza, M.L., Zucca, F.A., Deuschl, G., Sievers, J., and Lucius, R. (2008c) Human neuromelanin induces neuroinflammation and neurodegeneration in the rat substantia nigra: implications for Parkinson's disease. *Acta Neuropathol.* *116*, 47-55.

Zhang, W., Phillips, K., Wielgus, A. R., Liu, J., Albertini, A., Zucca, F. A., Faust, R., Qian, S.Y., Miller, D.S., Chignell, C.F., Wilson, B., Jackson-Lewis, V., Przedborski, S., Joset, D., Loike, J., Hong, J. S., Sulzer, D., and Zecca, L. (2011) Neuromelanin activates microglia and induces degeneration of dopaminergic neurons: implications for progression of Parkinson's disease. *Neurotox Res.* *19*, 63-72.

Zhang, W., Dai, M., Fridberger, A., Hassan, A., Degagne, J., Neng, L., Zhang, F., He, W., Ren, T., Trune, D., Auer, M., and Shi, X. (2012) Perivascular-resident macrophage-like melanocytes in the inner ear are essential for the integrity of the intrastrial fluid-blood barrier. *Proc Natl Acad Sci U S A.* *109*, 10388-93

Zucca, F.A., Basso, E., Cupaioli, F.A., Ferrari, E., Sulzer, D., Casella, L., and Zecca, L. (2014) Neuromelanin of the human substantia nigra: an update. *Neurotox Res.* *25*, 13-23.

Legends to figures

Figure 1. Melanins and melanogenesis: elucidating chemical and physical properties is central to unravel biological roles and functions and to translate clues from pigment cells to innovative applications.

Figure 2. Structures of main melanin precursors in plants, fungi and microorganisms

Figure 3. Biosynthetic pathways leading to eumelanin and pheomelanin production (Ito and Wakamatsu, 2008). Note that the activities of tyrosinase, Tyrp1 and Tyrp2 are involved in the production of eumelanin, while only tyrosinase (and the amino acid cysteine) is necessary for the production of pheomelanin.

Figure 4. Casing model for mixed melanogenesis (Agrup et al., 1982; Bush et al., 2006; Ito, 2006; Greco et al., 2011). Note that in the process of mixed melanogenesis, pheomelanin pigment is produced first, followed by the deposit of eumelanin pigment. In the granule with the eumelanin surface, the side was intentionally cut away to reveal the inner pheomelanin core. Eumelanin is believed to act as a photoprotective anti-oxidant and pheomelanin as a phototoxic pro-oxidant

Figure 5. Comproportionation reaction of the basic units within the eumelanin polymer. The scheme shows how water-dependent, pH-driven comproportionation between 5,6-dihydroxyindole and 5,6-indolequinone units generates protons and free radicals (electrons) as free charge carriers.

Figure 6. The electrical conductivity of solid-state synthetic eumelanin measured (a) in a simple sandwich configuration, and (b) in a surface electrode van der Pauw configuration. The sandwich geometry does not allow the system to come into equilibrium during the measurement and produces an erroneous conductivity isotherm, whilst the surface electrode configuration does and demonstrates a “percolation-like” transition at ~ 10-15% by weight water. The MDAS model is the Mott Davies Amorphous Semiconductor model previously (and wrongly) used to describe the behaviour. (Adapted from Mostert et al., 2012b)

Figure 7. Overall view of main differences between synthetic DHI and DHICA melanins (adapted from Panzella et al., 2013). The scheme shows how differences in the mode of coupling of indole units determine the extent of electron delocalization, absorption and paramagnetic properties, and the mode of aggregation at the supramolecular level.

Figure 8. Main reaction pathways involved in polydopamine formation. (adapted from Della Vecchia et al., 2013). The model was supported by a subsequent experimental and computational investigation showing that PDA consists of mixtures of oligomers in which indole units with different degrees of (un)saturation and open-chain dopamine units give rise to charge transfer interactions between o-quinoid and catechol units. (Liebscher et al., 2013)




Fig 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7

Table 1. Relationship between melanin properties and the proposed biological roles.

Organism/organ//biological context	Role/function	Property	References
Mammalian fur, avian feathers, reptile scales	Camouflage, sociosexual display	Diverse color patterns	Galván and Alonso-Alvarez (2009)
Mammalian and avian testicles	Protection against oxidative stress caused by mutation	Antioxidant, free radical scavenger	Galván et al. (2011a)
Skin, eyes	Photoprotection, visual acuity, antioxidant action.	Light absorption and scattering, excited state deactivation, sequestration of redox-active metal ions, free radical scavenging	Meredith and Sarna, 2006
Substantia nigra, inner ear	Energy transducer	Water-dependent hybrid ionic-electronic semiconductor properties	Mostert et al, 2012a
Inner ear	Calcium homeostasis regulation, metal reservoir and buffering system	Metal chelation, drug binding	Hong and Simon, 2007
Inner ear, substantia nigra	Redox buffering and control	Redox behavior	Samokhvalov et al., 2005
Insect cuticle, avian feathers	Structure strengthening	Chemical reactivity, quinone cross-linking with proteins and polysaccharides	Andersen, 2010
Insects	Innate immune system	Melanin bioadhesion and coating properties	Liu et al., 2014
	Wound healing	Quinone intermediates of melanogenesis	Ito and Wakamatsu, 2008
Bacteria	Virulence	Shield against host immune response	Shao et al., 2012
Eye, substantia nigra	Protection	Antioxidant, free radical scavenger	Panzella et al., 2013; Jiang et al., 2010; Zecca et al., 2008b
	Detoxification	Drug binding and covalent incorporation, metal binding	Zecca et al., 2008a; Bridelli et al., 2006
Cephalopod ink	Defense against predator	Toxic activity of melanogenic enzymes, catecholamine binding	Cooksey et al., 1997; Solano, 2014
Reptiles, amphibians, cold-blooded animals	Thermoregulation	Energy dissipation into heat	Meredith and Sarna, 2006
Browning of fruits	Response to injury	Catechol oxidation	Korkina, 2007
Fungi	Radioprotection	Free radical quenching, spatial arrangement	d'Ischia et al., 2009