Red tails are effective decoys for avian predators

Belén Fresnillo^a, Josabel Belliure^a, José Javier Cuervo^b

^a Department of Life Sciences, Ecology Section, University of Alcalá, Madrid, Spain;

^b Department of Evolutionary Ecology, Museo Nacional de Ciencias Naturales (CSIC),
Madrid, Spain

Corresponding author: Belén Fresnillo

Department of Life Sciences, Ecology Section, University of Alcalá, University
Campus, 28871 Alcalá de Henares, Madrid, Spain

E-mail: belen.fresnillo@edu.uah.es

Running title: Red tails are effective decoys for avian predators

Key words: *Acanthodactylus erythrurus*, attack diversion, predator-prey interaction,
evolution of signals

Total words: 4 365

Introduction: 692, Material and Methods: 1 967, Results: 644, Discussion: 1 064

Cited references: 61

Figures: 3

Tables: 2

Abstract

The decoy or deflection hypothesis, which states that conspicuous colouration is present in non-vital parts of the body to divert attacks from head and trunk, thus increasing survival probability, is a possible explanation for the presence of such colouration in juveniles of non-aposematic species. To test this hypothesis we made plasticine and plaster lizard models of two colour morphs, red or dark-and-light striped tails, based on the colouration of spiny-footed lizard (*Acanthodactylus erythrurus*) hatchlings, which naturally show a dark-and-light striped dorsal pattern and red tail. Lizard models were placed in the field and also presented to captive common kestrels (*Falco tinnunculus*), a common avian lizard predator. The number of attacks and the body part attacked (tail or rest-of-body) were recorded, as well as the latency to attack. Our results suggest that models of both colour morphs were recognized as prey and attacked at a similar rate, but in the field, red-tailed models were detected, and thus attacked, sooner than striped-tailed. Despite this increase in detection rate by predators, red-tailed models effectively diverted attacks to the tail from the more vulnerable body parts, thus supporting the decoy hypothesis. Greater fitness benefits of attack diversion to the tail compared to the costs of increased detection rate by predators would explain the evolution and maintenance of red tail colouration in lizards.

Introduction

Understanding the function of conspicuous colouration in prey has attracted much interest because it may also increase detection by predators (Arnold 1984; Endler 1980; Haskell 1996). Conspicuous colouration may be a consequence of sexual selection in adults (Darwin 1871; Kemp and Rutowski 2011; Stuart-Fox et al. 2003) because it may improve mating success. Because conspicuous colouration is sometimes also present in juveniles (Booth 1990), and sexual selection cannot be driving colour evolution in such cases, at least three alternative hypotheses have been proposed to explain its evolution. The inter-age-class signalling hypothesis proposes that conspicuous colouration in juveniles evolves to reduce aggression from conspecific adults (Clark and Hall 1970; Ochi and Awata 2009; VanderWerf and Freed 2003). Conspicuous colouration in juveniles may also advertise to predators prey toxicity or unpalatability (aposematism hypothesis), or may mimic an aposematic prey model (Booth 1990; Kraus and Allison 2009). Lastly, conspicuous colouration of non-vital body parts may divert attacks from vital parts (decoy or deflection hypothesis; Bateman et al. 2014; Johnson et al. 2008; Kodandaramaiah et al. 2013; Telemeco et al. 2011; Watson et al. 2012). This function as a lure can be reinforced by stereotyped movements of the conspicuously coloured body part (Telemeco et al. 2011).

The decoy hypothesis is more plausible in species where conspicuous colouration is present in expendable body parts. This is taken to the extreme in reptiles, where juveniles and/or adults of many lizard species show conspicuous colouration in the tail (Castilla et al. 1999; Clark and Hall 1970; Hawlena et al. 2006; Watson et al. 2012), which can be easily detached from the rest of the body, thus facilitating the escape from predators, although its loss might eventually lead to increased risk of

mortality (Arnold 1998; Fox and McCoy 2000; Wilson 1992; review in Bateman and Fleming 2009). Although conspicuous colouration very often increases detection by predators (Arnold 1984; Bateman et al. 2014; Endler 1980; Haskell 1996; but see Castilla et al. 1999; Watson et al. 2012), this signal may still be selected if the benefits of escaping an attack exceed the costs of increased attack rate (Cooper and Vitt 1991). The first studies addressing the decoy hypothesis in lizards were conducted in captivity (Clark and Hall 1970; Cooper and Vitt 1985; Vitt and Cooper 1986), but the use of plasticine (Mochida 2011; Valkonen et al. 2011; Watson et al. 2012) or plaster (Baylis et al. 2012; Ruiz-Rodríguez et al. 2013) models has allowed experiments in the field with different vertebrate species, obtaining information on predator attacks either by video recording or by inferring the attacks by the marks left on the models.

The efficiency of blue and green lizard tails in diverting attacks has already been tested (Castilla et al. 1999; Cooper and Vitt 1985; Watson et al. 2012), but there are a number of species from several lizard families (e.g., Scincidae, Gymnophthalmidae, Lacertidae) that have red tails. Avian predators are known to discriminate colours (Bowmaker 2008), and different colours might cause different responses in bird predators, either because of differences in conspicuousness (e.g., Schaefer et al. 2006) or because of a sensory bias for certain colour (e.g., Møller and Erritzøe 2010). Consequently, results obtained when testing the decoy hypothesis in blue and green tails might not apply to red tails. The effectiveness of red tails as decoys has not yet been addressed. In the present study, our aim was to answer two specific questions: (1) do red tails make lizards more detectable to avian predators? and (2) are red tails effective decoys for diverting bird attacks from vital body parts? To answer these questions we used plasticine and plaster models based on the morphology of spiny-footed lizards (*Acanthodactylus erythrurus*, Schinz 1833), a species in which hatchlings have red tails

and dark-and-light striped bodies. We checked whether birds attacked red-tailed models (striped body and red tail) more and/or sooner than those with striped tails (striped body and tail), and whether red-tailed models received more attacks on the tail than striped-tailed. Models were placed in the field and also presented to captive common kestrels (*Falco tinnunculus*, Linnaeus 1758), one of the main avian predators of *A. erythrurus* (Martín and López 1990).

Materials and methods

Study species

A. erythrurus is a medium-sized lizard (snout-vent length (SVL) and total length up to around 80 and 230 mm, respectively, although size can vary among the populations (Carretero and Llorente 1993; Seva Román 1982)) present in southern and central Iberian Peninsula and northern Africa (Belluore 2006). In populations in central Spain, new-born lizards appear in mid-August and are active until November. At the beginning of April the following spring, these lizards become active again, and they do not reach sexual maturity until their second spring (Castilla et al. 1992; Pollo and Pérez-Mellado 1990). Therefore, all individuals can be classified in three age classes, hereafter referred to as hatchlings (from hatching until the first winter), juveniles (from first to second winter) and adults (from the second winter onwards).

Colouration in this species undergoes ontogenetic changes. Hatchlings show a dorsal pattern of strongly contrasting dark and light bands that run the length of their bodies. This pattern starts reticulating in juveniles, and becomes dark and light patches

in adults (Seva Román 1982). The ventrolateral parts of hatchling and juvenile tails are red (Carretero and Llorente 1993; Seva Román 1982), and this colour is clearly visible from above in hatchlings and small juveniles (personal observation). The rear part of the hind limbs of juveniles is also red. Juvenile males lose their red colouring when they are around one year old, whereas juvenile females retain it through adulthood until they become gravid (Cuervo and Belliure 2013; Seva Román 1982).

Lizard models

We made some lizard models with plasticine and some with plaster, because both types of models have been widely used in similar studies in various taxa (Castilla et al. 1999; Castilla and Labra 1998; Mochida 2011; Ruiz-Rodríguez et al. 2013; Valkonen et al. 2011; Vervust et al. 2007; Watson et al. 2012). Plaster models were made using plastic lizard toys which we covered with a thin layer of plaster (Aquaplast Standard, Beissier S.A., Guipúzcoa, Spain). These models were larger than real hatchlings (plastic lizard toys: mean SVL $\pm$ SD = 56.6 ± 7.9 mm, mean total length $\pm$ SD = 131.8 ± 31.4 mm, $N = 10$; hatchlings: mean SVL $\pm$ SD = 38.3 ± 2.6 mm, mean total length $\pm$ SD = 107.7 ± 10.7 mm, $N = 23$; t -Student test, $t_{31} \geq 3.32$, $P \leq 0.002$ in the two tests), but both juveniles and adults are common prey of avian predators (Belliure 2006), so we think these models could still be easily recognized as prey. Plasticine models were made by hand from black oil-based modelling clay (Plastilina Jovi, Barcelona, Spain) and attempted to mimic the size of real hatchlings (random sample of models: mean SVL $\pm$ SD = 38.4 ± 2.8 mm, mean total length $\pm$ SD = 107.7 ± 7.9 mm, $N = 15$; comparison with hatchlings: t -Student test, $0.00 \leq t_{36} \leq 0.09$, $P \geq 0.928$ in the two tests).

Model colouring was based on real *A. erythrurus* hatchlings. We used non-toxic paints (Satin Paint, La Pajarita, Manises, Spain) to create two colour morphs that differed in tail colouration: red-tailed models and striped-tailed models (Fig. 1). In both colour morphs, the back of the body consisted of a contrasting pattern of dark and light bands (Fig. 1). Neither the paints used in the experiment nor real lizard dorsal colouration reflect UV light (B. Fresnillo, J. Belliure and J. J. Cuervo, unpublished data), so differences in colour discrimination by birds and humans was probably not an issue in this study.

In both plaster and plasticine models, it was possible to identify the type of predator by the marks left on the model based on previous studies (Baylis et al. 2012; Brodie 1993; Valkonen et al. 2011; see Fig. 2). As plaster is harder than plasticine, we are aware that some avian attacks might not have left any mark on the plaster models. We only recorded avian attacks in this study because birds are mainly visually-directed predators, whereas mammals and reptiles also rely on chemical cues (Greene 1988). Materials used to make the models are not odourless and certainly do not smell like lizards, which might have affected the predatory behaviour of mammals and reptiles.

Field study

We distributed lizard models when hatchlings were active (August-November) in two different years (2010 and 2011) and in two localities in central Spain where *A. erythrurus* is common. One study area was located in Chapinería, in south-western Region of Madrid (N40°22'; W4°13'), a Mediterranean oak forest with a meadow structure, where oaks (*Quercus ilex* L.) and lavender (*Lavandula stoechas* Lam.)

dominated vegetation patches surrounded by open areas. The other study area was located in Aranjuez, in southern Region of Madrid (N40°1'; W3°33'), a steppe dominated by esparto grass (*Stipa tenacissima* L.) where trees were scarce and the space surrounding plants was mainly bare soil. The study areas were 67 km apart in a straight line, and were selected because vegetation cover differed dramatically (many fewer trees in Aranjuez), which may have influenced prey detectability by visually-directed predators (Cuadrado et al. 2001; Vásquez et al. 2002). Moreover, both study areas might also have different predator communities. The study was replicated in two different habitats because the effect of model colouration on avian predation rate might depend on the habitat (a stronger effect was expected in more open habitats or where the community of avian predators was more abundant).

In both localities and years, models were placed in open areas near bushes (as if they were basking lizards), at least 100 m apart from one another, and alternating both colour morphs. When both plaster and plasticine models were present in the same study area, locations of the two model types were randomized. In 2010, we worked only in Chapinería and only with plaster models. We distributed an equal number of models of both colour morphs and inspected them every 4-5 days to record the number of models attacked by birds and the body parts attacked (tail and/or rest-of-body). A total of ten models were always present in the field, as a new model was added in a new location when any model was removed due to an avian attack or after four visits without avian attacks. This makes a maximum of 20 days of exposition in the field for these models. In 2011, we worked in both localities, using plasticine and plaster models in Chapinería but only plasticine models in Aranjuez. Both study areas were visited every 3 days, and 10-16 models were left in each area, always evenly distributed between both colour

morphs. The protocol for recording attacks and replacing models was the same as in 2010, so models were exposed to predators a maximum of 12 days in both localities.

A total of 189 lizard models were placed in the field, but 85 of them (two plaster and 83 plasticine models) disappeared. We considered these models as attacked (in both body parts) if there were bird tracks (Fig. 2g) on the ground (11 models), or as “disappeared” if there were no clear bird tracks (74 models). Lizard models considered as disappeared were excluded from statistical analyses. Most plasticine models that disappeared were very probably eaten by ants.

Captivity study

F. tinnunculus is a bird of prey in the Falconidae family that is widely distributed across Eurasia and Africa (BirdLife International 2014). It commonly preys on lizards, at least in some areas (Carrillo and González-Dávila 2009; Carrillo et al. 1994; Vanzyl 1994), and is a typical predator of *A. erythrurus* in Spain (Martín and López 1990), what makes this species a good model organism for studying predatory interactions with our lizard models.

For this study, we used captive *F. tinnunculus* from the Santa Faz Wildlife Rehabilitation Centre (Alicante, Spain) from August to October in three consecutive years (2011, 2012 and 2013). Only birds that were able to fly were involved in the experiment, although some of them had sustained wing injuries. We conducted a total of 31 trials in which an *F. tinnunculus* (which had not been fed for 24 hours before the trial) was presented with a plasticine lizard model that was either red-tailed ($N = 15$) or striped-tailed ($N = 16$). Each bird was used only once. Trials were conducted in

individual cages (4.80 x 1.65 x 3.20 m). The lizard model was placed in the centre of the concrete cage floor at the same time that the bird was placed on a perch about 3 m from the lizard model and 1.3 m off the floor. Models were checked for attacks from outside the cage two or three times per day. The observer only entered the cage when an attack on the model was suspected. The trial ended when the model was attacked or after four days without attacks. If the model was attacked, the body part attacked, i.e., with marks (tail or rest-of-body), and the latency to attack in days were recorded.

Statistical analyses

In the field study, Logistic Regression Models (GLZs) with binomial error distribution were used to test for any effect of model colour morph (red or striped tail) on (1) the probability of being attacked and (2) the probability of being attacked on each body part (tail or rest-of-body). In the first analysis, all the models not counted as disappeared ($N = 115$) were included, although the effect of model colour morph was qualitatively identical if models disappearing without the presence of bird tracks were considered attacked and included in the analysis ($N = 189$). For the second analysis, we included only lizard models that had been attacked by birds on a single body part ($N = 38$), because when there were marks on both parts, it was impossible to know which part was attacked first. To test for the possible effect of model colour morph on the latency to attack, a Generalized Linear Model (GLM) with Poisson error distribution was used. Only models attacked by birds ($N = 63$) were included in this analysis. All analyses in the field study simultaneously tested for differences between study areas (Chapinería and Aranjuez) and types of models (plaster and plasticine), and also checked for any

interaction between these two factors and model colour morph. Year was excluded from our statistical analyses because it was strongly associated with the type of model in all data subsets used here (Fisher's exact test; $P < 0.001$ in the three tests), but when year was included instead of type of model, results were qualitatively identical (see Results).

In the captivity study, GLZs with binomial error distribution were used to test for the possible effects of model colour morph (red or striped tail) on (1) the probability of being attacked, and (2) the probability of being attacked on each body part (tail or rest-of-body). All models ($N = 31$) were used in the first analysis, whereas in the second, only lizard models that had been attacked on a single body part were included ($N = 16$). To test for the possible effect of model colour morph on the latency to attack, a GLM with Poisson error distribution was used. Only attacked models ($N = 22$) were included in this analysis. All analyses in the captivity study simultaneously tested for differences in years (2011, 2012 and 2013) and checked the interaction between model colour morph and year.

In both studies, a backward stepwise procedure was used in all analyses, retaining only terms associated with P values below 0.10 in the final statistical models. If Akaike's information criterion (Burnham and Anderson 2002) was used to select final models, results were qualitatively identical in all analyses (results not shown for brevity). All statistical analyses were carried out using R (R Development Core Team 2013) and lme4 package (Bates et al. 2012). All tests were two-tailed and the significance level was set to 0.05.

Results

Field study

Excluding models that were counted as disappeared, we recorded data for 115 models, 31 plaster (14 with red tails and 17 with striped tails) and 84 plasticine (48 with red tails and 36 with striped tails). A total of 63 models were attacked by birds. Of the attacked models, 38 were attacked on one body part, whereas the rest (25 models) were attacked in both body parts (tail and rest-of-body). Taking into account the total number of models placed in the field (not including models replacing those attacked by non-avian predators), 39.16% of them were counted as disappeared, 33.33% were attacked by birds, and 27.51% were not attacked by birds.

The probability of being attacked by birds was not significantly different for red and striped-tailed lizard models (Likelihood test; $\chi^2 = 0.15$, $N = 115$, $P = 0.697$; Table 2), types of models (plaster or plasticine), or study areas (Chapinería or Aranjuez), as none of these factors was retained in the final model. However, red-tailed models were attacked sooner than striped-tailed, and plasticine models were attacked sooner than plaster (Table 1, Fig. 3a). The interaction between model colour morph and type of model was statistically significant (Table 1), implying that the difference between colour morphs depended on the type of model. In separate analyses of plaster and plasticine models, we found that red-tailed plaster models were attacked sooner than striped-tailed (Likelihood test; $\chi^2 = 36.51$, $N = 19$, $P < 0.001$), but for plasticine models, the difference was not statistically significant (Likelihood test; $\chi^2 = 2.44$, $N = 44$, $P = 0.118$; Fig. 3a). When year was included in this analysis instead of type of model, red-tailed models were also attacked sooner than striped-tailed (Likelihood test; $\chi^2 = 24.82$, $N = 63$, $P < 0.001$). Models were attacked sooner in 2011 than in 2010 (Likelihood test; $\chi^2 = 150.78$, $N = 63$, $P < 0.001$), and the statistically significant interaction between

model colour morph and year (Likelihood test; $\chi^2 = 8.81$, $N = 63$, $P = 0.003$) implied that the difference between colour morphs depended on the year. We analysed data for both years separately and found that red-tailed models were attacked sooner than striped-tailed in 2010 (Likelihood test; $\chi^2 = 31.75$, $N = 16$, $P < 0.001$), but the difference was not statistically significant in 2011 (Likelihood test; $\chi^2 = 1.88$, $N = 47$, $P = 0.170$).

The proportion of attacks to each body part (tail or rest-of-body) was different for red and striped-tailed models, as the probability of being attacked on the tail was higher for red-tailed models (Likelihood test; $\chi^2 = 3.91$, $N = 38$, $P = 0.048$; Table 2). Neither type of model nor study area had a significant effect on the part of the body that was attacked, as none of these factors or their interactions was retained in the final model.

Captivity study

A total of 31 models were presented to the birds, and 22 of them were attacked. Of the attacked models, 16 were attacked on one body part, whereas the rest (6 models) were attacked on both body parts. The probability of being attacked by a captive *F. tinnunculus* was not significantly different for red or striped-tailed lizard models (Likelihood test; $\chi^2 = 0.08$, $N = 31$, $P = 0.779$; Table 2), and did not differ significantly among years either, as no factor was retained in the final model. Nor did we find significant differences in the latency to attack between model colour morphs (Likelihood test; $\chi^2 = 0.17$, $N = 22$, $P = 0.680$; Fig. 3b), or among years, as no factor was retained in the final model. The part of the body (tail or rest-of-body) that was

318 attacked in red and striped-tailed lizard models did not differ significantly either
319 (Likelihood test; $\chi^2 = 2.08$, $N = 16$, $P = 0.150$; Table 2).

322 **Discussion**

323
324 Our first goal was to find out whether red tails increase lizard detection rate by avian
325 predators. The field study showed that red-tailed models were attacked sooner than
326 striped-tailed, suggesting that the red colouring did indeed make models more easily
327 detected by avian predators. Previous studies checking the decoy hypothesis with green
328 (Castilla et al. 1999) and blue (Watson et al. 2012) tails concluded that they were not
329 significantly more detected by predators than brown or black, respectively. However,
330 recent studies found that lizard models with blue tails were attacked sooner and more
331 often than models with brown tails (Bateman et al. 2014). Blue and green are relatively
332 common colours among lizards (Arnold 1984; Pianka and Vitt 2006), possibly because
333 they are inconspicuous at long distances against a background of vegetation, although
334 they might serve as a lure at closer distances (Arnold 1984, Bateman et al. 2014). Red,
335 however, is very conspicuous against a background of vegetation (e.g., Schaefer et al.
336 2006), which may explain why red-tailed models in our study were more easily
337 detected, although other mechanisms such as a possible pre-existing sensory bias in
338 avian predators towards red objects (e.g., Møller and Erritzøe 2010) could also explain
339 the results. Determining the precise mechanism underlying the behaviour observed
340 (avian predators attacking red-tailed models sooner) is beyond the scope of this study,
341 but our results suggest that red colouration attracts bird attention and probably increase
342 predatory pressure on red-tailed lizards. A significant interaction between colour morph

and type of model was also found, but, unfortunately, our experimental design does not allow us to elucidate whether latency to attack differed between types of models or between years.

In free-ranging lizards, dorsal colour pattern is very important for crypsis (Calsbeek and Cox 2012, Lancaster et al. 2007), and we assumed that striped-tailed models matched background colour better than red-tailed. Therefore, a higher rate of attacks on red-tailed models was expected. Our results did not confirm this expectation, as models of both colour morphs were subject to a similar number of attacks in the field study. The fact that red-tailed models were attacked sooner but not more often than striped-tailed is counterintuitive, but the long periods between consecutive model inspections (3-5 days) might have contributed to the mismatch between timing and number of attacks. The similar attack rate for both colour morphs suggests that both were recognized as prey by avian predators. The captivity study showed similar attack rates and latency to attack for lizard models of both colour morphs, but this was expected due to the experimental setup (models of both colour morphs were probably very easily detected in relatively small cages with bare concrete floor).

Our second aim was to check whether red tails are effective decoys for diverting bird attacks from vital body parts. The field study corroborated this hypothesis, as red-tailed models received a larger proportion of attacks to the tail than striped-tailed models. Birds usually attack the head and body of their prey (Greene 1988; Shepard 2007; Vervust et al. 2011), so red tails seem to cause a change in avian predatory behaviour, thus increasing lizard probability of surviving an attack, particularly when the tail can easily be detached from the rest of the body (even though loss of the tail may imply later costs; see the Introduction). The benefits of having red tails might then outweigh the costs of increased detection by predators (Cooper and Vitt 1991), what

might have driven the evolution and maintenance of this trait in *A. erythrurus*. In the captivity study, a similar pattern of a larger proportion of attacks on the tail was observed for red-tailed models, but the difference between both colour morphs was not statistically significant, maybe in part because of the small sample size.

The present study suggests that red tails in lizards are effective decoys for avian predators, a function that has also been suggested for green and blue tails (Bateman et al. 2014; Castilla et al. 1999; Watson et al. 2012). The tail colour developed in different lizard species does not seem to be strongly phylogenetically constrained, as the same colour may be present in species of different families (e.g., red tails are found in *Morethia ruficauda*, Scincidae family, *Vanzosaura rubricauda*, Gymnophthalmidae family, or *Acanthodactylus erythrurus*, Lacertidae family), whereas different colours may be present in species of the same genus (e.g., *Acanthodactylus erythrurus* and *A. schreiberi* juveniles have red tails, and *A. boskianus* and *A. beershebensis* juveniles have blue tails). Therefore, the specific tail colouration of each species might be more influenced by environmental variables (e.g., contrast with background (Håstad et al. 2005; Rosenblum 2006)), other lizard traits (e.g., contrast with the rest of the body (Arnold 1984)), or even the predator's visual characteristics (e.g., different avian taxa can discriminate wavelengths at slightly different spectral ranges (Håstad et al. 2005)).

If conspicuously coloured lizard tails are an effective decoy for avian predators, we wonder why some lizard species lose this anti-predatory mechanism during the ontogenetic process (Booth 1990). *A. erythrurus* lizards, for example, lose the red colouration when they reach sexual maturity in the case of males, or when they become gravid in the case of females (Cuervo and Belliure 2013; Seva Román 1982). This ontogenetic colour change might be due to differences in behaviour between age classes in *A. erythrurus* (B. Fresnillo, J. Belliure, and J. J. Cuervo, unpublished data) and other

lizard species (Hawlena 2009; Hawlena et al. 2006), as hatchlings and juveniles are usually more active than adults and, therefore, more vulnerable to predators (Jackson et al. 1976). For age classes with high activity rate, the development of an anti-predatory mechanism such as a lure to divert attacks to expendable body parts may be advantageous. However, adults might increase their survival rates by developing more cryptic colouration (Arnold 1984; Vitt and Cooper 1986).

In conclusion, this study suggests that lizards with conspicuous red tails are more easily detected by avian predators, but also that red tails effectively divert avian predator attacks from vital body parts to the expendable tail. If the benefits of diverting attacks to the tail exceed the costs of being more detectable by predators, red tails have a positive net effect on survival and, thus, on fitness, then favouring the evolution and maintenance of this conspicuous colouration. Therefore, our results support the decoy or deflection hypothesis as an explanation for conspicuous red colouration in juvenile lizards.

Acknowledgments

We thank C. Esteban, J. Calatayud, M. Cruz, M. Almarcha and C. Zaragoza for their help in the field study, and the staff of the Santa Faz Wildlife Rehabilitation Centre (Alicante, Spain) for their help in the captivity study. Deborah Fuldauer revised English language usage. These studies were funded by the Spanish Ministry of Education and Science and the European Regional Development Fund (Grant CGL2008-00137/BOS).

References

- Arnold EN (1984) Evolutionary aspects of tail shedding in lizards and their relatives. *Journal of Natural History* 18:127-169
- Arnold EN (1988) Caudal autotomy as a defense. In: Gans C, Huey RB (eds) *Biology of the Reptilia*. Vol. 16. Ecology B: Defense and Life History. Alan R. Liss, Inc., New York, USA, pp 235-274
- Bateman PW, Fleming PA (2009) To cut a long tail short: a review of lizard caudal autotomy studies carried out over the last 20 years. *Journal of Zoology* 277:1-14
- Bateman PW, Fleming PA, Rolek B (2014) Bite me: blue tails as a ‘risky-decoy’ defense tactic for lizards. *Current Zoology* 60:333-337
- Bates D, Maechler M, Bolker B (2012) lme4: Linear mixed-effects models using S4 classes, R package version 0.999999-0 edn. <http://CRAN.R-project.org/package=lme4>
- Baylis SM, Cassey P, Hauber ME (2012) Capsaicin as a deterrent against introduced mammalian nest predators. *The Wilson Journal of Ornithology* 124:518-524
- Belliure J (2006) Lagartija colirroja – *Acanthodactylus erythrurus*. In: Carrascal LM, Salvador A (eds) *Enciclopedia Virtual de los Vertebrados Españoles*. <http://www.vertebradosibericos.org/>, Museo Nacional de Ciencias Naturales, Madrid
- BirdLife International (2014) Species factsheet: *Falco tinnunculus*. Available from <http://www.birdlife.org> (accessed January 2014)
- Booth CL (1990) Evolutionary significance of ontogenic colour-change in animals. *Biological Journal of the Linnean Society* 40:125-163

440 Bowmaker JK (2008) Evolution of vertebrate visual pigments. *Vision Research*
 441 48:2022-2041
 442 Brodie ED III (1993) Differential avoidance of coral snake banded patterns by free-
 443 ranging avian predators in Costa Rica. *Evolution* 47:227-235
 444 Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a
 445 practical information-theoretic approach, 2nd edn. Springer-Verlag New York,
 446 Inc., New York
 447 Calsbeek R, Cox RM (2012) An experimental test of the role of predators in the
 448 maintenance of a genetically based polymorphism. *Journal of Evolutionary*
 449 *Biology* 25:2091-2101
 450 Carretero MA, Llorente GA (1993) Morphometry in a community of Mediterranean
 451 lacertid lizards, and its ecological relationships. *Historia Animalium* 2:77-99
 452 Carrillo J, González-Dávila E (2009) Latitudinal variation in breeding parameters of the
 453 common kestrel *Falco tinnunculus*. *Ardeola* 56:215-228
 454 Carrillo J, Hernández EC, Nogales M, Delgado G, García R, Ramos T (1994)
 455 Geographic variation in the spring diet of *Falco tinnunculus* L. on the islands of
 456 Fuerteventura and El Hierro (Canary Islands). *Bonner Zoologische Beiträge*
 457 45:39-48
 458 Castilla AM, Labra A (1998) Predation and spatial distribution of the lizard *Podarcis*
 459 *hispanica atrata*: an experimental approach. *Acta Oecologica* 19:107-114
 460 Castilla AM, Barbadillo LJ, Bauwens D (1992) Annual variation in reproductive traits
 461 in the lizard *Acanthodactylus erythrurus*. *Canadian Journal of Zoology* 70:395-
 462 402

463 Castilla AM, Gosá A, Galán P, Pérez-Mellado V (1999) Green tails in lizards of the
 464 genus *Podarcis*: do they influence the intensity of predation? *Herpetologica*
 465 55:530-537

466 Clark DRJ, Hall RJ (1970) Function of the blue tail coloration of the five-lined skink
 467 *Eumeces fasciatus*. *Herpetologica* 26:271-274

468 Cooper WE Jr, Vitt LJ (1985) Blue tails and autotomy: enhancement of predation
 469 avoidance in juvenile skinks. *Zeitschrift für Tierpsychologie* 70:265-276

470 Cooper WE Jr, Vitt LJ (1991) Influence of detectability and ability to escape on natural
 471 selection of conspicuous autotomous defenses. *Canadian Journal of Zoology*
 472 69:757-764

473 Cuadrado M, Martín J, López P (2001) Camouflage and escape decisions in the
 474 common chameleon *Chamaeleo chamaeleon*. *Biological Journal of the Linnean*
 475 *Society* 72:547-554

476 Cuervo JJ, Belliure J (2013) Exploring the function of red colouration in female spiny-
 477 footed lizards (*Acanthodactylus erythrurus*): patterns of seasonal colour change.
 478 *Amphibia-Reptilia* 34:525-538

479 Darwin C (1871) *The descent of man, and selection in relation to sex*. D. Appleton and
 480 Company, New York

481 Endler JA (1980) Natural selection on color patterns in *Poecilia reticulata*. *Evolution*
 482 34:76-91

483 Fox SF, McCoy JK (2000) The effects of tail loss on survival, growth, reproduction,
 484 and sex ratio of offspring in the lizard *Uta stansburiana* in the field. *Oecologia*
 485 122:327-334

486 Greene H (1988) Antipredator mechanisms in reptiles. In: Gans C, Huey RB (eds)
 487 Biology of the Reptilia. Vol. 16. Ecology B: Defense and Life History. Alan R.
 488 Liss, Inc., New York, USA, pp 1-152

489 Haskell DG (1996) Do bright colors at nests incur a cost due to predation? *Evolutionary*
 490 *Ecology* 10:285-288

491 Hawlena D (2009) Colorful tails fade when lizards adopt less risky behaviors.
 492 *Behavioral Ecology and Sociobiology* 64:205-213

493 Hawlena D, Bochnik R, Abramsky Z, Bouskila A (2006) Blue tail and striped body:
 494 why do lizards change their infant costume when growing up? *Behavioral*
 495 *Ecology* 17:889-896

496 Håstad O, Victorsson J, Ödeen A (2005) Differences in color vision make passerines
 497 less conspicuous in the eyes of their predators. *Proceedings of the National*
 498 *Academy of Sciences of the United States of America* 102:6391-6394

499 Jackson JF, Ingram W III, Campbell HW (1976) The dorsal pigmentation pattern of
 500 snakes as an antipredator strategy: a multivariate approach. *The American*
 501 *Naturalist* 110:1029-1053

502 Johnson J, Burt DB, DeWitt TJ (2008) Form, function, and fitness: pathways to
 503 survival. *Evolution* 62:1243-1251

504 Kemp DJ, Rutowski RL (2011) The role of coloration in mate choice and sexual
 505 interactions in butterflies. *Advances in the Study of Behavior* 43:55-92

506 Kodandaramaiah U, Lindenfors P, Tullberg BS (2013) Deflective and intimidating
 507 eyespots: a comparative study of eyespot size and position in butterflies.
 508 *Ecology and Evolution* 3:4518-4524

509 Kraus F, Allison A (2009) A remarkable ontogenetic change in color pattern in a new
510 species of *Oreophryne* (Anura: Microhylidae) from Papua New Guinea. *Copeia*
511 2009:690-697

512 Lancaster LT, McAdam AG, Wingfield JC, Sinervo BR (2007) Adaptive social and
513 maternal induction of antipredator dorsal patterns in a lizard with alternative
514 social strategies. *Ecology Letters* 10:798-808

515 Martín J, López P (1990) Amphibians and reptiles as prey of birds in Southwestern
516 Europe. *Smithsonian Herpetological Information Service* 82:1-43

517 Mochida K (2011) Combination of local selection pressures drives diversity in
518 aposematic signals. *Evolutionary Ecology* 25:1017-1028

519 Møller AP, Erritzøe J (2010) Why birds eat colourful grit: colour preferences revealed
520 by the colour of gizzard stones. *Journal of Evolutionary Biology* 23:509-517

521 Ochi H, Awata S (2009) Resembling the juvenile colour of host cichlid facilitates
522 access of the guest cichlid to host territory. *Behaviour* 146:741-756

523 Pianka ER, Vitt LJ (2006) *Lizards: Windows to the Evolution of Diversity*. University
524 of California Press, Berkeley, California

525 Pollo CJ, Pérez-Mellado V (1990) Biología reproductora de tres especies mediterráneas
526 de Lacertidae. *Mediterránea, Serie de Estudios Biológicos* 12:149-160

527 R Development Core Team (2013) R: a language and environment for statistical
528 computing. Version 2.15.3. R Foundation for Statistical Computing, Vienna,
529 Austria. <http://www.R-project.org/>

530 Rosenblum EB (2006) Convergent evolution and divergent selection: lizards at the
531 White Sands ecotone. *The American Naturalist* 167:1-15

532 Ruiz-Rodríguez M, Avilés JM, Cuervo JJ, Parejo D, Ruano F, Zamora-Muñoz C, Sergio
 533 F, López-Jiménez L, Tanferna A, Martín-Vivaldi M (2013) Does avian
 534 conspicuous colouration increase or reduce predation risk? *Oecologia* 173:83-93
 535 Schaefer HM, Levey DJ, Schaefer V, Avery ML (2006) The role of chromatic and
 536 achromatic signals for fruit detection by birds. *Behavioral Ecology* 17:784-789
 537 Seva Román E (1982) Taxocenosis de lacértidos en un arenal costero alicantino. Ph.D.
 538 Thesis, Universidad de Alicante, Alicante
 539 Shepard DB (2007) Habitat but not body shape affects predator attack frequency on
 540 lizard models in the Brazilian Cerrado. *Herpetologica* 63:193-202
 541 Stuart-Fox DM, Moussalli A, Marshall NJ, Owens IPF (2003) Conspicuous males
 542 suffer higher predation risk: visual modelling and experimental evidence from
 543 lizards. *Animal Behaviour* 66:541-550
 544 Telemeco RS, Baird TA, Shine R (2011) Tail waving in a lizard (*Bassiana duperreyi*)
 545 functions to deflect attacks rather than as a pursuit-deterrent signal. *Animal*
 546 *Behaviour* 82:369-375
 547 Valkonen J, Niskanen M, Björklund M, Mappes J (2011) Disruption or aposematism?
 548 Significance of dorsal zigzag pattern of European vipers. *Evolutionary Ecology*
 549 25:1047-1063
 550 VanderWerf EA, Freed LA (2003) 'Elepaio subadult plumages reduce aggression
 551 through graded status-signaling, not mimicry. *Journal of Field Ornithology*
 552 74:406-415
 553 Vanzyl AJ (1994) Comparison of the diet of the common kestrel *Falco tinnunculus* in
 554 South Africa and Europe. *Bird Study* 41:124-130

555 Vázquez RA, Ebensperger LA, Bozinovic F (2002) The influence of habitat on travel
 556 speed, intermittent locomotion, and vigilance in a diurnal rodent. Behavioral
 557 Ecology 13:182-187
 558 Vervust B, Grbac I, Van Damme R (2007) Differences in morphology, performance and
 559 behaviour between recently diverged populations of *Podarcis sicula* mirror
 560 differences in predation pressure. Oikos 116:1343-1352
 561 Vervust B, Van Loy H, Van Damme R (2011) Seeing through the lizard's trick: do
 562 avian predators avoid autotomous tails? Central European Journal of Biology
 563 6:293-299
 564 Vitt LJ, Cooper WE Jr (1986) Tail loss, tail color, and predator escape in
 565 *Eumeces* (Lacertilia: Scincidae): age-specific differences in costs and benefits.
 566 Canadian Journal of Zoology 64:583-592
 567 Watson CM, Roelke CE, Pasichnyk PN, Cox CL (2012) The fitness consequences of
 568 the autotomous blue tail in lizards: an empirical test of predator response using
 569 clay models. Zoology 115:339-344
 570 Wilson BS (1992) Tail injuries increase the risk of mortality in free-living lizards (*Uta*
 571 *stansburiana*). Oecologia 92:145-152
 572

Table 1. Generalized linear model with Poisson error distribution analysing the effects of type of model (plaster or plasticine) and model colour morph (red or striped tail) on the latency to attack. The interaction between type of model and colour morph is also shown. Study area and interactions including this factor were not retained in the final model after a backward stepwise procedure (see Statistical analyses for details). Only models attacked by birds were included in the analysis ($N = 63$).

	d.f.	χ^2	<i>P</i>
Type of model	1	124.61	< 0.001
Colour morph	1	30.47	< 0.001
Type of model x Colour morph	1	8.49	0.004

588 **Table 2.** Percentage of lizard models that were attacked by birds and percentage of attacks on the tail for red- and striped-tailed models in the
589 field and captivity studies. Percentage of attacks on the tail was calculated considering only models attacked by birds on a single body part (tail
590 or rest-of-body).

591

592

593

594

595

596

597

598

599

600

	Type of study	<i>N</i>	Red-tailed models	Striped-tailed models
Percentage of models attacked	Field	115	56.45	52.83
	Captivity	31	69.23	58.33
Percentage of attacks on the tail	Field	38	47.62	17.65
	Captivity	16	77.88	42.68

Figure legends

Fig. 1 Types of models used in the study: (a) red-tailed plaster model (tail completely red), (b) striped-tailed plaster model (dark and light banded tail pattern), (c) red-tailed plasticine model (red tail with a light band on the back), and (d) striped-tailed plasticine model (dark and light banded tail pattern).

Fig. 2 Types of marks left on the models by predators: (a) V-shaped mark left by a bird on a plasticine model, (b) U-shaped mark left by a bird on a plasticine model, (c) puncture mark left by a bird on a plaster model, (d) V-shaped mark with tooth marks left by lizards on a plasticine model, (e) tooth marks left by rodents on a plasticine model, (f) tooth marks left by rodents on a plaster model, and (g) tracks left on the ground after an avian attack when the model disappeared.

Fig. 3 Mean ($\pm$ SE) latency to attack for red and striped-tailed models in (a) the field study by type of model (plaster or plasticine) ($N = 38$) and (b) the captivity study ($N = 16$). Only models attacked by birds are included.

Fig. 1

Fig. 2

Fig. 3