1
2
3
4
5
6 Effect of an icing medium containing the alga *Fucus spiralis* on
7 the microbiological activity and lipid oxidation in chilled megrim
8 (*Lepidorhombus whiffiagonis*)
9
10
11
12

13 José M. Miranda¹, Marcos Trigo², Jorge Barros-Velázquez¹, Santiago P. Aubourg^{2,*}
14
15
16
17
18
19

20 ¹ Department of Analytical Chemistry, Nutrition and Food Science, School of
21 Veterinary Sciences, University of Santiago de Compostela, Lugo (Spain).

22 ² Department of Food Science and Technology, Marine Research Institute (CSIC), Vigo
23 (Spain).

24 * Corresponding author. Phone: +34986231930; Fax: +34986292762; e-mail:
25 saubourg@iim.csic.es
26
27
28

ABSTRACT

The present study provides a first approach on the employment of an icing medium including *Fucus spiralis*, an alga exhibiting antimicrobial and antioxidant activities, for the preservation of fish quality during chilled storage. For it, two different concentrations of a *F. spiralis* extract (0.67 and 2.50 g lyophilized alga L⁻¹ aqueous solution; F-1 and F-2 batches, respectively) were tested as icing medium for the chilled storage of megrim (*Lepidorhombus whiffiagonis*) for 14 days. The effects of the alga were compared with a counterpart batch stored in traditional ice prepared only from water (F-0 batch). A significant ($p<0.05$) inhibition of microbial activity (aerobes, psychrotrophs, proteolytic bacteria, lipolytic bacteria; pH and trimethylamine formation) in F-1 batch and, especially in F-2 batch, was concluded. Concerning lipid oxidation development, a significantly ($p<0.05$) lower formation of interaction compounds (fluorescence assessment) in samples corresponding to the F-2 batch was also observed, proving the inhibitory effect of *F. spiralis* on the formation of tertiary lipid oxidation compounds in chilled megrim. The icing medium proposed in this study may open the way to the development of a natural biopreservation strategy for chilled seafood based on algae.

Keywords: *Lepidorhombus whiffiagonis*; *Fucus spiralis*; chilling; microbiological activity; lipid oxidation; quality.

Head Tittle: Biopreservation of chilled megrim with *Fucus spiralis*.

1. INTRODUCTION

With a view to slow down fish damage during chilled storage, thus extending its shelf life, a wide number of preservation strategies have been combined with flake ice. Among them, several chemical (phenolic compounds, bisulphites, polyphosphates, etc.) and physical (namely, irradiation and high-pressure processing) treatments (Ashie, Smith, & Simpson, 1996; Briones, Reyes, Tabilo-Munizaga, & Pérez-Won, 2010) and the combination with packaging (Sivertsvik, Jeksrud, & Rosnes, 2002; González-Fandos, Villarino-Rodríguez, García-Linares, García-Arias, & García-Fernández, 2005) have been reported. Additionally, recent studies accounted for the incorporation of preservative compounds such as ozone (Pastoriza, Bernárdez, Sampedro, Cabo, & Herrera, 2008), vegetable extracts (namely, thyme, rosemary and oregano) (Oral, Gülmez, Vatansever, & Güven, 2008; Quitral, Donoso, Ortiz, Herrera, Araya, & Aubourg, 2009; Özyurt, Kuley, Balıkçı, Kaçar, Gökdogan, Etyemez, & Özogul, 2012) and natural low-molecular weight organic acids (Sallam, 2007; García-Soto, Fernández-No, Barros-Velázquez, & Aubourg, 2014a) in the icing medium.

Marine algae are known to be part of the diet in different Asiatic countries (namely, Japan, China and Korea) and constitute a source of beneficial nutrients, such as vitamins, trace minerals, lipids, amino acids and dietary fibres (Díaz-Rubio, Pérez-Jiménez, & Saura-Calixto, 2009; Paiva, Lima, Ferreira Patarra, Neto, & Baptista, 2014). Their major use in Western countries has concentrated on the extraction of compounds used in pharmaceutical, cosmetics and food industries (Smit, 2004; Gyawali & Ibrahim, 2014). Recently, red, green and brown macroalgae have offered the possibility of exploring a wide variety of natural compounds with potential antioxidant (Halldorsdóttir, Sveinsdóttir, Gudmundsdóttir, Thorkelsson, & Kristinsson, 2014) and antimicrobial (Sandsdalen, Haug, Stensvag, & Styrvold, 2003) activities susceptible to

be applied to seafood. In this sense, a wide number of preservative metabolites such as polyphenols, terpenes, phlorotannins, steroids, halogenated ketones and alkanes, fucoxanthin, polyphloroglucinol or bromophenols have been isolated from macroalgae (Fleurence, Moránçais, Dumay, Decottignies, Turpin, Munier, García-Bueno, & Jaouen, 2012; Peinado, Girón, Koutsidis, & Ames, 2014). As a result, previous research accounts for a preservative effect of alga extracts on seafood such as chilled cod muscle (Wang, Jónsdóttir, Kristinsson, Thorkelsson, Jacobsen, Yuca Hamaguchi, & Olafsdóttir, 2010), canned Atlantic salmon (Ortiz, Vivanco, & Aubourg, 2014) and fish-oil enriched food (Hermund, Iltas, Honold, Jónsdóttir, Kristinsson, & Jacobsen, 2015).

Flatfish (e.g., flounder, sole, turbot, plaice and halibut) species represent a very important seafood group. Among them, megrim (*Lepidorhombus whiffiagonis*) is abundant in the Northeast Atlantic waters. This species, which is considered one of the most fished species in the Grand Sole North Atlantic Fishing bank, has been exploited by a wide number of European countries, these including the United Kingdom, France, Ireland and Spain (FAO, 2007). Previous studies on megrim have evaluated the effects of different on-board storage conditions on chilled fish quality (Jehanno, Thuault, Larnaud, Alaume, & Bourgeois, 1996; Aubourg, Losada, Gallardo, Miranda, & Barros-Velázquez, 2006). Moreover, the evolution of different damage pathways in megrim has been analysed, these including volatile amine formation during refrigeration (1-3°C) (Civera, Turi, Bisio, Parisi, & Fazio, 1995), microbial activity (Sanjuás-Rey, García-Soto, Fuertes-Gamundi, Aubourg, & Barros-Velázquez, 2012) and lipid oxidation (García-Soto, Sanjuás-Rey, Barros-Velázquez, Fuertes-Gamundi, & Aubourg, 2011) during chilled storage. These reports underline the need of additional studies for the development of natural refrigeration systems to meet the increasing consumer demand for natural, high quality and safer seafood products.

Accordingly, this study evaluates the inclusion, to the best of our knowledge for the first time, of an alga extract in the icing medium employed for the chilled storage of a marine species. According to European Council Regulation (1997) algae are considered food or food ingredients, so their use in the icing medium should not constitute any hazard to health. *F. spiralis* was chosen for this study in agreement with its abundance in the Galician Atlantic coast (North-western Spain) and its promising biopreservation properties revealed recently by other authors (Andrade, Barbosa, Pedro Matos, Lopes, Vinholes, Mouga, & Valentão, 2013; Tierney, Smyth, Hayes, Soler-Vila, Croft, & Brunton, 2013a). For it, aqueous solutions including ethanolic extracts of lyophilised *F. spiralis* were employed at two different concentrations as icing media. The effects of the alga extracts on microbial activity inhibition and lipid oxidation stability were monitored in megrim muscle for up to 14 d of chilled storage by means of microbiological and biochemical analyses.

2. MATERIAL AND METHODS

2.1. Preparation of *Fucus spiralis* extracts and icing systems

The lyophilised alga *F. spiralis* was provided by Porto-Muiños (Cereda, A Coruña, Spain). Fifteen g of lyophilised alga were mixed with absolute ethanol (2 x 120 mL), stirred for 30 s and centrifuged at 3500 rpm for 10 min at 4°C. Then, the supernatant was recovered and diluted to 6 L with distilled water (2.50 g lyophilised alga L⁻¹ aqueous solution). This solution was packaged in polyethylene bags, kept frozen at -18°C and later used as icing medium (F-2 batch). In the same way, 4 g of lyophilised alga were extracted with ethanol as described above in order to provide a more diluted alga-icing medium (0.67 g lyophilised alga L⁻¹ aqueous solution; F-1 batch). Finally, traditional ice was prepared from distilled water that was packaged and kept frozen in

the same way as the two other ices (F-0, control batch). Before addition to individual fish specimens, the different icing systems were ground to obtain ice flakes.

Experimental conditions (content of lyophilised alga extract in the ice) employed in the present study were based on several preliminary tests carried out at our laboratory in the 0.2-10.0 g lyophilised alga L⁻¹ aqueous solution range. Thus, an increasing presence of alga in the icing medium provided better sensory acceptance. However a concentration of 2.50 g L⁻¹ resulted in the highest concentration without modifying the sensory descriptors (external odour and colour, as well as flesh odour and flavour) of the fish (data not shown). Consequently, this concentration was considered in the F-2 batch, together with a less concentrated concentration (0.67 g L⁻¹, F-1 batch).

2.2. Fish material, processing and sampling

Fresh megrim (117 specimens) were caught near the Galician Atlantic coast (North-Western Spain) and transported to the laboratory. Throughout this process (10 h), the fish were maintained in ice. The length and weight of the fish specimens ranged from 23 to 27 cm and from 110 to 137 g, respectively.

Upon arrival to the laboratory, nine individual fish specimens were separated and analysed as initial fish (day 0). These fish specimens were divided into three different groups (three individuals per group) that were analysed independently to achieve the statistical analysis (n=3). The remaining fish specimens were divided into three batches (36 individuals in each batch), that were placed in independent boxes and directly surrounded by different kinds of ice (F-0, F-1 and F-2 batches, respectively), prepared as previously described. Ice was added at a 1:1 fish:ice ratio, and all batches were placed inside a refrigerated room (2±1°C). Boxes that allowed draining of melted ice were used for fish storage. The ice of all batches was renewed when required in

order to maintain the 1:1 fish:ice ratio. Fish samples from all of the batches were stored for a 14-day period, being sampled and analysed on days 4, 7, 11 and 14. At each sampling time, nine specimens were taken from each batch for analysis and divided into three groups (three individuals in each group) that were studied independently (n=3).

2.3. Microbiological analyses

Samples of 10 g of fish white muscle were taken aseptically from chilled fish specimens, mixed with 90 mL of 0.1% peptone water (Merck, Darmstadt, Germany) and homogenised in sterilised stomacher bags (AES, Combourg, France) as previously described (Ben-Gigirey, Vieites Baptista de Sousa, Villa, & Barros-Velázquez, 1998; Ben-Gigirey, Vieites Baptista de Sousa, Villa, & Barros-Velázquez, 1999). Serial dilutions from the microbial extracts were prepared in 0.1% peptone water.

Total aerobes were investigated by plating on plate count agar (PCA, Oxoid Ltd., London, UK) after incubation at 30°C for 48 h. The anaerobe counts were also determined in PCA at 30±0.5°C, except that an anaerobic atmosphere kit (Oxoid Ltd.) was placed inside the anaerobiosis jar. Psychrotrophs were also investigated in PCA, being the incubation carried out at 7-8°C for 7 days. Enterobacteriaceae were investigated by pour plating using Violet Red Bile Agar (VRBA) (Merck, Darmstadt, Germany) after an incubation period of 24 h at 37±0.5°C. Microorganisms exhibiting a proteolytic or lipolytic phenotype were determined on casein-agar medium or tributyrine-agar, respectively, after incubation at 30°C for 48 h, as previously described (Ben-Gigirey, Vieites Baptista de Sousa, Villa, & Barros-Velázquez, 2000).

In all cases, microbial counts were transformed into log CFU g⁻¹ muscle before undergoing statistical analysis. All of the analyses were conducted in triplicate.

2.4. Chemical analyses

Total polyphenols content of lyophilised *F. spiralis* was assessed by means of the Folin-Ciocalteu colorimetric method (Cary 3E UV–Visible spectrophotometer, Varian; Mulgrave, Victoria, Australia) as described previously (Rodríguez-Bernaldo de Quirós, Frecha-Ferreiro, Vidal-Pérez, & López-Hernández, 2010). Measurements were made in triplicate. Gallic acid (GA) was used as standard. Results were expressed as mg GA g⁻¹ lyophilised alga.

Chemical analyses related to fish quality were carried out on the white muscle of megrim. All solvents and chemical reagents used were of reagent grade (Merck, Darmstadt, Germany).

The evolution of pH values in megrim muscle along storage time was determined using a 6-mm diameter insertion electrode (Crison, Barcelona, Spain).

Trimethylamine-nitrogen (TMA-N) values were determined using the picrate colorimetric (Beckman Coulter, DU 640; London, UK) method, as previously described by Tozawa, Erokibara, & Amano (1971). This method involves the preparation of a 5% trichloroacetic acid extract of fish muscle (10 g/25 mL). The results are expressed as mg TMA-N kg⁻¹ muscle.

Lipids were extracted from the fish muscle by the Bligh & Dyer (1959) method, which employs a single-phase solubilisation of the lipids using a chloroform-methanol (1:1) mixture. The results were calculated as g lipid kg⁻¹ muscle.

Free fatty acid (FFA) content was determined in the lipid extract of the fish muscle by the Lowry & Tinsley (1976) method based on complex formation with cupric acetate-pyridine followed by spectrophotometric (715 nm) assessment. Results were expressed as g FFA kg⁻¹ lipids and as mg FFA kg⁻¹ muscle.

The peroxide value (PV) was determined spectrophotometrically (Beckman Coulter, DU 640; London, UK) using the lipid extract via previous peroxide reduction with ferric thiocyanate according to the Chapman & McKay (1949) method. The results are expressed as meq active oxygen kg⁻¹ lipids.

The thiobarbituric acid index (TBA-i) was determined according to Vyncke (1970). This method is based on the reaction between a trichloroacetic acid extract of the fish muscle and thiobarbituric acid. Content of thiobarbituric acid reactive substances (TBARS) was spectrophotometrically measured at 532 nm and calculated from a standard curve using 1,1,3,3-tetraethoxy-propane (TEP); results were expressed as mg malondialdehyde kg⁻¹ muscle.

Tertiary lipid oxidation compounds resulting from the interaction between oxidised lipids and nucleophilic compounds (namely, protein-like molecules) were measured by fluorescence spectroscopy. Such measurement has shown to be a valuable tool for assessing the fish quality changes during processing (Losada, Piñeiro, Barros-Velázquez, & Aubourg, 2004). Thus, the formation of fluorescent compounds (Fluorimeter LS 45; Perkin Elmer España; Tres Cantos, Madrid, Spain) was determined by measurements at 393/463 nm and 327/415 nm. The relative fluorescence (RF) was calculated as follows: $RF = F/F_{st}$, where F is the fluorescence measured at each excitation/emission maximum and F_{st} is the fluorescence intensity of a quinine sulphate solution (1 µg mL⁻¹ in 0.05 M H₂SO₄) at the corresponding wavelength. The fluorescence ratio (FR) was calculated as the ratio between the two RF values: $FR = RF_{393/463\text{ nm}}/RF_{327/415\text{ nm}}$. The FR value was determined using the aqueous phase that resulted from the lipid extraction of the fish muscle (Bligh & Dyer, 1959).

2.5. Statistical analysis

Data obtained from the different microbiological and chemical analyses were subjected to the ANOVA method to explore differences resulting from the effects of both the icing condition and the chilling time; the comparison of means was performed using the least-squares difference (LSD) method. In all cases, analyses were carried out using the PASW Statistics 18 software for Windows (SPSS Inc., Chicago, IL, USA); differences among batches and among refrigeration times were considered significant for a confidence interval at the 95% level ($p < 0.05$) in all cases.

Correlation analysis among parameters (chilling storage time, microbiological values and chemical indices) was also carried out. In it, linear fittings are mentioned; otherwise, the kind of fitting is expressed.

3. RESULTS AND DISCUSSION

3.1. Analysis of the microbial development in megrim

Preliminary plate bioassays carried out at our laboratory showed that ethanolic extracts of *F. spiralis* exhibited antimicrobial activity against *Bacillus cereus*, *Bacillus subtilis*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Pseudomonas fluorescens*, *Escherichia coli*, *Aeromonas hydrophila*, *Vibrio alginolyticus* and *Vibrio parahaemolyticus* (data not shown). Additionally, a high polyphenol content (53.3 ± 5.0 GA g⁻¹ lyophilised alga) was determined in ethanolic extracts of *F. spiralis*.

The results of the investigation of aerobic mesophiles in all three batches are compiled in Table 1. A progressive increase was observed for all kinds of samples throughout the chilling storage time ($r^2 = 0.93$ - 0.95 , quadratic fitting). The inclusion of *F. spiralis* extract in the icing medium provided a better control of aerobes as compared with the control batch (F-0), the microbial inhibition reaching its maximum (1.25 log

units) at advanced storage periods. This was especially remarkable in the case of the F-2 batch, which included a higher concentration of *F. spiralis* extract. Thus, statistically significant ($p < 0.05$) differences were observed between the control batch (F-0) and F-2 batch after 11 and 14 d of storage. In contrast, F-1 batch provided an intermediate protection against aerobes growth as compared with F-2 batch, and no significant ($p > 0.05$) differences were observed between F-1 and F-0 batches (Table 1).

The development of psychrotrophs in megrim batches is also presented in Table 1. Psychrotrophs normally include bacteria belonging to genera *Pseudomonas*, *Moraxella*, *Acinetobacter*, *Shewanella* and *Flavobacterium*, some of them being of concern for the preservation of microbial fish quality. The evolution of psychrotrophs followed a similar behaviour as that observed for aerobes. Thus, a progressive increase was observed for all kinds of samples throughout the chilling storage time ($r^2 = 0.93-0.95$). The batches including *F. spiralis* extracts showed slight decreases of psychrotrophs counts as compared with the control batch, these reductions being more remarkable for F-2 batch at advanced storage times. Thus, significantly ($p < 0.05$) lower psychrotrophs counts were determined in the F-2 batch on day 11, such differences reaching 0.89 log units at that storage time. This result, as in the case of aerobes, indicates a positive effect derived from the inclusion of *F. spiralis* extract in the icing medium in terms of microbial inhibition in megrim muscle.

The investigation of Enterobacteriaceae (Table 1) provided similar results in all three batches. Remarkably, the numbers of this microbial group were very low in all batches at all storage times, a result that confirms the good initial microbial quality of megrim specimens. Thus, numbers below 1 log CFU g⁻¹ were determined in all batches even after 11 d of storage. As a consequence of this, no significant ($p > 0.05$) differences among batches could be concluded.

The evolution of proteolytic microorganisms in all three batches is presented in Figure 1. The results obtained for this microbial group were in agreement with those observed for aerobes and psychrothrophs. Thus, a progressive increase ($p < 0.05$) was observed for all kinds of samples throughout the chilling storage time ($r^2 = 0.92-0.94$). The presence of *F. spiralis* extracts in the icing medium, especially in the case of the F-2 batch, provided a significant ($p < 0.05$) protection against this microbial group at advanced storage times. The microbial inhibition reached its maximum (1.01 log units) on day 11 (Figure 1). The crucial role of proteolytic microorganisms in the degradation of fish muscle has been previously reported (Rodríguez, Barros-Velázquez, Ojea, Piñeiro, & Aubourg, 2003). Accordingly, the partial inhibition of the action of this microbial group due to the presence of *F. spiralis* extracts in the icing medium tested in this study is another remarkable result in terms of fish quality.

Figure 2 compiles the results of the investigation of lipolytic bacteria in all three batches. A progressive increase ($p < 0.05$) was observed for all kinds of samples throughout the chilling storage time ($r^2 = 0.89-0.93$). Significant ($p < 0.05$) differences among F-2 and F-0 batches were observed after 4 d of storage. Remarkably, such differences reached maximum values of 1.54 log CFU g⁻¹ after 11 d of storage, these being the most marked differences among batches determined for any microbial group.

The antimicrobial effect observed in this study on megrim for aerobes, psychrothrophs, proteolytic and lipolytic bacteria, especially at advanced storage times, is a consequence of the incorporation of *F. spiralis* ethanolic extracts in the icing medium. Chemicals responsible for preservative activities are widespread in macroalgae. The antimicrobial effect of *Fucus* spp. and other algae has been linked to the presence of terpenes and polyphenols (Sandsdalen et al., 2003) and oligomeric phlorotannins (Wang et al., 2010), although other compounds might also be involved

(Fleurence et al., 2012; Peinado et al., 2014). In a closely related alga species (*Fucus vesiculosus*), Sandsdalen et al. (2003) identified by electrospray ionization mass spectrometry a polyhydroxylated fucophlorethol as being responsible for antimicrobial activity against both the Gram-positive and Gram-negative bacteria tested. Interesting antimicrobial substances in particular are the halogenated compounds such as haloforms, halogenated alkanes and alkenes, alcohols, aldehydes, hydroquinones and ketones (Smit, 2004). Some of them have been applied for their effectiveness as active ingredient in bacterial antifouling (Kjelleberg & Steinberg, 2001).

Other authors have also reported that the inclusion of natural compounds in the icing system can reduce the microbial activity. Among them, the following two should be remarked: (i) the inclusion of a rosemary extract in the icing medium and its application to the chilled storage of sardines (*Sardinella aurita*) (Özyurt et al., 2012), and (ii) the incorporation of wild-thyme hydrosol extract to the chilling medium of Transcaucasian barb (*Capoeta capoeta capoeta*) (Oral et al., 2008). However, the present study represents, to our knowledge, the first report of *F. spiralis* extract in the icing medium for the preservation of fish.

This work, carried out on megrim, also complements previous reports focused on the preservation of this fish species under other advanced refrigeration systems, such as slurry ice (Aubourg et al., 2006) or addition of ozone to the icing medium (Pastoriza et al., 2008). Slurry ice slowed down microbial growth in fish muscle due to the washing effect of the salt solution on the fish surface during ice melting (Losada et al., 2004; Campos, Rodríguez, Losada, Aubourg, & Barros-Velázquez, 2005). In the present work, the melting of the flake ice crystals containing *F. spiralis* extract might exert a similar washing effect that may reduce the surface microbial load, thus limiting its diffusion towards megrim muscle.

Natural organic acids, such as citric acid and lactic acid, have previously been used for the preparation of flake ice used for megrim storage with good results (García-Soto, Böhme, Barros-Velázquez, & Aubourg, 2014b). Likewise, the inclusion of ascorbic acid, combined with citric and lactic acids, has also been reported to provide control of aerobic mesophiles in megrim as well as in hake (*Merluccius merluccius*) and angler (*Lophius piscatorius*) (Sanjuás-Rey et al., 2012).

3.2. Analysis of the chemical changes

Two chemical quality parameters (namely, pH and TMA-N values) closely related to the microbial activity development were studied.

Concerning the pH value, a significant ($p < 0.05$) and progressive increase was observed for all kinds of samples along chilling storage time ($r^2 = 0.90-0.95$, quadratic fitting) (Figure 3). An inhibitory effect on pH increase was observed as a result of including the lyophilised alga in the icing medium. At all storage times, analysis of the mean values of the different fish-sample groups led to the following pH increasing sequence: F-2 < F-1 < F-0, so that a 7.18 score was obtained for control fish at the end of the study. Differences between control and alga-treated fish were found significant ($p < 0.05$) at days 7 and 14. Comparison among both alga concentrations showed lower mean values in the F-2 batch, being the difference significant ($p < 0.05$) at day 4.

Increases in the pH value of fish muscle indicate the accumulation of alkaline compounds, such as ammonia, TMA and other nitrogen-including compounds, which are mainly derived from microbial activity. Thus, pH values showed good correlation values with microbiological parameters ($r^2 = 0.76-0.95$), the best scores being observed for aerobe counts ($r^2 = 0.89-0.95$). In agreement with these results, previous studies have reported an inhibition of pH increase as a result of using other natural preservative

compounds in ice during the chilled storage of other marine species. These studies included the use of oregano and rosemary extracts during the chilled storage of Chilean jack mackerel (*Trachurus murphyi*) (Quitral et al., 2009), a rosemary extract applied to sardine (*Sardinella aurita*) (Özyurt et al., 2012) and a wild-thyme hydrosol extract employed for the chilled storage of Transcaucasian barb (Oral et al., 2008).

A progressive formation of TMA was observed as storage time progressed ($r^2 = 0.88-0.94$, quadratic fitting) in all batches (Table 2). As a result of this, notably high TMA contents were determined at the end of the experiment, this being in agreement with previous studies on megrim chilled storage (Civera et al., 1995; Aubourg et al., 2006; Sanjuás-Rey et al., 2012). Comparison among batches allowed to conclude a significant ($p < 0.05$) inhibitory effect at days 7 and 11 in the F-2 batch.

Volatile amine compounds are produced partially by means of endogenous enzyme activity but mostly as a result of microbial development (Ashie et al., 1996; Campos, Gliemmo, Aubourg, & Barros-Velázquez, 2012). Among them, TMA is one of the most commonly employed quality methods to assess microbial activity in marine species kept under refrigerated conditions. In agreement with this close relationship with microbial activity, TMA values showed good correlation values with aerobe, psychrotroph and proteolytic counts ($r^2 = 0.77-0.91$), showing the best scores when compared with the aerobe counts ($r^2 = 0.88-0.91$). Additionally, a good correlation was obtained with the pH value ($r^2 = 0.84-0.90$).

A previous report (Sanjuás-Rey et al., 2012) showed that the employment of a commercial formula including natural organic acids (lactic, citric and ascorbic) in the icing system did not produce any significant improvement in terms of TMA formation. However, if citric and lactic acids were directly included in the icing medium, a small inhibitory effect on TMA formation was observed (García-Soto et al., 2014a; García-

Soto et al., 2014b). Additionally, the application of other preservative strategies such as slurry ice (Aubourg et al., 2006) and ozonised ice (Pastoriza et al., 2008) has shown to inhibit TMA formation in chilled fish.

Lipid hydrolysis was measured by means of the FFA content (Table 2). This parameter revealed a progressive formation in all cases throughout storage time ($r^2 = 0.89-0.93$, quadratic fitting). Comparison among the different samples under study did not provide a definite trend concerning the effect of *F. spiralis* on lipid hydrolysis. Thus, a higher FFA formation ($p < 0.05$) was observed at advanced storage (11-14 days) in F-1 batch as compared to F-0 batch. On the other hand, higher mean values were determined in the control batch in the 7-14-day period as compared with the alga-concentrated fish group (F-2 batch). Similar conclusions were obtained when FFA content is considered on muscle basis (mg kg^{-1} muscle). In this case, FFA formation also showed a good correlation value with the storage time in all cases ($r^2 = 0.88-0.92$, quadratic fitting).

Although the formation of FFA itself does not lead to nutritional losses, accumulation of FFA has been related to some extent to lack of acceptability, since FFAs are known to have detrimental effects on protein properties (Sikorski & Kolakowska, 1994) and oxidise faster than higher-molecular weight lipid classes (namely, triglycerides and phospholipids) by providing a greater accessibility (lower steric hindrance) to oxygen and other pro-oxidant molecules. FFA formation during chilling storage has been reported to be produced as a result of both endogenous enzyme activity and microbial activity (Ashie et al., 1996; Campos et al., 2012). Before the end of the microbial lag phase (up to 6-9 days, depending on several factors), FFA formation should be produced mostly as a result of endogenous enzyme (namely, lipases and phospholipases) activity. Later on, microbial activity should gain importance, so that

FFA formation should be produced especially as a result of bacterial catabolic processes. The fact that a definite trend of the effect of *F. spiralis* on FFA content was not found in the present work could be explained by the fact that different FFA formation mechanisms are involved. FFA values determined in the present study showed good correlation values with some indices related to microbial activity, such as aerobe counts ($r^2 = 0.86-0.93$), pH value ($r^2 = 0.86-0.92$) and TMA-N content ($r^2 = 0.92-0.94$).

In agreement with the present results, an inhibitory effect of a rosemary extract on lipid hydrolysis in chilled sardine was not observed by Özyurt et al. (2012). In contrast, an inhibitory effect on fish FFA formation could be observed by the application of a commercial formula including organic acids (citric, lactic and ascorbic) in the icing medium during megrim storage (García-Soto et al., 2011) as well as by including a rosemary and oregano extract to the preservation of chilled Chilean jack mackerel (Quitral et al., 2009). In addition, a lower FFA formation was observed in blue whiting (*Micromesistius poutassou*) muscle after a two-step process that consisted of dipping and icing fish with the above-mentioned formula (Sanjuás-Rey, García-Soto, Barros-Velázquez, Fuertes-Gamundi, & Aubourg, 2011).

Primary and secondary lipid oxidation development in megrim muscle (3.9-5.1 g lipids kg^{-1} muscle) was measured by assessment of the peroxide and TBARS content, respectively (Table 3). Formation of peroxides throughout storage time provided, with some exceptions, an increase with storage time, although correlation values were not satisfactory. This increase was especially marked in all cases at the end of the storage time. Comparison among batches showed higher values in fish specimens corresponding to the F-2 batch throughout the 4-14-day period. In contrast, the less-concentrated alga batch (F-1) showed higher mean values in the 11-14-day period when

compared with control fish. Consequently, an enhancement effect on peroxide content can be concluded by the presence of alga extract in the icing medium.

Concerning secondary lipid oxidation, TBARS formation can be considered as very low, being included all scores in the 0.22-0.44 range. TBARS content increased with storage time although correlation was poor. Comparison among batches provided only slight differences, so that a general trend of the effect of *F. spiralis* on TBARS formation in megrim muscle could not be concluded.

Tertiary lipid oxidation was assessed by means of fluorescent compounds formation (Table 3). With some exceptions, a general increase was observed as storage time progressed ($r^2 = 0.80-0.90$, quadratic fitting). Such increase was especially relevant in all batches at day 11. Comparison among the different samples showed lower mean FR values in samples corresponding to the F-2 batch in the 11-14-day period, being differences significant ($p < 0.05$) at the end of the storage time. Consequently, an inhibitory effect of *F. spiralis* on tertiary lipid oxidation compounds formation could be concluded. This inhibitory effect is in agreement with the high level of polyphenol compounds (53.3 ± 5.0 GA g⁻¹ lyophilised alga) determined in the ethanolic extracts of *F. spiralis*.

Lipid oxidation has been recognised as a complex process where different kinds of molecules are produced, most of them unstable, susceptible to breakdown and to originate lower-weight compounds, or to react with other molecules (nucleophilic-type, mostly) present in the fish muscle. As a result of this, determination of each kind of compound cannot always provide an accurate method for the quality loss assessment in fish. In the present study, primary and secondary lipid oxidation compounds did not provide a satisfactory correlation with storage time. The electrophilic nature of such compounds led them to interact with food constituents possessing nucleophilic

functions (Losada et al., 2004; Campos et al., 2012), so that FR assessment provided the best correlation value with storage time, this being in agreement with previous studies on chilled megrim (Aubourg et al., 2006; García-Soto et al., 2011). Additionally, FR values showed good correlation scores with aerobe counts ($r^2 = 0.81-0.90$), and fair with other chemical parameters such as pH value ($r^2 = 0.76-0.85$), TMA-N value ($r^2 = 0.61-0.92$) and FFA content ($r^2 = 0.63-0.93$).

The antioxidant capacity and reducing ability of ethanolic extracts of alga *F. spiralis* were previously reported by means of in-vitro DPPH and FRAP tests (Andrade et al., 2013; Peinado et al., 2014). This alga showed a marked content on polyphenols (90-205 μg phloroglucinol equivalents mg^{-1}) (Tierney et al., 2013a) and α -tocopherol (511.4 mg kg^{-1}) (Paiva et al., 2014). Additionally, a preliminary identification of active compounds was carried out by quadrupole time-of-flight mass spectrometry (Q-TOF-MS) (Tierney, Smyth, Rai, Soler-Vila, Croft, & Brunton, 2013b); as a consequence, the results supported the assumption that phlorotannins were present and were probably involved in the antioxidant activity.

Previous studies have shown that including natural preservative compounds in the icing system provides an inhibitory effect on fish lipid oxidation. Directly related to the present work, the employment of a commercial formula including citric, lactic and ascorbic acids led to a lower FR score during the chilled storage of megrim and hake (García-Soto et al., 2011) and blue whiting (Sanjuás-Rey et al., 2011). The same result was obtained when a laboratory-scale solution of citric and lactic acids was employed as icing medium during megrim storage (García-Soto et al., 2014b). Additionally, primary and secondary lipid oxidation compounds formation was inhibited in Chilean jack mackerel as a result of including oregano and rosemary extracts in the icing system (Quitral et al., 2009) and rosemary extract in sardine icing system (Özyurt et al., 2012).

4. CONCLUSIONS

The development and application of an icing system including an ethanolic extract of the alga *F. spiralis* has shown a protective effect on refrigerated megrim quality, according to the assessment of microbial activity and lipid oxidation development in the fish muscle. The present study provides a first approach focused on the employment of this abundant alga species, whose preservative properties and components have already been described in *in vitro* experiments. The results presented in this work constitute a promising basis in order to apply algae extracts in the icing system to enhance fish quality retention during storage in the different steps (on-board, transport, retail market, etc.) included in the commercialisation of refrigerated marine species.

ACKNOWLEDGEMENTS

The authors thank Mrs. Monserrat López for her excellent technical assistance. Porto-Muiños (Cerdeira, A Coruña, Spain) is greatly acknowledged for kindly providing the lyophilised alga. This work was supported by the Consejo Superior de Investigaciones Científicas (CSIC; Spain) through the project PIE 201370E001.

REFERENCES

- Andrade, P., Barbosa, M., Pedro Matos, R., Lopes, G., Vinholes, J., Mouga, T., & Valentão, P. (2013). Valuable compounds in macroalgae extracts. *Food Chemisrty*, 138, 1819-1828.
- Ashie, I., Smith, J., & Simpson, B. (1996). Spoilage and shelf-life extension of fresh fish and shellfish. *Critical Reviews in Food Science and Nutrition*, 36, 87-121.
- Aubourg, S., Losada, V., Gallardo, J., Miranda, M., & Barros-Velázquez, J. (2006). On-board quality preservation of megrim (*Lepidorhombus whiffiagonis*) by a novel ozonised-slurry ice system. *European Food Research Technology*, 223, 232-237.
- Ben-Gigirey, B., Vieites Baptista de Sousa, J., Villa, T., & Barros-Velázquez, J. (1998). Changes in biogenic amines and microbiological analysis in albacore (*Thunnus alalunga*) muscle during frozen storage. *Journal of Food Protection*, 61, 608-615.
- Ben-Gigirey, B., Vieites Baptista de Sousa, J., Villa, T., & Barros-Velázquez, J. (1999). Histamine and cadaverine production by bacteria isolated from fresh and frozen albacore (*Thunnus alalunga*). *Journal of Food Protection*, 62, 933-939.
- Ben-Gigirey, B., Vieites Baptista de Sousa, J., Villa, T., & Barros-Velázquez, J. (2000). Characterization of biogenic amine-producing *Stenotrophomonas maltophilia* strains isolated from white muscle of fresh and frozen albacore tuna. *International Journal of Food Microbiology*, 57, 19-31.
- Bligh, E., & Dyer, W. (1959). A rapid method of total extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37, 911-917.
- Briones, L., Reyes, J., Tabilo-Munizaga, G., & Pérez-Won, M. (2010). Microbial shelf-life extension of chilled Coho salmon (*Oncorhynchus kisutch*) and abalone

519 (*Haliotis rufescens*) by high hydrostatic pressure treatment. *Food Control*, 21,
520 1530-1535.

521 Campos, C., Gliemmo, M., Aubourg, S., & Barros-Velázquez, J. (2012). Novel
522 technologies for the preservation of chilled aquatic food products. In A.
523 McElhatton, & P. Amaral Sobral (Eds.), *Novel Technologies in Food Science*
524 (pp. 299-323), chapter 13. New York, USA: Springer.

525 Campos, C., Rodríguez, O., Losada, V., Aubourg, S., & Barros-Velázquez, J. (2005).
526 Effects of storage in ozonised slurry ice on the sensory and microbial quality of
527 sardine (*Sardina pilchardus*). *International Journal of Food Microbiology*, 103,
528 121–130.

529 Chapman, R., & McKay, J. (1949). The estimation of peroxides in fats and oils by the
530 ferric thiocyanate method. *Journal of the American Oil Chemists' Society*, 26,
531 360-363.

532 Civera, T., Turi, R., Bisio, C., Parisi, E., & Fazio, G. (1995). Sensory and chemical
533 assessment of marine teleosteans. Relationship between total volatile basic
534 nitrogen, trimethylamine and sensory characteristics. *Science des Aliments*, 13,
535 109-117.

536 Díaz-Rubio, M^aE., Pérez-Jiménez, J., & Saura-Calixto, F. (2009). Dietary fiber and
537 antioxidant capacity in *Fucus vesiculosus* products. *International Journal of*
538 *Food Science and Nutrition*, 60, 23-34.

539 European Council Regulation (1997). European Community (EC), No 258/97, 27
540 January 1997 concerning novel foods and novel food ingredients. *CELEX-EUR*
541 *Official Journal* L-43, 14/02/1997, pp. 1-7.

542 FAO (2007). Fishery statistics. Capture production. In *Food and Agriculture*
543 *Organization of the United Nations* (p. 97) vol. 101/1. Rome, Italy: Yearbook
544 2005.

545 Fleurence, J., Moranchais, M., Dumay, J., Decottignies, P., Turpin, V., Munier, M.,
546 García-Bueno, N., & Jaouen, P. (2012). What are the prospects for using
547 seaweed in human nutrition and for marine animals raised through aquaculture?
548 *Trends in Food Science and Technology*, 27, 57-61

549 García-Soto, B., Böhme, K., Barros-Velázquez, J., & Aubourg, S. (2014b). Inhibition of
550 quality loss in chilled megrim (*Lepidorhombus whiffiagonis*) by employing citric
551 and lactic acid icing. *International Journal of Food Science and Technology*, 49,
552 18-26.

553 García-Soto, B., Fernández-No, I., Barros-Velázquez, J., & Aubourg, S. (2014a). Use of
554 citric and lactic acids in ice to enhance quality of two fish species during on-
555 board chilled storage. *International Journal of Refrigeration*, 40, 390-397.

556 García-Soto, B., Sanjuás-Rey, M., Barros-Velázquez, J., Fuertes-Gamundi, R., &
557 Aubourg, S. (2011). Preservative effect of an organic acid-icing system on
558 chilled fish lipids. *European Journal of Lipid Science and Technology*, 113, 487-
559 496.

560 González-Fandos, E., Villarino-Rodríguez, A., García-Linares, M. C., García-Arias, M.
561 T., & García-Fernández, M. C. (2005). Microbiological safety and sensory
562 characteristics of salmon slices processed by the sous vide method. *Food*
563 *Control*, 16, 77-85.

564 Gyawali, R., & Ibrahim, S. (2014). Natural products as antimicrobial agents. *Food*
565 *Control*, 46, 412-429.

- Halldorsdóttir, S., Sveinsdóttir, H., Gudmundsdóttir, A., Thorkelsson, G., & Kristinsson, H. (2014). High quality fish protein hydrolysates prepared from by-product material with *Fucus vesiculosus* extract. *Journal of Functional Foods*, 9, 10-17.
- Hermund, D., Iltas, B., Honold, P., Jónsdóttir, R., Kristinsson, H., & Jacobsen, C. (2015). Characterisation and antioxidant evaluation of Icelandic *F. vesiculosus* extracts *in vitro* and in fish-oil-enriched milk and mayonnaise. *Journal of Functional Foods*, doi: 10.1016/j.jff.2015.02.020.
- Jehanno, D., Thuault, D., Larnaud, P., Alaume, P., & Bourgeois, C. (1996). Effect of the method of storage on-board fishing vessels on the quality of fish. *Science des Aliments*, 16, 167-174.
- Kjelleberg, S., & Steinberg, P. (2001). Surface warfare in the sea. *Microbiology Today*, 28, 134-135.
- Losada, V., Piñeiro, C., Barros-Velázquez, J., & Aubourg, S. (2004). Effect of slurry ice on chemical changes related to quality loss during European hake (*Merluccius merluccius*) chilled storage. *European Food Research and Technology*, 219, 27-31.
- Lowry, R., & Tinsley, I. (1976). Rapid colorimetric determination of free fatty acids. *Journal of the American Oil Chemists' Society*, 53, 470-472.
- Oral, N., Gülmez, M., Vatansever, L., & Güven, A. (2008). Application of antimicrobial ice for extending shelf life of fish. *Journal of Food Protection*, 71, 218-222.
- Ortiz, J., Vivanco, J., & Aubourg, S. (2014). Lipid and sensory quality of canned Atlantic salmon (*Salmo salar*): Effect of the use of different seaweed extracts as covering liquids. *European Journal of Lipid Science and Technology*, 116, 596-605.

591 Özyurt, G., Kuley, E., Balikçi, E., Kaçar, Ç., Gökdoğan, S., Etyemez, M., & Özogul, F.
592 (2012). Effect of the icing with rosemary extract on the oxidative stability and
593 biogenic amine formation in sardine (*Sardinella aurita*) during chilled storage.
594 *Food and Bioprocess Technology*, 5, 2777-2786.

595 Paiva, L., Lima, E., Ferreira Patarra, R., Neto, A., & Baptista, J. (2014). Edible Azorean
596 macroalgae as source of rich nutrients with impact on human health. *Food*
597 *Chemistry*, 164, 128-135.

598 Pastoriza, L., Bernárdez, M., Sampedro, G., Cabo, M., & Herrera, J. (2008). The use of
599 water and ice with bactericide to prevent onboard and onshore spoilage of
600 refrigerated megrim (*Lepidorhombus whiffiagonis*). *Food Chemistry*, 110, 31-38.

601 Peinado, I., Girón, J., Koutsidis, G., & Ames J. M. (2014). Chemical composition,
602 antioxidant activity and sensory evaluation of five different species of brown
603 edible seaweeds. *Food Research International*, 66, 36-44.

604 Quiral, V., Donoso, M^aL., Ortiz, J., Herrera, M^aV., Araya, H., & Aubourg, S. (2009).
605 Chemical changes during the chilled storage of Chilean jack mackerel
606 (*Trachurus murphyi*): Effect of a plant extract-icing system. *LWT-Food Science*
607 *and Technology*, 42, 1450-1454.

608 Rodríguez, O., Barros-Velázquez, J., Ojea, A., Piñeiro, C., & Aubourg, S. (2003).
609 Evaluation of sensory and microbiological changes and identification of
610 proteolytic bacteria during the iced storage of farmed turbot (*Psetta maxima*).
611 *Journal of Food Science*, 68, 2764-2771.

612 Rodríguez-Bernaldo de Quirós, A., Frecha-Ferreiro, S., Vidal-Pérez, A., & López-
613 Hernández, J. (2010). Antioxidant compounds in edible brown seaweeds.
614 *European Food Research Technology*, 231, 495-498.

- Sallam, K. (2007). Antimicrobial and antioxidant effects of sodium acetate, sodium lactate, and sodium citrate in refrigerated sliced salmon. *Food Control*, 18, 566-575.
- Sandsdalen, E., Haug, T., Stensvag, K., & Styrvold, O. (2003). The antibacterial effect of a polyhydroxylated fucophlorethol from the marine brown alga, *Fucus vesiculosus*. *World Journal of Microbiology and Biotechnology*, 19, 777-782.
- Sanjuás-Rey, M., García-Soto, B., Barros-Velázquez, J., Fuertes-Gamundi, J., & Aubourg, S. (2011). Effect of a two-step natural organic acid-treatment on microbial activity and lipid damage during blue whiting (*Micromesistius poutassou*) chilling. *International Journal of Food Science and Technology*, 46, 1021-1030.
- Sanjuás-Rey, M., García-Soto, B., Fuertes-Gamundi, R., Aubourg, S., & Barros-Velázquez, J. (2012). Effect of a natural organic acid-icing system on the microbiological quality of commercially relevant chilled fish species. *LWT-Food Science and Technology*, 46, 217-223.
- Sikorski, Z., & Kolakowska, A. (1994). Changes in protein in frozen stored fish. In Z. Sikorski, B. Sun Pan, & F. Shahidi (Eds.), *Seafood proteins* (pp. 99-112). New York, USA: Chapman and Hall.
- Sivertsvik, M., Jeksrud, W., & Rosnes, T. (2002). A review of modified atmosphere packaging of fish and fishery products- significance of microbial growth, activities and safety. *International Journal of Food Science and Technology*, 37, 107-127.
- Smit, A. (2004). Medicinal and pharmaceutical uses of seaweed natural products: A Review. *Journal of Applied Phycology*, 16, 245-262.

- Tierney, M., Smyth, T., Hayes, M., Soler-Vila, A., Croft, A., & Brunton, N. (2013a). Influence of pressurised liquid extraction and solid-liquid extraction methods on the phenolic content and antioxidant activities of Irish macroalgae. *International Journal of Food Science and Technology*, 48, 860-869.
- Tierney, M., Smyth, T., Rai, D., Soler-Vila, A., Croft, A., & Brunton, N. (2013b). Enrichment of phenol contents and antioxidant activities of Irish brown macroalgae using food-friendly techniques based on polarity and molecular size. *Food Chemistry*, 139, 753-761.
- Tozawa, H., Erokibara, K., & Amano, K. (1971). Proposed modification of Dyer's method for trimethylamine determination in codfish. In R. Kreuzer (Ed.), *Fish Inspection and Quality Control* (pp. 187-190). London, UK: Fishing News Books Ltd.
- Vyncke, W. (1970). Direct determination of the thiobarbituric acid value in trichloroacetic acid extracts of fish as a measure of oxidative rancidity. *Fette Seifen Anstrichmittel*, 72, 1084-1087.
- Wang, T., Jónsdóttir, R., Kristinsson, H., Thorkelsson, G., Jacobsen, C., Yuca Hamaguchi, P., & Olafsdóttir, G. (2010). Inhibition of haemoglobin-mediated lipid oxidation in washed cod muscle and cod protein isolates by *Fucus vesiculosus* extract and fractions. *Food Chemistry*, 123, 321-330.

FIGURE LEGENDS

Figure 1: Proteolytic bacteria (log CFU g⁻¹ muscle) assessment* in chilled megrim muscle stored under different icing conditions.

* Initial value: 3.39±0.18. Mean values of three replicates (n = 3); standard deviations are indicated in brackets. For each chilling time, mean values followed by different capital letters (A-B) indicate significant (p<0.05) differences as a result of the icing condition. No letters are indicated when significant differences are not found (p>0.05).

Figure 2: Lipolytic bacteria (log CFU g⁻¹ muscle) assessment* in chilled megrim muscle stored under different icing conditions.

* Initial value: 2.16±0.28. Mean values of three replicates (n = 3); standard deviations are indicated in brackets. For each chilling time, mean values followed by different capital letters (A-B) indicate significant (p<0.05) differences as a result of the icing condition. No letters are indicated when significant differences are not found (p>0.05).

Figure 3: Assessment* of the pH value in chilled megrim muscle stored under different icing conditions.

* Initial value: 6.42±0.09. Mean values of three replicates (n = 3); standard deviations are indicated in brackets. For each chilling time, mean values followed by different capital letters (A-B) indicate significant (p<0.05) differences as a result of the icing condition. No letters are indicated when significant differences are not found (p>0.05).

TABLE 1**Microbiological count assessment (log CFU g⁻¹)* in chilled megrim muscle stored under different icing conditions****

Chilling time (days)	Aerobes			Psychrotrophs			Enterobacteriae		
	F-0	F-1	F-2	F-0	F-1	F-2	F-0	F-1	F-2
0	3.25 a (0.24)	3.25 a (0.24)	3.25 a (0.24)	3.76 a (0.31)	3.76 a (0.31)	3.76 a (0.31)	< 1 a	< 1 a	< 1 a
4	3.80 ab (0.56)	3.90 ab (0.61)	3.23 a (0.58)	4.71 a (0.70)	5.00 b (0.06)	4.93 b (0.22)	< 1 a	1.10 a (0.70)	< 1 a
7	5.11 bc (0.72)	4.22 b (0.28)	4.19 b (0.24)	6.15 b (0.32)	5.77 bc (0.81)	5.67 c (0.34)	< 1 a	1.11 a (0.17)	< 1 a
11	6.35 cB (0.65)	5.96 cB (0.21)	5.25 cA (0.21)	7.28 cB (0.62)	6.99 cdB (0.37)	6.39 dA (0.04)	< 1 a	< 1 a	< 1 a
14	7.39 dB (0.11)	7.26 dAB (0.26)	6.72 dA (0.36)	7.42 c (0.47)	7.30 d (0.36)	7.25 e (0.21)	2.37 b (0.13)	2.73 b (0.38)	2.73 b (0.38)

* Mean values of three replicates (n = 3); standard deviations are indicated in brackets. For each parameter and for each chilling time, mean values followed by different capital letters (A-B) indicate significant (p<0.05) differences as a result of the icing condition. For each parameter and for each icing condition, values followed by different low-case letters (a-e) denote significant (p<0.05) differences as a result of the chilling time. No letters are indicated when significant differences are not found (p>0.05).

** Abbreviations of icing conditions: F-0 (Control; no seaweed extract presence in ice), F-1 (ice prepared from a 0.67 g L⁻¹ aqueous solution of *F. spiralis* extract) and F-2 (ice prepared from a 2.50 g L⁻¹ aqueous solution of *F. spiralis* extract).

TABLE 2

Assessment* of trimethylamine-nitrogen (TMA-N) and free fatty acids contents in chilled megrim muscle stored under different icing conditions**

Chilling time (days)	TMA-N (mg TMA-N kg⁻¹ muscle)			Free fatty acids content (g kg⁻¹ lipids)		
	F-0	F-1	F-2	F-0	F-1	F-2
0	1.5 a (0.7)	1.5 a (0.7)	1.5 a (0.7)	10.4 a (1.6)	10.4 a (1.6)	10.4 a (1.6)
4	5.4 b (2.0)	4.1 ab (2.3)	7.2 b (2.9)	10.4 a (0.1)	14.3 a (5.7)	12.0 ab (1.7)
7	45.2 cB (14.4)	7.2 bA (8.9)	18.1 bA (7.9)	26.3 b (8.4)	14.1 a (3.9)	17.1 bc (3.5)
11	100.8 dB (29.3)	180.4 cC (18.1)	58.3 cA (11.5)	27.5 bA (7.7)	63.7 bB (8.4)	21.1 cA (3.7)
14	198.3 e (17.5)	193.4 c (48.2)	240.7 d (50.6)	62.8 cA (12.4)	95.3 cB (11.9)	48.1 dA (13.7)

* Mean values of three replicates (n = 3); standard deviations are indicated in brackets. For each parameter and for each chilling time, mean values followed by different capital letters (A-C) indicate significant (p<0.05) differences as a result of the icing condition. For each parameter and for each icing condition, values followed by different low-case letters (a-e) denote significant (p<0.05) differences as a result of the chilling time. No letters are indicated when significant differences are not found (p>0.05).

** Abbreviations of icing conditions as expressed in Table 1.

TABLE 3**Lipid oxidation assessment* in chilled megrim muscle stored under different icing conditions****

Chilling time (days)	Peroxide value (meq active oxygen kg ⁻¹ lipids)			Thiobarbituric acid index (mg malondialdehyde kg ⁻¹ muscle)			Fluorescence ratio		
	F-0	F-1	F-2	F-0	F-1	F-2	F-0	F-1	F-2
0	0.45 a (0.27)	0.45 a (0.27)	0.45 a (0.27)	0.22 a (0.03)	0.22 a (0.03)	0.22 a (0.03)	2.46 a (0.64)	2.46 a (0.64)	2.46 ab (0.64)
4	1.55 bA (0.62)	2.69 bcA (0.55)	5.33 cB (1.26)	0.24 ab (0.04)	0.35 ab (0.11)	0.24 a (0.05)	2.18 a (0.42)	1.91 a (0.35)	1.80 a (0.30)
7	1.38 bA (0.17)	0.92 aA (0.41)	2.67 bB (1.11)	0.34 abAB (0.15)	0.22 aA (0.02)	0.33 bB (0.01)	2.18 a (0.19)	2.25 a (0.50)	2.66 bc (0.27)
11	0.87 abA (0.72)	1.99 bB (0.27)	5.08 cC (1.05)	0.36 b (0.07)	0.34 b (0.05)	0.38 b (0.07)	4.06 b (0.41)	4.15 b (0.43)	3.66 d (0.09)
14	2.53 bA (1.54)	5.02 cA (2.00)	8.96 dB (0.84)	0.31 b (0.03)	0.44 b (0.10)	0.42 b (0.06)	4.77 bB (0.44)	5.03 bB (0.57)	3.45 cdA (0.49)

* Mean values of three replicates (n = 3); standard deviations are indicated in brackets. For each parameter and for each chilling time, mean values followed by different capital letters (A-C) indicate significant (p<0.05) differences as a result of the icing condition. For each parameter and for each icing condition, values followed by different low-case letters (a-d) denote significant (p<0.05) differences as a result of the chilling time. No letters are indicated when significant differences are not found (p>0.05).

** Abbreviations of icing conditions as expressed in Table 1.

Figures 1-3

Chilling time (days)	Proteolitics (Fig 1)			Lipolitics (Fig 2)			pH (Fig 3)		
	F-0	F-1	F-2	F-0	F-1	F-2	F-0	F-1	F-2
4	3.98 a (0.26)	4.38 a (0.20)	4.10 a (0.23)	3.43 a (0.90)	3.84 a (0.11)	3.54 ab (0.52)	6.61 aAB (0.14)	6.58 aB (0.03)	6.44 aA (0.06)
7	5.13 bAB (0.60)	5.05 bB (0.25)	4.36 aA (0.21)	4.36 aB (0.47)	4.20 abB (0.42)	3.49 aA (0.25)	6.75 aB (0.02)	6.54 aA (0.05)	6.51 abA (0.07)
11	6.19 cB (0.13)	6.79 cB (0.49)	5.18 bA (0.22)	5.82 bB (0.12)	4.60 bA (0.24)	4.28 abA (0.48)	6.78 a (0.10)	6.71 b (0.03)	6.65 b (0.05)
14	6.61 dB (0.21)	6.53 cAB (0.58)	5.99 cA (0.24)	5.90 bB (0.10)	5.50 cB (0.54)	4.36 bA (0.42)	7.18 bB (0.06)	6.96 cA (0.08)	6.83 cA (0.07)