Post-print of: Mellado-Ortega, E., Hornero-Méndez, D. Carotenoids in cereals: an ancient resource with present and future applications. Phytochemistry Reviews 14 (2015) 873-890
[http:// dx.doi.org/10.1007/s11101-015-9423-3](http://dx.doi.org/10.1007/s11101-015-9423-3)

Carotenoids in cereals. An ancient resource with present and future applications.

Elena Mellado-Ortega & Dámaso Hornero-Méndez*

Group of Chemistry and Biochemistry of Pigments. Food Phytochemistry Department.
Instituto de la Grasa (CSIC). Pablo de Olavide University Campus, Ctra. de Utrera km.
1. 41013 - Seville (Spain).

* Corresponding author

Telephone: +34 954611550

Fax: +34 954616790

e-mail: hornero@ig.csic.es

Abstract

Cereals are considered to be a major source of carbohydrates and proteins as well as minor micronutrients such as mineral, vitamins and antioxidants, including carotenoids. Carotenoids are natural lipophilic pigments biosynthesized mainly by plants, and certain bacteria and fungi. Cereals, although having a low carotenoid content when compared with the majority of fruits and vegetables, may have an important impact in the nutritional status of consumers. The daily consumption of cereals, and products derived from them, by a large part of the population, especially in under-developed and developing countries, makes cereals a contributor of carotenoids which should not be overlooked and must be taken into consideration in biofortification strategies. In the present manuscript, we revise the existing information about the composition and distribution of carotenoids in cereals, highlighting factors which alter their profile, such as domestication of wild varieties, genotype, storage, milling and processing techniques. Strategies for stimulating the carotenoid content in cereals, either by means of traditional breeding methods or by genetic manipulation, are also discussed.

Keywords. xanthophylls; carotenes; lutein; wheat; staple food

Abbreviations

MAS: marker assisted selection

PSY: phytoene synthase gene

QTL: quantitative trait locus

XAT: xanthophyll acyltransferase

YPC: yellow pigment content

1. Introduction

Carotenoid pigments are molecules of lipophilic nature and are responsible for some of the yellow, orange and red colour of a wide range of fruits and some animal teguments (Britton and Hornero-Méndez 1997). Animals are not able to synthesize them *de novo* and so they must be acquired through the diet. Carotenoids are located in oil droplets, crystalloids and membrane structures within specialized vegetable organelles, the so-called plastids, being the most common the chloroplasts (in green tissues) and the chromoplasts (yellow, orange and red tissues), and are found in all parts of the plant: roots, leaves, flowers, fruits and seeds (Howitt and Pogson 2006). Carotenoids carry out their most basic functions in vegetables as pigment collectors of light in the photosynthetic apparatus, and preventing oxidative damage acting as antioxidants.

When carotenoids are ingested, they exert important biological activities; antioxidant, inhibition of carcinogenesis, enhancement of the immune response and cell defence against reactive oxygen species (ROS) and free radicals, and the reduction on the risk of developing cardiovascular and other degenerative diseases (Nishino 1997; recently reviewed by Britton et al. 2009). In addition some carotenoids (β -carotene, α -carotene, β -cryptoxanthin, etc) have provitamin A activity (Baker and Günther 2004; Olson 1989). Of particular interest are the epidemiological studies showing an inverse correlation between the progression of age-related macular degeneration (AMD) and cataracts and the high intake of lutein and zeaxanthin rich-vegetables, being both pigments present in high concentration at the macula in the retina humans and primates (Ahmed et al. 2005; Alves-Rodrigues and Shao 2004; Calvo 2005; Landrum and Bone, 2001).

Cereals are basic food stuffs of vegetable origin, also known as staple food,

which are characterised mainly as sources of carbohydrates and proteins, and contain a relatively small amount of carotenoids compared to the majority of fruits and vegetables. However, the daily consumption of cereals and products derived from them by a large part of the population, especially in under-developed and developing countries, makes cereals a contributor of carotenoids which must be taken into consideration (Graham and Rosser 2000). Therefore, cereals can be considered ideal elements for use in biofortification strategies (Bai et al. 2011).

An essential step for the development of carotenoid biofortified crops is the complete characterization of the carotenoids profile contained in the vegetables, allowing this an adequate knowledge of their metabolism, which can be used for the selection and breeding of new cultivars. The success of such processes lies in the existence of a more or less extensive variation of carotenoid contents among different cereals. This variability is mainly due to three factors: a) the genotype of the cereal, b) the varietal selection, the pressure exerted by man and its domestication, and c) the growing conditions and post-harvest storage and processing (industrial and domestic) practices. Regarding the last factor, the evaluation of such changes is an active area of study in food science research, which allows the prediction and selection of the optimal conditions for preserving these phytochemicals, driven by their nutritional properties as well as the applications derived from their natural role as being responsible for the colour of foods. Thermal treatments and others which include homogenisation and modification of the food matrix are factors which have a beneficial effect on the bioaccessibility of carotenoids since it facilitates their solubility. The term bioaccessibility is used to evaluate the amount of a nutrient that is released from a food during the digestion process. The bioaccessibility of lipophilic compounds, such as carotenoids, in natural foods is frequently fairly low and is constrained by various

factors, particularly the degree of food processing and matrix composition. There are evidences suggesting that homogenisation and thermal treatment have positive effects on the bioaccessibility of these compounds. Therefore, foods such as cereals, which are normally consumed as processed foods, could present some advantages in comparison to fresh food.

Up to now, there have been several studies regarding the qualitative and quantitative composition of carotenoids in fruits and vegetables, but very few with respect to cereals and therefore studies which determine the carotenoid profile of these foodstuffs are necessary. Undoubtedly, investigations aimed at improving the bioaccessibility of carotenoids in cereals will benefit from such knowledge and it will aid in the information and recommendation provided to the consumer about the way of processing and consumption of these basic elements.

In the present manuscript, we revise the existing information about the composition and distribution of carotenoids present in cereals, highlighting factors which alter their profile, such as storage and processing. At the same time, the strategies for stimulating the carotenoid content in cereals, either by means of traditional breeding methods or by genetic manipulation, are also discussed.

2. Composition and distribution of carotenoids in cereals.

Cereals and their derivatives are good examples of foods that contain a complex mixture of minor components such as phenols, folates, vitamin E, phytic acid, sterols and carotenoids, some of them with antioxidants properties (McKevith 2004). In general, the content of bioactive compounds in cereals is mainly concentrated in the germ and bran portions and most of them are located in the aleurone layer. At the same time, the content in phytochemicals is subjected to important variations due to factors

such as the type of cereal, the cultivar, cultivation conditions, stage in development, and storage and processing conditions (Fardet et al. 2008; Liu 2007;), with the greatest importance lying in the interactions between the genotype and the environment (Hidalgo et al. 2009; Lv et al. 2013; Van Hung and Hatcher 2011). The carotenoid profile in cereals is mainly composed of xanthophylls, with lutein as the most abundant, followed by zeaxanthin and β -cryptoxanthin in addition to carotenes such as α - and β -carotene in small amounts (**Figure 1**). The largest concentration of carotenoids can be found in the embryo, although this part of the seed only represents 3-5% of its total weight. On the contrary, the contribution of the endosperm, around 80-85% of the cereal grain, is the most influential in the total content of carotenoids in the grain. The distribution of this carotenoid profile seems to vary among genotypes of the same type of cereal (Siebenhandl et al. 2007) and within the same grain since α -, β -carotene and zeaxanthin are concentrated in the bran and the germ, while lutein is distributed more homogeneously (Borrelli et al. 2008; Konopka et al. 2004; Ndolo and Beta 2013; Panfili et al. 2004). In relation to the composition of carotenoids in cereals, the bibliography is fairly scarce, especially for barley (*Hordeum vulgare*), rye (*Secale cereale*) and millet (*Panicum miliaceum*) (Choi et al. 2007; Kandlakunta et al. 2008; Mamatha et al. 2011).

In light of the fact that the level of carotenoids in cereals is low in comparison to fruits and vegetables, corn (*Zea mays*) of the yellow genotype may be cited as the cereal which presents the greatest levels (up to 63 $\mu\text{g/g}$) of these phytochemicals. Most of the cultivars present zeaxanthin as the major pigment but also show small amounts of β -cryptoxanthin α -carotene and β -carotene (Moros et al. 2002; Panfili et al. 2004). From the quantitative studies carried out on this cereal, a large variation in the contents of carotenoids has been determined among varieties (Berardo et al. 2004, 2009; Egesel et al. 2003; Kurilich and Juvik 1999; Quackenbush et al. 1961), which serves as proof that

genotype is the determining factor for this variety (Ibrahim and Juvik 2009; Menkir and Maziya-Dixon 2004;), followed by storage and processing conditions (Burt et al. 2010; De Oliveira and Rodríguez-Amaya 2007; Scott and Eldridge 2005).

In the case of rice (*Oryza sativa*), it is obvious that the pigmented varieties are the ones which result in some interest from a quantitative point of view. In the grains of conventional rice the carotenoid content is concentrated almost exclusively in the bran and therefore the common practices of milling and stripping of the grain result in a drastic reduction in this component in the final product (Tan et al. 2005). Similar to most cereals, lutein is the main pigment (Belefant-Miller and Grace 2010), followed by zeaxanthin and β -carotene in crops with dark or black pigmentation (Frei and Becker 2005; Kim et al. 2010; Lamberts and Delcour 2008; Nakornriab et al. 2008). In some varieties of black rice from Thailand, the concentration of β -carotene found in its bran is significant (up to 40 $\mu\text{g/g}$) (Nakornriab et al. 2008).

A recent study carried out on glabrous canary seed (*Phalaris canariensis* L.) has shown a content of β -carotene ranging from 5.2 to 6.3 $\mu\text{g/g}$, which is the major pigment in this cereal (Li and Beta 2012). Although sorghum (*Sorghum bicolor*) has been considered one of the cereals with the least amount of carotenoids, the pioneer findings of Blessin et al. (1958) and Suryanarayana Rao et al. (1968) concerning varieties of yellow endosperm, as well as the recent studies of Kean et al. (2007, 2011), offer better perspectives for this cereal. These authors found a carotenoid content between 0.11-0.32 mg/kg, which although lower than the content corresponding to yellow corn, these data must not be overlooked, given the importance of this cereal in the diet of some populations (mainly in Asia and Africa).

Oats (*Avena sativa*) are also cited in the literature as one of the cereals with the lowest level of carotenoids, with contents of lutein and carotenes (α - and β -carotene) in

the order of 0.20 mg/kg and 0.01 mg/kg, respectively (Panfili et al. 2004). Wheat (genus *Triticum*), along with corn and rice, are the most popular cereals due to their generalised consumption. This is the most consistently seen aspect in the literature, with special interest being paid to the study of their antioxidant profile and contents of healthy phytochemicals as well as their inter- and intra-varietal distribution (Adom et al. 2003, 2005; Di Silvestro et al. 2012; Lier and Lacroix 1974; Okarter et al. 2010; Velioglu et al. 1998;; Zhou et al. 2004; Zhou and Yu 2004a, b). Similar to the case of corn, the genetic variability among wheat genotypes concerning the contents of carotenoids, offers an interesting tool which can be effectively used in programs aimed at augmenting the contents of bioactive compounds with properties which are beneficial to health (Digesù et al. 2009; Leenhardt et al. 2006a).

Wheat varieties with diploid genome (AA), in particular the Einkorn wheat (*Triticum monococcum*), are considered those presenting the highest levels, in the order of approximately 2 to 4 times higher with respect to other wheat varieties. Einkorn also stands out as possessing the greatest antioxidant activity (Lavelli et al. 2009). Followed by these, tetraploid wheats (AABB) can be found and has been determined to contain the highest levels of carotenoids (around 5-6 µg/g) among the durum wheat cultivars (*Triticum turgidum* conv. *durum*), although others have been mentioned such as the ancestral Emmer (*T. turgidum* subsp. *dicoccoides*) with similar levels (approximately 3-5 µg/g). Finally, wheat varieties belonging to the hexaploid genome (AABBDD) are considered to have the lowest carotenoid content, with almost exclusive attention paid to the common bread wheat (*T. aestivum*), which shows an average content around 2 µg/g (Abdel-Aal et al. 2007; Hidalgo et al. 2006; Hidalgo and Brandolini 2008a; Serpen et al. 2008).

These quantitative data indicate that the selection and domestication of cereal

species carried out by man has derived in a wide variation of their carotenoid levels (both low and high) (Digesù et al. 2009). The case of durum wheat constitutes an example which has been extensively analysed due to the fact that the yellow colour of its semolina (also referred to as *yellow pigment content*; YPC, determined as lutein equivalents by the official method 125 of the International Association for Cereal Science and Technology and method 14-50 of the American Association of Cereal Chemists) is considered one of the principal criteria of quality of processed derived foods, mainly pasta (Blanco et al. 2011; Fratianni et al. 2005; Hentschel et al. 2002; Humphries et al. 2004). An inverse situation occurs with bread wheat destined to flour production, which presents lower levels of carotenoids due to the pressure exerted by industries motivated by a consumer preference for whiter flour for the further preparation of flour-containing products according to market demand.

The *Triticeae* tribe, which includes wheat, barley and rye species, is a series of polyploids which are closely related, being possible to generate fertile hybrid amphiploids among the different cultivated members of this tribe and its wild parents. Among the hybrid cereals, tritordeum, an innovative cereal with interesting properties and applications, must be pointed out. Tritordeum is a cereal which has been obtained from the cross-breeding between a wild barley, *Hordeum chilense*, with diploid genome ($H^{ch}H^{ch}$), and diploid and polyploid wheats (Martín and Chapman 1977). In addition to the development of amphiploids such as tritordeum, the use of *H. chilense* in the enhancement programs has also been focused on the inclusion of wheat with new characteristics of interest. Among the highlighted attributes, a high content of carotenoids, determined by $7H^{ch}$ chromosome, stands out. The first fertile hybrids with the optimal characteristics for becoming a possible crop were only obtained following to the crossing of *H. chilense* with *T. turgidum* (a tetraploid durum wheat, with genomic

denotation AABB), generating those denominated hexaploid tritordeums ($2n=6x=42$, $H^{ch}H^{ch}AABB$) which showed a low frequency of aneuploids, a wide variation in growth rate and an optimal level of fertility (Martín and Sánchez-Monge Laguna 1982).

Most of the estimations on the carotenoid content made on tritordeum did not provide detailed information about the individual composition of the pigment profile due to the lack of specificity of the traditional analytical methods employed for the characterisation of carotenoids in cereals. By means of estimations such as YPC, levels of carotenoids in the order of 5 to 6 times higher (in the range of 11-13 $\mu\text{g/g}$) have been found in tritordeum with respect to common wheat (2 $\mu\text{g/g}$) and 2 to 3 times higher with respect to durum wheat (5-6 $\mu\text{g/g}$) (Álvarez et al. 1995; Martín et al. 1999), which is comparable to the levels reported for Einkorn wheat. In an extensive study which evaluated a total of 35 primary lines of tritordeum together with their respective parental lineage, 27 *H. chilense* accessions and 19 durum wheat cultivars, the average levels of carotenes ($\mu\text{g/g}$ of β -carotene equivalents, determined as YPC) for the amphiploid continued to be in the order of 2 times higher than durum wheat parent, but 3 times lower than the average of *H. chilense* (Álvarez et al. 1999). Among the advanced lines of tritordeum, HT621 was registered as an elite germplasm line due to its high content of carotenoid pigments (19 μg of β -carotene equivalents/g, YPC) (Ballesteros et al. 2005). Recent studies carried out in our laboratory (Atienza et al. 2007a; Mellado-Ortega and Hornero-Méndez 2012) have contributed substantially to the characterisation and quantification of individual carotenoid pigments in tritordeum. Similar to most cereals, lutein is also the major pigment (>85%), with a high esterification degree (monoesters and diesters), along with minor quantities of β -carotene (**Figure 2**). On average, the total carotenoid content of tritordeum (6.5 $\mu\text{g/g}$ of fresh weight) was significantly higher, approximately 8 times higher, when compared to

durum wheat (0.7 µg/g of fresh weight). For the first time, the regioisomers of the monoesters and diesters of lutein in a cereals, in this case in tritordeum, have been isolated and characterized (Mellado-Ortega and Hornero-Méndez 2012) (**Figure 3**). The fraction of monoesters is made up of the regioisomers, namely lutein-3'-*O*-linoleate, lutein-3-*O*-linoleate, lutein-3'-*O*-palmitate and lutein-3-*O*-palmitate, while the fraction of diesters is composed of two homoesters, lutein dilinoleate and lutein dipalmitate, and by the two regioisomers of a heterodiester, lutein-3'-*O*-linoleate-3-*O*-palmitate and lutein-3'-*O*-palmitate-3-*O*-linoleate. The esterification of lutein with only two different fatty acids (linoleic and palmitic acids) suggests a high-specificity degree for the *in vivo* process, which should be further investigated. The ability to distinguishing the regioisomers of mono- and diesters of lutein provides a powerful tool contributing to a better deciphering of this poorly known biochemical pathway, for which tritordeum grains could be used as an excellent plant model. Evidences for a preferential xanthophyll acyltransferase activity regarding the position (3 or 3') and the acyl moiety are discussed. Further studies should be carried out in order to identify the acyltransferase enzymes (XAT) and the acyl donor molecules involved in the xanthophylls esterification process.

Tritordeum is currently the subject of an intense breeding program being developed at the Institute of Sustainable Agriculture (CSIC; Cordoba, Spain) in order to optimise its use as a new cereal and its incorporation into the formulation of functional foods. Its cultivation areas are concentrated in Spain (Andalusia, Castilla and Catalonia), the south of Italy and the south of Portugal. In 2008 it was registered, under the name Aucan, the first line of tritordeum in the European Union Plant Variety Registration (CPVO), with other advanced lines in their final phases of development and evaluation. As a result of these efforts, at the beginning of 2013, the Spanish

company Agrasys (www.agrasys.es) began to commercialise tritordeum flour under the brand name Vivagran®.

3. Cereal technology and its influence on the carotenoid contents in grains and derived products

Conservation and storage conditions, as well as different processing treatments, result in overall changes in the composition of foods, being this the main reason for the numerous studies carried out in the field of food technology which are crucial for determining some important sensorial and nutritional quality attributes (Kalt 2005; Mínguez-Mosquera et al. 1997; Nicoli et al. 1999; Rodriguez-Amaya 2003). In this way, it is important to distinguish between the effects of processing itself, and other environmental factors such as partial oxygen pressure, temperature, exposure to light and humidity as well as the interactions with other antioxidant and pro-oxidant molecules which are present (Lindley 1998; Nicoli et al. 1999). In the case of the carotenoids occurring in cereals, the storage of the natural and processed grain normally results in a decrease in the pigment contents, which is directly proportional to the processing time and the increase in the intensity of other variables such as temperature or the degradation process (Cristobal 1965; Weber 1987). It is interesting to note that the literature concerning this aspect is especially scarce since cereals are characterised as undergoing prolonged storage periods as part of their industrial and technological treatments. On the other hand, it is not easy to define a set of general guidelines in the behaviour of pigments in processes such as peeling, grinding, drying, roasting, fermentation, etc., due to the fact that positive, negative and even indifferent results can be found in the literature regarding the level of carotenoids present. In general, these processes would affect the internal structure of the cereal matrix and the processed

product with substantial implications regarding accessibility and further bio-availability of the carotenoids.

The rate at which these changes occur has been a generalised subject of food science. The bibliography concerning the kinetics of carotenoid degradation and/or retention in foods is abundant as well as the disparity of the results which have often been contradictory (Mínguez-Mosquera et al. 1997). The profusion of such studies reflects the importance of carotenoid pigments for the food industry, not only from a nutritional point of view, but from a technological standpoint, with interest in not just quantifying losses in carotenoids but also analysing the conditions which provide their greater stability and subsequent retention. Colour is considered one of the main consumer acceptance criteria for processed foods. As mentioned before, in the flour and bread as well as the semolina and pasta this factor is important, especially for products derived from durum wheat. Therefore the research efforts in this field try to reproduce the processing and storage conditions of food, including cereals, with the aim of comparing different industrial treatments and monitoring the incidences of individual or joint factors such as water, temperature, light, oxygen and pH activities along with pro-oxidants and antioxidants (Fish and Davis 2003; Mínguez-Mosquera and Gandul-Rojas 1994; Ouchi et al. 2010; Saxena et al. 2012; Selim et al. 2000; Tsimidou 1997), allowing for an estimate of the shelf-life of foods and their bioactive compounds. Most of the studies have been carried out on vegetables such as carrots (Koca et al. 2007; Lemmens et al. 2010; Wagner and Warthesen 1995), tomato (Sharma and Le Maguer 1996; Tonon et al. 2007); peppers and pepper derived products (Carbonell et al. 1986), potatoes (Bechoff et al. 2010), and citrus fruit juices (Dhuique-Mayer et al. 2007; Zepka et al. 2009) where the authors commonly describe a degradation reaction which occurs according to a zero or first order kinetics. However, the literature is quite limited in the

particular case of cereals, probably due to the low level of pigments that these foods present and there are very few studies which carry out a detailed kinetic evaluation (Guzmán-Tello and Cheftel 1990; Hidalgo and Brandolini 2008b; Mellado-Ortega 2013). Nevertheless, although such alterations have not been quantified, most authors agree in the fact that changes generally occur more rapidly at the beginning of processing and storage and as they progress the rate at which they are occurring diminishes (Burt et al. 2010; Quackenbush 1963).

4. Stability of carotenoids during the post-harvest storage of grains and flours.

The storage process produces a loss in carotenoids mainly caused by oxidation, either of an enzymatic or non-enzymatic nature. The oxygen present in the medium is considered the major factor affecting the stability of carotenoids (Britton and Khachik 2009). Other modifications of the carotenoid molecule associated with these processes are the geometric isomerisation (*cis/trans*) produced by temperature and/or light and cause alterations in the composition rather than net losses in the pigments. The mechanisms of both processes have been extensively studied (El-Agamey and McGarvey 2008; Liaaen-Jensen and Lutnaes 2008).

While the oxidation of pigments in the storage of cereal grains comes mainly from oxygen, in processed foods such as flour the alteration of the matrix produces direct oxidative processes as well as those caused by enzymes which come into contact with carotenoids (Doblado-Maldonado et al. 2012). As previously mentioned, the results obtained in these studies depend on the conditions applied, mostly evaluating the effects of time and temperature applied during the treatments. The literature contains examples in which the effects of these two variables are clearly observed, which allow for the prediction of the consequences of these variables in the stability of the

carotenoids present (Arya and Parihar 1981; Calucci et al. 2004; Farrington et al. 1981; Hidalgo and Brandolini 2008b; Nghia et al. 2006; Pinzino et al. 1999).

On occasion, in addition to storage, the experimental designs additionally cover another factors corresponding to common practices in cereal technology (such as drying, grinding, shelling, etc). This widens the study but at the same time difficult the analysis of the obtained results. Belefant-Miller and Grace (2010) evaluate the behaviour of carotenoids during the prolonged storage of rice which was previously shelled. A common practice in cereals prior to storage is the drying of the grains since the reduction in humidity prevents deterioration due to mould, acarus, etc., as well as a premature germination of the grain (Chelowski 1994; Jood and Kapoor 1994). Some authors, such as Burt et al. (2010) did not obtain a significant reduction in pigment content when comparing drying at a high temperature (90 °C) to the corresponding ambient temperature while maintain the rest of the storage conditions for corn. On the contrary, other studies have found significant differences in the pigment content which are attributed not only to the drying of the grain but also to duration of the process (Quackenbush 1963). In spite of this controversy, the optimal conditions for the preservation of cereals, and particularly to preserve the carotenoids present in them are always reached by employing an exhaustive method of water removal (dehydration) and/or storage at low temperatures.

Studies on the storage of grains and processed grains also allow for the evaluation of aspects relating to the post-carotenogenic metabolism such as esterification with fatty acids which mediate its accumulation and stability in these foods (Kaneko and Oyanagi 1995; Kaneko et al. 1995). The increase in the esterification of lutein as the storage of grain progresses, including storage under controlled temperature, is one aspect that has been characterised by several authors

(Ahmad et al. 2013; Mellado-Ortega et al 2015). This provides information about the ideal conditions for the storage of cereals which preserves the contribution of these phytochemicals. In addition, a specific evaluation has been applied in order to characterise the enzymatic systems involved in esterification of the xanthophylls during the storage of grains and flours of tritordeum and durum wheat. This has resulted in important differences attributed not only to the nature of esterification in each case, but also to the enzymatic regional selectivity of the lutein and fatty acid molecules participating in such reactions (Mellado-Ortega 2013). Regardless of the type of plant material, either grain or flour, the stability of the esterified and free pigment has been studied and it has been found that the degradation rate is always greater for the latter case. This provides valuable information which could be used in crop enhancement programs of cereals. In agreement with these results the studies aimed to increase the cereal contents in carotenoids should be based on the selection of varieties with greater concentrations of esterified xanthophylls with the objective of increasing their ability to store them within the grain tissues, along with improving their stability in the seeds.

On the other hand, the influence of thermal pre-treatments on the storage of flour has been studied to evaluate the generation of free radicals and their impact on endogenous antioxidants (Andersen et al. 2011), as well as the possible intervention of degrading enzymes (Rodríguez-Amaya 1997).

5. Effect of processing on the carotenoid contents of cereals.

As explained above, the different processing techniques undoubtedly modify the carotenoid content present in the final product of processed cereals, due to, among other factors, an uneven distribution of the pigments in the different parts of the grain (Konopka et al. 2004; Zhou et al. 2004). This has led to the modification and

development of new processing techniques with the aim of preserving or stimulating the carotenoid contents along with other phytochemicals of nutritional importance. These emerging techniques are being incorporated into the traditional enhancement programs and genetic manipulation for obtaining bio-fortified grain crops (Fardet 2010; Hemery et al. 2007). Cereals are generally processed in two ways: dry fractioning followed by baking (at different temperature conditions, water contents and pressure) to produce product such as pasta, biscuits, breakfast cereal, etc., and fermentation, which produce alcoholic beverages. The production of the most popular cereal-derived products such as bread and baked goods include both treatments.

The fractionation and grinding processes of the cereal grains have allowed for the detailed analysis of carotenoids in the different layers and industrial fractions of the seeds (Ndolo and Beta 2013). The analyses of whole-grain products and those containing the germ (Kean et al. 2008), flour without bran and its formulations (Žilić et al. 2012), whole and shelled grains (Kean et al. 2011), grains, flour and semolina (Fares et al. 2008; Luterotti and Kljak 2010), are among the most common comparisons. One of the first studies carried out in this way was done by Blessin et al. (1963) on corn. These authors analysed the effect of manual and industrial processing on the pigment content found in different layers of the grain. Recently, Sellappan et al. (2009) has described the losses in β -carotene, iron and zinc in genetically modified rice varieties subjected to polishing and have estimated them to be higher than 70%. Polishing is a common industrial practice for rice grains, which involves the near complete elimination of the aleurone layer and the embryo (Juliano 1994). The classification of the genotype of cereals, by means of the characterisation of its pigment contents due to specific formulations of the grains can be a good example of the use of these crop enhancement techniques. Descriptive analyses like those of Siebenhandl et al. (2007)

contribute to this aspect, evaluating formulations of different bran particle sizes and flour for the classification and selection of diverse genotypes of wheat and barley. The size of the particle generated in these processes has a great influence from the technical and nutritional viewpoints. A reduction in the particle size facilitates the liberation of vitamins and other compounds from the outer layers of the grain (Fратиanni et al. 2005; Kahlon et al. 1986; Zhou et al. 2004). Zhou et al. (2004) showed that micronizing the aleurone layer resulted in greater antioxidant activity compared to the non-micronized aleurone layers of bran and grain, possibly due to a greater availability of antioxidants.

The diverse thermal treatments which are applied with the objective of prolonging the shelf-life of processed cereal products have also been reproduced in the laboratory to determine their effect on carotenoid contents. Toasting the cereal seems to have a rather pronounced repercussion which causes a significant reduction in pigment contents. This was reported by De Oliveira and Rodríguez-Amaya (2007), who analysed a series of fresh and processed corn products and estimated the loss in zeaxanthin to be about 53% after toasting the corn flour. Similar results have been described as a result of cooking rice (Lamberts and Delcour 2008). Scott and Eldridge (2005) performed a comparison of diverse thermal treatments and their effects on the carotenoid profiles of fresh, frozen and canned corn and found that the thermal treatments applied to corn prior to canning did not produce a relevant decrease in its carotenoid content. The freezing and scalding pre-treatments applied to cereals and vegetables in general may result in an increase in the bioavailability of carotenoids (Selman 1994). The heating by microwave of rice bran produced similar results without a reduction in the pigment levels which makes it a good alternative for favouring the stability of this cereal formulation (Abdul-Hamid et al. 2007). A more complex treatment consists of an extrusion, a multi-step treatment which is widely applied in the breakfast cereal

industry. The mild conditions applied during the extrusion (high moisture content, short treatment time and low temperature) have a clearly positive effect on the retention of vitamins and carotenoids and the decrease in lipid peroxidation, among others (Cheftel 1986; Singh et al. 2007). Guzmán-Tello and Cheftel (1990) studied the changes in the concentration of β -carotene under the most severe extrusion conditions of wheat flour and estimated losses ranging from 38 to 73% for an applied temperature interval of 125 to 200 °C. Other treatments such as those involved in the malting process of barley are characterised for generating contradictory results concerning their effect on pigment levels. For example, Goupy et al. (1999) registered losses (of approximately 76%) as well as gains in carotenoid contents in a study with different varieties of barley.

The analyses carried out during the different steps involved in the processing of semolina and making of pasta indicate that it is during the kneading phase when the greatest decrease in carotenoid content is registered while the subsequent drying and maturation periods turn out to be the least aggressive (Fратиanni et al. 2012; Hidalgo et al. 2010; Panfili et al. 2005). At the same time, important losses (up to 66%) in carotenoids during the kneading phase are associated with the production of bread, biscuits and baked goods (Leenhardt et al. 2006b). Other authors, however, point out different production phases, especially the baking step as being influential in the reduction in pigments. In a study of the evolution of pigments during the production of bread from wheat flour and einkorn, Hidalgo et al. (2010) registered averaged losses of 47% for the crust compared to 21% for the centre of the bread. After comparing the two types of flour, einkorn was found to suffer the greatest loss in pigments. Degradation during kneading is governed by an enzymatic oxidation of the pigments which has led some authors to relate both aspects (degradative enzymatic activity and carotenoid loss) in the enhancement programs for the selection of optimal cereal genotypes for the bread

making industry (Borrelli et al. 2003; Fu et al. 2013; Leenhardt et al. 2006a,b; Trono et al. 1999).

In the case of tritordeum, the properties of this new cereal make it one of the most suitable for bread applications, similar to common wheat, but on the contrary, less appropriate for formulations of semolina. As mentioned before, tritordeum flour has recently began to be commercialised with the aim of several applications such as bread, sandwich bread, biscuits, muffins and other cereal products (www.agrasys.es). In the sense, even though the elaboration process of bread and other products results in losses in carotenoid contents, as in any other cereal, tritordeum bread presents superior levels of lutein (6 times greater) than bread made from traditional wheat (according to the analyses carried out in our laboratory for the Agrasys company).

6. Applications of the biotechnology of carotenoids to cereal selection and breeding.

Biotechnology, as a vegetable enhancement technique, applied to the increase in the carotenoid content in cereals has been used fundamentally on three species: corn, wheat and sorghum. In the case of rice the grains do not produce any carotenoid in the endosperm, however a new rice variety, *Golden Rice*, was developed at the beginning of this century by engineering grains to produce β -carotene (provitamin A) in order to help combating vitamin A deficiency in populations with rice-based traditional diets (this applies to countries such as India, Vietnam, Bangladesh, the Philippines, and Indonesia) (Paine et al. 2005; Ye et al. 2000). The relevance of these cereals and their impact on the nutrition of the populations of under-developed countries is reflected in biofortification programs such as the one of Harvestplus, belonging to the Consultive Group on International Agricultural Research (CGIAR) which is focused on the

501 augmentation of zinc, iron and provitamin A contents in basic crops producing staple
502 foods (<http://www.harvestplus.org/>).

503 The last developments in the exploration of the genomes of the main cereals
504 such as wheat has been recently reported (Eversole et al. 2014), and has been made
505 thanks to the existence of genomic models such as rice and sorghum (Mace and Jordan
506 2011; Paterson et al. 2009). A large part of the investigations has been based, in the first
507 instance, on the study of the variability in the feature of being susceptible to traditional
508 selection and breeding, in this case, for the enhancement of carotenoid content in a
509 given population and in the second place on the ability to inherit such a trait (Clarke et
510 al. 2006; Santra et al. 2005). Rice, sorghum and wheat genomes have shown diverse
511 genetic importance for this characteristic as reflected in the existence of elite germplasm
512 in these species (Chander et al. 2008; Ibrahim and Juvik 2009). Up to now, the study
513 has been focused on the identification of QTLs (Quantitative Trait Loci) whose allelic
514 variation is associated with a variation of quantitative trait such as the carotenoid
515 content, which varies continuously. Numerous studies can be found which describe such
516 locus and their relation to the phenotypic variation in the carotenoid contents of the
517 endosperm of cereals (Chander et al. 2008; Patil et al. 2008; Pozniak et al. 2007; Salas-
518 Fernández et al. 2008; Wong et al. 2004). Studies at this level go beyond the
519 justification of the overall carotenoid content and identify specific QTLs for certain
520 carotenoids such as lutein and β -carotene. Salas-Fernández et al. (2008) detected up to
521 five QTLs for β -carotene content in sorghum located at the 1, 2, and 10 chromosomes.
522 Howitt et al. (2009) identified three QTLs for lutein contents in the 3B, 5B and 7A
523 chromosomes of durum wheat. More recently, Blanco et al. (2011) have mapped QTLs
524 for β - and α -carotene in the chromosomes 2A, 3B and 7A.

525 In wheat, the most determinant QTLs for the pigment contents have been

repeatedly mapped onto the 7A and 7B chromosomes, both on durum wheat and common wheat. The PSY1 codifying genes (phytoene synthase, EC 2.5.1.32) generally co-segregate with these QLTs. The allelic variations of these genes are numerous and their continuous appearance is very probable. This demonstrates a large number of genetic polymorphisms as well as many genomes to be explored (Ravel et al. 2013). One of the strategies in crop improvement is the search for new variation sources in related wild species such as *Lophopyrum ponticum* (Zhang et al. 2005) and *H. chilense* (Rodríguez-Suárez et al. 2011; Rodríguez-Suárez and Atienza 2012). The small grains of *H. chilense* are characterised as having a large amount of carotenoids and present at least two loci for the content of pigments located in the 2H^{ch} and 7H^{ch} chromosomes (Álvarez et al. 1998 1999; Atienza et al. 2004, 2008). Knowledge about the chromosomal location was first studied by Álvarez et al. (1998) using additional lines of *H. chilense* in wheat. These authors located genes responsible for the level of pigments (determined as YPC) in *H. chilense* at chromosome 7 and mapped this feature onto the α arm of this chromosome. However, the possibility of an interaction between *H. chilense* genes and the durum wheat genes of tritordeum promoted a search for new locus. Subsequently, by means of the construction of the first genetic map of *H. chilense*, a new QTL entitled *carot 1* was identified at chromosome 2 (Atienza et al. 2004; Hernández et al. 2001). Currently, the generation of maps of a wider range for these purposes is now available, and therefore there is a greater chance of finding genomic regions of interest in *H. chilense* (Rodríguez-Suárez et al. 2012).

This information constitute the first steps towards the development of a “Marker Assisted Selection” (MAS) program for the content of carotenoids in tritordeum, which is interesting for the identification and further transference of particular genes related with the carotenoid content from tritordeum to wheat lines subjected to improvement

programs. In this way, the amphiploid tritordeum will also become a useful species for the enhancement of cereals acting as bridge species between barley and wheat (Atienza et al. 2005, 2007a,b Martín et al. 1999). The tritordeum lines, regardless of their ploidy level, show carotenoid contents which are higher than their wheat relatives (Atienza et al. 2007a). Therefore, the study of the genetic variation of this feature in tritordeum and its relation to the level of pigments of the parental species *H. chilense* and durum wheat has become a very active research topic. From these studies it appears that even though the genome H^{ch} is clearly responsible for the level of pigments in tritordeum there can be interactions between the genetic backgrounds of both parental species that may not be obvious (Álvarez et al. 1999). Recently the first exhaustive analysis of the carotenoid profile of *H. chilense* was carried out, discovering that more than half of the lutein was esterified, with a similar esterification pattern to tritordeum. This corroborates the previous results and indicates that the esterification pattern of tritordeum should result from the genetic background of this wild barley (Mellado-Ortega et al. 2015). Recent studies have demonstrated that lutein esterification in wheat and *H. chilense* is controlled by a loci located at the chromosomes 7D and 7H^{ch}, respectively (Ahmad et al. 2015; Mattera et al. 2015).

The utility of the metabolic engineering to improve human nutrition reaches its best objectives in the basic foods field (staple foods) which have an important impact on the nutrition of under-developed populations. The new generation of crops with the *Golden* label such as *Golden Potato* or *Golden Canola* are increasing with time leading up to the well-known *Golden Rice* (Bai et al. 2011; Beyer 2010). The success of last case is encouraging similar studies and improvement programs with other species. The intense investigation from the first attempts made in rice by Burkhardt et al. (1997) to the most recent ones with the *Golden Rice 2* generation (Paine et al. 2005) have

provided stimulating ideas for this purpose. Another example is the introduction and expression of the *yl* gene (gene coding for phytoene synthase in yellow endosperm corn) from corn into the hexaploid common wheat and its positive effect on increasing the carotenoid content in the endosperm (Cong et al. 2009). In the case of corn, the improved use of the γ -zein promotor (Marzábal et al. 1998), highly specific for the expression in endospermic tissue (super γ -zein), derived into an increase in carotenoids of up to 34 times with a preferential accumulation of β -carotene (Aluru et al. 2008). A more complex strategy was the one designed by Zhu et al. (2008) with the generation of an entire transgenic corn plant battery that resulted from the multiple combinations possible derived from a multi-gene transformation. More recently, the simultaneous increase in the vitamin contents (vitamins A, B and C) involving the simultaneous manipulation of three different metabolic routes (Naqvi et al. 2009), establishes corn as a likely crop to maximise carotenoid provitamin A contents, showing less restrictions than others such as rice or wheat (Wurtzel et al. 2012). The most recent *Golden Rice* investigations focus now on the study of the stability and transference of the transgenes for the selection and breeding of rice crops with a higher efficiency through introgression and other enhancement techniques (Datta et al. 2006, 2007). Other less studied grains such as sorghum and millet, but having a great impact on under-developed populations such as the African, are increasingly adapting to these techniques making its way into the biofortification mediated by genetic manipulation (O’Kennedy et al. 2006; www.grandchallenges.org/ImproveNutrition/Challenges/NutrientRichPlants/Pages/Sorghum.aspx).

599

600 **Acknowledgements**

601 This work was supported by funding from the Ministerio de Ciencia e
602 Innovación (Spanish Government, Projects AGL2010-14850/ALI and AGL2014-
603 53195R). EMO was the recipient of a JAE-Predoctoral grant (CSIC) co-financed by the
604 ESF. Authors are members of the IBERCAROT Network, funded by CYTED (ref.
605 112RT0445).

606

607 **Conflict of interest.** The authors declare that they have no conflict of interest.

References

- Abdel-Aal EM, Young JC, Rabalski I et al (2007) Identification and quantification of seed carotenoids in selected wheat species. *J Cereal Sci* 55:787–794
- Abdul-Hamid A, Sulaiman RRR, Osman A et al (2007) Preliminary study of the chemical composition of rice milling fractions stabilized by microwave heating. *J Food Comp Anal* 20:627–637
- Adom KK, Sorrells, ME, Liu RH (2003) Phytochemical profiles and antioxidant activity of wheat varieties. *J Agric Food Chem* 51:7825–7834
- Adom KK, Sorrells ME, Liu RH (2005) Phytochemicals and antioxidant activity of milled fractions of different wheat varieties. *J Agric Food Chem* 53:2297–2306
- Agrasys. <http://www.agrasys.es/es/index.html> (accessed December 2014).
- Ahmad FT, Asenstorfer RE, Soriano IR et al (2013) Effect of temperature on lutein esterification and lutein stability in wheat grain. *J Cereal Sci* 58:408–413
- Ahmad FT, Mather DE, Law H-Y et al (2015) Genetic control of lutein esterification in wheat (*Triticum aestivum* L.) grain. *J Cereal Sci* 64:109–115
- Ahmed SS, Lott MN, Marcus DM (2005) The macular xanthophylls. *Surv Ophthalmol* 50:183–193
- Aluru M, Xu Y, Guo R et al (2008) Generation of transgenic maize with enhanced provitamin A content. *J Exp Bot* 59:3551–3562
- Álvarez JB, Ballesteros J, Arriaga HO et al (1995) Rheological properties and baking performances of flours from hexaploid tritordeums. *J Cereal Sci* 23:291–299

631 Álvarez JB, Martín LM, Martín A (1998) Chromosomal localization of genes for
 632 carotenoid pigments using addition lines of *Hordeum chilense* in wheat. Plant
 633 Breeding 117:287–289
 634 Álvarez JB, Martín LM, Martín A (1999) Genetic variation for carotenoid pigment
 635 content in the amphiploid *Hordeum chilense* × *Triticum turgidum* conv. *durum*.
 636 Plant Breeding 118:187–189
 637 Alves-Rodrigues A, Shao A (2004) The science behind lutein. Toxicol Lett 150:57–83
 638 Andersen ML, Erichsen HR, Skibsted LH et al (2011) Heat induced formation of free
 639 radicals in wheat flour. J Cereal Sci 54:494–498
 640 Arya SS, Parihar DB (1981) Effect of moisture and temperature on storage changes in
 641 lipids and carotenoids of atta (wheat flour). Food Nahrung 25:121–126
 642 Atienza S, Ballesteros J, Martín A et al (2007a) Genetic variability of carotenoid
 643 concentration and degree of esterification among tritordeum (×*Tritordeum*
 644 Ascherson et Graebner) and durum wheat accessions. J Agric Food Chem 55:4244–
 645 4251
 646 Atienza SG, Avila CM, Ramírez MC et al (2005) Application of near infrared
 647 reflectance spectroscopy to the determination of carotenoid content in tritordeum
 648 for breeding purposes. Aust J Agric Res 56:85–89
 649 Atienza SG, Martín A, Pecchioni N et al (2008) The nuclear-cytoplasmic interaction
 650 controls carotenoid content in wheat. Euphytica 159:325–331
 651 Atienza SG, Martín AC, Ramírez MC et al (2007b) Effects of *Hordeum chilense*
 652 cytoplasm on agronomic traits in common wheat. Plant Breeding 126:5–8
 653 Atienza SG, Ramírez CM, Hernández P et al (2004) Chromosomal location of genes for
 654 carotenoid pigments in *Hordeum chilense*. Plant Breeding 123:303–304

655 Bai C, Twyman RM, Farré G et al (2011) A golden era- pro-vitamin A enhancement in
 656 diverse crops. *In Vitro Cell Dev Biol Plant* 47:205–221
 657 Baker R, Günter C (2004) The role of carotenoids in consumer choice and the likely
 658 benefits from their inclusion into products for human consumption. *Trends Food*
 659 *Sci Tech* 15:484–488
 660 Ballesteros JB, Ramírez MC, Martínez C et al (2005) Registration of HT621, a high
 661 carotenoid content tritordeum germplasm line. *Crop Sci* 45:2662–2663
 662 Bechoff A, Dhuique-Mayer C, Dornier M et al (2010) Relationship between the kinetics
 663 of β -carotene degradation and formation of norisoprenoids in the storage of dried
 664 sweet potato chips. *Food Chem* 121:348–357
 665 Belefant-Miller H, Grace SC (2010) Variations in bran carotenoid levels within and
 666 between rice subgroups. *Plant Food Hum Nutr* 65:358–363
 667 Berardo N, Brenna OV, Amato A et al (2004) Carotenoids concentration among maize
 668 genotypes measured by near infrared reflectance spectroscopy (NIRS). *Innov Food*
 669 *Sci Emerg* 5:393–398
 670 Berardo N, Manzzinelli G, Valoti P et al (2009) Characterization of maize germplasm
 671 for the chemical composition of the grain. *J Agric Food Chem* 57:2378–2384
 672 Beyer P (2010) Golden Rice and ‘Golden’ crops for human nutrition. *New Biotechnol*
 673 27:478–481
 674 Blanco A, Colasuonno P, Gadaleta A et al (2011) Quantitative trait loci for yellow
 675 pigment concentration and individual carotenoid compounds in durum wheat. *J*
 676 *Cereal Sci* 54:255–264
 677 Blessin CW, Brecher JD, Dimler RJ (1963) Carotenoids of corn and sorghum: V.
 678 Distribution of xanthophylls and carotenes in hand-dissected and dry-milled
 679 fractions of yellow dent corn. *Cereal Chem* 40:582–590

680 Blessin CW, VanEtten H, Wiebe R (1958) Carotenoid content of the grain from yellow
681 endosperm-type sorghums. *Cereal Chem* 35:359–365

682 Borrelli GM, De Leonardis AM, Fares C et al (2003) Effects of modified processing
683 conditions on oxidative properties of semolina dough and pasta. *Cereal Chem*
684 80:225–231

685 Borrelli GM, De Leonardis AM, Platani C et al (2008) Distribution along durum wheat
686 kernel of the components involved in semolina colour. *J Cereal Sci* 48:494–502

687 Britton G, Hornero-Méndez D (1997) Carotenoids and colour in fruit and vegetables.
688 In: Tomás-Barberán FA, Robins RJ (eds) *Phytochemistry of fruit and vegetables*,
689 Clarendon Press, Oxford (United Kingdom), Ch. 2, pp11–27

690 Britton G, Liaaen-Jensen S, Pfander H (2009) Carotenoids. vol 5: Nutrition and Health.
691 Basel: Birkhäuser Verlag.

692 Britton G, Khachik F (2009) Carotenoids in Food. In: Britton G, Liaaen-Jensen S,
693 Pfander H (eds) *Carotenoids*, vol 5: Nutrition and Health. Birkhäuser Verlag Basel,
694 pp45–66

695 Burkhardt PK, Beyer P, Wünn J et al (1997) Transgenic rice (*Oryza sativa*) endosperm
696 expressing daffodil (*Narcissus pseudonarcissus*) phytoene synthase accumulates
697 phytoene, a key intermediate of provitamin A biosynthesis. *Plant J* 11:1071–1078

698 Burt A, Grainger C, Young J et al (2010) Impact of postharvest handling on carotenoid
699 concentration and composition in high-carotenoid maize (*Zea mays* L.) kernels. *J*
700 *Agric Food Chem* 58:8286–8292

701 Calucci L, Capocchi A, Galleschi L et al (2004) Antioxidants, free radicals, storage
702 proteins, puroindolines, and proteolytic activities in bread wheat (*Triticum*
703 *aestivum*) seeds during accelerated aging. *J Agric Food Chem* 52:4274–4281

704 Calvo MM (2005) Lutein: a valuable ingredient of fruit and vegetables. *Crit Rev Food*

705 Sci 45:1–26

706 Carbonell JV, Piñaga F, Yusá V et al (1986) The dehydration of paprika with ambient
 707 and heated air and the kinetics of colour degradation during storage. J Food Eng
 708 5:179–193

709 Chander S, Guo YQ, Yang XH et al (2008) Using molecular markers to identify two
 710 major loci controlling carotenoid contents in maize grain. Theor Appl Genet
 711 116:223–233

712 Cheftel JC (1986) Nutritional effects of extrusion-cooking. Food Chem 20:263–283

713 Chelowski J (ed) (1994) Cereal grain. Mycotoxins, fungi and quality in drying and
 714 storage. Developments in food Science. Elsevier, Amsterdam

715 Choi Y, Jeong H, Lee J (2007) Antioxidant activity of methanolic extracts from some
 716 grains consumed in Korea. Food Chem 103:130–138

717 Clarke FR, Clarke JM, McCaig TN et al (2006) Inheritance of yellow pigment
 718 concentration in seven durum wheat crosses. Can J Plant Sci 86:133–141

719 Cong L, Wang C, Chen L et al (2009) Expression of phytoene synthase 1 and carotene
 720 desaturase crtI genes result in an increase in the total carotenoids content in
 721 transgenic elite wheat (*Triticum aestivum* L.). J Agric Food Chem 57:8652–8660

722 Cristobal, JAR (1965) Variation of the content of carotenes in cereals in relation to the
 723 time of storage. An Inst Invest Vet 15:85–92

724 Datta K, Rai M, Parkhi V et al (2006) Improved “Golden” rice and post-transgeneration
 725 enhancement of metabolic target products of carotenoids (β -carotene) in transgenic
 726 elite cultivars (IR64 and BR29). Current Sci 91:935–939

727 Datta SK, Datta K, Parkhi V et al (2007) Golden rice: introgression, breeding, and field
 728 evaluation. Euphytica 154:271–278

729 De Oliveira G, Rodriguez-Amaya D (2007) Processed and prepared corn products as
730 sources of lutein and zeaxanthin: compositional variation in the food chain. *J Food*
731 *Sci* 72:S79–S85

732 Dhuique-Mayer C, Tbatou M, Carail M et al (2007) Thermal degradation of antioxidant
733 micronutrients in citrus juice: kinetics and newly formed compounds. *J Agric Food*
734 *Chem* 55:4209–4216

735 Di Silvestro R, Marotti LL, Bosi S et al (2012) Health-promoting phytochemicals of
736 italian common wheat varieties grown under low-input agricultural management. *J*
737 *Sci Food Agric* 92:2800–2810

738 Digesù AM, Platani C, Cattivelli L et al (2009) Genetic variability in yellow pigment
739 components in cultivated and wild tetraploid wheats. *J Cereal Sci* 50:210–218

740 Doblado-Maldonado AF, Pike OA, Sweley JC et al (2012) Key issues and challenges in
741 whole wheat flour milling and storage. *J Cereal Sci* 56:119–126

742 Egesel CO, Wong JC, Lambert RJ et al (2003) Combining ability of maize inbreds for
743 carotenoids and tocopherols. *Crop Sci* 43:818–823

744 El-Agamey A, McGarvey DJ (2008) Carotenoid Radicals and Radical Ions. In: Britton
745 G, Liaaen-Jensen S, Pfander H (eds) Carotenoids, vol 4: Natural Functions.
746 Birkhäuser Verlag Basel, pp119–154

747 Eversole K, Feuillet C, Mayer KFX et al (2014) Slicing the wheat genome. *Science*
748 18:285–287

749 Fardet A (2010) New hypotheses for the health-protective mechanisms of whole-grain
750 cereals: what is beyond fibre? *Nutr Res Rev* 23:65–134

751 Fardet A, Rock E, Rémésy C (2008) Is the *in vitro* antioxidant potencial of whole-grain
752 cereals and cereal products well reflected *in vivo*? *J Cereal Sci* 48:258–276

753 Fares C, Codianni P, Nigro F et al (2008) Processing and cooking effects on chemical,
 754 nutritional and functional properties of pasta obtained from selected emmer
 755 genotypes. *J Sci Food Agric* 88:2435–2444

756 Farrington FF, Warwick MJ, Shearer G (1981) Changes in the carotenoids and sterol
 757 fractions during the prolonged storage of wheat flour. *J Sci Food Agric* 32:948–950

758 Fernandez-Orozco R, Gallardo-Guerrero L, Hornero-Méndez D (2013) Carotenoid
 759 profiling in tubers of different potato (*Solanum* sp) cultivars: Accumulation of
 760 carotenoids mediated by xanthophyll esterification. *Food Chem* 141:2864–2872

761 Fish WW, Davis AR (2003) The effects of frozen storage conditions on lycopene
 762 stability in watermelon tissue. *J Agric Food Chem* 51:3582–3585

763 Fratianni A, Di Criscio T, Mignogna R et al (2012) Carotenoids, tocots and retinols
 764 evolution during egg pasta-making processes. *Food Chem* 131:590–595

765 Fratianni A, Irano M, Panfili G et al (2005) Estimation of color of durum wheat.
 766 Comparison of WSB, HPLC, and reflectance colorimeter measurements. *J Agric*
 767 *Food Chem* 53:2373–2378

768 Frei M, Becker K (2005) Fatty acids and all-trans- β -carotene are correlated in
 769 differently colored rice landraces. *J Sci Food Agric* 85:2380–2384

770 Fu BX, Schlichting L, Pozniak CJ et al (2013) Pigment loss from semolina to dough:
 771 rapid measurement and relationship with pasta colour. *J Cereal Sci* 57:560–566.

772 Goupy P, Hugues M, Boivin P, Amiot MJ (1999) Antioxidant composition and activity
 773 of barley (*Hordeum vulgare*) and malt extracts and of isolated phenolic compounds.
 774 *J Sci Food Agric* 79:1625–1634

775 Graham RD, Rosser JM (2000) Carotenoids in staple foods: their potential to improve
 776 human nutrition. *Food Nutr Bull* 21:404–409

777 Grand Challenges in Global Health.
 778 [www.grandchallenges.org/ImproveNutrition/Challenges/NutrientRichPlants/Pages/](http://www.grandchallenges.org/ImproveNutrition/Challenges/NutrientRichPlants/Pages/Sorghum.aspx)
 779 [Sorghum.aspx](http://www.grandchallenges.org/ImproveNutrition/Challenges/NutrientRichPlants/Pages/Sorghum.aspx) (accessed May 2015).

780 Guzman-Tello R, Cheftel JC (1990) Colour loss during extrusion cooking of β -
 781 carotene- wheat flour mixes as an indicator of the intensity of thermal and oxidative
 782 processing. *Int J Food Sci Technol* 25:420–434

783 Harvestplus. Breeding Crops for Better Nutrition. <http://www.harvestplus.org/> (accessed
 784 May 2015).

785 Hemery Y, Rouau X, Lullien-Pellerin V et al (2007) Dry processes to develop wheat
 786 fractions and products with enhanced nutritional quality. *J Cereal Sci* 46:327–347

787 Hentschel V, Kranl K, Hollmann J et al (2002) Spectrophotometric determination of
 788 yellow pigment content and evaluation of carotenoids by high-performance liquid
 789 chromatography in durum wheat grain. *J Agric Food Chem* 50:6663–6668

790 Hernández P, Dorado G, Prieto P et al (2001) A core genetic map of *Hordeum chilense*
 791 and comparisons with maps of barley (*Hordeum vulgare*) and wheat (*Triticum*
 792 *aestivum*). *Theor Appl Genet* 102:1259–1264

793 Hidalgo A, Brandolini A (2008a) Protein, ash, lutein and tocopherols distribution in einkorn
 794 (*Triticum monococcum* L. ssp. *monococcum*) seed fractions. *Food Chem* 107:444–
 795 448

796 Hidalgo A, Brandolini A (2008b) Kinetics of carotenoids degradation during the storage
 797 of einkorn (*Triticum monococcum* L. ssp. *monococcum*) and bread wheat (*Triticum*
 798 *aestivum* L. ssp. *aestivum*) flours. *J Agric Food Chem* 56:11300–11305

799 Hidalgo A, Brandolini A, Pompei C (2010) Carotenoids evolution during pasta, bread
 800 and water biscuit preparation from wheat flours. *Food Chem* 121:746–751

801 Hidalgo A, Brandolini A, Ratti S (2009) Influence of genetic and environmental factors
802 on selected nutritional traits of *Triticum monococcum*. J Agric Food Chem
803 57:6342–6348

804 Hidalgo A, Brandolini A, Pompei C et al (2006) Carotenoids and tocopherols of einkorn
805 wheat (*Triticum monococcum* ssp. *monococcum* L.). J Cereal Sci 44:182–193

806 Howitt CA, Pogson BJ (2006) Carotenoid accumulation and function in seeds and non-
807 green tissues. Plant Cell Environ 29:435–445

808 Howitt CA, Cavanagh CR, Bowerman AF et al (2009) Alternative splicing, activation
809 of cryptic exons and amino acid substitutions in carotenoid biosynthetic genes are
810 associated with lutein accumulation in wheat endosperm. Funct Integr Genomics
811 9:363–376

812 Humphries JM, Graham RD, Mares DJ (2004) Application of reflectance colour
813 measurement to the estimation of carotene and lutein content in wheat and triticale.
814 J Cereal Sci 40:151–159

815 Ibrahim K, Juvik J (2009) Feasibility for improving phytonutrient content in vegetable
816 crops using conventional breeding strategies: case study with carotenoids and
817 tocopherols in sweet corn and broccoli. J Agric Food Chem 57:4636–4644

818 ICC (International Association for Cereal Science and Technology), 1990. Method 152.
819 Determination of the yellow pigment content of durum wheat semolina and flour.
820 In: Standard methods of the International Association for Cereal Science and
821 Technology. Verlag Moritz Schäfer: Detmold, Germany

822 Jood S, Kapoor AC (1994) Vitamin contents of cereal grains as affected by storage and
823 insect infestation. Plant Food Hum Nutr 46:237–243

824 Juliano, B.O. 1994. *Rice: Chemistry and Technology*, Juliano, B.O. (ed.), 2nd ed., St.
825 Paul, EUA, pp. 17–160 and 647–680.

826 Kahlon TS, Chow FI, Hoefer JL et al (1986) Bioavailability of vitamins A and E as
827 influenced by wheat bran and bran particle size. *Cereal Chem* 63:490–493

828 Kalt, W (2005) Effects of production and processing factors on major fruit and
829 vegetable antioxidants. *J Food Sci* 70:R11–R19

830 Kandlakunta B, Rajendran A, Thingnganing L (2008) Carotene content of some
831 common (cereals, pulses, vegetables, spices and condiments) and unconventional
832 sources of plant origin. *Food Chem* 106:85–89

833 Kaneko S, Oyanagi A (1995) Varietal differences in the rate of esterification of
834 endosperm lutein during the storage of wheat seeds. *Biosci Biotechnol Biochem*
835 59:2312–2313

836 Kaneko S, Nagamine T, Yamada T (1995) Esterification of endosperm lutein with fatty
837 acids during the storage of wheat seeds. *Biosci Biotechnol Biochem* 59:1–4

838 Kean EG, Bordenave N, Ejeta G et al (2011) Carotenoid bioaccessibility from whole
839 grain and decorticated yellow endosperm sorghum porridge. *J Cereal Sci* 54:450–
840 459

841 Kean EG, Ejeta G, Hamaker BR et al (2007) Characterization of carotenoid pigments in
842 mature and developing kernels of selected yellow-endosperm sorghum varieties. *J*
843 *Agric Food Chem* 55:2619–2626

844 Kean EG, Hamaker BR, Ferruzzi MG (2008) Carotenoid bioaccessibility from whole
845 grain and degermed maize meal products. *J Agric Food Chem* 56:9918–9926

846 Kim JK, Lee SY, Chu SM et al (2010) Variation and correlation analysis of flavonoids
847 and carotenoids in korean pigmented rice (*Oryza sativa* L.) cultivars. *J Agric Food*
848 *Chem* 58:12804–12809

849 Koca N, Burdurlu HS, Karadeniz F (2007) Kinetics of color changes in dehydrated
850 carrots. *J Food Eng* 78:449–455

851 Konopka I, Kozirok W, Rotkiewicz D (2004) Lipids and carotenoids of wheat grain and
 852 flour and attempt of correlating them with digital image analysis of kernel surface
 853 and cross-sections. Food Res Int 37:429–438
 854 Kurilich AC, Juvik JA (1999) Quantification of carotenoid and tocopherol antioxidants
 855 in *Zea mays*. J Agric Food Chem 47:1948–1955
 856 Lamberts L, Delcour JA (2008) Carotenoids in raw and parboiled brown and milled
 857 rice. J Agric Food Chem 56:11914–11919
 858 Landrum JT, Bone RA (2001) Lutein, zeaxanthin, and the macular pigment. Arch
 859 Biochem Biophys 385:28–40
 860 Lavelli V, Hidalgo A, Pompei C et al (2009) Radical scavenging activity of einkorn
 861 (*Triticum monococcum* L. subsp. *monococcum*) wholemeal flour and its
 862 relationship to soluble phenolic and lipophilic antioxidant content. J Cereal Sci
 863 49:319–321
 864 Leenhardt F, Lyan B, Rock E et al (2006a) Genetic variability of carotenoid
 865 concentration, and lipoxygenase and peroxidase activities among cultivated wheat
 866 species and bread wheat varieties. Eur J Agron 25:170–176
 867 Leenhardt F, Lyan B, Rock E et al (2006b) Wheat lipoxygenase activity induces greater
 868 loss of carotenoids than vitamin E during breadmaking. J Agric Food Chem
 869 54:1710–1715
 870 Lemmens L, De Vleeschouwer K, Moelants KRN et al (2010) β -Carotene isomerization
 871 kinetics during thermal treatments of carrot puree. J Agric Food Chem 58:6816–
 872 6824
 873 Li W, Beta T (2012) An evaluation of carotenoid levels and composition of glabrous
 874 canaryseed. Food Chem 133:782–786

875 Liaaen-Jensen S, Lutnaes BF (2008) E/Z Isomers and Isomerization. In: Britton G,
 876 Liaaen-Jensen S, Pfander H (eds) Carotenoids, vol 4: Natural Functions. Birkhäuser
 877 Verlag Basel, pp15–36

878 Lier JB, Lacroix LJ (1974) Carotenoids of durum wheat: induced high pigment levels
 879 obtained by treatment of the growing plant with chlorophenylthiotriethylamine
 880 (CPTA) hydrochloride. *Cereal Chem* 51:188–194

881 Lindley MG (1998) The impact of food processing on antioxidants in vegetable oils,
 882 fruits and vegetables. *Trends Food Sci Tech* 9:336–340

883 Liu RH (2007) Whole grain phytochemicals and health. *J Cereal Sci* 46:207–219

884 Luterotti S, Kljak K (2010) Spectrophotometric estimation of total carotenoids in cereal
 885 grain products. *Acta Chim Slov* 57:781–787

886 Lv J, Lu Y, Niu Y et al (2013) Effect of genotype, environment, and their interaction on
 887 phytochemical compositions and antioxidant properties of soft winter wheat flour.
 888 *Food Chem* 138:454–462

889 Mace ES, Jordan DR (2011) Integrating sorghum whole genome sequence information
 890 with a compendium of sorghum QTL studies reveals uneven distribution of QTL
 891 and of gene-rich regions with significant implications for a crop improvement.
 892 *Theor Appl Genet* 123:169–191

893 Maiani G, Periago-Castón MJ, Catasta G et al (2009) Carotenoids: actual knowledge on
 894 food sources, intakes, stability and bioavailability and their protective role in
 895 humans. *Mol Nutr Food Res* 53:S194–S218

896 Mamatha B, Sangeetha R, Baskaran V (2011) Provitamin-A and xanthophyll
 897 carotenoids in vegetables and food grains of nutritional and medicinal importance.
 898 *Int J Food Sci Tech* 46:315–323

899 Martín A, Álvarez JA, Martín LM et al (1999) The development of tritordeum: a novel
900 cereal for food processing. *J Cereal Sci* 30:85–95

901 Martín A, Chapman V (1977) A hybrid between *Hordeum chilense* and *Triticum*
902 *aestivum*. *Cereal Res Commun* 5:365–368

903 Martín A, Sanchez-Monge Laguna E (1982) Cytology and morphology of the
904 amphiploid *Hordeum chilense* x *Triticum turgidum* conv. *durum*. *Euphytica*
905 31:261–267

906 Marzábal P, Busk PK, Ludevid MD et al (1998) The bifactorial endosperm box of γ -
907 zein gene: characterisation and function of the Pb3 and GZM cis-acting elements.
908 *Plant J* 16:41–52

909 Mattera G, Cabrera A, Hornero-Méndez D et al (2015) Lutein esterification in wheat
910 endosperm is controlled by the homoeologous group 7, and is increased by the
911 simultaneous presence of chromosomes 7D and 7Hch from *Hordeum chilense*.
912 *Crop Pasture Sci* 66:912–921

913 McKeivith B (2004) Nutricional aspects of cereals. *Nutrition Bulletin* 29:111–142

914 Mellado-Ortega E (2013) Biosynthesis, accumulation and stability of carotenoids in
915 cereals. Comparative study of durum wheat (*Triticum turgidum*) and tritordeum
916 (*Hordeum chilense* x *T. turgidum* conv. *durum*). PhD Thesis Dissertation.
917 University of Seville, Spain.

918 Mellado-Ortega E, Hornero-Méndez D (2012) Isolation and identification of lutein
919 esters, including their regioisomers, in tritordeum ($\times$ *Tritordeum* Ascherson et
920 Graebner) grains: evidence for a preferential xanthophyll acyltransferase activity.
921 *Food Chem* 135:1344–1352

922 Mellado-Ortega E, Atienza SG, Hornero-Méndez D (2015) Carotenoid evolution during
923 postharvest storage of durum wheat (*Triticum turgidum* conv. *durum*) and
924 tritordeum ($\times$ *Tritordeum* Ascherson et Graebner) grains. J Cereal Sci 62:134–142

925 Mellado-Ortega E, Hornero-Méndez D (2015) Carotenoid profiling of *Hordeum*
926 *chilense* grains: the parental proof for the origin of the high carotenoid content and
927 esterification pattern of tritordeum. J Cereal Sci 62:15–21

928 Menkir A, Maziya-Dixon B (2004) Influence of genotype and environment on β -
929 carotene content of tropical yellow-endosperm maize genotypes. *Maydica* 49:313–
930 318

931 Mínguez-Mosquera MI, Jarén-Galán M, Gandul-Rojas B, Hornero-Méndez D, Garrido-
932 Fernández J, Gallardo-Guerrero L (eds) (1997) Clorofilas y carotenoides en
933 tecnología de alimentos. Servicio de Publicaciones de la Universidad de Sevilla,
934 Sevilla

935 Mínguez-Mosquera MI, Gandul-Rojas B (1994) Mechanism and kinetics of carotenoid
936 degradation during the processing of green table olives. J Agric Food Chem
937 42:1551–1554

938 Moros EE, Darnoko D, Cheryan M, Perkins EG, Jerrell J (2002) Analysis of
939 xanthophylls in corn by HPLC. J Agric Food Chem 50:5787–5790

940 Nakornriab M, Sriseadka T, Wongpornchai S (2008) Quantification of carotenoid and
941 flavonoid components in brans of some Thai black rice cultivars using supercritical
942 fluid extraction and high-performance liquid chromatography-mass spectrometry. J
943 Food Lipids 15:488–503

944 Naqvi S, Zhu C, Farre G et al (2009) Transgenic multivitamin corn through
945 biofortification of endosperm with three vitamins representing three distinct
946 metabolic pathways. P Natl Acad Sci USA 106:7762–7767

947 Ndolo VU, Beta T (2013) Distribution of carotenoids in endosperm, germ, and aleurone
 948 fractions of cereal grain kernels. *Food Chem* 139:663–671

949 Nghia PT, Liem DT, Hai TV et al (2006) Effect of storage conditions on total
 950 carotenoid content in golden rice grains. *Omonrice* 14:18–27

951 Nicoli MC, Anese M, Parpinel M (1999) Influence of processing on the antioxidant
 952 properties of fruit and vegetables. *Trends Food Sci Tech* 10:94–100

953 Nishino H (1997) Cancer prevention by natural carotenoids. *J Cell Biochem* 27:86–91

954 O’Kennedy MM, Grootboom A, Shewry PR (2006) Harnessing sorghum and millet
 955 biotechnology for food and health. *J Cereal Sci* 44:224–235

956 Okarter N, Liu C, Sorrells M et al (2010) Phytochemical content and antioxidant
 957 activity of six diverse varieties of whole wheat. *Food Chem* 119:249–257

958 Olson JA (1989) Biological actions of carotenoids. *J Nutr* 119:94–95

959 Ouchi A, Aizawa K, Iwasaki Y et al (2010) Kinetic study of the quenching reaction of
 960 singlet oxygen by carotenoids and food extracts in solution. Development of a
 961 singlet oxygen absorption capacity (SOAC) assay method. *J Agric Food Chem*
 962 58:9967–9978

963 Paine JA, Shipton CA, Chaggner S et al (2005) Improving the nutritional value of
 964 Golden Rice through increased pro-vitamin A content. *Nat Biotechnol* 23:482–487

965 Panfili G, Fratianni A, Distaam M (2004) Improved normal-phase high-performance
 966 liquid chromatography procedure for the determination of carotenoids in cereals. *J*
 967 *Agric Food Chem* 52:6373–6377

968 Panfili G, Fratianni A, Irano M (2005) Change in tocochromanol and carotenoid content
 969 during technological processes of cereals. *Tecnica Molitoria* 5:493–498

970 Paterson AH, Bowers JE, Bruggmann R et al (2009) The sorghum bicolor genome and
 971 the diversification of grasses. *Nature* 457:551–556

972 Patil RM, Oak MD, Tamhankar SA et al (2008) Mapping and validation of a major
 973 QTL for yellow pigment content on 7AL in durum wheat (*Triticum turgidum* L.
 974 ssp. *durum*). Mol Breeding 21:485–496
 975 Pinzino C, Nanni B, Zandomenighi M (1999) Aging, free radicals, and antioxidants in
 976 wheat seeds. J Agric Food Chem 47:1333–1339
 977 Pozniak CJ, Knox RE, Clarke FR et al (2007) Identification of QTL and association of a
 978 phytoene synthase gene with endosperm colour in durum wheat. Theor Appl Genet
 979 114:525–537
 980 Quackenbush FW (1963) Corn carotenoids: effects of temperature and moisture on
 981 losses during storage. Cereal Chem 40:266–269
 982 Quackenbush F, Firch J, Rabourn W et al (1961) Composition of corn. Analysis of
 983 carotenoids in corn grain. J Agric Food Chem 9:132–135
 984 Ravel C, Dardevet M, Leenhardt F et al (2013) Improving the yellow pigment content
 985 of bread wheat flour by selecting the three homoeologous copies of Psy1. Mol
 986 Breeding 31:87–99
 987 Rodríguez-Amaya DB (1997) Carotenoids and Food Preparation: The Retention of
 988 Provitamin A Carotenoids in Prepared, Processed, and Stored Foods. Opportunities
 989 for Micronutrient Intervention (OMNI), Arlington, USA.
 990 Rodríguez-Amaya DB (2003) Food carotenoids: analysis, composition and alterations
 991 during storage and processing of foods. Forum Nutr 56:35–37
 992 Rodríguez-Suárez C, Atienza SG (2012) *Hordeum chilense* genome, a useful tool to
 993 investigate the endosperm yellow pigment content in the *Triticeae*. BMC Plant Biol
 994 12:200.

995 Rodríguez-Suárez C, Giménez MJ, Gutiérrez N et al (2012) Development of wild barley
 996 (*Hordeum chilense*)-derived DArT markers and their use into genetic and physical
 997 mapping. *Theor Appl Genet* 124:713–722

998 Rodríguez-Suárez C, Giménez MJ, Ramírez MC et al (2011) Exploitation of nuclear
 999 and cytoplasm variability in *Hordeum chilense* for wheat breeding. *Plant Genet*
 1000 *Resour* 9:313–316

1001 Salas-Fernandez MG, Hamblin MT, Li L et al (2008) Quantitative trait loci analysis of
 1002 endosperm color and carotenoid content in sorghum grain. *Crop Sci* 48:1732–1743

1003 Santra M, Rao VS, Tamhankar SA (2003) Modification of AACC procedure for
 1004 measuring β -carotene in early generation durum wheat. *Cereal Chem* 80:130–131

1005 Santra M, Santra DK, Rao VS et al (2005) Inheritance of β -carotene concentration in
 1006 durum wheat (*Triticum turgidum* L. ssp. *durum*). *Euphytica* 144:215–221

1007 Saxena A, Maity T, Raju PS et al (2012) Degradation kinetics of colour and total
 1008 carotenoids in jackfruit (*Artocarpus heterophyllus*) bulb slices during hot air
 1009 drying. *Food Bioprocess Tech* 5:672–679

1010 Scott C, Eldridge A (2005) Comparison of carotenoid content in fresh, frozen and
 1011 canned corn. *J Food Comp Anal* 18:551–559

1012 Selim K, Tsimidou M, Biliaderis CG (2000) Kinetic studies of degradation of saffron
 1013 carotenoids encapsulated in amorphous polymer matrices. *Food Chem* 71:199–206

1014 Sellappan K, Datta K, Parkhi V et al (2009) Rice caryopsis structure in relation to
 1015 distribution of micronutrients (iron, zinc, β -carotene) of rice cultivars including
 1016 transgenic indica rice. *Plant Sci* 177:557–562

1017 Selman JD (1994) Vitamin retention during blanching of vegetables. *Food Chem*
 1018 49:137–147

1019 Serpen A, Gökmen V, Karagöz A et al (2008) Phytochemical quantification and total
 1020 antioxidant capacities of Emmer (*Triticum dicoccon* Schrank) and Einkorn
 1021 (*Triticum monococcum* L.) wheat landraces. J Agric Food Chem 56:7285–7292
 1022 Sharma SK, Le Maguer M (1996) Kinetics of lycopene degradation in tomato pulp
 1023 solids under different processing and storage conditions. Food Res Int 29:309–315
 1024 Siebenhandl S, Grausgruber H, Pellegrini N et al (2007) Phytochemical profile of main
 1025 antioxidants in different factions of purple and blue wheat, and black barley. J
 1026 Agric Food Chem 55:8541–8547
 1027 Singh S, Gamlath S, Wakeling L (2007) Nutritional aspects of food extrusion: a review.
 1028 Int J Food Sci Tech 42:916–929
 1029 Suryanarayana Rao K, Rukmini C, Mohan VS (1968) Carotene content of some yellow-
 1030 endosperm varieties of sorghum. Indian J Agr Sci 38:368–372
 1031 Tan J, Baisakh N, Oliva N et al (2005) The screening of rice germplasm, including
 1032 those transgenic rice lines which accumulate β -carotene in their polished seeds, for
 1033 their carotenoid profile. Int J Food Sci Technol 40:563–569
 1034 Tonon RV, Baroni AF, Hubinger MD (2007) Osmotic dehydration of tomato in ternary
 1035 solutions: influence of process variables on mass transfer kinetics and an evaluation
 1036 of the retention of carotenoids. J Food Eng 82:509–517
 1037 Trono D, Pastore D, Di-Fonzo N (1999) Carotenoid dependent inhibition of durum
 1038 wheat lipoxygenase. J Cereal Sci 29:99–102
 1039 Tsimidou M (1997) Kinetic studies of saffron (*Crocus sativus* L.) quality deterioration.
 1040 J Agric Food Chem 45:2890–2898
 1041 Van Hung P, Hatcher DW (2011) Ultra-performance liquid chromatography (UPLC)
 1042 quantification of carotenoids in durum-wheat: influence of genotype and

1043 environment in relation to the colour of yellow alkaline noodles (YAN). Food
 1044 Chem 125:1510–1516
 1045 Velioglu YS, Mazza G, Gao L et al (1998) Antioxidant activity and total phenolics in
 1046 selected fruits, vegetables, and grain products. J Agric Food Chem 46:4113–4117
 1047 Wagner LA, Warthesen JJ (1995) Stability of spray-dried encapsulated carrot carotenes.
 1048 J Food Sci 60:1048–1053
 1049 Weber EJ (1987) Carotenoids and tocopherols of corn grain determined by HPLC. J Am Oil
 1050 Chem Soc 8:1129–1134
 1051 Wong JC, Lambert RJ, Wurtzel ET et al (2004) QTL and candidate genes phytoene
 1052 synthase and ζ -carotene desaturase associated with the accumulation of carotenoids
 1053 in maize. Theor Appl Genet 108:349–359
 1054 Wurtzel ET, Cuttriss A, Vallabhaneni R (2012) Maize provitamin A carotenoids,
 1055 current resources, and future metabolic engineering challenges. Front Plant Sci 3:29
 1056 Ye X, Al-Babili S, Klöti A, Zhang J, Lucca P, Beyer P, Potrykus I (2000) Engineering
 1057 the provitamin A (beta-carotene) biosynthetic pathway into (carotenoid-free) rice
 1058 endosperm. Science 287:303–305
 1059 Zepka LQ, Borsarelli CD, Azevedo MA et al (2009) Thermal degradation kinetics of
 1060 carotenoids in a cashew apple juice model and its impact on the system color. J
 1061 Agric Food Chem 57:7841–7845
 1062 Zhang W, Lukaszewski AJ, Kolmer J et al (2005) Molecular characterization of durum
 1063 and common wheat recombinant lines carrying leaf rust resistance (Lr19) and
 1064 yellow pigment (Y) genes from *Lophopyrum ponticum*. Theor Appl Genet
 1065 111:573–582
 1066 Zhou K, Yu L (2004a) Antioxidant properties of bran extracts from Trego wheat grown
 1067 at different locations. J Agric Food Chem 52:1112–1117

1068 Zhou K, Yu L (2004b) Effects of extraction solvent on wheat bran antioxidant activity
 1069 estimation. *Lebensm Wiss Technol* 37:717–721
 1070 Zhou K, Laux JJ, Yu L (2004) Comparison of swiss red wheat grain fractions for their
 1071 antioxidant properties. *J Agric Food Chem* 52:1118–1123
 1072 Zhou K, Su L, Yu L (2004) Phytochemicals and antioxidant properties in wheat bran. *J*
 1073 *Agric Food Chem* 52:6108–6114
 1074 Zhu C, Naqvi S, Breitenbach J et al (2008) Combinatorial genetic transformation
 1075 generates a library of metabolic phenotypes for the carotenoid pathway in maize. *P*
 1076 *Natl Acad Sci USA* 105:18232–18237
 1077 Žilić S, Serpen A, Akillioğlu G et al (2012) Distribution of phenolic compounds, yellow
 1078 pigments and oxidative enzymes in wheat grains and their relation to antioxidant
 1079 capacity of bran and debranned flour. *J Cereal Sci* 56:562–568
 1080

all-trans-Lutein (3,3'-diol- β , ϵ -carotene)

all-trans-Zeaxanthin (3,3'-diol- β , β -carotene)

all-trans- β -Cryptoxanthin (3-ol- β , β -carotene)

all-trans- β -Carotene (β , β -carotene)

all-trans- α -Carotene (β , ϵ -carotene)

Figure 1. Chemical structures of the most common carotenoid pigment (carotenes and xanthophylls) present in cereal grains.

Figure 2. HPLC chromatograms corresponding to direct carotenoid extracts obtained from tritordeum and durum wheat. Peaks identity: 1. all-*trans*-zeaxanthin; 2. all-*trans*-lutein; 3. 9-*cis*-lutein; 4. 13-*cis*-lutein; 5. lutein monolinoleate; 6. lutein monopalmitate; 7. all-*trans*- α -carotene; 8. all-*trans*- β -carotene; 9. lutein dilinoleate; 10. lutein linoleatopalmitate; 11. lutein dipalmitate.

Lutein-3-*O*-palmitate

Lutein-3'-*O*-palmitate

Lutein-3-*O*-linoleate

Lutein-3'-*O*-linoleate

Lutein dipalmitate

Lutein dilinoleate

Lutein-3'-*O*-linoleate-3-*O*-palmitate

Lutein-3'-*O*-palmitate-3-*O*-linoleate

Figure 3. Chemical structure of the monoesters and diesters, including the regioisomers, of lutein with fatty acids identified in grains of tritordeum (*×Tritordeum* Ascherson et Graebner).