**Genetic and Functional Analysis of Biogenic Amine Production Capacity Among
Starter and Non-Starter Lactic Acid Bacteria Isolated from Artisanal Cheeses**

**Victor Ladero^{1*}, María Cruz Martín, Begoña Redruello, Baltasar Mayo, Ana
Belén Flórez, María Fernández, and Miguel A. Alvarez**

Instituto de Productos Lácteos de Asturias (IPLA-CSIC), Villaviciosa, Spain.

Short title: BA production capacity among dairy LAB

***Corresponding author: Victor Ladero**

**Mailing address: Instituto de Productos Lácteos de Asturias (IPLA-CSIC), Paseo
del Río Linares s/n, 33300 Villaviciosa, Spain.**

Phone: +34 985 89 21 31

Fax: +34 985 89 22 32

E-mail: ladero@ipla.csic.es

Abstract

This work reports the capacity of 137 strains of starter and non-starter LAB belonging to nine species of the genera *Lactobacillus*, *Lactococcus*, *Streptococcus* and *Leuconostoc* (all isolated from artisanal cheeses) to produce histamine, tyramine, putrescine and β -phenylethylamine, the biogenic amines (BA) most commonly found in dairy products. Production assays were performed in liquid media supplemented with the appropriate precursor amino acid; culture supernatants were then tested for BA by (U)HPLC. In addition, the presence of key genes involved in the biosynthetic pathways of the target BA, including the production of putrescine via the agmatine deiminase pathway, was assessed by PCR. Twenty strains were shown to have genes involved in the synthesis of BA; these belonged to the species *Lactobacillus brevis* (4), *Lactobacillus curvatus* (3), *Lactococcus lactis* (11) and *Streptococcus thermophilus* (2). With the exception of the two *S. thermophilus* strains, all those possessing genes involved in BA production synthesized the corresponding compound. Remarkably, all the putrescine-producing strains used the agmatine deiminase pathway. Four *Lb. brevis* and two *Lb. curvatus* strains were found able to produce both tyramine and putrescine. There is increasing interest in the use of autochthonous LAB strains in starter and adjunct cultures for producing dairy products with ‘particular geographic indication’ status. Such strains should not produce BA; the present results show that BA-production capacity should be checked by (U)HPLC and PCR.

Key words: Biogenic amines, tyramine, putrescine, tyrosine decarboxylase, agmatine deiminase.

Introduction

Lactic acid bacteria (LAB) play an essential role in the production of fermented dairy products, with *Lactococcus lactis* and *Streptococcus thermophilus* being the species most commonly used as primary fermentation starters [1]. Their major function is the rapid production of lactic acid from lactose, resulting in a lowering of the pH.

The so-called non-starter lactic acid bacteria (NSLAB) participate in the development of the final organoleptic properties of fermented dairy products [2]. NSLAB may be present in the milk itself, be part of the flora of dairy facilities, or be added to fermentations as adjunct cultures [3]. These bacteria are frequently facultative, heterofermentative lactobacilli belonging to the species *Lactobacillus casei/paracasei*, *Lactobacillus plantarum* or *Lactobacillus curvatus* [4,5]. *Leuconostoc* may be involved in the development of aroma components [6]. There is increasing interest in the characterization and use of NSLAB from artisanal products for use in tailored cultures to be employed in the manufacture of dairy products with ‘protected geographic indication’ (PGI) status. Their use would help maintain their typical organoleptic characteristics [6-9].

The long and safe history of the use of LAB in dairy products has resulted in the assignment of Qualified Presumption of Safety (QPS) status (awarded by the European Food Safety Authority [EFSA]) to the majority of LAB. However, some properties and enzymatic activities can generate undesirable flavours [10] or even toxic compounds such as biogenic amines (BA) [11], the presence of which should be avoided in dairy products.

BA are low molecular weight nitrogenous compounds formed by the decarboxylation of certain amino acids that may be present in foods. The consumption of foods with high

BA concentrations may cause intoxications manifested as headache, nausea or vomiting, alterations in blood pressure, rashes, etc. [12]. Cheese is the fermented food most commonly associated with BA poisoning; indeed, the term *cheese reaction* was coined to refer to it [13]. Tyramine, putrescine and histamine are the most commonly encountered and abundant BA; in cheese [11,14,15]. Certainly, cheese provides an ideal matrix for the production and accumulation of BA since the amino acid substrates required are made easily available by casein proteolysis, and the low pH favours decarboxylase gene transcription and enzyme activity [11]. Further, cheese naturally contains milk-derived Gram positive LAB, generally of the genera *Lactobacillus* and *Enterococcus*, which possess decarboxylating activity [11,16]. BA-producing strains have also been described among the species most commonly used as dairy starters, such as *Lactococcus lactis*, *Streptococcus thermophilus* and *Lactobacillus delbrueckii* [17-19]. BA producers may also enter dairy products via contamination [20,21].

The selection of starter strains with no BA-producing capacity would be a good starting point for reducing BA accumulation in dairy products [22]. Different methods have been devised for assessing the capacity of LAB to produce BA, including the use of differential media and pH indicators [23]. Unfortunately, the strong acidification of the medium occasioned by harmless LAB can result in false negatives. Moreover, these methods target the presence of amino acid decarboxylases and do not test the presence of deamination routes involved in the production of some BA such as putrescine [11].

Analytical methods that directly detect BA compounds in culture supernatants after incubation with an amino acid precursor have also been commonly used [24,25]. However, culture-independent methods based on PCR techniques, aimed to detect the genetic determinants involved in the synthesis of BA, are now regarded as the most suitable for screening collections of isolates [26]. Agreement between the results

obtained by analytical and molecular methods strengthens the case for the use of the latter [27,28].

In the present work, U(H)PLC and PCR methods were used to examine the capacity of 137 LAB strains (four genera, nine species), isolated from artisanal cheeses, and all with potential for use in dairy starter or adjunct cultures designed for the production of artisanal cheeses with PGI status, to produce histamine, tyramine, putrescine and β -phenylethylamine.

2. Materials and Methods

Bacterial strains

One hundred and thirty seven strains isolated from different artisanal cheeses [29,30], identified by comparison of partial 16S rRNA gene sequences, and belonging to four different genera - *Lactococcus*, *Streptococcus*, *Leuconostoc*, and *Lactobacillus* - were assessed for their capacity to produce BA (Table 1). *Lactococcus lactis*, *Streptococcus thermophilus* and *Leuconostoc mesenteroides* strains were grown statically in M17 (Oxoid) supplemented with 0.5% glucose and 0.5% lactose (w/v) at either 30°C (*Lc. lactis*, *Le. mesenteroides*) or 37°C (*S. thermophilus* strains). All *Lactobacillus* strains, which belonged to six species (Table 1), were grown statically in MRS (Oxoid) at 30°C, except those belonging to *Lb. delbrueckii* which were grown at 37°C.

***In vivo* BA-production capacity**

BA production was assessed in triplicate in culture supernatants of the LAB strains grown for 24 h in 10 ml M17 or MRS broth supplemented with 1 mM tyrosine (M17/MRS-T), 1 mM histidine (M17/MRS-H), 1 mM ornithine (M17/MRS-O) or 1 mM agmatine (M17/MRS-A). Both ornithine and agmatine are precursors of putrescine,

although via different pathways. Tyramine, histamine and putrescine detection was performed as previously described [31] after the centrifugation of the cultures (2000 x g for 15 min) and filtering of the supernatant through a 0.2 µm pore diameter membrane (Pall, USA), followed by derivatization of 100 µl with diethyl ethoxymethylene malonate. Derivatized samples were analyzed by (U)HPLC in a Waters H-Class Acquity UPLC apparatus with a UV detector (Waters, USA) controlled by Empower 2.0 software (Waters), under the conditions described by Redruello et al. [32].

Detection of BA-producing genes

The presence of the tyrosine decarboxylase gene *tdcA*, the histidine decarboxylase gene *hdcA*, the ornithine decarboxylase gene *odc*, and the *aguA* and *aguD* genes from the agmatine deiminase cluster (AgdI), was checked by PCR using the primer pairs P2-for and P1-rev [33], JV16HC and JV17HC [34], ODC3 and ODC16 [35], and Seq1 and Seq2 [17], respectively. The PCR conditions were those described in [33-35,17], respectively and were performed in a MyCycler™ thermal cycler (Bio-Rad, Spain) using DreamTaq polymerase (Fermentas, Lithuania). Total DNA from the strains was obtained as previously described [36] and used as a template in PCR. Total DNA from the tyramine- and putrescine-producing strain *Enterococcus faecalis* V583 [27], from the ODC⁺ strain *Lactobacillus saerimneri* 30A [37], and from the histamine-producer *Lactobacillus buchneri* B301 [38], were used to provide positive controls.

PCR products were separated in 0.8% (w/v) agarose gels in 1XTAE buffer, and visualized after staining with ethidium bromide using a GelDoc 2000 system (Bio-Rad, Hercules, USA). The Gene Ruler DNA ladder mix (Fermentas, Lithuania) was used as molecular weight marker.

Compliance with Ethical Requirements

This study does not involve animal or human subjects.

Results and Discussion

The selection of starter strains with no BA-producing capacity is an important step towards reducing the presence of these toxins in dairy products [22]. In this work, 137 LAB strains, previously isolated from artisanal cheeses made from raw milk, were evaluated for their BA-producing capability.

Twenty (14.69%) of the 137 examined strains were found to possess genes involved in BA production, including 4 strains of *Lb. brevis*, 3 of *Lb. curvatus*, 11 of *Lc. lactis* (8 belonging to *Lc. lactis* subsp. *lactis* and 3 to *Lc. lactis* subsp. *cremoris*), and 2 of *S. thermophilus* (Table 1 and Fig. 1).

Eighteen (13.1%) of the tested strains showed the capacity to produce at least one BA in a supplemented medium. These corresponded to all the strains in which the presence of genes involved in BA production were detected by PCR, except for the two *S. thermophilus* strains (see below). Six strains (4.4%), four *Lb. brevis* and two *Lb. curvatus* strains, produced both tyramine and putrescine.

Similar percentages of BA-producing strains have been reported by other authors [19,39]. During their analysis of dairy isolates, Bunkova et al. [19] found 20% of the strains tested to produce tyramine and to be positive for the *tdcA* gene. In some studies, higher percentages of BA-producers have been reported [39,40], but in most of these investigations strains of *Enterococcus* were analyzed. The capacity to produce BA is widespread among enterococci and has been shown as a species-specific trait in some enterococcal species [27,41]; thus increasing the occurrence of BA producing strains. It was, therefore decided not to include enterococcal strains in the present work.

All of the strains that gave a positive PCR result for the presence of genes involved in BA production were able to synthesize the corresponding BA (Table 1), except for two strains of *S. thermophilus*. Both of these strains possessed the histidine decarboxylase gene *hdcA*, but no histamine was found in the culture supernatant after 24 or even after 48 h of culture in M17-H. This might be due to a non-functionality of the HDC cluster or because the conditions assayed were not optimal for histamine production in these strains since the production of BA can be affected by different cultures conditions [11]. Certainly, some authors report that all *S. thermophilus* strains with the capacity to produce histamine from histidine produce small amounts of histamine in broth but not in milk etc. [18,42]. In any event, the present work highlights a good correlation between the results of molecular and functional analysis of BA-producing capacity. All the BA-producing strains returned positive PCR results, indicating that this culture-independent technique is suitable for assessing this property in potential LAB starter strains [28].

Even though two *S. thermophilus* strains were negative for the *in vivo* production of BA, their possession of genes involved in BA production must be seen as a risk. These genes could be horizontally transferred to other LAB present in the culture or dairy product [43-45], conferring the ability to produce histamine upon them.

Of the 137 strains tested, seven produced tyramine from tyrosine in broth, and were positive for *tdcA* in PCR tests (Table 1). All these strains belonged to *Lb. brevis* or *Lb. curvatus*. Tyramine-producing strains of these species have been isolated from cheeses by other authors [44,46,47]. In *Lb. brevis*, tyramine production has been described as a strain level trait - perhaps horizontally acquired [44,48]. For *Lb. curvatus* there are insufficient data to confirm whether it is a species- or strain-dependent trait. The majority of *Lb. curvatus* strains isolated from meat, however, have been reported

tyramine producers [49-51]. All the present tyramine producers, independent of their species, were 'strong tyramine producers' (Table 2). *Lb. curvatus* strains have been described as strong tyramine producers by other authors [47,47], showing high conversion rates in broth media supplemented with tyrosine. *Lb. brevis* has also been described as a strong tyramine producer, although different media and conditions were assayed and variation in tyramine production capacity was observed [52].

None of the tested strains was able to produce β -phenylethylamine under the present assay conditions. No specific phenylalanine decarboxylases have been described, but several authors have reported that certain tyrosine decarboxylases can use phenylalanine as an alternative substrate, converting it into β -phenylethylamine [53]. In the present work, only the *E. faecalis* positive control was able to produce β -phenylethylamine in medium supplemented with tyrosine (data not shown).

Putrescine is produced from arginine via a decarboxylation and a deimination reaction [11,54]. However, the order of these reactions can differ, and, depending on that order, two different pathways are recognized: (i) the ornithine decarboxylation pathway (ODC) (in which arginine is first deiminated to form ornithine, which is then decarboxylated to form putrescine), and (ii) the agmatine deimination pathway (AGDI) (in which arginine is first decarboxylated to form agmatine, which is then deiminated to form putrescine) [11,54]. To distinguish which pathway was being used, the tested strains were grown in media supplemented with ornithine or agmatine. No strain produced putrescine from ornithine. Although the ODC pathway has been described in several LAB strains, including strains of *Lb. brevis* [25,43], it is not a pathway commonly used by dairy bacteria [11,54,55]. Thus, the lack of strains with ODC pathway capacity among those tested in the present work was expected. Seventeen strains of the 137 examined were, however, able to produce putrescine from agmatine

(Table 1). Putrescine is the most commonly found BA in dairy products [14]. It is not surprising, therefore, that the largest number of BA-producing strains detected should be putrescine producers. It is important to highlight that all the putrescine producers detected in the present survey have the AGDI and not the ODC pathway. Although the prevalence of the AGDI pathway in dairy strains has been previously suggested [11], a test to determine the presence of the AGDI pathway is not usually done. In fact, as far as we are aware, this is the first study to include screening for the AGI pathway when testing for BA-producing capacity in dairy LAB.

The production of putrescine via the AGDI pathway has, however, been described in *Lb. brevis* of non-dairy origin by other authors [48,56]. All the present strains of *Lb. brevis* shown to be putrescine producers were also tyramine producers. It has been suggested that, in this species, the AGDI genetic determinants are linked to those of the TDC pathway, producing a *locus* of acid resistance mechanisms probably acquired by horizontal gene transfer [48,43]. Two of the three *Lb. curvatus* strains tested produced putrescine from agmatine and also returned positive PCR results (Table 1), both strains were also able to produce tyramine. As in *Lb. brevis*, these two BA-producing capacities have been related to acid resistance. The corresponding genes have been described as lying adjacent to one another in the chromosome of some dairy isolates of *Lb. curvatus* [43].

Among the *Lc. lactis* strains tested, i.e., of both subspecies *lactis* and *cremoris*, 11 were shown to produce putrescine from agmatine. Such a capacity had already been reported for some *Lc. lactis* strains [17], and putrescine-producing *Lc. lactis* can be found in large numbers in cheeses with high putrescine concentrations [55]. Not all the *Lc. lactis* strains tested were able to produce putrescine, although the capacity to produce it from agmatine has been described as a species level trait [17]. Traditionally, BA-producing

pathways have been thought horizontally acquired [44,48]. The present *Lc. lactis* strains unable to synthesize putrescine may have lost this capacity during their use in the dairy environment: putrescine would negatively affect acidification and/or final flavour and such putrescine-producing strains would have been avoided [17]. The loss of this capacity seems to have occurred in two ways: (i) via the loss of the AGDI pathway genes, as has been shown for strains of *Lc. lactis* subsp. *cremoris*, and (ii) the inactivation of the cluster by an insertion element (IS) in *Lc. lactis* subsp. *lactis* strains [17]. To differentiate between putrescine and non-putrescine producers, a specific PCR test is available [17] in which non-putrescine-producing *Lc. lactis* subsp. *cremoris* strains do not render a PCR band, while *Lc. lactis* subsp. *lactis* non-putrescine-producing strains do, although the amplification product is 1 kb larger than expected due to the presence of an IS element. In the present work, none of the non-putrescine-producing strains of *Lc. lactis* subsp. *cremoris* was associated with any positive amplification, while those of *Lc. lactis* subsp. *lactis* rendered the expected enlarged amplicon (Fig. 1).

Variation in the efficiency of putrescine production was observed among the producing strains of *Lc. lactis*; this allowed their classification as 'strong' or 'medium putrescine producers' (Table 2). Variation in the capacity to produce putrescine from agmatine has been described before among *Lc. lactis* subsp. *cremoris* strains [31]. In the present work, however, the greatest variation was observed among the *Lc. lactis* subsp. *lactis* producers (Table 2).

One of the most effective measures for reducing the presence of BA in dairy products is the use of starter cultures that have been properly tested and selected as non-BA-producers [22]. The present results show that both culture-dependent and culture-independent methods are appropriate for ruling out BA-producing strains for use as

starters and adjunct cultures. The culture-independent methods based on PCR testing not only detected BA producers but also non-producer strains that possessed genes involved in BA production; these pose a risk since they might be spread by horizontal gene transfer.

There is increasing interest in the use of autochthonous LAB strains in starter and adjunct cultures for producing dairy products with PGI status. [8,57]. Strains intended for use in their manufacture should be systematically monitored for BA production capacity to avoid the accumulation of these toxins and thus produce safer dairy products.

Conclusions

This work shows that some of the strains belonging to the species most frequently used in the manufacture of dairy products can produce BA, highlighting the importance of adequately selecting indigenous strains for inclusion in starter and adjunct cultures. The prevalence of putrescine-producing strains (which use the AGDI pathway) is noteworthy. The literature contains little on this, even though putrescine is one of the commonest BA in dairy products and the AGDI pathway is the main route of its synthesis. Tests for the presence of the AGDI pathway should be included when examining the BA production capability in dairy strains. As shown in this work, the capacity to produce BA can be tested by either chromatographic or molecular methods, although PCR testing affords many advantages.

Acknowledgments

This work was funded by the Ministry of Economy and Competitiveness, Spain (AGL2013-45431-R), the Spanish National Research Council (CSIC201270E144), and the INIA (RM2011-00005-00-00). The authors thank Adrian Burton for language assistance.

Conflict of interest

None.

REFERENCES

1. Parente E, Cogan TM (2004) Starter cultures: general aspects. In: Fox PF, McSweeney PLH, Cogan TM, Guinee TP (eds) Cheese: Chemistry, Physics and Microbiology, vol 1. 3rd edn. Elsevier, Oxford, UK, pp 123-147
2. Poveda JM, Cabezas L, McSweeney PLH (2004) Free amino acid content of Manchego cheese manufactured with different starter cultures and changes throughout ripening. Food Chem 84 (2):213-218.
3. Coeuret V, Dubernet S, Bernardeau M, Gueguen M, Vernoux JP (2003) Isolation, characterisation and identification of lactobacilli focusing mainly on cheeses and other dairy products. Lait 83 (4):269-306.
4. Antonsson M, Molin G, Ardo Y (2003) *Lactobacillus* strains isolated from Danbo cheese as adjunct cultures in a cheese model system. Int J Food Microbiol 85 (1-2):159-169.
5. Pisano MB, Patrignani F, Cosentino S, Guerzoni ME, Franz CMAP, Holzapfel WH (2011) Diversity and functional properties of *Lactobacillus plantarum*-group strains isolated from Italian cheese products. Dairy Sci Technol 91 (1):65-76.

6. Nieto-Arribas P, Sesena S, Poveda JM, Palop L, Cabezas L (2010) Genotypic and technological characterization of *Leuconostoc* isolates to be used as adjunct starters in Manchego cheese manufacture. Food Microbiol 27 (1):85-93.
7. Nieto-Arribas P, Poveda JM, Sesena S, Palop L, Cabezas L (2009) Technological characterization of *Lactobacillus* isolates from traditional Manchego cheese for potential use as adjunct starter cultures. Food Control 20 (12):1092-1098.
8. Capozzi V, Spano G (2011) Food microbial biodiversity and "microbes of protected origin". Front Microbiol 2:237.
Doi 10.3389/Fmicb.2011.00237
9. Florez AB, Lopez-Diaz TM, Alvarez-Martin P, Mayo B (2006) Microbial characterisation of the traditional Spanish blue-veined Cabrales cheese: identification of dominant lactic acid bacteria. Eur Food Res Technol 223 (4):503-508.
10. Smit G, Smit BA, Engels WJM (2005) Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. FEMS Microbiol Rev 29 (3):591-610.
11. Linares DM, Del Rio B, Ladero V, Martinez N, Fernandez M, Martin MC, Alvarez MA (2012) Factors influencing biogenic amines accumulation in dairy products. Front Microbiol 3:180.
12. Ladero V, Calles-Enríquez M, Fernández M, Alvarez MA (2010) Toxicological effects of dietary biogenic amines. Curr Nutr Food Sci 6 (2):145-156.
13. ten Brink B, Damink C, Joosten HMLJ, Tveld JHJHI (1990) Occurrence and formation of biologically-active amines in foods. Int J Food Microbiol 11 (1):73-84.
14. Fernandez M, Linares DM, del Rio B, Ladero V, Alvarez MA (2007) HPLC quantification of biogenic amines in cheeses: correlation with PCR-detection of tyramine-producing microorganisms. J Dairy Res 74 (3):276-282.

15. Spano G, Russo P, Lonvaud-Funel A, Lucas P, Alexandre H, Grandvalet C, Coton E, Coton M, Barnavon L, Bach B, Rattray F, Bunte A, Magni C, Ladero V, Alvarez M, Fernandez M, Lopez P, de Palencia PF, Corbi A, Trip H, Lolkema JS (2010) Biogenic amines in fermented foods. *Eur J Clin Nutr* 64:S95-S100.
16. Ladero V, Fernandez M, Cuesta I, Alvarez MA (2010) Quantitative detection and identification of tyramine-producing enterococci and lactobacilli in cheese by multiplex qPCR. *Food Microbiol* 27 (7):933-939.
17. Ladero V, Rattray FP, Mayo B, Martin MC, Fernandez M, Alvarez MA (2011) Sequencing and transcriptional analysis of the biosynthesis gene cluster of putrescine-producing *Lactococcus lactis*. *Appl Environ Microbiol* 77 (18):6409-6418.
18. Calles-Enriquez M, Eriksen BH, Andersen PS, Rattray FP, Johansen AH, Fernandez M, Ladero V, Alvarez MA (2010) Sequencing and transcriptional analysis of the *Streptococcus thermophilus* histamine biosynthesis gene cluster: Factors that affect differential *hdcA* expression. *Appl Environ Microbiol* 76 (18):6231-6238.
19. Bunkova L, Bunka F, Hlobilova M, Vanatkova Z, Novakova D, Drab V (2009) Tyramine production of technological important strains of *Lactobacillus*, *Lactococcus* and *Streptococcus*. *Eur Food Res Technol* 229 (3):533-538.
20. Ladero V, Fernandez M, Alvarez MA (2009) Effect of post-ripening processing on the histamine and histamine-producing bacteria contents of different cheeses. *Int Dairy J* 19 (12):759-762.
21. Novella-Rodriguez S, Veciana-Nogues MT, Roig-Sagues AX, Trujillo-Mesa AJ, Vidal-Carou MC (2002) Influence of starter and nonstarter on the formation of biogenic amine in goat cheese during ripening. *J Dairy Sci* 85 (10):2471-2478.

22. EFSA (2011) Scientific Opinion on risk based control of biogenic amine formation in fermented foods. EFSA Panel on Biological Hazards (BIOHAZ). EFSA J 9(10):2393-2486.
23. Bover-Cid S, Holzapfel WH (1999) Improved screening procedure for biogenic amine production by lactic acid bacteria. Int J Food Microbiol 53 (1):33-41.
24. Garcia-Moruno E, Carrascosa AV, Munoz R (2005) A rapid and inexpensive method for the determination of biogenic amines from bacterial cultures by thin-layer chromatography. J Food Protec 68 (3):625-629.
25. Coton M, Romano A, Spano G, Ziegler K, Vetrana C, Desmarais C, Lonvaud-Funel A, Lucas P, Coton E (2010) Occurrence of biogenic amine-forming lactic acid bacteria in wine and cider. Food Microbiol 27 (8):1078-1085.
26. Landete JM, de las Rivas B, Marcobal A, Munoz R (2007) Molecular methods for the detection of biogenic amine-producing bacteria on foods. Int J Food Microbiol 117 (3):258-269.
27. Ladero V, Fernandez M, Calles-Enriquez M, Sanchez-Llana E, Canedo E, Martin MC, Alvarez MA (2012) Is the production of the biogenic amines tyramine and putrescine a species-level trait in enterococci? Food Microbiol 30 (1):132-138.
28. Bhardwaj A, Gupta H, Iyer R, Kumar N, Malik RK (2009) Tyramine-producing enterococci are equally detected on tyramine production medium, by quantification of tyramine by HPLC, or by *tdc* gene-targeted PCR. Dairy Sci Technol 89 (6):601-611.
29. Lopez S, Mayo B (1997) Identification and characterization of homofermentative mesophilic *Lactobacillus* strains isolated from artisan starter-free cheeses. Lett Appl Microbiol 25 (4):233-238.

30. Estepar J, Sanchez MD, Alonso L, Mayo B (1999) Biochemical and microbiological characterization of artisanal 'Penamellera' cheese: analysis of its indigenous lactic acid bacteria. *Int Dairy J* 9 (10):737-746.
31. Linares DM, del Rio B, Ladero V, Redruello B, Martin MC, Fernandez M, Alvarez MA (2013) The putrescine biosynthesis pathway in *Lactococcus lactis* is transcriptionally regulated by carbon catabolic repression, mediated by CcpA. *Int J Food Microbiol* 165 (1):43-50.
32. Redruello B, Ladero V, Cuesta I, Alvarez-Buylla JR, Martin MC, Fernandez M, Alvarez MA (2013) A fast, reliable, ultra high performance liquid chromatography method for the simultaneous determination of amino acids, biogenic amines and ammonium ions in cheese, using diethyl ethoxymethylenemalonate as a derivatising agent. *Food Chem* 139 (1-4):1029-1035.
33. Lucas P, Lonvaud-Funel A (2002) Purification and partial gene sequence of the tyrosine decarboxylase of *Lactobacillus brevis* IOEB 9809. *FEMS Microbiol Lett* 211 (1):85-89.
34. Le Jeune C, Lonvaud-Funel A, ten Brink B, Hofstra H, van der Vossen JM (1995) Development of a detection system for histidine decarboxylating lactic acid bacteria based on DNA probes, PCR and activity test. *J Appl Bacteriol* 78 (3):316-326.
35. Marcobal A, De las Rivas B, Moreno-Arribas MV, Munoz R (2005) Multiplex PCR method for the simultaneous detection of histamine-, tyramine-, and putrescine-producing lactic acid bacteria in foods. *J Food Protec* 68 (4):874-878.
36. Ruiz-Barba JL, Maldonado A, Jimenez-Diaz R (2005) Small-scale total DNA extraction from bacteria and yeast for PCR applications. *Anal Biochem* 347 (2):333-335.

37. Romano A, Trip H, Campbell-Sills H, Bouchez O, Sherman D, Lolkema JS, Lucas PM (2013) Genome sequence of *Lactobacillus saerimneri* 30a (Formerly *Lactobacillus* sp. Strain 30a), a reference lactic acid bacterium strain producing biogenic amines. *Genome announce* 1 (1): e00097-12.
38. Martin MC, Fernandez M, Linares DM, Alvarez MA (2005) Sequencing, characterization and transcriptional analysis of the histidine decarboxylase operon of *Lactobacillus buchneri*. *Microbiol* 151:1219-1228.
39. Pircher A, Bauer F, Paulsen P (2007) Formation of cadaverine, histamine, putrescine and tyramine by bacteria isolated from meat, fermented sausages and cheeses. *Eur Food Res Technol* 226 (1-2):225-231.
40. Lorencova E, Bunkova L, Matoulkova D, Drab V, Pleva P, Kuban V, Bunka F (2012) Production of biogenic amines by lactic acid bacteria and bifidobacteria isolated from dairy products and beer. *Int J Food Sci Technol* 47 (10):2086-2091.
41. Capozzi V, Ladero V, Beneduce L, Fernandez M, Alvarez M, Benoit B, Laurent B, Grieco F, Spano G (2011) Isolation and characterization of tyramine-producing *Enterococcus faecium* strains from red wine. *Food Microbiol* 28 (3):434-439.
42. Rossi F, Gardini F, Rizzotti L, La Gioia F, Tabanelli G, Torriani S (2011) Quantitative analysis of histidine decarboxylase gene (*hdcA*) Transcription and histamine production by *Streptococcus thermophilus* PRI60 under conditions relevant to cheese making. *Appl Environ Microbiol* 77 (8):2817-2822.
43. Romano A, Ladero V, Alvarez MA, Lucas PM (2014) Putrescine production via the ornithine decarboxylation pathway improves the acid stress survival of *Lactobacillus brevis* and is part of a horizontally transferred acid resistance locus. *Int J Food Microbiol* 175:14-19.

44. Coton E, Coton M (2009) Evidence of horizontal transfer as origin of strain to strain variation of the tyramine production trait in *Lactobacillus brevis*. Food Microbiol 26 (1):52-57.
45. Lucas PM, Wolken WAM, Claisse O, Lolkema JS, Lonvaud-Funel A (2005) Histamine-producing pathway encoded on an unstable plasmid in *Lactobacillus hilgardii* 0006. Appl Environ Microbiol 71 (3):1417-1424.
46. Komprda T, Burdychova R, Dohnal V, Cwikova O, Sladkova P, Dvorackova H (2008) Tyramine production in Dutch-type semi-hard cheese from two different producers. Food Microbiol 25 (2):219-227.
47. Bunkova L, Bunka F, Mantlova G, Cablova A, Sedlacek I, Svec P, Pachlova V, Kracmar S (2010) The effect of ripening and storage conditions on the distribution of tyramine, putrescine and cadaverine in Edam-cheese. Food Microbiol 27 (7):880-888.
48. Lucas PM, Blancato VS, Claisse O, Magni C, Lolkema JS, Lonvaud-Funel A (2007) Agmatine deiminase pathway genes in *Lactobacillus brevis* are linked to the tyrosine decarboxylation operon in a putative acid resistance locus. Microbiol 153:2221-2230.
49. Aymerich T, Martin B, Garriga M, Vidal-Carou MC, Bover-Cid S, Hugas M (2006) Safety properties and molecular strain typing of lactic acid bacteria from slightly fermented sausages. J Appl Microbiol 100 (1):40-49.
50. Bover-Cid S, Hugas M, Izquierdo-Pulido M, Vidal-Carou MC (2001) Amino acid-decarboxylase activity of bacteria isolated from fermented pork sausages. Int J Food Microbiol 66 (3):185-189.
51. Pereira CI, Crespo MTB, Romao MVS (2001) Evidence for proteolytic activity and biogenic amines production in *Lactobacillus curvatus* and *L. homohiochii*. Int J Food Microbiol 68 (3):211-216.

52. Landete JM, Pardo I, Ferrer S (2007) Tyramine and phenylethylamine production among lactic acid bacteria isolated from wine. *Int J Food Microbiol* 115 (3):364-368.
53. Marcobal A, de las Rivas B, Munoz R (2006) First genetic characterization of a bacterial beta-phenylethylamine biosynthetic enzyme in *Enterococcus faecium* RM58. *FEMS Microbiol Lett* 258 (1):144-149.
54. Wunderlichova L, Bunkova L, Koutny M, Jancova P, Bunka F (2014) Formation, degradation, and detoxification of putrescine by foodborne bacteria: A review. *Compr Rev Food Sci Food Safety* 13 (5):1012-1030.
55. Ladero V, Canedo E, Perez M, Cruz Martin M, Fernandez M, Alvarez MA (2012) Multiplex qPCR for the detection and quantification of putrescine-producing lactic acid bacteria in dairy products. *Food Control* 27 (2):307-313.
56. Arena MP, Romano A, Capozzi V, Beneduce L, Ghariani M, Grieco F, Lucas P, Spano G (2011) Expression of *Lactobacillus brevis* IOEB 9809 tyrosine decarboxylase and agmatine deiminase genes in wine correlates with substrate availability. *Lett Appl Microbiol* 53 (4):395-402.
57. Tristezza M, Vetrano C, Bleve G, Spano G, Capozzi V, Logrieco A, Mita G, Grieco F (2013) Biodiversity and safety aspects of yeast strains characterized from vineyards and spontaneous fermentations in the Apulia Region, Italy. *Food Microbiol* 36 (2):335-342.

Fig 1: Results of PCR tests for the presence of genes involved in BA production (*tdcA*, *hdcA*, *odc* and *aguD-AguA*). A representative of each positive species is shown. For *Lc. lactis* subsp. *lactis* and *cremoris*, a representative of the negative strains is also shown (see text for details). For each BA cluster, the negative (-) and positive (+) controls (*E. faecalis* V583 for TDC and AGDI, *Lb. buchneri* B301 for HDC and *Lb. saerimneri* 30A for ODC) are indicated. MW: Molecular weight standard Gene Ruler (Fermentas). TDC: Tyramine-producing cluster; HDC: Histamine-producing cluster; ODC: Putrescine-producing cluster (via the ornithine decarboxylase pathway); AGDI: putrescine producing cluster (via the agmatine deiminase pathway).

487 **Table 1: BA-producing strains among the LAB tested.** Number of strains with the capacity to produce tyramine (Tym), β -phenylethylamine
488 (β -phe), histamine (Him) or putrescine (Put) in supplemented broth, as determined by (U)HPLC, and the presence of the corresponding genes, as
489 shown by PCR. N: number of strains tested. ODC: ornithine decarboxylase pathway. AGDI: agmatine deiminase pathway.

490

Species	N	Tym		β -phe	Him		Put (ODC)		Put (AGDI)	
		PCR	(U)HPLC	(U)HPLC	PCR	(U)HPLC	PCR	(U)HPLC	PCR	(U)HPLC
<i>Lactobacillus brevis</i>	4	4	4	0	0	0	0	0	4	4
<i>Lactobacillus casei</i>	12	0	0	0	0	0	0	0	0	0
<i>Lactobacillus curvatus</i>	3	3	3	0	0	0	0	0	2	2
<i>Lactobacillus delbrueckii</i>	9	0	0	0	0	0	0	0	0	0
<i>Lactobacillus fermentum</i>	10	0	0	0	0	0	0	0	0	0
<i>Lactobacillus plantarum</i>	19	0	0	0	0	0	0	0	0	0
<i>Lactococcus lactis</i> subsp. <i>lactis</i>	16	0	0	0	0	0	0	0	8	8
<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	7	0	0	0	0	0	0	0	3	3
<i>Leuconostoc mesenteroides</i>	14	0	0	0	0	0	0	0	0	0
<i>Streptococcus thermophilus</i>	43	0	0	0	2	0	0	0	0	0
Total	137	7	7	0	2	0	0	0	17	17

491

492

493

494

495

496

497 **Table 2: Classification of tyramine- and putrescine-producing strains based on their production capacity.** The strains were classified as
498 'strong' (more than 90% of the substrate present [1 mM tyrosine or agmatine] converted after 24 h of incubation), 'medium' (between 40 and 90%
499 converted) or 'poor' (<40% converted) producers. N: number of strains tested. Tym: tyramine. Put: putrescine. AGDI: agmatine deiminase
500 pathway. CR: conversion rate.

501

Species	N	Tym			Put (AGDI)		
		CR>90	90>CR>40	CR<40	CR>90	90>CR>40	CR<40
<i>Lactobacillus brevis</i>	4	4	0	0	1	2	1
<i>Lactobacillus curvatus</i>	3	3	0	0	2	0	0
<i>Lactococcus lactis</i> subsp. <i>lactis</i>	8	0	0	0	5	3	0
<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	3	0	0	0	0	3	0
Total	18	7	0	0	8	8	1

502

503

504