Bacterioplankton responses to riverine and atmospheric inputs in
a coastal upwelling system (Ría de Vigo, NW Spain)

By

E. Teira¹, Hernández-Ruiz M¹, Barber-Lluch E¹, Sobrino C¹, Teixeira IG², Álvarez-
Salgado XA², Nieto-Cid M², Martínez-García S^{1,3}, Figueiras FG², Fernández E¹

¹ Departamento Ecoloxía e Bioloxía Animal, Universidade de Vigo, 36310 Vigo, Spain

² Instituto de Investigaciones Marinas, CSIC, 36208 Vigo, Spain

³ Present address: Centre for Ecology and Evolution in Microbial model Systems –
EEMiS, Linnaeus University, SE-39182 Kalmar, Sweden.

Keywords: bacterioplankton, production, biomass, riverine water, atmospheric
deposition, Spain, Galicia, Ría de Vigo

Running head: Bacterioplankton response to inputs of matter

Abstract

Anthropogenic pressures are changing the magnitude and nature of matter inputs into the ocean. The Ría de Vigo (NW Iberian Peninsula) is a highly productive and dynamic coastal system, likely affected by such alterations. Previous nutrient-addition microcosm experiments conducted during contrasting hydrographic conditions suggested that heterotrophic bacteria appear to be limited by organic carbon (C) and occasionally co-limited by inorganic nutrients in this coastal area. In order to assess short-term responses in biomass, production and respiration of heterotrophic bacteria from the Ría de Vigo to increasing amounts of natural inputs of matter, we conducted 6 microcosm experiments, where surface seawater collected in spring, summer and autumn was mixed with increasing amounts of dissolved natural matter concentrates from riverine and atmospheric origin. Simultaneous experiments with controlled inorganic and/or organic additions indicated that bacteria were co-limited by inorganic nutrients and C in spring and summer and primarily limited by C in autumn. Production responded more than biomass to increasing inputs of matter whereas respiration did not change. The bacterial production response to increasing dissolved organic C load associated with riverine and atmospheric inputs was strongly related with the relative phosphorous (P) content of the dissolved matter concentrates. Our data suggest that bacterial production might decrease with the increase of P-deficient allochthonous matter inputs, which would have important biogeochemical consequences for C cycling in coastal areas.

Introduction

Nutrient inputs to the ocean have increased over the past decades as a result of human activity (Galloway & Cowling 2002) and are expected to further increase in the future (Galloway et al. 2004, Duce et al 2008). Atmospheric deposition, surface run-off and ground-water effluents introduce inorganic and organic nutrients and pollutants from anthropogenic origin into the coastal ocean (Jickells 1998, Doney 2010, Statham 2012, Jickells et al. 2014). The flux of reactive nitrogen to the coastal oceans through atmospheric deposition and riverine discharge is expected to increase 10-20% by 2050, primarily due to the intensification of fertilizer and energy production (Howarth et al. 2012, Moore et al. 2013). Anthropogenic alterations of global biogeochemical cycles are changing not only the magnitude but also the nature of matter inputs into the ocean. For instance, the relatively higher increase in anthropogenic carbon (C) and nitrogen (N) compared to phosphorus (P) inputs appears to result in a global increase of the C:P and N:P supply ratios to the global biosphere (Peñuelas et al. 2012, 2013). These altered nutrient inputs will likely affect microbial plankton dynamics (Grover 2000, Danger et al. 2007, Hitchcock & Mitrovic 2013) and suggest a global scenario of increasing P limitation in marine ecosystems (Peñuelas et al. 2012), where N is known to be currently the major limiting nutrient (Elser et al. 2007).

When bacteria are not limited by organic C, they are predicted to outcompete phytoplankton for mineral nutrient uptake (Grover 2000). Many studies indicate that bacteria are better competitors than phytoplankton for P uptake (Pengerud et al. 1987, Jansson 1993, Guerrini et al. 1998, Joint et al. 2002, Danger et al. 2007, Vadstein et al. 2012), particularly at very low concentrations (Thingstad et al. 1993). Therefore, the distinct nutrient requirements and uptake efficiencies of bacteria and phytoplankton, as

well as the magnitude and composition of allochthonous matter inputs, may determine the responses of the microbial communities to nutrient enrichment.

The Ría de Vigo is an eutrophic embayment located in the coastal system of the North-West Iberian Peninsula, characterized by the intermittent upwelling of inorganic nutrient-rich water (Fraga 1981, Tenore et al. 1995). Water exchange between this embayment and the adjacent shelf is determined by the balance between river discharge and on-shelf wind stress (Álvarez-Salgado et al. 2000). Nutrient delivery in this area has been reported to be in the order of $1500 \text{ mg N m}^{-2} \text{ yr}^{-1}$ (Gago et al. 2005) associated with riverine discharge, and about $100\text{-}250 \text{ mg N m}^{-2} \text{ yr}^{-1}$ associated with wet atmospheric deposition (Rodríguez & Macías 2006). Significant inputs of organic C in the Ría de Vigo have been also measured associated to riverine (Gago et al. 2005) and atmospheric (Teira et al. 2013) matter. Previous experimental studies on the effect of controlled inorganic and organic nutrient additions (Martínez-García et al. 2010) and natural additions of rainwater (Teira et al. 2013, Martínez-García et al. 2015) on coastal microbial planktonic communities in this coastal area showed that (1) microbial plankton responses to nutrient enrichment are highly variable; (2) phytoplankton is more responsive to natural rainwater additions than heterotrophic bacteria; (3) heterotrophic bacteria are primarily limited by organic C and occasionally co-limited by inorganic nutrients.

In the present study we aimed at further investigating the response of microbial plankton to dissolved natural matter inputs in the Ría de Vigo. The type of inputs studied included not only atmospheric but also riverine dissolved matter entering through fluvial discharge, as the latter introduces higher amounts of inorganic and organic nutrients into this highly productive ecosystem (Gago et al. 2005, Rodríguez & Macías 2006). In order to improve our predictive capability we conducted three

experiments using different microbial plankton communities collected under contrasting hydrographic conditions (spring, summer and autumn) where we evaluated the response of heterotrophic bacterial biomass, production and respiration to increasing amounts of matter inputs from riverine and atmospheric sources. As riverine discharge and atmospheric deposition introduce organic C to the Ría de Vigo, and bacteria are primarily limited by C in this region, we hypothesize that increasing amounts of riverine or atmospheric inputs will increasingly stimulate bacterial production, biomass and respiration in surface waters from the Ría de Vigo.

Methods

Natural seawater for the experiments was taken in the middle sector of the Ría de Vigo (42°14.09' N, 8°47.18' W) in spring, summer and autumn 2013. Vertical profiles of water column temperature, salinity and in situ fluorescence down to 25 m depth were obtained with a SBE 9/11 CTD probe and a Seatech fluorometer attached to a rosette sampler. Then, sub-surface seawater (3-4 m) was collected in 12 L acid-clean Niskin bottles and filtered through a 200 µm pore size mesh to remove larger zooplankton. Subsequently, 4 L UV transparent whirlpak bags were gently filled with 2 L of seawater under dim light conditions.

Preparation of natural concentrates for the addition experiments. River and rainwater samples and <10 µm atmospheric particles were collected and processed to obtain natural concentrates of the riverine and atmospheric inputs to the Ría de Vigo. Our aim was to reduce the volume of the original water samples 10-fold without altering their chemical composition, i.e. trying to avoid the addition or the loss of any component in the natural samples.

The River Oitabén-Verdugo was sampled in April, July and October 2013, a week before the addition experiments. We collected the water samples just upstream of the freshwater-seawater interface, to ensure that the chemical composition of the river samples were representative of the riverine water that mixes with the seawater of the Ría de Vigo. Five liters of each sample were gravity filtered through a pre-washed (with 10 L of ultrapure water) dual-stage (0.8 and 0.2 μm) filter cartridge (Pall-Acropak supor Membrane) and the filtrate was concentrated 10-fold by using rotatory evaporation with a Buchi R215 evaporator. This concentration was performed under mild conditions (bath temperature: 25 °C, vacuum: 13 mbar, condenser: acetone/ CO_2) to avoid breakage of any organic compound present in the original water samples. Measurements of the concentration of inorganic (ammonium, nitrite, nitrate and phosphate) and organic (dissolved organic carbon and nitrogen) substrates confirmed that the samples were concentrated quantitatively without any significant loss or gain.

A MTX rainwater sampler (model FAS005AB) and a high volume PM10 MCV PM1025 particle sampler (model CAV-A/MS) were installed at the rooftop of the Instituto de Investigaciones Marinas (CSIC) to collect samples of wet and dry deposition to the Ría de Vigo. The MTX sampler was equipped with a humidity sensor to open the system only when it was raining, allowing the sampling of just the wet fraction of the atmospheric deposition, the rainwater. Rainwater was collected from four weeks to one week before the addition experiment. Samples were taken on a daily basis and frozen immediately after collection. A week before the experiment, the daily samples were thawed at ambient temperature, mixed in one volume (6 L), and quantitatively concentrated following the same procedure as for the riverine samples. Rainwater was collected only for the experiments conducted in April and October 2013 because wet deposition was very scarce the weeks before the July experiments (36 mm

accumulated from 11 June to 10 July collected in the meteorological station at the Vigo city hall). The high volume sampler was used to gather atmospheric particles (1–10 μm) on precombusted (450 $^{\circ}\text{C}$, 4 h) 140 mm GF/F filters. The particles were collected the week before the three addition experiments, operating during 48 h at a rate of 30 $\text{m}^3 \text{h}^{-1}$. Then, the water-soluble fraction (WSF) of one eighth of the filter was extracted in 400 mL of the corresponding rainwater concentrate by mechanic stirring during 40 min. For the case of the July experiment, the WSF was extracted in milli-Q water. These proportions (1/8 of the filter in 400 mL of water) were decided to obtain 10-fold the expected concentrations according to previous existing information on the composition of wet and dry deposition to the Ría de Vigo (Teira et al. 2013, Martínez-García et al. 2015). Final mixed extracts were filtered through precombusted (450 $^{\circ}\text{C}$, 4 h) 47 mm diameter Whatman GF/F filters in an acid-cleaned glass filtration system, under low N_2 flow pressure, to be chemically characterized and used in the experiments as atmospheric concentrate. As for the riverine concentrates, quantitative concentration was observed except for the silicate since the reduction of the rainwater volume was carried out in a glass rotary evaporator and the atmospheric particles were collected onto a glass fibber filter.

Natural and controlled addition experiments. Increasing aliquots of riverine and atmospheric concentrates were added to surface seawater collected in the middle Ría de Vigo in spring, summer and autumn 2013. The concentrates were supplemented in proportions ranging from 1% (0.1% concentrate: 0.9% ultrapure water: 99% seawater) to 10% (1% concentrate: 99% seawater) of the original (previous to concentration) riverine and atmospheric materials ensuring that the final salinity of the samples was kept constant independently of the amount of extract added. Seawater was

167 mixed with 1%, 2.5%, 4%, 5%, 7.5% and 10% of natural matter from riverine (riverine
168 discharge) and atmospheric (dry and wet deposition) origin. A control treatment (no
169 addition) was included for each type of input. Three replicates were included for the
170 control, 1%, 5% and 10% treatments, and one replicate for the 2.5%, 4% and 7.5%
171 treatments.

172 With this procedure we aimed to test the impact of natural additions of riverine
173 and atmospheric materials over a wide range of realistic concentrations. To calculate the
174 current average riverine and atmospheric inputs to the Ría de Vigo we have considered
175 the average river flow of the River Oitabén-Verdugo, $17 \text{ m}^3 \text{ s}^{-1}$ (Gago et al. 2005) and
176 the average precipitation to the Ría de Vigo, 7.7 mm d^{-1} . Considering that the surface
177 area of the ría is 174 km^2 , a mean surface mixing layer of 2 m and an average flushing
178 time of this layer of 5 days, it results that the surface mixing layer of the ría contains
179 about 2% of riverine water and 2% of rainwater. Therefore, the additions up to 10%
180 would serve as a test for the response of the Ría de Vigo to future global change
181 scenarios in which human activities increase the quantity without modifying the quality
182 of riverine and atmospheric substrates.

183 Controlled nutrient addition experiments were also conducted in order to
184 describe the limiting nutrient for bacterial growth during each sampled season. We used
185 the same controlled nutrient addition treatments as in Martínez-García et al. (2010): (1)
186 No addition treatment; (2) Inorganic nutrient treatment: $5 \text{ } \mu\text{mol L}^{-1}$ nitrate (NO_3^-), 5
187 $\mu\text{mol L}^{-1}$ ammonium (NH_4^+) and $1 \text{ } \mu\text{mol L}^{-1}$ phosphate (HPO_4^{2-}); (3) Organic nutrient
188 treatment: $5 \text{ } \mu\text{mol L}^{-1}$ glucose and $5 \text{ } \mu\text{mol L}^{-1}$ equimolar mix of 18 aminoacids (all the
189 protein amino acids except cysteine and tyrosine); and (4) Mixed treatment: Inorganic
190 and organic nutrient treatments. Three replicates were included for each treatment. All
191 the experimental bags were incubated for 48 h under natural irradiance and temperature

conditions. Bacterial biomass, production and respiration were measured at time zero and after 48 h incubations. In the nutrient controlled experiments only biomass and production were measured after 48 h.

Inorganic and organic nutrients. Aliquots for inorganic nutrients determination (ammonium, nitrite, nitrate and phosphate) were collected in 50 mL polyethylene bottles and frozen at -20°C until analysis by standard colorimetric methods with an Alliance Futura segmented flow analyzer (Hansen and Grasshoff 1983). Water for the analysis of dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) was filtered through $0.2\ \mu\text{m}$ filters (Pall, Supor membrane Disc Filter) in an all-glass filtration system under positive pressure of N_2 and collected into pre-combusted (450°C , 12 h) 10 mL glass ampoules acidified with H_3PO_4 to $\text{pH} < 2$. Samples were measured with a Shimadzu TOC-V total organic carbon analyzer fitted with a Shimadzu TNM-1 total nitrogen measurement unit. Dissolved organic nitrogen (DON) was obtained by subtracting ammonium + nitrite + nitrate from TDN.

Dissolved organic matter (DOM) fluorescence. The fluorescence of dissolved protein-like (FDOM_T) and humic-like substances (FDOM_A) was also determined on the concentrates. Measurements were performed in a Perkin Elmer LS 55 luminescence spectrometer at two fixed pairs of excitation/emission wavelengths: 280/350 nm for FDOM_T and 250/435 nm for FDOM_A (Coble 1996). Calibration was done using a mixed standard of quinine sulfate and tryptophan in 0.1 N sulphuric acid following Nieto-Cid et al (2006) to convert fluorescence units to normalized fluorescence intensity units (NFIU).

Bacterial biomass. The abundance of heterotrophic bacteria was determined with a BD FACSCalibur flow cytometer equipped with a laser emitting at 488 nm. Picoplankton samples (1.8 mL) were preserved with 1% paraformaldehyde + 0.05% glutaraldehyde and frozen at -80°C until analysis. Prior to analysis, heterotrophic bacteria were stained with 2.5 mM SybrGreen DNA fluorochrome. The empirical calibrations between side scatter and mean cell diameter described in Calvo-Díaz & Morán (2006) were used to estimate biovolume (BV) of heterotrophic bacteria. BV was finally converted into biomass by using the allometric relationship of Norland (1993): $\text{fg C cell}^{-1} = 120 \times \text{BV}^{0.72}$.

Heterotrophic Bacterial Production. The [^3H]leucine incorporation method (Kirchman et al. 1985), modified as described by Smith & Azam (1992), was used to determine leucine (Leu) incorporation rates. Leucine was added at 40 nM final concentration. Samples were incubated for 1 h simulating in situ temperature conditions in a dark incubation chamber. We used the empirical leucine to carbon conversion derived by Martínez-García et al. (2010) ($2.6 \pm 1.1 \text{ kg C mol Leu}^{-1}$).

Bacterial Respiration. INT ((iodo-phenyl)-3(nitrophenyl)-5(phenyl) tetrazolium chloride) reduction rate was used as estimator of bacterial respiration. Size-fractionated in vivo INT reduction rates were measured as described in Martínez-García et al. (2009). Four 100 mL dark bottles were filled from each microcosm bottle. One bottle was immediately fixed by adding formaldehyde (2% w/v final concentration) and used as killed-control. Fifteen minutes later, all the replicates were inoculated with a sterile solution of 7.9 mM INT to a final concentration of 0.2 mM. Samples were incubated at the same temperature as the microcosm bottles in dark conditions during 1

h. After incubation, samples were filtered sequentially through 0.8 and 0.2- μm pore size polycarbonate filters, which were stored at $-20\text{ }^{\circ}\text{C}$ in 1.5 mL cryovials until further processing. The formed insoluble formazan crystals (INT-F) were extracted from the filters by adding 1 mL of propanol and sonicating for 20-30 minutes in $50\text{ }^{\circ}\text{C}$ water using an ultrasonic bath. One mL of the propanol extract containing the INT-F was transferred to 1.5 mL microfuge vials and then centrifuged at $13200 \times g$ during 10 minutes at $18\text{ }^{\circ}\text{C}$. The absorbance at 485 nm was then measured using a spectrophotometer (Beckman model DU640). Bacterial respiration was operationally defined as INT reduction rates in the $<0.8\text{ }\mu\text{m}$ size fraction (Robinson 2008). In order to transform INT reduction rates into carbon respiration an R/ETS ratio of 12.8 (Martínez-García et al 2009) and a respiratory quotient (RQ) of 0.8 (Williams and del Giorgio 2005) were used.

Chlorophyll a (Chl a) concentration. Chl *a* concentrations were measured in 100 mL water samples which were filtered through $0.2\text{ }\mu\text{m}$ polycarbonate filters. The filters were immediately frozen at $-20\text{ }^{\circ}\text{C}$ until pigment extraction in 90% acetone at $4\text{ }^{\circ}\text{C}$ overnight in the dark. Chl *a* concentrations were determined, with a 10-AU Turner Designs fluorometer calibrated with pure Chl *a*.

Primary production. Five 75 mL Corning tissue flasks (3 light and 2 dark) were filled with seawater and spiked with 185 kBq ($5\text{ }\mu\text{Ci}$) $\text{NaH}^{14}\text{CO}_3$. Samples were incubated for 2 h in a temperature-controlled incubation chamber illuminated with cool white light from fluorescent tubes providing an average PAR of $240\text{ }\mu\text{E m}^{-2}\text{ s}^{-1}$. After the incubation period, samples were filtered through $0.2\text{ }\mu\text{m}$ polycarbonate filters at

very low vacuum (<50 mm Hg). Filters were exposed to HCl fumes for 24 h to remove unincorporated inorganic ^{14}C and radioassayed.

Results

Initial conditions and chemical composition of natural inputs

Initial conditions for each experiment are summarized in Table 1. Different hydrographic conditions were found during each survey. In spring (May 2013), high dissolved inorganic N (DIN= nitrate + nitrite + ammonium) and Chl *a* concentrations occurred while in autumn (October 2013) lower Chl *a* concentration were observed regardless of higher nutrient concentrations. In the summer experiment (July 2013) inorganic nutrient and Chl *a* concentrations were the lowest. Primary production rates were the highest in spring and lower values were measured in summer and autumn (Table 1). Bacterial biomass was lower in summer than in spring and autumn, whereas bacterial production and respiration rates did not notably varied among seasons (Table 1). The chemical composition of riverine and atmospheric (wet and dry) matter inputs collected during the three studied seasons showed a high temporal variability (Table 2). In general, riverine water collected in October 2013 contained higher DIN, phosphate and DOC than that collected in spring and summer (Table 2). On the other hand, atmospheric inputs in May 2013 contained higher DIN and DON but lower DOC concentrations than that collected in autumn. Atmospheric deposition in summer contained relatively low N concentrations (Table 2). Higher DOC and DON concentrations and lower DOC:DIN ratios (except in spring) were measured in riverine water compared to atmospheric inputs (Table 2). The relative contribution of nitrate to total DIN was, on average, significantly higher in riverine (98%) than in atmospheric (58%) inputs (t-test, $p < 0.001$). DON accounted on average 20% of total dissolved

nitrogen (Table 2) both in riverine and atmospheric inputs. Phosphate concentration was relatively low in both atmospheric and riverine inputs resulting in P:DIN ratios largely below the Redfield values (0.0625) (Table 2). Considering that in general protein-like substances are labile and humic-like compounds are recalcitrant, the ratio $FDOM_T/FDOM_A$ can be used as a proxy for DOM bioavailability. The mean value of this ratio in the DOM concentrates was significantly higher for atmospheric (7.4 ± 2.5) than for riverine (3.3 ± 0.7) inputs (t-test, $p = 0.05$) (Table 2). We also confirmed that the annual average nutrient concentrations of riverine and atmospheric inputs (mean value $\pm$ standard deviation of the three different seasons) extrapolated from the concentrates were within the limits of previous reported values obtained in this area (Table 2), except for the riverine nitrate, which is very unpredictable and variable.

Bacterial response to controlled nutrient additions

The response of heterotrophic bacteria to controlled nutrient additions differed among experiments. Enhanced bacterial production relative to the control (up 34-fold) was observed after mixed additions in the 3 experiments and also after organic addition in autumn (Fig. 1A). Bacterial biomass significantly increased after mixed addition (5 to 13-fold) and to a lesser extent after organic addition (3.8 to 6-fold) in the three experiments (Fig. 1B). Neither biomass nor production responded to inorganic addition alone (Fig. 1A, 1B).

Response of bacterial biomass and production to natural matter additions

In order to describe changes in biomass and production associated to increasing amounts of riverine or atmospheric inputs we represented the response, calculated as the ratio between the mean value in the treatment and the corresponding mean value in the control, versus the amount of DOC load associated to the different treatments (Fig. 2, 3). In spring, the bacterial biomass response significantly decreased with increasing

amounts of DOC associated to the riverine inputs (Fig. 2A), but did not change with increasing DOC load associated to atmospheric inputs (Fig. 2B). In summer, the biomass response did not change with increasing DOC load from riverine inputs (Fig. 2C), but was positively related to the total DOC load associated to atmospheric inputs (Fig. 2D), which explained 46% of the observed variability. Mean ($\pm$ SE) biomass response was 1.06 ± 0.02 -fold, ranging from 0.76-fold, for atmospheric inputs in spring, to 1.29-fold for riverine inputs in summer.

Bacterial production showed a different pattern of response. The production response associated to increasing amounts of riverine inputs significantly decreased in spring and summer (Fig. 3A, 3C), explaining 34-36% of the variability. By contrast, production significantly increased with increasing DOC load associated to atmospheric inputs in summer and autumn, explaining 37-52% of the observed variability. The ordinate intercepts of the significant regressions did not significantly differed from 1 (t-test, $p > 0.05$). Mean production response was 1.28 ± 0.05 -fold, ranging from 0.76- to 1.97-fold, for riverine inputs in summer, and was significantly higher than the mean biomass response (t-test, $p < 0.001$). Mean production response was significantly higher for atmospheric (1.37 ± 0.07 -fold) than for riverine inputs (1.19 ± 0.07 -fold) (t-test, $p = 0.039$).

The slopes of the regressions between the production response (i.e., the production response rates) and the DOC load (a zero value was assigned for the atmospheric inputs in spring and the riverine inputs in autumn) were significantly correlated with the P:DIN ratio of the inputs ($r = 0.87$, $p = 0.025$, $n = 6$). The P:DIN ratio of the inputs explained 75% of the response rate variability (Fig. 4).

We did not find any significant response in bacterial respiration to increasing matter additions for in any of the 6 experiments (data not shown).

Bacterial response compared to phytoplankton response

In order to interpret the response of bacteria to the different natural matter additions within the context of the microbial food web, we compared the response of bacterial production to that of primary production (Fig. 5). A detailed description of phytoplankton responses is described elsewhere (Fernández et al, in prep). Overall, the mean bacterial production response was significantly lower (0.83-fold) than primary production response for riverine inputs (t-test, $p = 0.044$) while no significant differences were observed for atmospheric inputs. The bacterial to primary production response decreased as the % of riverine addition increased in spring and summer (Fig. 5B), although the observed trend was not significant.

Discussion

The impact of natural matter inputs of riverine or atmospheric origin promoted variable responses of heterotrophic bacteria depending on the initial physico-chemical and ecological conditions of the water samples and the chemical composition of the inputs.

Overall, the mean production response was higher than the mean biomass response, as previously observed for natural rainwater additions (Teira et al. 2013), natural dust additions (Bonnet et al. 2005, Marañón et al. 2010) and controlled additions (Mills et al. 2008, Martínez-García et al. 2010), which is likely associated to top-down processes (i.e. predation) (Zubkov et al. 2000) which prevent bacterial biomass accumulation regardless of increments in bacterial production (Jürgens & Massana 2008). Discrepancies between biomass and production responses may also result from a lag period between changes in production and biomass due to the heterogeneity of population growth rates; if only a fraction of the total bacterial assemblage is growing

or actively incorporating leucine, production rates will increase faster than the total biomass because the cells are “diluted” by the inactive or non-growing fraction (Ducklow 2000). Previous studies in the same region showed that the effect of controlled dissolved matter addition on bacterial production and biomass was most apparent 24 h and 48 h after the amendments, respectively (Martínez-García et al. 2010). Nevertheless, due to the great number of replicates and different treatments in the present study (i.e a total of 42 experimental units per experiment) we chose to sample only after 48 h in order to capture the response of phytoplankton which typically occur after 48-72 h (Martínez-García et al. 2010, 2015).

Overall, respiration did not significantly change with increasing nutrient loads in agreement with previous experimental rainwater additions in the same sampling site (Teira et al. 2013).

Bacterial production response to atmospheric and riverine natural inputs

The production response rate, estimated as the slope of the linear regression between production response and DOC-load, varied from -0.10 to 0.14 $\mu\text{M DOC}^{-1}$ (Figures 3, 4) across the six experiments. As the ordinate intercepts did not significantly differ from 1, a response rate of 0.14 implies that the production will increase 14% per each μM of DOC-load increment. Even though production response rates were negative for riverine and positive for atmospheric additions, both types of inputs had the greatest effect on production when initial DIN concentrations were the lowest (summer) (Fig. 3, Table 1), as shown before in rainfall addition experiments (Zou et al. 2000, Teira et al. 2013). As previously observed in the sampling area (Martínez-García et al. 2010), the response of bacteria to controlled nutrient amendment experiments indicate that bacteria were primarily limited by organic carbon during the three experiments as neither

biomass nor production responded to the addition of inorganic nutrients alone (Fig. 1A, 1B).

Since bacteria in our system are primarily limited by organic C, a higher production would be expected associated to increasing DOC-load. However, such response pattern was only observed in summer and autumn for the atmospheric inputs (Fig. 3D, 3F). A lower response of bacterial production associated to riverine compared to atmospheric inputs could be expected due to differences in DOM bioavailability. Previous studies indicate that riverine DOM is largely refractory (Søndergaard & Middelboe 1995, Moran et al. 1999) compared to rainwater DOM (Avery et al. 2003). The significantly higher $FDOM_T/FDOM_A$ ratio (used as a proxy for DOM bioavailability) of atmospheric inputs compared to those of riverine inputs (Table 2) suggest a higher availability of atmospheric than riverine DOC, however the $FDOM_T/FDOM_A$ ratio did not significantly explain the variability observed in the response rates.

Bacterial response to the additions might be also constrained by the mineral components of the inputs. Particularly, the lack of bacterial production response to our controlled organic additions, containing C and N but not P (Figure 1A), strongly suggests a P deficiency for bacterial growth in spring and summer, as previously observed on certain occasions in the sampling area (Martínez-García et al. 2010). Despite the response rate was not correlated with the phosphorous load, the strong and significant relationship observed between the response rate and the P:DIN ratio of the inputs (Fig. 4), suggest that the balance between inorganic P and N forms largely regulate the response of bacteria to riverine and atmospheric nutrient fluxes. Furthermore, the intriguing negative bacterial production response rate associated with riverine matter inputs observed in spring and summer (Fig. 3A, 3C), suggests that

bacteria may be competing with phytoplankton for inorganic nutrients. The higher response of primary production compared to bacterial production after riverine inputs, particularly in spring and summer (Fig 5), partially support this possibility.

Considering that bacteria typically have lower C:N and C:P ratios than primary producers (Chrzanowski et al. 1996, Cotner et al. 2000, Cotner & Biddanda 2002, Vrede et al. 2002, Carlsson et al. 2012), they may exhibit high inorganic nutrient demands and, thus, directly compete with phytoplankton for the uptake of mineral N and P. Bacteria, due to their smaller size, are expected to outcompete phytoplankton for the uptake of limiting inorganic nutrients (Thingstad et al. 1993, Cotner & Biddanda 2002, Joint et al. 2002). In accordance to this, bacterial production response relative to phytoplankton production response to dust inputs has been shown to increase as the degree of oligotrophy increases in the Atlantic Ocean (Marañón et al. 2010), likely due to the superior ability of bacteria to take up inorganic nutrients at very low concentrations. We also observed that the production response of bacteria relative to that of phytoplankton after the natural matter inputs is higher for treatments with low percentage of addition (Fig. 5B).

Even though the competitive advantage of bacteria over phytoplankton has been seen to hold for dissolved inorganic P uptake (Pengerud et al. 1987, Jansson 1993, Guerrini et al. 1998, Joint et al. 2002), there are no equally clear evidences for bacteria as better competitors than phytoplankton for DIN (Danger et al. 2007, Vadstein et al. 2012). Although heterotrophic bacteria significantly contribute to both ammonium and nitrate uptake (Kirchman and Wheeler 1998, Zehr & Ward 2002, Fouilland et al. 2007), they are not expected to outcompete phytoplankton for nitrate due to the higher energetic cost associated to nitrate uptake compared to ammonium or dissolved organic nitrogen (Vallino et al. 1996, Joint et al. 2002). Likewise, contrary to phytoplankton,

bacteria appear to be more commonly limited by P than by N in marine ecosystems (Cotner et al. 1997, Church 2008, Zohary et al. 2005, Carlsson et al. 2012, Vadstein et al. 2012), as it has been also suggested in our sampling area (Martínez-García et al. 2010). If bacteria are secondarily limited by inorganic P rather than by inorganic N, and phytoplankton primarily limited by DIN, the negative bacterial production response rate observed for riverine inputs in spring and summer (Fig. 3A, 3C), when P supply is extremely low relative to DIN (Table 2, Fig. 4), suggests that the bacterial response might be modulated by the phytoplankton response, which seems to profit from the large nitrate concentration associated to the riverine inputs (Table 2, Fig. 5). A very low P supply associated to riverine inputs have been previously reported elsewhere (Labry et al. 2001).

The hypothesis of a P-mediated bacterial response to natural matter inputs in the Ría de Vigo is further supported by controlled nutrient addition experiments conducted by our research group in the same sampling site. When pooling all the available data from such experiments, including the present and two previous studies (Martínez-García et al. 2010, Prieto et al. 2015) we found a significant and positive correlation between the production response to organic additions (containing C and N but not P) and the ambient phosphate concentration in the Ría de Vigo ($r^2 = 0.75$, $p = 0.011$, $n = 7$; data not shown).

In conclusion, we have shown that bacterial production response to increasing DOC-load associated with atmospheric and riverine additions largely depends on the P:DIN ratio of the inputs. Negative production response rates are associated to riverine inputs that show extremely low P:DIN ratios, probably due to phytoplankton outcompeting bacteria for P uptake when nitrate concentration is high. In a future global change scenario, where both the riverine and atmospheric nutrient fluxes are expected to

increase and their associated relative P content is expected to decrease as a consequence of anthropogenic activities (Galloway et al. 2004, Peñuelas et al. 2013), autotrophic production would likely benefit more than heterotrophic bacterial production. Moreover, the limited response of bacteria to increasing P-limited inputs to coastal waters may have further implications for ocean C cycling as the unused allochthonous DOC might be eventually transported to open ocean waters where it could be utilized by bacteria or exported to the ocean interior.

Acknowledgements

We thank all the people involved in the project REIMAGE who helped with the preparation and sampling of the experiments and the analytical work. We thank Estación de Ciencias Mariñas de Toralla (ECIMAT) for the technical support during the experiments as well as the crew of the R/V Mytilus for their help during the work at sea. This research was supported by the Spanish Ministry of Science and Innovation through project REIMAGE (grants CTM2011-30155-C03-01 and CTM2011-30155-C03-02) and by the Xunta de Galicia through project DIMENSION (grant EM2013/023). E. B-L was funded by a F.P.I. MICINN fellowship. M.N.-C. was funded by the CSIC Program "Junta para la Ampliación de Estudios" co-financed by the ESF. S.M.G. was funded by a Marie Curie fellowship.

References

- Álvarez-Salgado XA, Gago J, Míguez BM, Gilcoto M, Pérez FF (2000) Surface waters of the NW Iberian margin: Upwelling on the shelf versus outwelling of upwelled waters from the Rias Baixas. *Estuar Coast and Shelf Sci* 51(6): 821–837
- Avery GB, Willey JD, Kieber RJ, Shank GC, Whitehead RF (2003) Flux and bioavailability of Cape fear River and rainwater dissolved organic carbon to Long Bay, southeastern United States. *Global Biogeochem Cycles* 17: article number 1042
- Bonnet S, Guieu C, Chiaverini J, Ras J, Stock A (2005) Effect of atmospheric nutrients on the autotrophic communities in a low nutrient, low chlorophyll system. *Limnol Oceanogr* 50: 1810-1819
- Calvo-Díaz A, Morán XAG (2006) Seasonal dynamics of picoplankton in shelf waters of the southern Bay of Biscay. *Aquat Microb Ecol* 42: 159-174
- Carlsson P, Granéli E, Granéli W, Gonzales Rodriguez E, Fernandes de Carvalho W, Brutemark A, Lindehoff E (2012) Bacterial and phytoplankton nutrient limitation in tropical marine waters, and a coastal lake in Brazil. *J Exp Mar Biol Ecol* 418-149: 37-45
- Charzanowski TH, Kyle M, Elser JJ, Sterner RW (1996) Element ratios and growth dynamics of bacteria in an oligotrophic Canadian shield lake. *Aquat Microb Ecol* 11: 119-125
- Coble PG (1996) Characterization of marine and terrestrial DOM in seawater using excitation-emission matrix spectroscopy. *Marine Chem* 51, 325–346
- Cotner JB, Ammerman JW, Peele ER, Bentzen E (1997) Phosphorous-limited bacterioplankton growth in the Sargasso Sea. *Aquat Microb Ecol* 13: 141-149
- Cotner JB, Sada RH, Bootsma H, Johengen T, Cavaletto JF, Gardner WS (2000) Nutrient limitation of heterotrophic bacteria in Florida Bay. *Estuaries* 23: 611-620

- Cotner JB, Bidanda BA (2002) Small players, large role: microbial influence on biogeochemical processes in pelagic aquatic ecosystems. *Ecosystems* 5: 105-121
- Danger M, Leflaive J, Oumarou C, Ten-Hage L, Lacroix G (2007) Control of phytoplankton-bacteria interactions by stoichiometric constraints. *Oikos* 116: 1079-1086
- Doney SC (2010) The growing human footprint on coastal and open-ocean biogeochemistry. *Science* 328: 1512-1516
- Duce RA, LaRoche J, Altieri K, Arrigo KR and others (2008) Impacts of atmospheric anthropogenic nitrogen on the open ocean. *Science* 320:893-897
- Ducklow HW (2000) Bacterial production and biomass in the oceans. In: *Microbial Ecology of the Oceans* (1st edition). Edited by Kirchman D.L., pp: 85–120. Wiley-Liss, New York
- Elser JJ, Bracken MES, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, Ngau JT, Seabloom EW, Shurin JB, Smith JE (2007) Global analysis of nitrogen and phosphorous limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecol Lett* 10: 1135-1142
- Fraga F (1981) Upwelling off the Galician coast, Northwest Spain in *Coastal Upwelling*. *Coast and Estuar Sci*, Vol. 1, FA Richards, ed, AGU, Washington, DC, 176-182
- Fouilland E, Gosselin M, Rivkin RB, Vasseur C, Mostajir B (2007) Nitrogen uptake by heterotrophic bacteria and phytoplankton in Arctic surface waters. *J Plankton Res* 29: 369-376
- Gago J, Álvarez-Salgado XA, Nieto-Cid M, Brea S, Piedracoba S (2005) Continental inputs of C, N, P and Si species to the Ría de Vigo (NW Spain). *Estuar, Coastal and Shelf Sci* 65(1): 74-82

Galloway JN, Cowling B (2002) Reactive Nitrogen and The World: 200 Years of Change. *Ambio* 31 (2): 64-71

Galloway JN, Dentener FJ, Capone DG, and other 12 (2004) Nitrogen cycles: past, present, and future. *Biogeochem* 70: 153-226

Grover JP (2000) Resource competition and community structure in aquatic microorganisms: experimental studies of algae and bacteria along a gradient of organic carbon to inorganic phosphorus supply. *J Plankton Res* 22: 1591-1610

Guerrini F, Mazotti A, Boni L, Pistocchi R (1998) Bacterial-algal interactions in polysaccharide production. *Aquat Microb Ecol* 15: 247-253

Hansen HP, Grasshoff K (1983) Automated chemical analysis. In: Grasshoff, K., Ehrhardt, M., Kermling, K. (ed), *Methods of seawater analysis* (2nd edition). Verlag Chemie, Weinheim, p 347–395

Havskum H, Thingstad TF, Scharek R, Peters F, Berdalet E, Sala MM et al (2003) Silicate and labile DOC interfere in structuring the microbial food web via algal-bacterial competition for mineral nutrients: results from a mesocosm experiment. *Limnol Oceanogr* 48: 129-140

Hitchcock JN, Mitrovic SM (2013) Different resource limitation by carbon, nitrogen and phosphorus between base flow and high flow conditions for estuarine bacteria and phytoplankton. *Est Coast and Shelf Sci* 135: 10-115

Howarth R, Swaney D, Billen G, Garnier J, Hong B, Humborg C, Johnes P, Mörrth C-M, Marino R (2012) Nitrogen fluxes from the landscape are controlled by net anthropogenic nitrogen inputs and climate change. *Front Ecol Environ* 10: 37-43

Jansson M (1993) Uptake, exchange and excretion of orthophosphate in phosphate-starved *Scenedesmus quadricauda* and *Pseudomonas K7*. *Limnol Oceanogr* 38: 1162-1178.

561 Jickells TD, 1998. Nutrient biogeochemistry of the coastal zone. *Science* 281:
562 217-222.

563 Jickells TD, Andrews JE, Parkes DJ, Suratman S, Aziz AA, Hee YY (2014)
564 Nutrient transport through estuaries: the importance of the estuarine geography. *Est*
565 *Coast and Shelf Sci* 150: 215-229

566 Joint I, Henriksen P, Fonnes GA, Bourne D, Thingstad TF, Riemann B (2002)
567 Competition for inorganic nutrients between phytoplankton and bacterioplankton in
568 nutrient manipulated mesocosms. *Aquat Microb Ecol* 29: 145-159

569 Jürgens, K and Massana, R (2008) Protistan grazing on marine bacterioplankton,
570 in: *Microbial Ecology of the Oceans* (2nd edition), edited .by: Kirchman, D L, Wiley-
571 Liss, New York, 383–441.

572 Kirchman DL, Knees E, Hodson R (1985) Leucine incorporation and its
573 potential as a measure of protein synthesis by bacteria in natural aquatic systems. *Appl*
574 *Environ Microbiol* 49: 599-607

575 Kirchman DL and Wheeler PA (1998) Uptake of ammonium and nitrate by
576 heterotrophic bacteria and phytoplankton in the sub-Arctic Pacific. *Deep-Sea Res I* 45:
577 347-365

578 Labry C, Herbland A, Delmas D (2001) The role of phosphorus on planktonic
579 production of the Gironde plume waters in the Bay of Biscay. *J Plankton Res* 24: 97-
580 117

581 Marañón E, Fernandez A , Mouriño-Carballido B, Martinez-Garcia S, Teira E et
582 al (2010) Degree of oligotrophy controls the response of microbial plankton to Saharan
583 dust. *Limnol Oceanogr* 55: 2339-2352

Martínez-García S, Fernández E, Aranguren-Gassis M, Teira E (2009) In vivo electron transport system activity: a method to estimate respiration in marine microbial planktonic communities. *Limnol Oceanogr Methods* 7, 459–469

Martínez-García S, Fernández E, Álvarez-Salgado XA, González J, Lønborg C, Marañón E, Morán XAG, Teira E (2010). Differential responses of phytoplankton and heterotrophic bacteria to organic and inorganic nutrient additions in coastal waters off the NW Iberian Peninsula. *Mar Ecol Prog Ser* 416, 17–33

Martínez-García S, Arbones B, García-Martín EE, Teixeira IG, Serret P, Fernández E, Figueiras FG, Teira E, Álvarez-Salgado XA (2015) Impact of atmospheric deposition on the metabolism of coastal microbial communities. *Est Coast Shelf Sci* 153: 18-28

Mills MM, Moore CM, Langlois R, Milne A, Achterberg E, Nachtigall K, Lochte K, Geider RJ, La Roche J (2008) Nitrogen and phosphorus co-limitation of bacterial productivity and growth in the oligotrophic subtropical North Atlantic. *Limnol Oceanogr* 53: 824-834

Moore CM, M. M. Mills, K. R. Arrigo, I. Berman-Frank, L. Bopp, P. W. Boyd, E. D. and other 17 (2013) Processes and patterns of oceanic nutrient limitation. *Nature Geoscience* 6: 701-710

Moran MA, Sheldon Jr WM, Sheldon JE (1999) Biodegradation of riverine dissolved organic carbon in five estuaries of the southeastern United States. *Estuaries* 22: 55-64

Nieto-Cid M, Álvarez-Salgado XA, Perez FF (2006) Microbial and photochemical reactivity of fluorescent dissolved organic matter in a coastal upwelling system. *Limnol Oceanogr* 51:1381-1390

Nogueira E, Pérez FF, Ríos AF (1997) Seasonal Patterns and Long-term Trends in an Estuarine Ecosystem (Ría de Vigo, NW Spain). *Estuar and Coast Shelf Sci* 44, 285–300

Norland S (1993) The relationship between biomass and volume of bacteria In: Kemp PF, Sherr BF, Sherr EB, Cole JJ (ed) *Handboook of methods in aquatic microbial ecology*. Lewis Publishers, Boca Raton, FL, pp 303–307

Pengerud B, Skjoldal EF, Thingstad TF (1987) The reciprocal interaction between degradation of glucose and ecosystem structure. *Studies in mixed chemostat cultures of marine bacteria, algae and bacterivorous nanoflagellates*. *Mar Ecol Prog Ser* 25: 111-117

Peñuelas J, Sardans J, Rivas-Ubach A, Janssens IA (2012) The human-induced imbalance between C, N and P in Earth’s life system. *Global Change Biol* 18: 3-6

Peñuelas J, Poulter B, Sardans J, Ciais P, van der Velde M and other 9 (2013) Human-induced nitrogen–phosphorus imbalances alter natural and managed ecosystems across the globe. *Nature Communications* 4, Article number: 2934

Prieto A, Barber-Lluch E, Hernández-Ruiz M, Martínez-García S, Fernández E, Teira E (2015) Assessing the role of phytoplankton-bacterioplankton coupling in the response of microbial plankton to nutrient additions (submitted to *J Plankton Res*)

Robinson C (2008) Heterotrophic bacterial respiration In Kirchman DL (ed) *Microbial ecology of the oceans* (2nd edition). Wiley-Liss, New York, pp 299-336.

Rodríguez L, Macías F (2006) Eutrophication trends in forest soils in Galicia (NW Spain) caused by the atmospheric deposition of nitrogen compounds. *Chemosphere* 63, 1598–1609

631 Smith DC, Azam F (1992) A simple, economical method for measuring bacterial
632 protein synthesis rates in seawater using ³H-leucine. *Mar Microb Food Webs* 6: 107-
633 114

634 Søndergaard M, Middelboe A (1995) A cross-system analysis of labile dissolved
635 organic carbon. *Mar Ecol Prog Ser* 118: 283-294

636 Statham PJ (2012) Nutrients in estuaries-An overview and the potential impacts
637 of climate change. *Science of the Total Environment* 434: 213-227

638 Teira E, Hernando-Morales V, Martínez-García S, Figueiras FG, Arbones B,
639 Álvarez-Salgado XA (2013) Response of bacterial community structure and function to
640 experimental rainwater additions in a coastal eutrophic embayment. *Estuar Coast and*
641 *Shelf Sci* 119, 44–53

642 Tenore KR, Alonso-Noval M, Álvarez-Ossorio M and other 16 (1995) Fisheries
643 and oceanography off Galicia, NW Spain (FOG): Mesoscale spatial and temporal
644 changes in physical processes and resultant patterns in biological productivity. *J*
645 *Geophys Res* 100 (C6): 10943-10966

646 Thingstad TF, Skjoldal EF, Bohne RA (1993) Phosphorus cycling and algal-
647 bacterial competition in Sandsfjord, western Norway. *Mar Ecol Prog Ser* 99: 239-259

648 Thingstad TF, Øvreas L, Egge JK, Løvdal T, Heldal M (2005) Use of non-
649 limiting substrates to increase size; a generic strategy to simultaneously optimize uptake
650 and minimize predation in pelagic osmotrophs? *Ecol Lett* 8: 675-682

651 Vadstein O, Andersen T, Reinertsen HR, Olsen Y (2012) Carbon, nitrogen and
652 phosphorus resource supply and utilization for coastal planktonic heterotrophic bacteria
653 in a gradient of nutrient loading. *Mar Ecol Prog Ser* 447: 55-75

Vallino JJ, Hokinson CS, Hobbie JE (1996) Modeling bacterial utilization of dissolved organic matter: optimization replaces Monod growth kinetics. *Limnol Oceanogr* 41: 1591-1609

Vrede K, Heldal M, Norland S, Bratbak G (2002) Elemental composition (C, N, P) and cell volume of exponentially growing and nutrient-limited bacterioplankton. *Appl Environ Microbiol* 68: 2965-2971

Williams PJ le B, del Giorgio PA (2005) Respiration in aquatic ecosystems: history and background In: del Giorgio PA, Williams PJ le B (ed), *Respiration in aquatic ecosystems*, Oxford University Press, London, p 1-18

Zehr JP, Ward BB (2002) Nitrogen cycling in the ocean: new perspectives on processes and paradigms. *Appl Environ Microbiol* 68: 1015-1024

Zohary T, Herut B, Krom MD, Mantoura RFC, Pitta P, Psarra S, Rassoulzadegan F, Stambler N, Tanaka T, Thingstad F, Woodward EMS (2005) P-limited bacteria but N and P co-limited phytoplankton in the Eastern Mediterranean- a microcosm experiment. *Deep Sea Res II* 52: 3011-3023

Zou L, Chen HT, Zhang J (2000) Experimental examination of the effects of atmospheric wet deposition on primary production in the Yellow Sea. *Journal of Experimental Marine Biology and Ecology* 259: 111-121

Zubkov MV, Sleigh MA, Burkill PH, Leakey RJG (2000) Bacterial growth and grazing loss in contrasting areas of North and South Atlantic. *J Plank Res* 22: 685-711

Figure legends

Figure 1. Response (mean value in treatment relative to mean value in control after 48 h of incubation) of (A) bacterial production and (B) bacterial biomass to controlled nutrient additions in spring, summer and autumn. Error bars represent the standard error; where error bars are not visible, they are smaller than the size of the symbol. A response equal to 1 means no change relative to control. Asterisks indicate a response significantly >1 (t-test; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

Figure 2. Response (mean value in treatment relative to mean value in control after 48 h of incubation) of bacterial biomass to increasing concentrations of natural riverine (A, C, E) and atmospheric (B, D, F) matter inputs (expressed as total DOC-load) in spring (A, B), summer (C, D) and autumn (E, F). Error bars represent the standard error; where error bars are not visible, they are smaller than the size of the symbol. A biomass response equal to 1 means no change relative to control. Regression line, slope value (b) and determination coefficient (r^2) are represented if a significant increase or decrease of the biomass response with increasing DOC-load was found.

Figure 3. Response (mean value in treatment relative to mean value in control after 48 h of incubation) of bacterial production to increasing concentrations of natural riverine (A, C, E) and atmospheric (B, D, F) matter inputs (expressed as total DOC-load) in spring (A, B), summer (C, D) and autumn (E, F). Error bars represent the standard error; where error bars are not visible, they are smaller than the size of the symbol. A production response equal to 1 means no change relative to control. Regression line, slope value (b) and determination coefficient (r^2) are represented if a significant increase or decrease of the production response with increasing DOC-load was found.

Figure 4. Relationship between the production response rate (slope $\pm$ SE of the regression between production response and DOC-load) and the P:DIN ratio of the inputs.

Figure 5. (A) Response (mean value in treatment relative to mean value in control after 48 h of incubation) of primary production versus response of bacterial production to natural continental and atmospheric inputs. The diagonal represents the 1:1 line where bacterial and primary production equally responded to the inputs. (B) Bacterial to primary production response along the gradient of increasing additions. Dashed line indicates where bacterial and primary production equally responded to the inputs.

Table 1 Summary of the physical-chemical and biological conditions of seawater at the sampling station in spring, summer and autumn. DOC, dissolved organic carbon; DON, dissolved organic nitrogen; DIN, dissolved inorganic nitrogen.

Variable	May-13	July-13	October-13
Temperature °C	14.3	16.6	18.5
Salinity	35.35	35.62	33.40
Nitrate (μM)	2.80	0.22	3.48
Nitrite (μM)	0.11	0.05	0.31
Ammonium (μM)	1.05	0.96	2.02
Phosphate (μM)	0.13	0.16	0.33
DOC (μM)	65.2	76.2	84.9
DON (μM)	4.7	6.3	7.6
P:DIN	0.033	0.130	0.057
Chlorophyll-a (mg m^{-3})	13.5	0.6	1.4
Primary production ($\text{mg C m}^{-3} \text{ h}^{-1}$)	17.4	1.2	3.6
Bacterial Biomass (mg C m^{-3})	26.8	11.1	27.0
Bacterial Production ($\text{mg C m}^{-3} \text{ h}^{-1}$)	0.17	0.18	0.18
Bacterial Respiration ($\text{mg C m}^{-3} \text{ h}^{-1}$)	0.87	0.60	0.93

Table 2. Summary of the chemical characteristics of dissolved matter concentrates from riverine water and atmospheric wet (spring and autumn) or dry deposition (summer) collected in spring, summer and autumn. DOC, dissolved organic carbon; DON, dissolved organic nitrogen; DIN, dissolved inorganic nitrogen; FDOM_T and FDOM_A, protein-like and humic-like fluorescence of dissolved organic matter, respectively. *The annual mean composition was extrapolated from the composition of the concentrates by dividing by the concentration factor (10). **Reference for riverine inputs is Gago et al 2005 (values for Eiras station during year 2002). Reference for atmospheric inputs is project IMAN (values for station Bouzas wet, during years 2008-2009).

		May-13	July-13	October-13	Calculated Annual Mean*	Reference Annual Mean**
Riverine	DOC (μM)	795	788	1492	102 ± 40	89 ± 12
	DON (μM)	64	50	39	5.1 ± 1.2	7.7 ± 1.5
	Nitrate (μM)	151	186	330	22.3 ± 9.5	6.0 ± 4.2
	Nitrite (μM)	0.43	0.50	1.28	0.07 ± 0.05	0.11 ± 0.04
	Ammonium (μM)	1.5	4.6	8.9	0.50 ± 0.37	0.62 ± 0.34
	Phosphate (μM)	0.14	0.48	2.46	0.10 ± 0.13	0.15 ± 0.08
	DOC:DIN	5.2	4.1	4.4		
	P:DIN	0.0009	0.0025	0.0072		
	FDOM _T /FDOM _A	2.5	3.7	3.8		
Atmospheric	DOC (μM)	369	552	817	58 ± 23	56 ± 34
	DON (μM)	41	18	22	2.7 ± 1.2	7.5 ± 26
	Nitrate (μM)	105	28	71	6.8 ± 3.9	9.1 ± 10.5
	Nitrite (μM)	0.04	0.03	0.61	0.02 ± 0.03	0.05 ± 0.07
	Ammonium (μM)	74.7	22.5	49.2	4.9 ± 2.6	9.7 ± 8.4
	Phosphate (μM)	0.66	0.70	1.52	0.10 ± 0.05	0.09 ± 0.18
	DOC:DIN	2.1	11.0	6.8		
	P:DIN	0.0037	0.0139	0.0127		
	FDOM _T /FDOM _A	6.3	5.8	10.3

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5