An improved method to obtain pure α -galactosides from lupin seeds

Cristina Martínez-Villaluenga¹, Juana Frias¹, Krzysztof Gulewicz² and Concepción Vidal-Valverde^{1*}

¹Instituto de Fermentaciones Industriales (CSIC), Juan de la Cierva 3, Madrid 28006, Spain

²Institute of Bioorganic Chemistry (PAS), Noskowskiego 12/14, Poznan 61-704, Poland.

*Autor to whom correspondence should be addressed

Tel.: + 34 915622900 Ext 241

Fax: 34 915644873

Email: ificv12@ifi.csic.es

ABSTRACT

Improvement of a previous described method of purification of α -galactosides, raffinose family oligosaccharides (RFOs) has been developed for lupins. The considerable amount of sucrose present in the RFO preparations obtained by previous method has been removed by modifying the purification stage on diatomaceous earth and charcoal. The present method allows for the preparation of high purity RFOs containing about 99.4% of RFOs in form of white fine powder which provides new perspectives for the production of pure α -galactoside preparations for their use as prebiotics in functional foods.

Keywords: α -galactosides, raffinose family oligosaccharides (RFOs), lupins

1 INTRODUCTION

2 α -Galactosides, called also raffinose family oligosaccharides (RFOs) are
3 widely distributed in the plant kingdom. Large amounts of RFOs occur in the
4 generative parts of higher plants where perform protective physiological functions (1-
5 4). In legumes, RFOs accumulate during seed development and raffinose and
6 stachyose are formed *de novo* during seed maturation (5, 6) and are carbon reserves
7 utilised during germination (7, 8). Legumes seeds contain a large α -galactoside
8 content and their consumption has been associated with the production of flatulence,
9 since these oligosaccharides are not hydrolysed in the small intestine, due to the
10 absence of α -galactosidases in monogastrical animals, and they are fermented in the
11 lower intestine (9, 10). This problem has been considered to be the single most
12 important factor that deters people from eating this nutritious food (10).

13 It has been considered recently that pure RFOs are prebiotics. RFOs promote
14 the growth of bifidobacteria population (11,12) and, consequently, there is a great
15 deal of interest to use prebiotic oligosaccharides as ingredients in functional foods in
16 order to manipulate the composition of colonic microflora, contributing to human
17 health in many ways.

18 There are many prebiotic oligosaccharides known, principally in the Japanese
19 market and they are added as ingredients in a large number of products such as soft
20 drinks, cookies, cereals and candies (12). α -Galactosides from soybean are the only
21 legume oligosaccharides on the market and the main producer is Japan (13). In the
22 USA and Europe RFOs are also available on the market, however the market leaders
23 are fructo-oligosaccharides in these regions.

1   These low-molecular weight oligosaccharides are present in different seeds,
2 but lupins are one of the legumes with the most of α -galactoside content (7-15%) (8;
3 14-17). The sweet lupins, containing very low levels of alkaloids, can be a good
4 source of RFOs. In 2000 a paper was published describing a simple method for the
5 isolation and purification of α -galactosides from lentils and peas (18), but the
6 obtained RFO preparations contained sucrose in considerable amounts (11 and
7 12%, respectively). In the present work, a modification of the previous published
8 method is presented that allows obtaining lupin RFOs free of sucrose. It creates new
9 perspectives for production pure α -galactoside preparations for their use as
10 prebiotics in functional foods.

12   **MATERIALS AND METHODS**

13   **Samples and Chemicals.** Sweet lupin seeds of *Lupinus albus* L. cv.
14 Multolupa, were obtained from the Agricultural Research and Technology
15 Development Service of the Agriculture and Commerce Council of the Junta de
16 Extremadura (Spain). Seeds were cleaned and stored in polyethylene container at
17 4°C until used. Diatomaceous earth, charcoal, and naphthoresorcinol were
18 purchased from Sigma, Darmstadt (Germany). Dowex 50WX8, 100-200 mesh was
19 purchased from BDH Laboratories (England). Ethanol, 2-propanol and acetic acid
20 were supplied by Scharlau (Germany). Acetonitrile (HPLC grade) was purchased
21 from ACROS-ORGANIC (Belgium). Silica gel 60 F₂₅₄ TLC plates, sucrose, raffinose,
22 and stachyose were obtained from Merck, Darmstadt (Germany). Milipore FH (0.45
23 μ m) membranes were obtained from Millipore (Bedford, MA).

Isolation of RFOs. Lupins seeds were submitted to different selective extractions according to a previous procedure described by Gulewicz et al. (18), (Figure 1).

Purification of RFOs. The purification scheme of RFOs is shown in Figure 1. RFOs precipitate was dissolved in 25 mL of distilled water and suspended in a glass funnel (pore size G4, 7cm x 5 cm i.d.) with diatomaceous earth and charcoal (1:1 w/w) and connected to vacuum. The funnel was washed with 6% ethanol to remove sucrose. The absence of sucrose in the eluat was confirmed by thin-layer chromatography (TLC) according to Dey, (7). TLC of carbohydrates was performed on silica gel 60 F254 plates with 2-propanol:acetic acid:water (5:2:3 v/v). Carbohydrates were visualised with naphthoresorcinol. After confirming the absence of sucrose, RFOs were eluted with 70% ethanol until negative naphthoresorcinol reaction. The ethanol RFOs fraction was concentrated to dryness on a vacuum evaporator at 50°C. The RFOs were dissolved in 10 mL of distilled water, transferred into a Dowex 50WX8 column (12 x 1.5 cm i.d.), and washed with distilled water (50 mL) until negative naphthoresorcinol reaction. The acid solution of RFOs (pH 2) was adjusted to pH 7 with 4% freshly prepared Ca(OH)_2 . The solution was then boiled for 2 minutes and centrifuged for 10 minutes at 10.000 rpm. Supernatant containing a high purity RFOs was then evaporated to dryness on a vacuum evaporator at 50°C.

Determination of total soluble carbohydrate content. Sucrose and α -galactosides content were determined by HPLC following the procedure described by Granito et al. (19).

RESULTS AND DISCUSSION

Figure 2 shows chromatograms of RFO preparations from lupins obtained as previously described by Gulewicz *et al.*, (18) (method A) and improved method described in this work (method B). It can be observed that chromatographic peak of sucrose was sharply reduced using method B, whilst raffinose, stachyose and verbascose increased. These peaks were quantified and results are shown in Table 1. The improved method described in this paper increased to 99.4% the purity of RFO preparations, compared with a 75.1 % obtained using method A. The RFO powder obtained presented a 12.6% of raffinose, a 62.3% of stachyose and a 24.5% of verbascose, and only a trace amount of sucrose (0.8%) was detectable (Table 1).

There are five stages in the improved method of isolation and purification α -galactosides from legume: (i) imbibition of seeds, (ii) extraction of RFOs, (iii) RFO precipitation, (iv) purification of RFOs on diatomaceous earth and charcoal, and (v) cation-exchange chromatography. In the purification stage with diatomaceous earth and charcoal (iv) percolation of RFOs with water used in method described by Gulewicz *et al.*, (18) was not efficient for total removal of monosaccharides, sucrose and other impurities. Using the 6% ethanol solution makes possible entire elution of sucrose, that has been confirmed by HPLC analysis. Finally, using a higher concentration of ethanol (70%) conducted to the total elution of RFOs and the obtained preparation shows very high purity (> 99%).

The removal of RFOs from legume seeds has been widely studied from the nutritional point of view, in order to obtain legume flour with low flatulent-causing compounds (20, 21, 22), but little attention has been paid to the extraction and purification of these α -galactosides as prebiotic substances. Lupin seeds have been previously used as source of RFOs but in those studies purity of the isolated have

not been reported (15). Kim et al. (23) optimized the extraction of oligosaccharides from soybean meal with either water at 50°C or 10% ethanol and purification was carried out by ultrafiltration. These authors found fructose, sucrose, raffinose and stachyose in their concentrates. Gulewicz *et al.*, (18) obtained soluble sugars preparations containing 11.5% sucrose, 7.2% raffinose, 29.3% stachyose and 40.3% verbascose from peas and 12.2% sucrose, 5.6% raffinose, 9.3% ciceritol, 39.5% stachyose and 11.5% verbascose from lentils. The improved method described in this paper offer a simple and rapid procedure for obtaining high purity RFO preparations free on monosaccharides and disaccharides from lupin seeds and it could be apply to other legumes. The pure RFOs can be offered as functional food ingredient like prebiotic, which can contribute to human health.

ACKNOWLEDGMENT

This work is a result of a bilateral scientific cooperation joint project between Poland (PAS) and Spain (CSIC), it has been supported by CICYT AGL-2001-2302 and it is part of C. Martínez-Villaluenga Thesis.

LITERATURE CITED

- (1) Dey, P. M. D-Galactose containing oligosaccharides. In: Biochemistry of storage carbohydrates in green plants; Dey, P. M. Eds.; Academic Press: London, **1985**, pp 5-129.

- (2) Kuo, T. M.; Van Middlesworth, F. K.; Wolf, W. J. Content of raffinose oligosaccharides and sucrose in various plant seeds. *J. Agric. Food Chem.*, **1988**, 36, 32-36.
- (3) Larsson, S.; Johansson, L. A.; Svenningsson, M. Soluble sugars and membrane lipids in winter wheats (*Triticum aestivum* L.) during cold acclimation. *Eur. J. Agron.* **1993**, 1, 85-90.
- (4) Horbowicz, M.; Obendorf, R. L. Seed desiccation tolerance and storability: Dependence on flatulence-producing oligosaccharides and cyclitols-review and survey *Seed Sci. Res.* **1994**, 4, 385-405.
- (5) Lowell, C. A.; Kuo, T. M. Oligosaccharide metabolism and accumulation in developing soybean seeds. *Crop Sci.*, **1989**, 29, 459-465.
- (6) Frias, J.; Vidal-Valverde, C.; Kozłowska, H.; Gorecki, R.; Honke, J.; Hedley, C. L. Evolution of soluble carbohydrates during the development of pea, faba bean and lupin seeds. *Z. Lebensm. Unters. Forsch.*, **1996**, 203, 27-32.
- (7) Dey, P. M. Oligosaccharides. In: *Methods in plant biochemistry*. Dey, P. M., Eds. Academic Press: New York, **1990**, Vol. 2, pp 189-218.
- (8) Frias, J.; Diaz-Pollan, C.; Hedley, C. L.; Vidal-Valverde, C. Evolution and kinetics of monosaccharide, disaccharide and α -galactosides during germination of lentils. *Z. Lebens. Unters. Forsch.* **1996**, 202, 35-39.

- (9) Cristofaro, E.; Mottli, F.; Whurmann, J. J. Involvement of raffinose family oligosaccharides in flatulence. *Sugars in nutrition*; Sepple, H. I., McNut, K. W., Eds.; Academic Press; New York, **1974**, pp 313-363.
- (10) Price, K. R.; Lewis, J.; Wyatt, G. M.; Fenwick, R. G. Flatulence-causes, relation to diet and remedies. *Nahrung* **1988**, 32, 609-626.
- (11) Gibson, G. R.; Roberfroid, M. D. Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J. Nutr.* **1995**, 125, 1401-1412.
- (12) Tomomatsu, H. Health effects of oligosaccharides. *Food Technology* **1994**, 48, 61-65
- (13) Crittenden, R. G.; Playne, M. J. Production, properties and applications of food-grade oligosaccharides. *Trends in Food Science and Technology*, **1996**, 7, 353-361.
- (14) Van Kempen, R. J.; Hughes, R. J. The nutritive value of lupin for pigs and poultry. *Proccedings of the first Australian lupin technical symposium*, **1994**, pp. 49-57.
- (15) Muzquiz, M.; Burbano, C.; Pedrosa, M. M.; Folkman, W. Y.; Gulewicz, K. Lupin as a potential source of raffinose family oligosaccharides. Preparative method for their isolation and purification. *Ind. Crops Prod.* **1999**, 9, 183-188.

- (16) Ruiz-López, M. A.; García-López, P. M.; Castañeda-Vazquez, H.; Zamora, N. J. F.; Garzón-De la Mora, P.; Bañuelos Pineda, J.; Burbano, C.; Pedrosa, M. M.; Cuadrado, C.; Muzquiz, M. Chemical composition and antinutrient content of three *Lupinus* species from Jalisco, Mexico. *J. Food Comp. Anal.*, **2000**, *13*, 193-199.
- (17) Saini, H. S.; Gladstones J. S. Variability in the total and component galactosyl sucrose oligosaccharides of *Lupinus* species. *Aust. J. Agric. Res.* **1986**, *37*, 157-166.
- (18) Gulewicz, P.; Ciesiolka, D.; Frías, J.; Vidal-Valverde, C.; Frejnagel, S.; Trojanowska, K.; Gulewicz, K. Simple method of isolation and purification of α -galactosides from legumes. *J. Agric. Food Chem.*, **2000**, *48*, 3120-3123.
- (19) Granito, M.; Frías, J.; Doblado, R.; Guerra, M.; Champ, M.; Vidal-Valverde, C. Nutritional improvement of beans (*Phaseolus vulgaris*) by natural fermentation. *Eur. Food Res. Technol.*, **2002**, *214*, 226-231.
- (20) Frías, J.; Vidal-Valverde, C.; Sotomayor, C.; Díaz-Pollan, C.; Urbano, G. Influence of processing on available carbohydrate content and antinutritional factors of chickpeas. *Eur. Food Res. Technol.*, **2000**, *210*, 340-345

1 (21) Sanz, M.A.; Blazquez, I.; Sierra, I.; Medrano, M. A.; Frias, J.; Vidal-Valverde,
2 C.; Hernandez, A. Nutritional evaluation of ethanol extracted lentil flours. *J.*
3 *Agric. Food Chem.*, **2001**, 49, 1854-1860.

4
5 (22) Vidal-Valverde, C.; Sierra, I.; Frias, J.; Prodanov, M.; Sotomayor, C.; Hedley,
6 C.L.; Urbano, G. Nutritional evaluation of lentil flours obtained after short time
7 soaking processes. *Eur. Food Res. Technol.*, **2002**, 215, 138-144.

8
9 (23) Kim, S.; Kim, W.; Hwang, K. Optimization of the extraction and purification of
10 oligosaccharides from defatted soybean meal. *Int. J. Food Sci. Technol.*, **2003**,
11 38, 337-342.

Table 1. Percentage of α -galactosides and sucrose in the RFO preparations from lupin obtained by method A and method B

Sugars	Method A*	Method B**
Sucrose	20.22 $\pm$ 0.11	0.82 $\pm$ 0.01
Raffinose	9.65 $\pm$ 0.33	12.56 $\pm$ 0.91
Stachyose	52.99 $\pm$ 2.13	62.33 $\pm$ 0.48
Verbascose	12.50 $\pm$ 1.01	24.47 $\pm$ 0.05
Total of α-galactosides	75.14 $\pm$ 2.86	99.36 $\pm$ 1.43

*) Method A: According to Gulewicz et al. (18)

**) Method B: Present improved method