[Click here to view linked References](#)

Allometric size-scaling of biometric growth parameters and metabolic and excretion rates. A comparative study of intertidal and subtidal populations of mussels (*Mytilus galloprovincialis*)

Kristina Arranz^a, Uxío Labarta^{a*}, M. José Fernández-Reiriz^a and Enrique Navarro^b

^aConsejo Superior de Investigaciones Científicas (CSIC), Instituto de Investigaciones Marinas (IIM), C/Eduardo Cabello 6, 36208 Vigo, Spain

^bDepartamento GAFFA (Animal Physiology), Facultad de Ciencia y Tecnología, Universidad del País Vasco/Euskal Herriko Unibertsitatea, Apartado 644, 48080 Bilbao, Spain

Abstract

Allometric relationships between biometric parameters (i.e., soft body and shell weights and shell organic content vs. shell length) as well as for routine and standard metabolic and ammonia excretion rates related to flesh weight and shell length were estimated and compared for subtidal and intertidal populations of *Mytilus galloprovincialis* in Galicia (NW Spain). This is the first report on allometric size-scaling of excretion and metabolic (both routine and standard) rates in this species. No evidences of differences in size-exponent were found between physiological rates or between both populations for any physiological rate. Intercepts of regression lines were significantly higher in subtidal than in intertidal mussels, indicating greater levels of energy expenditure in the former. However, metabolic scope for feeding and growth was about two-fold in intertidal mussels, pointing to a reduced growth efficiency compared with subtidal mussels. Evolution of biometric parameters of body components with size indicated that subtidal mussels allocated energy resources preferably into flesh growth, achieving higher condition indices, while intertidal mussels put more effort on shell calcification and thickening which resulted in heavier shells of reduced organic content. These

26 differentiated growth “strategies” of both populations could be related to their differences in
27 growth efficiencies.

28

29

30

31 *Corresponding author. Tel.: +34 986231930, fax: +34 986292762. E-mail address:
32 labarta@iim.csic.es (Uxío Labarta).

33 *Introduction*

34 The extensive culture of the mussel (*M. galloprovincialis*), with a production volume
35 that ranged between 200,000–300,000 tons, and a production value that exceeded 100
36 million euros in 2012 (www.pescadegalicia.com), is the main aquaculture industry in
37 Galicia (NW Spain). Mussels are cultured in floating systems (rafts) consisting of a
38 500m² wood structure anchored to the seafloor, from which culture ropes and/or seed
39 collectors are suspended. Nowadays, the number of ropes per raft is limited to 500 and
40 ca. 3300 rafts are located in the Galician Rias. Mussel culture is scheduled according to
41 the availability of natural resources for feeding and seed recruitment, the biological
42 cycle of mussels and the fluctuations of market demand (Labarta et al., 2004).

43 Producers collect mussel seeds for culture either from intertidal or subtidal habitats;
44 hence, the physiological and metabolic differences associated to the origin of the
45 individuals may constitute an important factor toward the optimization of mussel
46 production (Pérez-Camacho et al., 2013; 2014). Several factors have been invoked to
47 account for such differences in physiological behavior:

48 In the first place, intertidal populations are subjected to cycles of air exposure, which
49 implies intervals of hypoxic or anoxic conditions. Moreover, tidal cycles lead to
50 periodical shortages in feeding (Peterson & Black, 1988; Marsden & Weatherhead,
51 1999). Intertidal mussels cannot compensate for periods of starvation; even though
52 some advantages of the intertidal habitat have been identified, such as organic matter
53 resuspension and thermal fluctuations liable to improve energy gain at low tide by
54 increasing rates of digestion (Elvin & Gonor, 1979; Bayne et al., 1988). Storey &
55 Storey (1990) observed that organisms subjected to air exposure periods present a

reduced metabolic rate, considering this response as an energy saving mechanism acting to compensate for the lesser feeding or energy acquisition time (Shick et al., 1988).

In the second place, physiological responses associated with the origin of individuals have been considered as an indicator of the existence of genetic diversity (Rawson & Hilbish, 1991; Widdows et al., 1984; Dickie et al., 1984). However, some other authors suggest that these responses would reflect the persistence of original habitat influences in the form of an “ecological memory” (Mallet et al., 1987; Okumuş & Stirling, 1994; Pérez-Camacho et al., 1995; Labarta et al., 1997; Babarro et al., 2000a;b; 2003). In *M. galloprovincialis*, such ecological memory is found to be responsible for the differences in physiological condition between intertidal and subtidal populations. Hence, subtidal individuals exhibited higher values of growth rate, condition index (Pérez-Camacho et al., 1995; Babarro et al., 2003), energy reserves (i.e., triacylglycerol and phospholipid levels) (Freites et al., 2002), and SFG (“**Scope for growth**”, Labarta et al., 1997); since, despite of their higher values of oxygen consumption (Babarro et al., 2000b) and ammonia excretion rates, both clearance rate (Labarta et al., 1997; Babarro et al., 2000a) and absorption efficiencies (Labarta et al., 1997) were also higher in subtidal mussels.

Differentiated growth trends encompass most of these physiological differences between intertidal and subtidal mussel populations. Growth is frequently measured in bivalves as changes in shell length or weight, but this approach tends to disregard essential features of this phenomenon. For example, growth trajectories often differ for shell and soft tissues according to environmental factors or variations in the reproductive cycle (Hilbish, 1986; Borrero & Hilbish, 1988; Dame, 2012). Concerning shell growth itself, shell architecture and organic content are important attributes often subjected to variations between populations. Dynamics of shell formation includes growth in both circumference and thickness (Gosling, 2003) as variables simultaneously

contributing to determine size and shape of bivalves. Habitat can be responsible for much of the variation in the relationships between biometric parameters accounting for different aspects of growth in mussels (Rao, 1953; Seed, 1973; Brown & Seed, 1977; Aldrich & Crowley, 1986). Since these relationships are known also to change along the life-span of individuals, the characterization of allometric scaling of these parameters to body size (usually shell length) in different populations constitutes a useful approach in the comparative analysis of habitat effects.

Among the various physiological components of growth, metabolic rate is a key parameter determining rates of growth in two related ways: In the context of the energy budget of an individual, metabolic rate constitutes, together with excretion rate, the main component of energy expenditure. At the same time, it summarizes the metabolic energy demands to sustain maintenance and growth processes. In the literature on metabolic rates in bivalves it is common use to distinguish between measurements performed on active fed organisms representative of routine rates and standard or resting rates characteristics of starved organisms (Bayne & Newell, 1983). In sessile continuous feeders, such as bivalves, the difference between both metabolic measurements represents the energy in excess of basal requirements used in the various activities of feeding, digestion and biosynthesis involved in tissue growth. This metabolic component has been recently designed as metabolic scope for feeding and growth (MSFG) (Tamayo et al., 2013).

As stated for biometric parameters, the analysis of allometric scaling of physiological rates to body size is meaningful in interpreting growth processes. Allometric relationships have been formalized as power functions of the form:

$$Y = aX^b$$

where Y is the biological variable, a the intercept, X the body mass, and b the allometric scaling exponent. Concerning metabolism, one of the most important points of controversy in scientific discussion about power functions is focused on the value of the exponent (for review see Glazier, 2005; White, 2011). Along many years different authors have reported that mass scaling exponents fluctuate within a range of values of 0.5–1 (Prosser, 1973; Withers, 1992; White et al., 2006). On account of observed variability, the assumption of a common weight exponent for metabolism (the proposed $\frac{3}{4}$ scaling law) is no longer tenable (Riisgård, 1998; Atanasov & Dimitrov, 2002; Bokma, 2004; Glazier, 2005; Muller-Landau et al., 2006; Reich et al., 2006; White et al., 2006; Glazier, 2008; 2009a; b; c; 2010; White, 2011).

Empirical knowledge of allometric exponents is of particular importance in the parameterization of bioenergetics growth models, where metabolic expenditure corresponding to the size groups has forcibly been estimated indirectly in different ways (Duarte et al., 2010). Particularly, in the case of *M. galloprovincialis* lack of specific information on allometric scaling values for any metabolic level has compelled size-standardization to be based on values reported for related species of *Mytilus*, mainly *M. edulis* (Navarro et al., 1991; Labarta et al., 1997; Babarro et al., 2000b; Tamayo, 2012; Anestis et al., 2010) and eventually *M. chilensis* (Sarà & Pusceddu, 2008).

In the context of the energy balance rates of nitrogen excretion (as ammonia-N) constitute a minor component of total energy losses (10% on average: Bayne & Newell, 1983); however, its determination is important as an indicator of changes in metabolizable substrates mainly occurring along the seasonal cycle.

Summarizing, habitat variation has been shown to promote differentiated growth trends in intertidal and subtidal populations of mussels that are relevant in regards to

suspended culture of this species. Such differentiation involves changing relationships between biometric parameters representative of shell and soft tissue dynamics that can be conveniently approached by means of allometric functions. Consequently, the aims of this study were: (1) to compute allometric parameters for the scaling of flesh and shell weight (both total and organic) to body size represented by shell length, (2) to calculate allometric functions relating rates of energy loss (both metabolic and ammonia-N excretion rates) to body size for subsequent comparison between intertidal and subtidal populations, and (3) to analyze functional relationships of growth trends associated to body size and habitat with the metabolic scope for feeding and growth (MSFG).

Materials and methods

Collection and maintenance of mussels

Between September and October 2014 mussels (*Mytilus galloprovincialis*) were sampled from subtidal and intertidal habitats in Ria de Ares-Betanzos (Galicia, NW Spain), and brought to the laboratory where they were cleaned of epibionts and microbial biofilms with sterile scalpels and kept in open flow-through tanks of 20 L of capacity in seawater. The diet during the maintenance period consisted of a monoalgal suspension of *Rhodomonas lens* supplied in a continuous flow to each aquarium by a peristaltic pump (ISMATEC MPC Process). The concentration of food entering the tanks was established at 8000 cells ml⁻¹, at a flow rate of 10 L h⁻¹.

On the second day, shell lengths were measured to the nearest 0.5 mm with a caliper (Mitutoyo®) and individuals sorted in 8 size-classes in the range of 15–50 mm (Table 1). These groups were maintained in separate tanks at the above conditions for 15 days to let the mussels acclimate to laboratory conditions.

Physiological measurements

Metabolic rate

Metabolic rate was determined indirectly through the measure of oxygen consumption rate. Mussels were cleaned and placed in respirometers of about 780 ml of capacity, filled with filtered seawater (1µm), and maintained at a constant temperature (15°C). Two respirometers were left without animals as a control in order to correct for bacterial respiration, electronic drift, etc. (Labarta et al., 1997). The number of mussels constituting each group is reported in Table 1. This distribution was chosen in order to promote a uniform decrement in oxygen concentration. Determinations started 15

minutes after placing the mussels in the respirometers in order to let the mussels open their valves and start the normal respiratory activity. Dissolved oxygen concentration (mg L^{-1}) was registered by a LDO probe connected to a HATCH HQ40d oxymeter. Determinations were concluded before oxygen concentration had dropped below 70% of the initial concentration. Routine metabolic rate (RMR) was estimated after a continuous feeding period, while standard metabolic rate (SMR) was determined after 72 h of starvation, when a stable level of respiration had been attained as based on previous studies (data not published). Respiration rates were calculated following the formula used by Babarro et al. (2000b), being modified in order to correct the oxygen consumed with the control chambers:

where ΔO_2 is the difference between oxygen concentration registered in a respirometer with individuals from final to initial time, ΔO_2^c is the difference between oxygen concentration registered in control chambers from final to initial time, Vol represents the capacity (L) of the respirometer, t is the time (h) between final and initial oxygen registration and n means the number of individuals placed in the respirometer.

Ammonia excretion rate ($\text{VNH}_4\text{-N}$)

Ammonia ($\text{VNH}_4\text{-N}$) excretion rate was determined after placing the mussels cleaned of epibionts and biofilm in open Erlenmeyer flasks with 250 ml of filtered seawater (0.2 μm Millipore membranes). Temperature was maintained during the determinations by immersing the flasks in an isothermal bath. Two Erlenmeyer without animals were used as a control. After 120 min, water samples were collected from each Erlenmeyer flask

and frozen to -20°C until analysis in the laboratory, according to the phenol–hypochlorite method described by Solórzano (1969). Excretion rates were calculated as:

where VNH_4-N represents the ammonia excretion rate; μM and μM_c are the ammonia concentration estimated through the calibration curve in the sample and in the control chamber, respectively; Vol represents the capacity (ml) of the incubation chamber; and t is the incubation time (h).

Metabolic scope for feeding and growth

Metabolic scope for feeding and growth (MSFG) was computed as the difference between routine and standard metabolic rates and expressed as fraction of routine metabolic rate:

Biometry and condition index

After concluding the physiological determinations, individuals were dissected to determine flesh and shell dry weight (100°C for 24 h), as well as ash free dry weight (450°C for 24 h). Condition Index (CI) was calculated according to Freeman (1974) using the following equation:

Shell organic content (%) was estimated through the equation:

205 *Data analysis*

206 Allometric relationships of oxygen consumption and ammonia excretion rates vs. size
207 were determined on log-log transformed data by linear regressions using the least
208 squares method. Allometric equations were compared through a covariance analysis
209 (ANCOVA test) (Zar, 1996). Assumptions of ANCOVA were verified using residual
210 plots (linearity), Kolmogorov–Smirnov tests (normality of residuals), Levene tests
211 (homoscedasticity) and Durbin Watson tests (independence of residuals). The level of
212 **significance (α) for all analyses was set at $P = 0.05$** . Statistical analyses were performed
213 with the statistical package R 2.15.2 (<http://www.r-project.org/>), using a custom made R
214 script based on Zar (1996).

Results:

Biometry and condition index

Condition index (CI) increased with size in both subtidal and intertidal mussels (Fig. 1A); however, in subtidal individuals, CI was significantly higher than in those from intertidal habitat. Shell organic content (OC %) (Fig. 1B) was characterized by a marked decline with size in both populations. In addition, subtidal mussels had a greater percentage of shell organic content than intertidal individuals at each size class.

Relationships of flesh weight (FW) to shell length (SL) for intertidal and subtidal populations were fitted by linear regression after log-log transformation (Fig. 2A). As there were not statistically significant differences between slopes (see Table 2), a common slope was calculated and a values recalculated for each population:

$$\text{Subtidal: Log FW} = 3.053 \text{ Log SL} - 5.395$$

$$\text{Intertidal: Log FW} = 3.053 \text{ Log SL} - 5.455$$

Relationships of shell weight (SW) to shell length (SL) were estimated as described above for flesh weight/shell length ratio (Fig. 2B). Again, lack of statistically significant differences between slopes (see Table 2) allowed a common slope to be calculated and a values were recalculated for each population:

$$\text{Subtidal: Log SW} = 2.637 \text{ Log SL} - 4.052$$

$$\text{Intertidal: Log SW} = 2.637 \text{ Log SL} - 3.863$$

Total shell organic content (g) vs. shell length (mm) did not show significant differences between slopes (see Table 2), while intercepts were statistically significant. Therefore, equations were recalculated according to their common slope:

237 Subtidal: $\text{Log OC} = 2.415 \text{ Log SL} - 4.985$

238 Intertidal: $\text{Log OC} = 2.415 \text{ Log SL} - 4.926$

239 *Allometries of respiration rates*

240 Allometric equations for routine and standard metabolic rates were fitted as a function
241 of dry weight (Fig. 3) and length (Fig. 4). Due to the lack of significant differences (see
242 Table 3) between slopes for routine metabolic rate (RMR) or standard metabolic rate
243 (SMR) in relation to dry mass (dry flesh weight, FW) corresponding to intertidal and
244 subtidal groups (Fig. 3A and 3B), common weight exponents were calculated and
245 intercepts (a) recalculated according to these common slopes:

246 Subtidal: $\text{Log RMR} = 0.715 \text{ Log FW} - 0.428$

247 Intertidal: $\text{Log RMR} = 0.715 \text{ Log FW} - 0.485$

248 Subtidal: $\text{Log SMR} = 0.716 \text{ Log FW} - 0.512$

249 Intertidal: $\text{Log SMR} = 0.716 \text{ Log FW} - 0.661$

250 Similarly, covariance analyses performed on regression lines for metabolic rates (both
251 RMR and SMR) in relation to shell length (SL) (Figure 4 A,B) resulted in lack of
252 significant differences in slope but significant differences in intercepts between
253 intertidal and subtidal groups (Table 3). Therefore, common slopes were computed, and
254 a values recalculated for each population:

255 Subtidal: $\text{Log RMR} = 2.199 \text{ Log SL} - 4.308$

256 Intertidal: $\text{Log RMR} = 2.199 \text{ Log SL} - 4.410$

257 Subtidal: $\text{Log SMR} = 2.220 \text{ Log SL} - 4.424$

258 Intertidal: $\text{Log SMR} = 2.220 \text{ Log SL} - 4.618$

Metabolic scope for feeding and growth (MSFG)

MSFG was expressed as a percentage of routine metabolic rate since mass scaling exponents for both respiration rates were similar ($p > 0.05$) while intercepts were significantly different ($p < 0.005$) (Table 3). Thus, recalculated a values were used for this purpose. Therefore, MSFG represented 17.43% and 23.37% of routine rate in the subtidal population, in terms of flesh weight and shell length, respectively; whereas intertidal mussels showed a higher percentage of reduction, amounting to 33.33% and 38.05%, respectively.

Allometries of ammonia excretion rates

Relationships between ammonia excretion rate ($\text{VNH}_4\text{-N}$) and size were performed also by regression analyses (Fig. 5). Equations relating ammonia excretion to dry flesh weight (FW) (Fig. 5A) in intertidal and subtidal populations (Table 4) did not show significant differences between slopes. Hence, equations were recalculated according to their common slope:

$$\text{Subtidal: } \text{Log VNH}_4\text{-N} = 0.616 \text{ Log FW} + 1.217$$

$$\text{Intertidal: } \text{Log VNH}_4\text{-N} = 0.616 \text{ Log FW} + 1.138$$

Similarly, slopes of regressions relating ammonia excretion rate ($\text{VNH}_4\text{-N}$) to shell length (SL) (Fig. 5B) did not differ statistically between origins (Table 4). Thus, a common slope was calculated, recalculating then intercepts as the following form:

$$\text{Subtidal: } \text{Log VNH}_4\text{-N} = 1.910 \text{ Log SL} - 2.150$$

$$\text{Intertidal: } \text{Log VNH}_4\text{-N} = 1.910 \text{ Log SL} - 2.268$$

To assess a possible size-effect on the ratio of oxygen consumption to ammonia excretion (the O:N ratio), regression lines for RMR and VNH_4 vs. FW were compared

by ANCOVA (Table 5), resulting in absence of significant differences in slope for any population of mussels.

Discussion:

Biometry and condition index

The relevant amount of shell organics found in both populations suggests that the energy required for shell growth is not an insignificant portion of a bivalve's total energy budget, as stated previously by Jørgensen (1976); Rodhouse et al. (1984); Hawkins & Bayne (1985; 1992); Goulletquer & Wolowicz (1989); Wolowicz & Goulletquer (1999). Shell organic content (%) decreased with size in both subtidal and intertidal populations. Although subtidal mussels had higher levels of shell organic content (%), absolute shell organics (g) was only slightly lower in subtidal than in intertidal mussels due to the higher shell weight found in the latter group.

Subtidal mussels showed higher values of condition index (CI) than intertidal's. In some size classes (40–45 mm), subtidal values were about two-fold higher in relation to the intertidal ones. These results can be interpreted as indicative of a higher growth index (Smaal & Stralen, 1990; Pérez-Camacho et al., 1995). Pérez-Camacho et al. (1995) found similar results on *M. galloprovincialis*, which were attributed to the lesser feeding time in the intertidal population. In fact, use of energy reserves associated to reduced food availability have been reported in intertidal mussels (Freites et al., 2002), which could also explain the observed differences between populations concerning the CI in this experiment. On the other hand, broader fluctuations in flesh content of bigger individuals along the seasonal cycle are likely accounting for the increased variability recorded for CI in the largest size classes, since by the end of summer-early autumn some individuals are spawning while others are recovering from this event. The origin

of this variability would also account for greater CI fluctuation in subtidal mussels endowed with a thinner shell.

Both FW/SL and SW/SL ratios, as well as CI, increased with size in each population. However, subtidal mussels were characterized by a high flesh weight and low shell weight per unit shell length relative to those in the intertidal habitat. Thippeswamy & Joseph (1991; 1992) suggested that size of organisms is controlled by the ambient coupled with the population selection strategies. Thus, shell dimensions are influenced by the environmental conditions (Hemachandra & Thippeswamy, 2008). Our results confirm previous studies on habitat differences regarding condition and biometry in bivalves (Rao, 1953; Seed, 1973; Brown & Seed, 1977; Aldrich & Crowley, 1986). The higher shell thickness found in intertidal bivalves in contrast to their subtidal conspecifics could be explained as a protection strategy against the destructive effects of wave action (Fox & Coe 1943; Raubenheimer & Cook 1990; Akester & Martel 2000; Steffani & Branch, 2003). For instance, Akester & Martel (2000) found that mean shell thickness at a typical wave-exposed site was about 60% greater than at a sheltered site. It was also seen that some intertidal mussel species may increase shell thickness –subsequently decreasing growth rates– in response to predation (Leonard et al., 1999; Naddafi & Rudstam, 2014). The process of shell-thickening is thought to be mediated by increasing calcification (Brookes, 2006, Brookes & Rochette, 2007, Freeman, 2007) and would involve a decline in the percentage organic content of the shells (Brookes, 2006), as reported in the present work. Since 25 to 50% of the total body energy can be allocated to shell production (Jørgensen, 1976; Griffiths & King, 1979; Gardner & Thomas, 1987), thickening of shells can be considered to occur subjected to elevated metabolic costs. Thus, higher costs of shell production in intertidal mussels would account for slower growth whilst reduced condition (lower flesh weights) probably

reflects the poorer feeding conditions prevailing in this habitat (Aldrich & Crowley, 1986).

Allometric scaling of respiration rates to body size

Recorded values of size-scaling exponents for respiration in different species of the genus *Mytilus* (for review see Winter, 1978; Bayne & Newell, 1983) fall in the range 0.65 to 0.87, that encloses the average value (0.78) reported for bivalves (Glazier, 2005). Most these fluctuations in weight exponents are attributable to the experimental conditions under which determinations were performed, considering that some variables such as temperature or season differ among studies. Activity level of endogenous origin was an additional source of variation since these measurements combined routine as well as standard rates (for review see Bayne & Newell, 1983). This particular issue of a relationship between metabolic size-exponents and activity levels has been recently formalized by Glazier (2005) who put forward the metabolic level boundaries (MLB) hypothesis.

According to Griffiths & Griffiths (1987), allometric scaling exponents (b values) for metabolism in bivalves are subjected to minimal variations at the intraspecific level. MLB hypothesis, by contrast, indicates that b values would increase with activity level in ectothermic organisms (Glazier, 2009a). Results reported here revealed no differences in scaling exponents between routine and standard metabolic rates. Hence, it is possible that the increment in the activity level –from resting to active levels– is not high enough to achieve a significant change in scaling exponents, since many other studies (for review see Glazier, 2005; 2009a; Jensen et al., 2013) at the intraspecific level have proved significant differences in scaling exponents based on activity level.

As previously stated, this is the first report on allometries of respiration rates for *M. galloprovincialis* covering routine and standard levels and habitat differences. Scaling exponents obtained for routine (0.715) and standard metabolic rate (0.716) vs. dry flesh weight were similar to those estimated by Bayne et al. (1973) for *Mytilus edulis*. No comparable data have been reported for the allometric relationship of respiration rate and shell length (for review see Winter, 1978; Bayne & Newell, 1983; Glazier, 2005). As for weight exponents, scaling exponents for length were found similar between routine (2.199) and standard (2.220) metabolic rates. Regression analyses (Table 3) provided models that accounted for 88 and 92% of the variation on oxygen consumption based on shell length; which was exactly the same percentage of the variation than in models based on dry flesh weight. These results allow using shell length as an alternative to soft body weight in standardizing metabolic rates of mussels, which represents some advantages. As measuring shell length is neither an invasive nor destructive method, it makes possible to repeat measures throughout the time and allows researchers to work with endangered species, enabling reintroduction of the specimens in their habitat once measurements were concluded.

Regression intercepts (a values) have been reported to vary both among species and depending on experimental conditions; particularly temperature and activity level (Griffiths & Griffiths, 1987). Comparisons among different a values are frequently made under the assumption that these coefficients are mass-independent, and subsequently, considering that are independent of scaling exponents. As Carey et al. (2013) adequately described, “any alteration of the value of the slope b necessarily means that the intercept a will also change, and this confounds direct comparisons of metabolic level using this metric”; so in this report comparisons between intercepts of

regression lines are made on the assumption of lack of statistical significance of differences among allometric scaling exponents.

Unlike allometric exponents, a values exhibited significant effects associated to both habitat and activity level. Compared with subtidal mussels, intertidal specimens experienced an 11.8% reduction in routine metabolic rate and 22.5% in standard metabolic rate. Specific restrictions found in the intertidal habitat might account for these metabolic reductions: metabolic expenditure can amount up to 84% of the absorbed energy in bivalves (56% on average in *M. edulis*) (Bayne & Newell, 1983), so that adjustments in respiration rate can operate as an efficient mechanism for saving energy, especially under limiting conditions of food availability. In this respect, lower rates of respiration recorded in intertidal *M. galloprovincialis* has been associated to limitations in feeding time imposed by tidal cycles (Babarro et al. 2000b). Moreover, reduced resting demands are known to diminish resource to anaerobiosis during air exposure periods (Shick et al., 1988), and this may help understanding why metabolic depression reported here for intertidal mussel concerns mainly to the standard rather than routine rates (see above). In addition to restrictions imposed to metabolic expenditure in the intertidal habitat, differences in metabolic rate between mussels from both populations could be enhanced by the specific demands of an increased gametogenic activity in subtidal mussels, as can be inferred from their greater CI and higher lipid content (Freites et al., 2002).

Metabolic scope for feeding and growth (MSFG)

Intercepts from regression lines were significantly different for routine and standard rates in both populations of mussels and this allowed using recalculated a values for computing MSFG. When expressed in relative terms (as a fraction of RMR) this

metabolic scope can be considered a nearly constant amount size-independent on account of the lack of difference between the scaling exponents for both metabolic levels (either in terms of soft body weight or shell length). However, the magnitude of metabolic reduction experienced by starved mussels differed greatly between populations, from 17 to 33% of total metabolic costs (represented by RMR) in subtidal and intertidal mussels, respectively. These values are in the range reported for species of *Mytilus*: 26–45% in *M. edulis* (Bayne et al., 1989; Widdows & Hawkins, 1989) and 17–53% in *M. galloprovincialis* (Tamayo et al., submitted). Greater metabolic investments in growth processes exhibited by intertidal mussels contrasts with their reduced rates of growth suggesting that growth efficiency would be considerably reduced, a condition very likely associated to the elevated costs of shell production in the intertidal media that were previously considered.

Allometric scaling of ammonia excretion rates to body size

Information on allometric scaling of rates of ammonia excretion to body weight in bivalves is extremely scarce and no comparative data exist referred to shell length (for review see Griffiths & Griffiths, 1987). There was no evidence in this study of any effect of habitat on the scaling exponents of ammonia excretion rates *i.e.*, regression models obtained for subtidal and intertidal populations showed no effect on the scaling exponents. Bayne & Scullard (1977) reported variability both in habitat –rocky shore and estuary–and season on nitrogen excretion rates. Results reported in the present study concerning to habitat differences in a values are in agreement with Labarta et al. (1997), who also found higher ammonia excretion rates in the subtidal population under laboratory conditions.

Size exponents for metabolism and ammonia excretion were found no-significant for any metabolic level or habitat, implying that O:N indices were size-independent. This result agrees with reported data for *M. californianus* (Bayne et al., 1976a) but contrasts with previous information on *M. edulis* (Bayne et al., 1976b) where O:N index was found to increase or decrease with body size in resting or actively growing mussels, respectively.

Summarizing, results report neither effect of habitat nor effect of activity level on the allometric scaling exponent in both respiration and ammonia excretion rates. Allometric scaling exponents based on weight were 0.715 and 0.716 for the routine and standard respiration rates, respectively; while b values for the relationship between respiration rates and shell length amounted to 2.199 and 2.220, respectively. Allometric scaling exponents concerning ammonia excretion rates were 0.616 and 1.910 for weight and shell length, respectively. Origin differences found in respiration rates could reflect physiological compensations in the intertidal population for the lesser feeding time and air exposure. The higher CI registered in subtidal mussels suggests a greater energy budget than in intertidal mussels, despite the higher respiration and excretion rates of subtidal individuals. This suggestion is confirmed by higher feeding rates (Babarro et al., 2000a) and absorption efficiencies (Labarta et al., 1997) found for subtidal mussels in previous studies. The differences in MSFG between populations could explain differences in growth efficiencies; furthermore, the higher shell thickness found in the intertidal individuals suggests that energy resources are allocated as a priority to shell growth to the detriment of flesh growth.

Acknowledgements

448 The authors thank Lourdes Nieto and Beatriz González for their expert technical
449 assistance with the algae culture, physiological and chemical analyses. We thank P.
450 Markaide for his valuable contribution on the custom made R script. This work was
451 funded by the project FIGEBIV (AGL2013-49144-C3-2-R).

References

- Akester, R. J. & A. L. Martel, 2000. Shell shape, dysodont tooth morphology, and hinge–ligament thickness in the bay mussel *Mytilus trossulus* correlate with wave exposure. Canadian Journal of Zoology 78: 240–253.
- Aldrich, J. C. & M. Crowley, 1986. Condition and variability in *Mytilus edulis* (L.) from different habitats in Ireland. Aquaculture 52: 273–286.
- Anestis, A., H. O. Pörtner, D. Karagiannis, P. Angelidis, A. Staikou & B. Michaelidis, 2010. Response of *Mytilus galloprovincialis* (L.) to increasing seawater temperature and to marteliosis: Metabolic and physiological parameters. Comparative Biochemistry and Physiology, Part A 156: 57–66.
- Atanasov, A. T. & D. B. Dimitrov, 2002. Changes of the power coefficient in the ‘metabolism–mass’ relationship in the evolutionary process of animals. BioSystems 66: 65–71.
- Babarro, J. M. F., M. J. Fernández-Reiriz & U. Labarta, 2000a. Feeding behaviour of seed mussel *Mytilus galloprovincialis*: environmental parameters and seed origin. Journal of Shellfish Research 19: 195–201.
- Babarro, J. M. F., M. J. Fernández-Reiriz & U. Labarta, 2000b. Metabolism of the mussel *Mytilus galloprovincialis* from two origins in the Ría de Arousa (north–west Spain). Journal of the Marine Biological Association of the United Kingdom 80: 865–872.
- Babarro, J. M. F., U. Labarta, & M. J. Fernández-Reiriz, 2003. Growth patterns in biomass and size structure of *Mytilus galloprovincialis* cultivated in the “Ría de

474 **Arousa”** (north-west Spain). Journal of the Marine Biological Association of the United
 475 Kingdom 83: 151–158.

476 Bayne, B. L., C. J. Bayne, T. C. Carefoot & R. J. Thompson, 1976a. The physiological
 477 ecology of *Mytilus californianus* Conrad. Oecologia 22: 229–250.

478 Bayne, B. L. & R. C. Newell, 1983. Physiological energetics of marine molluscs. In:
 479 Saleuddin, A. S. M. & K. M. Wilbur (eds.), The Mollusca, Vol. 4. Physiology, Part 1.
 480 Academic Press, New York: 407–515.

481 Bayne, B. L. & C. J. Scullard, 1977. Rates of nitrogen excretion by species of *Mytilus*
 482 (Bivalvia: Mollusca). Journal of the Marine Biological Association of the United
 483 Kingdom 57: 355–369.

484 Bayne, B. L., R. J. Thompson & J. Widdows, 1973. Some effects of temperature and
 485 food on the rate of oxygen consumption by *Mytilus edulis* L. In: Weiser W. (ed.),
 486 Effects of temperature on ectothermic organisms, Berlin: Springer-Verlag: 181–193.

487 Bayne, B. L., A. J. S. Hawkins & E. Navarro, 1988. Feeding and digestion in
 488 suspension-feeding bivalve mollusks – the relevance of physiological compensations.
 489 American Zoologist 28: 147 –159.

490 Bayne, B. L., A. J. S. Hawkins, E. Navarro & J. I. P. Iglesias, 1989. Effects of seston
 491 concentration on feeding, digestion and growth in the mussel *Mytilus edulis*. Marine
 492 Ecology Progress Series 55: 47–54.

493 Bayne, B. L., J. Widdows & A. Thompson, 1976b. Physiological integrations. In:
 494 Bayne, B. L. (ed.), Marine mussels: their ecology and physiology. Cambridge
 495 University Press, Cambridge, first edition: 261–291.

496 Bokma, F., 2004. Evidence against universal metabolic allometry. *Functional Ecology*
 497 18: 184–187.

498 Borrero, F. J. & T. J. Hilbish, 1988. Temporal variation in shell and soft tissue growth
 499 of the mussel *Geukensia demissa*. *Marine Ecology Progress Series* 42: 9–15.

500 Brookes, J. I. & R. Rochette, 2007. Mechanism of a plastic phenotypic response:
 501 predator-induced shell thickening in the intertidal gastropod *Littorina obtusata*. *Journal*
 502 *of Evolutionary Biology* 20: 1015–1027.

503 Brookes, J. I., 2006. Proximate control and microstructure characteristics of predator–
 504 induced shell thickening in the intertidal gastropod *Littorina obtusata*. University of
 505 New Brunswick, Canada. 84 pp.

506 Brown, R. A. & R. Seed, 1977. *Modiolus modiolus*– an autecological study. In: Keegan,
 507 B. F., P. O’Ceidigh & P. J. S. Boaden, (eds) *Proc 11th Eur. Mar. Biol. Symp.*, Pergamon
 508 Press, Oxford: 93–100.

509 Carey, N., J. D. Sigwart & J. G. Richards, 2013. Economies of scaling: More evidence
 510 that allometry of metabolism is linked to activity, metabolic rate and habitat. *Journal of*
 511 *Experimental Marine Biology and Ecology* 439: 7–14.

512 Dame, R. F., 2012. *Ecology of marine bivalves: an ecosystem approach*. CRC Press.
 513 Boca Raton. 2nd edition. 260 pp.

514 Dickie, L. M., P. R. Boudreaup & F. R. Freeman, 1984. Influences of stock and site on
 515 growth and mortality in the blue mussel (*Mytilus edulis*). *Canadian Journal of Fisheries*
 516 *and Aquatic Sciences* 41: 225–236

517 Duarte, P., M. J. Fernández-Reiriz, R. A. Filgueira & U. Labarta, 2010. Modelling
 518 mussel growth in ecosystems with low suspended matter loads. *Journal of Sea Research*
 519 64: 273–286.

520 Elvin, O. W. & J. J. Gonor, 1979. The thermal regime of an intertidal *Mytilus*
 521 *californianus* Conrad population on the central Oregon coast. *Journal of Experimental*
 522 *Marine Biology and Ecology* 39: 265–279.

523 Fox, D. L. & W. R. Coe, 1943. Biology of the Californian sea mussel (*Mytilus*
 524 *californianus*). II. Nutrition, metabolism, growth and calcium deposition. *Journal of*
 525 *Experimental Zoology* 93: 203–249.

526 Freeman, K. R., 1974. Growth, mortality and seasonal cycle of *Mytilus edulis* in two
 527 Nova Scotian embayments. Department of the Environment, Fisheries and Marine
 528 Service, Canada, Technical Report N° 500: 1–112.

529 Freeman, A. S., 2007. Specificity of induced defenses in *Mytilus edulis* and
 530 asymmetrical predator deterrence. *Marine Ecology Progress Series* 334: 145–153.

531 Freitas, L., M. J. Fernández-Reiriz & U. Labarta, 2002. Lipid classes of mussel seeds
 532 *Mytilus galloprovincialis* of subtidal and rocky shore origin. *Aquaculture* 207: 97–111.

533 Gardner, J. P. A. & M. L. H. Thomas, 1987. Growth, mortality and production of
 534 organic matter by a rocky intertidal population of *Mytilus edulis* in the Quoddy Region
 535 of the Bay of Fundy. *Marine Ecology Progress Series* 39: 31–36.

536 Glazier, D. S., 2005. **Beyond the “3/4-power law”: Variation in the intra- and**
 537 **interspecific scaling of metabolic rate in animals.** *Biological Reviews of the Cambridge*
 538 *Philosophical Society* 80: 611–662.

539 Glazier, D. S., 2008. Effects of metabolic level on the body-size scaling of metabolic
540 rate in birds and mammals. *Proceedings of the Royal Society of London B* 275: 1405–
541 1410.

542 Glazier, D. S., 2009a. Activity affects intraspecific body-size scaling of metabolic rate
543 in ectothermic animals. *Journal of Comparative Physiology B* 179: 821–828.

544 Glazier, D. S., 2009b. Metabolic level and size-scaling of rates of respiration and
545 growth in unicellular organisms. *Functional Ecology* 23: 963–968.

546 Glazier, D. S., 2009c. Ontogenetic body-mass scaling of resting metabolic rate covaries
547 with species-specific metabolic level and body size in spiders and snakes. *Comparative*
548 *Biochemistry and Physiology. Part A* 153: 403–407.

549 Glazier, D. S., 2010. A unifying explanation for diverse metabolic scaling in animals
550 and plants. *Biological Reviews* 85: 111–138.

551 Gosling, E., 2003. *Bivalve molluscs: biology, ecology and culture*. Oxford, UK:
552 Blackwell Science. 443 pp.

553 Gouletquer, O. & M. Wollowitz, 1989. The shell of *Cardium edule*, *Cardium glaucum*
554 and *Ruditapes philippinarum*. Organic content, composition and energy value as
555 determined by different methods. *Journal of the Marine Biological Association of the*
556 *United Kingdom* 69: 563–572.

557 Griffiths, C. L. & R. J. Griffiths, 1987. Bivalvia. In: *Animal Energetics*, Vol. 2, Bivalvia
558 through Reptilia (eds T.J. Pandian & F. J. Vernberg), Academic Press, California: 1–87.

559 Griffiths, C. L. & J. A. King, 1979. Some relationship between size, food availability
560 and energy balance in the ribbed mussel *Aulacomya ater*. *Marine Biology* 51: 141–149.

561 Hawkins, A. J. S. & B. L. Bayne, 1985. Seasonal variation in the relative utilization of
 562 carbon and nitrogen by the mussel *Mytilus edulis*: Budgets, conversion efficiencies and
 563 maintenance requirements. Marine Ecology Progress Series 25: 181–188.

564 Hawkins, A. J. S. & B. L. Bayne, 1992. Physiological interrelations, and the regulation
 565 of production. In: Gosling, E. (ed.), The mussel *Mytilus*: ecology, physiology, genetics
 566 and culture. Elsevier Science, Amsterdam: 171–222.

567 Hemachandra, D. & S. Thippeswamy, 2008. Allometry and condition index in green
 568 mussel *Perna viridis* (L.) from St Mary's Island off Malpe, near Udupi, India.
 569 Aquaculture Research 39: 1747–1758.

570 Hilbish, T. J., 1986. Growth trajectories of shell and soft tissue in bivalves: seasonal
 571 variation in *Mytilus edulis* L. Journal of Experimental Marine Biology and Ecology 96:
 572 103–113.

573 Jensen, M., Q. P. Fitzgibbon, C. G. Carter & L. R. Adams, 2013. Effect of body mass
 574 and activity on the metabolic rate and ammonia-N excretion of the spiny lobster
 575 *Sagmariasus verreauxi* during ontogeny. Comparative Biochemistry and Physiology
 576 Part A 166: 191–198.

577 Jørgensen, C. B., 1976. Growth efficiencies and factors controlling size in some mytilid
 578 bivalves, especially *Mytilus edulis* L.: review and interpretation. Ophelia 15: 175–192.

579 Labarta, U., M. J. Fernandez-Reiriz & J. M. F. Babarro, 1997: Differences in
 580 physiological energetics between intertidal and raft cultivated mussels *Mytilus*
 581 *galloprovincialis*. Marine Ecology Progress Series 152: 167 –173.

582 Labarta U., M. J. Fernández-Reiriz, A. Pérez-Camacho & A. P. Corbacho, 2004.
 583 Bateiros, mar, mejillón. Una perspectiva bioeconómica. Centro de Investigaciones

584 Económicas y Financieras (CIEF). Fundación CaixaGalicia. Santiago de Compostela.
585 274 pp.

586 Leonard, G. H., M. D. Bertness & P. O. Yund, 1999. Crab predation, waterborne cues,
587 and inducible defenses in the blue mussel, *Mytilus edulis*. Ecology 80: 1–14.

588 Mallet, A. L., C. E. A. Carver, S. S. Coffen & K. R. Freeman, 1987. Winter growth of
589 the blue mussel *Mytilus edulis* L.: importance of stock and site. Journal of Experimental
590 Marine Biology and Ecology 108: 217–228.

591 Marsden, I. D. & M. A. Weatherhead, 1999. Shore level induced variations in condition
592 and feeding of the mussel *Perna canaliculus* from the east coast of the South Island,
593 New Zealand. N. Z. J. Marine Freshwater Research 33: 611–622.

594 Muller-Landau, H. C., R. S. Condit, J. Chave, S. C. Thomas, S. A. Bohlman, S.
595 Bunyavejchewin, S. Davies, R. Foster, S. Gunatilleke, K. E. Harms, T. Hart, S. P.
596 Hubbell, A. Itoh, A. R. Kassim, J. V. La Frankie, H. S. Lee, E. Losos, J. R. Makana, T.
597 Ohkubo, R. Sukumar, I. F. Sun, N. Supardi, S Tan, J. Thompson, R. Valencia, G. V.
598 Muñoz, C. Wills, T. Yamakura, G. Chuyong, H. S. Dattaraja, S. Esufali, P. Hall, C.
599 Hernandez, D. Kenfack, S. Kiratiprayoon, H. S. Suresh, D. Thomas, M. I. Vallejo, P.
600 Ashton, 2006. Testing metabolic ecology theory for allometric scaling of tree size,
601 growth and mortality in tropical forests. Ecology Letters 9: 575–588.

602 Naddafi, R. & L. G. Rudstam, 2014. Predator-induced morphological defences in two
603 invasive dreissenid mussels: implications for species replacement. Freshwater Biology
604 59: 703–713.

605 Navarro, E., J. I. P. Iglesias, A. P. Camacho, U. Labarta & R. Beiras, 1991. The
 606 physiological energetics of mussels (*Mytilus galloprovincialis* Lmk.) from different
 607 cultivation rafts in the Ria de Arosa (Galicia N.W. Spain). *Aquaculture* 94: 197–212.

608 Okumuş, I. & H. P. Stirling, 1994. Physiological energetics of cultivated mussel
 609 (*Mytilus edulis*) populations in 2 Scottish west coast sea lochs. *Marine Biology* 119:
 610 125–131.

611 Pérez-Camacho A., E. Aguiar, U. Labarta, V. Vinseiro, M. J. Fernández-Reiriz & X. A.
 612 Álvarez-Salgado, 2014. Ecosystem-based indicators as a tool for mussel culture
 613 management strategies. *Ecological Indicators* 45: 538–548.

614 Pérez-Camacho, A., U. Labarta & R. Beiras, 1995. Growth of mussels (*Mytilus edulis*
 615 *galloprovincialis*) on cultivation rafts: influence of seed source, cultivation site and
 616 phytoplankton availability. *Aquaculture* 138: 349–362.

617 Pérez-Camacho, A., U. Labarta, V. Vinseiro & M. J. Fernandez-Reiriz, 2013. Mussel
 618 production management: raft culture without thinning-out. *Aquaculture* 406–407: 172–
 619 179.

620 Peterson, C. H. & R. Black, 1988. Responses of growth to elevation fail to explain
 621 vertical zonation of suspension-feeding bivalves on a tidal flat. *Oecologia* 76: 423–429.

622 Prosser, C. L., 1973. Oxygen: respiration and metabolism. In: Prosser, C. L. (ed.),
 623 *Comparative Animal Physiology*, vol. 1. Saunders College, Philadelphia: 165–211.

624 Rao, K. P., 1953. Shell weight as a function of intertidal height in a littoral population
 625 of pelecypods. *Experientia* 9: 465–466.

626 Raubenheimer, D. & P. Cook, 1990. Effects of exposure to wave action on allocation of
 627 resources to shell and meat growth by the subtidal mussel, *Mytilus californianus*.
 628 Journal of Shellfish Research 9: 87–93.

629 Rawson, P. D. & T. J. Hilbish, 1991. Genotype–environment interaction for juvenile
 630 growth in the hard clam *Mercenaria mercenaria* (L.). Evolution 45: 1924–1935.

631 Reich, P. B., M. G. Tjoelker, J. L. Machado & J. Oleksyn, 2006. Universal scaling of
 632 respiratory metabolism, size and nitrogen in plants. Nature 439: 457–461.

633 Riisgård, H. U., 1998. **No foundation of a ‘3/4 power scaling law’ for respiration in**
 634 **biology.** Ecology Letters 1: 71–73.

635 Rodhouse, P. G., C. M. Roden, M. P. Hensey & T. H. Ryan, 1984. Resource allocation
 636 in *Mytilus edulis* on the shore and in suspended culture. Marine Biology 84: 27– 34.

637 Sarà, G. & A. Pusceddu, 2008. Scope for growth of *Mytilus galloprovincialis* (Lmk.,
 638 1819) in oligotrophic coastal waters (Southern Tyrrhenian Sea, Italy). Marine Biology
 639 156: 117–126.

640 Seed, R. 1973. Absolute and allometric growth in the mussel, *Mytilus edulis* L.
 641 (Mollusca Bivalvia). Proceedings of the Malacological Society of London 40: 343–357.

642 Shick, J. M., J. Widdows & E. Gnaiger, 1988. Calorimetric studies of behaviour,
 643 metabolism and energetics of sessile intertidal animals. American Zoologist 28: 161–
 644 181.

645 Smaal, A. C. & M. R. Stralen, 1990. Average annual growth and condition of mussel as
 646 a function of food source. Hydrobiologia 195: 179–188.

647 Solórzano, L., 1969. Determination of ammonia in natural waters by the
 648 phenolphthalein method. Limnology and Oceanography 14: 779–801.

649 Steffani, C. N. & G. M. Branch, 2003. Growth rate, condition, and shell shape of *Mytilus*
 650 *galloprovincialis*: responses to wave exposure. Marine Ecology Progress Series 246:
 651 197–209.

652 Storey, K. B. & J. M. Storey, 1990. Metabolic rate depression and biochemical
 653 adaptation in anaerobiosis, hibernation and estivation. Quarterly Review of Biology 65:
 654 145–174.

655 Tamayo, D., 2012. Physiological basis for inter-individual differences in growth among
 656 bivalve molluscs. PhD thesis. University of the Basque Country (UPV/EHU), Bilbao,
 657 Spain, 183 pp.

658 Tamayo D., K. Azpeitia, E. Navarro & I. Ibarrola, Submitted. Food regime modulates
 659 physiological processes underlying size-differentiation in juvenile mussels *Mytilus*
 660 *galloprovincialis* from an intertidal population.

661 Tamayo, D., I. Ibarrola & E. Navarro, 2013. Thermal dependence of clearance and
 662 metabolic rates in slow- and fast-growing spat of Manila clam *Ruditapes*
 663 *philippinarum*. Journal of Comparative Physiology B 183: 893–904.

664 Thippeswamy, S. & M. M. Joseph, 1991. Population selection strategies in the wedge
 665 clam *Donax incarnates* (Gmelin) from Panambur beach. Indian Journal of Marine
 666 Sciences 20: 147–151.

667 Thippeswamy, S. & M. M. Joseph, 1992. Allometry in the wedge clam *Donax*
 668 *incarnates* (Gmelin) from Panambur beach. Indian Journal of Marine Sciences 21: 147–
 669 149.

670 White, C. R., N. F. Phillips & R. S. Seymour, 2006. The scaling and temperature
671 dependence of vertebrate metabolism. *Biology Letters* 2: 125–127.

672 White, C. R., 2011. Allometric estimation of metabolic rates in animals. *Comparative*
673 *Biochemistry and Physiology Part A* 158: 346–57.

674 Widdows, J., P. Donkin, P. N. Salkeld, J. J. Cleary, D. M. Lowe, S. V. Evans & P. E
675 Thompson, 1984. Relative importance of environmental factors in determining
676 physiological differences between two populations of mussels (*Mytilus edulis*). *Marine*
677 *Ecology Progress Series* 17: 33–48.

678 Widdows, J. & A. J. S. Hawkins, 1989. Partitioning of rate of heat dissipation by
679 *Mytilus edulis* into maintenance feeding, and growth components. *Physiological*
680 *Zoology* 62: 764–784.

681 Winter, J. E., 1978. A review on the knowledge of suspension-feeding in
682 lamellibranchiate bivalves, with special reference to artificial aquaculture systems.
683 *Aquaculture* 13: 1–33.

684 Withers, P. C., 1992. *Animal Energetics. Comparative Animal Physiology.* Saunders
685 College Publishing, Fort Worth: 82–121.

686 Wolowicz, M. & P. Goulletquer, 1999. The shell organic content in the energy budget
687 of *Mytilus trossulus* from the South Baltic. *Haliotis* 28: 1–10.

688 Zar, J. H., 1996. *Biostatistical analysis.* 3rd ed. Prentice-Hall, Englewood Cliffs, USA.
689 718 pp.