TITLE: New large threatened populations of *Phengaris nausithous* discovered in the SW of Europe

AUTHORS: Fernando Jubete ¹ Jacinto Román ²

1. Asociación de Naturalistas Palentinos. C/ Vega, 7, E-34337, Fuentes de Nava, Palencia (Spain).

2. Department of Conservation Biology. Doñana Biological Station (CSIC), C/ Américo Vespucio
s/n, E-41092, Seville (Spain).

Corresponding author:

Jacinto Román

jroman@ebd.csic.es

ABSTRACT

The near threatened *Phengaris nausithous* is distributed in two large mountainous areas in the northern part of the Iberian Peninsula, living in small dispersed populations. During July and August of 2012-2015 we systematically searched for the species in the southeastern part of the Cordillera Cantábrica. We found 38 new populations grouped into 7 metapopulations. Two of the latter are the largest known, representing 34% of the species' occurrence range in Spain. The discovery of these new metapopulations does not improve the conservation status of *P. nausithous* in the SW of Europe because a land consolidation project currently underway and an established peat extraction plant could lead to the imminent extinction of the largest metapopulations. However, there is still time to reverse the present course of change. Knowledge of these recently discovered populations allows the establishment of management measures to protect them over the long term.

KEYWORDS

butterfly, conservation, near threatened, land consolidation, peat extraction

Introduction

Butterfly species within the genus *Phengaris* Doherty, 1891, have complex and highly specialized breeding systems, making them very susceptible to factors adversely affecting their habitat (Kotiaho et al. 2005). The near threatened *Phengaris nausithous* (Bergsträsser, 1779) (World Conservation Monitoring Centre 1996; van Swaay et al. 2010) lays eggs inside the flowerheads of *Sanguisorba officinalis*, its exclusive hostplant. When larvae subsequently leave the plant, they are adopted by foraging ants of the genus *Myrmica* and spend the autumn, winter and spring inside the ants' nest, feeding on ant larvae and pupae (Thomas 1984). In summer, during July and August, *P. nausithous* individuals fly away as imagoes (García-Barros et al. 2013). Suitable habitat patches include wet and periodically flooded meadows and riverbanks, in which their host plant and host ant are found (Munguira and Martín 1999).

P. nausithous is typically distributed in metapopulations, with small populations occupying discrete habitat patches (Thomas 1984; Nowicki et al. 2005a; 2007). The average life span is only 2-3 days in the field (Nowicki et al. 2005a). Their mobility is low (commonly within a few hundred metres), restricted by unsuitable habitat, mainly forests, croplands and urban areas (Hovestadt et al. 2011; Nowicki et al. 2014). The dispersal rate is very low, but regular disturbance by land use and high density might lead to increased dispersal (Nowicki et al. 2005a; 2007; Dover and Settele 2009).

In Europe, the southernmost populations are located in Spain, where it is listed as vulnerable in the Spanish Catalog of Threatened Species (Real Decreto 139/2011). The species is distributed in two large mountainous areas in the northern part of the Iberian Peninsula (García-Barros et al. 2013). In recent years species surveys have been conducted in many parts of the Spanish distribution range (Munguira et al. 2001; 2011; Manceñido-González and González-Estébanez 2013; Vicente et al. 2013). Previous data for Spain have been detailed in standard 10x10 km or 1x1 km UTM grids (10 000 ha and 100 ha grids, respectively), both at coarser size resolutions than the size of occupied patches (Vicente et al. 2013). Additionally, models on the potential distribution of the species are already available (Romo et al. 2006; 2015; Jiménez-Valverde et al. 2008), and according to several studies their current distribution in Spain will be strongly affected by climatic change (Settele et al. 2008; Schweiger et al. 2012; Romo et al. 2015).

Here, we present the recent discovery of the largest populations of *P. nausithous* in Spain, and show that there are imminent threats that hinder the conservation of these important populations.

Material and Methods

We searched for the species during July and August of 2012-2015 in the north of Palencia and Burgos provinces (south and east of the Cordillera Cantábrica, Fig. 1). We defined habitat patches on the basis of land use and vegetation composition: suitable meadows limited by forests, bushes or urban areas. These meadows are normally located at the low parts of mountain valleys. We used Google Earth and Google Street View to systematically identify potential sites that were later visited. Once in the patch, we searched for *S. officinalis*, which, in the study region, typically grows dispersed or in sparse stands. In each patch with the presence of flowering foodplants, we conducted transect surveys visiting all the identified stands of *S. officinalis*. We considered the species as absent if it was not detected. The detectability of *P. nausithous* in open habitats is high (Pellet et al. 2012) and it is considered that, in occupied patches, flying individuals are seen within the first five minutes of survey in *S. officinalis* stands (Jimenez-Valverde et al. 2008). We define a population as an occupied patch of continuous suitable habitat. We consider the number of records of *P. nausithous* found in a meadow as a proxy of population size. We define populations located less than 5 km apart as the same metapopulation because this is the maximum recorded displacement of *P. nausithous* (Nowicki et al. 2005b). We calculated the distance between populations as the Euclidean distance between the centroids of occupied patches. To allow comparison with previous studies made in Spain we also used grids of both 10x10 km and 1x1 km.

Results

We found several new locations where the species is present. In Palencia Province we obtained 568 individual records, distributed in 34 populations, grouped into five metapopulations (Figure 1; Table 1). The largest metapopulation is located in the upper basin of the Pisuerga River. This metapopulation has the lowest mean distance between populations (Table 1). In this province, the new records occupy 40 cells of the 1x1 km grid and 9 cells of the 10x10 km grid, in 4 of which the species had not been previously recorded. In Burgos Province, we obtained 53 individual records, distributed in four populations, grouped into two metapopulations. Nine of these records belong to the three populations of the Tozo-Loras metapopulation which shows the greatest mean inter-population distance (Table 1). In two of these populations we found only one individual, whereas the third population occupies three grid cells of 1x1 km. The remaining 44 records belong to the Nava River basin where we found a single large population occupying 10 grid cells of 1x1 km. In all populations, most records are confined to vegetation adjacent to streams and waterlogged peatlands.

Discussion

Our study provides an update to the distribution of *P. nausithous* in the southwest of Europe. This information indicates that the species is distributed in dispersed and small populations (Vicente et al. 2013). It is not easy to compare the new data with previously known sites because no common resolution and terminology were employed. Munguira et al. (2011) used a 1x1 km grid as reference, recording the study species in 41 grid cells. In the most recent review, Vicente *et al.* (2013) used both locality and colony alike, mentioning 15 new small and isolated populations. Only two were larger than 5 ha, while 8 were smaller than 1 ha (without specifying how area was calculated). Assuming that each population represents a single grid cell at a 1x1 km resolution, the previously known distribution of *P. nausithous* in Spain would cover 56 grid cells. The 55 new cells described in the present study nearly double the previous estimations.

Previous data also indicated a high level of isolation: the 56 known 1x1 km grid cells were distributed in 36 10x10 km grid cells (Munguira et al. 2011; Vicente et al. 2013). In our study, aggregation is higher in both the Pisuerga metapopulation, with 28 1x1 km grid cells occupying four 10x10 km grid cells, and the Nava basin, with 10 1x1 km grid cells present in a single 10x10 grid cell. Measured at the 1x1 km resolution the two metapopulations represent 34% of the area occupied by the species in Spain.

Unfortunately, we do not think that the discovery of these new populations represents an improvement in the conservation status of the species in Spain. There are serious threats affecting them that could lead to their imminent extinction. The Pisuerga, Carrión and Otero de Guardo metapopulations are completely included in Fuentes Carrionas and Fuente Cobre Natural Park, which is part of the Natura 2000 network. In this protected area a land consolidation plan has been approved for the whole area. If implemented, all the pastures and croplands will be rearranged at the landscape scale in order to increase their average size, including the removal of hedgerows, the levelling-off of the new patches of land, the eventual drainage of pastures and river banks, and the canalization of streams (by straightening them and lowering their beds), effectively destroying the habitat of *P. nausithous*. The project has already been implemented in San Salvador de Cantamuda municipality, where, in spite of its potential, the species is absent. The remaining localities are waiting for funds to be allocated, mainly by the European Regional Development Fund, to undertake the land consolidation.

Likewise, the habitat of the Nava basin population is being destroyed by the extraction of peat (permit granted in the 1990s) across a large area of the habitat patch occupied by the population. To date, about 40 ha of habitat have been directly destroyed, and the surrounding habitat belt is being drained.

These threats make the largest populations recently discovered at serious risk of extinction in the near future. However, we still have time to reverse the present course of change. This work should allow the rethinking of land consolidation projects as well as the inclusion of measures allowing for the conservation and management of populations of *P. nausithous*. The most important action is to avoid the drainage of pastures and surrounding areas, or other works that could adversely affect the conservation of *P. nausithous* populations (for example, the dumping of waste products resulting from work projects, repairs and new construction of service roads near populations, etc.). If this is not possible, a key action is to halt the funding of land consolidation by the European Union within the Natural Park of Fuentes Carrionas.

In the population of the Nava basin, it is necessary to check whether the peat extraction grant complies with the Habitats Directive as well as other Spanish and European legislation.

Knowledge of these recently discovered populations, allows the establishment of management measures to protect them over the long term.

Acknowledgments

Inventory work in Palencia was funded by the project “Colony inventory, conservation actions and stewardship for the dusky large butterfly (*Phengaris nausithous*) in the natural park of Fuentes Carrionas and Fuente Cobre-Montaña Palentina” (Fundación Biodiversidad; Ministry of Agriculture, Food and Environment). Permission for sampling was obtained from the Junta de Castilla y León. We would like to thank Carlos Rodríguez and Eloy Revilla for reviewing an earlier version of the manuscript. Piotr Nowicki and an anonymous referee reviewed a draft version of the manuscript. English corrections to the original text were made by P. James Macaluso.

References

- Dover J, Settele J (2009) The influences of landscape structure on butterfly distribution and movement: a review. *J Insect Conserv*, 13:3-27. doi: 10.1007/s10841-008-9135-8
- García-Barros E, Munguira ML, Stefanescu C, Vives-Moreno A (2013) *Lepidoptera Papilionoidea*. In: Ramos MA (Ed.) *Fauna Ibérica*, Vol 37. Museo Nacional de Ciencias Naturales. CSIC. Madrid. 1213 pp.

150 Hovestadt T, Binzenhöfer B, Nowicki P, Settele J (2011) Do all inter-patch movements represent
 151 dispersal? A mixed kernel study of butterfly mobility in fragmented landscapes. *J Anim Ecol*, 80: 1070-
 152 1077. doi: 10.1111/j.1365-2656.2011.01848.x
 153 Jiménez-Valverde A, Gómez JF, Lobo JM, Baselga A, Hortal J (2008) Challenging species distribution
 154 models: the case of *Maculinea nausithous* in the Iberian Peninsula. *Ann Zool Fennici* 45: 200–210. doi:
 155 10.5735/086.045.0305
 156 Kotiaho JS, Kaitala V, Komonen A, Päävinen J (2005) Predicting the risk of extinction from shared
 157 ecological characteristics. *PNAS*, 102 (6): 1963-1967. doi: 10.1073/pnas.0406718102
 158 Manceñido-González DC, González-Estébanez FJ (2013) Mariposas diurnas de la provincia de León.
 159 León. Ed. El Búho Viajero. 653 pp.
 160 Munguira ML, Martin J (1999) Action plan for *Maculinea* butterflies in Europe. Nature and environment,
 161 No. 97, Council of Europe Publishing. Strasbourg Cedex.
 162 Munguira ML, Martín J, Orueta D, Viejo JL, García-Barros E (2001) *Maculinea nausithous*
 163 (Bergsträsser, 1779). In: Ramos M, Bragado D, Fernández J (Eds.) Los invertebrados no insectos de la
 164 “Directiva Hábitat” en España. Organismo Autónomo de Parques Nacionales, Ministerio de Medio
 165 Ambiente, Madrid. pp.: 163-173.
 166 Munguira ML, Romo H, Martín J, García-Barros E (2011) *Phengaris nausithous* (Bergsträsser, 1779). In:
 167 Verdú JR, Numa C, Galante E (eds) Atlas y Libro Rojo de los Invertebrados amenazados de España
 168 (Especies Vulnerables). Dirección General de Medio Natural y Política Forestal, Ministerio de Medio
 169 Ambiente, Medio rural y Marino, Madrid. pp: 1258-1264.
 170 Nowicki P, Witek M, Skórka P, Settele J, Woyciechowski M (2005a) Population ecology of the
 171 endangered butterflies *Maculinea teleius* and *M. nausithous* and the implications for conservation. *Popul*
 172 *Ecol*, 47: 193-202. doi: 10.1007/s10144-005-0222-3
 173 Nowicki P, Settele J, Thomas JA, Woyciechowski M (2005b) A review of population structure of
 174 *Maculinea* butterflies. In: Settele J, Kühn E, Thomas JA (Eds) Studies on the Ecology and Conservation
 175 of Butterflies in Europe. Vol. 2: Species Ecology along a European Gradient: *Maculinea* Butterflies as a
 176 model. PENSOFT Publishers. Sofia-Moscow. pp: 144-149
 177 Nowicki A, Pepkowska A, Kudlek J, Skórka P, Witek M, Settele J, Woyciechowski M (2007) From
 178 metapopulation theory to conservation recommendations: Lessons from spatial occurrence and abundance
 179 patterns of *Maculinea* butterflies. *Biol Cons*, 140: 119-129. doi:10.1016/j.biocon.2007.08.001
 180 Nowicki P, Vrabec V, Binzenhöfer B, Feil J, Zaksek B, Hovestadt T, Settele J (2014) Butterfly dispersal
 181 in inhospitable matrix: rare, risky, but long-distance. *Landscape Ecol*, 29: 401-412. doi:10.1007/s10980-
 182 013-9971-0
 183 Pellet J, Bried JT, Parietti D, Gander A, Heer PO, Cherix D, Arlettaz R (2012) Monitoring Butterfly
 184 Abundance: Beyond Pollard Walks. *PLoS ONE* 7(7): e41396. doi:10.1371/journal.pone.0041396

- Romo H, García-Barros E, Munguira ML (2006) Distribución potencial de trece especies de mariposas diurnas amenazadas o raras en el área ibero-balear (Lepidoptera: Papilionoidea & Hesperioidea). Bol. Asoc. Esp. Ent., 30 (3-4): 25-49
- Romo H, Silvestre M, Munguira ML (2015) Potential distribution models and the effect of climatic change on the distribution of *Phengaris nausithous* considering its food plant and host ants. J Insect Conserv 19: 1101-1118. doi 10.1007/s10841-015-9825-y
- Schweiger O, Heikkinen RK, Harpke A, Hickler T, Klotz S, Kudrna O, Kühn I, Pöyry J, Settele J (2012) Increasing range mismatching of interacting species under global change is related to their ecological characteristics. Global Ecol. Biogeogr, 21, 88-99. doi: 10.1111/j.1466-8238.2010.00607.x
- Settele J, Kudrna O, Harpke A, Kühn I, van Swaay C, Verovnik R, Warren M, Wiemers M, Hanspach J, Hickler T, Kühn E, van Halder I, Veling K, Vliegenthart A, Wynhoff I, Schweiger O (2008) Climatic Risk Atlas of European Butterflies. PENSOFT Publishers. Sofia-Moscow. 710 pp.
- Thomas JA (1984) The behaviour and habitat requirements of *Maculinea nausithous* (the dusky large blue butterfly) and *M. teleius* (the scarce large blue) in France. Biol Cons, 28(4): 325-347
- Van Swaay C, Cuttelod A, Collins S, Maes D, López-Munguira M, Šašić M, Settele J, Verovnik R, Verstrael T, Warren M, Wiemers M, Wynhoff I (2010) European red list of butterflies. Publications Office of the European Union, Luxembourg. 58 pp.
- Vicente JC, Salvador V, Alcalde J, Parra B (2013) Ampliación de la distribución de *Phengaris nausithous* (Bergstrasser, 1779) (Lepidoptera: Lycaenidae) en la Península Ibérica, y algunas consideraciones para su conservación. Boletín de la Sociedad Entomológica Aragonesa, 52: 249-258
- World Conservation Monitoring Centre (1996) *Phengaris nausithous*. The IUCN Red List of Threatened Species 1996. doi: 10.2305/IUCN.UK.1996.RLTS.T12662A3371835. Downloaded on 23 September 2015.

Metapopulation		Province	Number of populations	Inter-population distance (m)	
				Mean± SD	Range (m)
A	Guardo	Palencia	2	1149	-
B	Otero de Guardo	Palencia	1	-	-
C	Carrión	Palencia	5	1626±202	1407-1821
D	Pisuerga	Palencia	25	837±338	360-1415
E	Quintanaluengos	Palencia	1	-	-
F	Tozo-Loras	Burgos	3	3607±986	3038-4745
G	Nava basin	Burgos	1	-	-

Table 1 Number of populations in each metapopulation and mean nearest distance between the nearest neighbouring populations. The range represents the shortest and the longest nearest distances in each metapopulation.

Fig. 1 a) Distribution of *P. nausithous* in Spain (standard 10x10 km UTM). Empty bold cells correspond to published localities where the species is present. Gray bold cells correspond to new locations found in this study (note that we also detected the species in cells where it was already recorded). b) Study area (shaded). Black dots correspond to the location of populations. Different metapopulations are encircled and identified with letters: A. Guardo, B. Otero de Guardo, C. Carrión, D. Pisuergra, E. Quintanaluengos, F. Tozo-Loras and G. Nava basin.