A new risk variant for multiple sclerosis at the immunoglobulin heavy chain locus associates with intrathecal IgG, IgM index and oligoclonal bands

Mercedes Delgado-García, Fuencisla Matesanz, Antonio Alcina, María Fedetz, María Isabel García-Sánchez, Juan Luis Ruiz-Peña, Óscar Fernández, María Jesús Pinto Medel, Laura Leyva, Carmen Arnal, Concepción Delgado, José Antonio López Guerrero, Antonio González-Pérez, María E Sáez, Luisa María Villar, José Carlos Álvarez-Cermeño, Carmen Picón, Rafael Arroyo, Jezabel Varadé, Elena Urcelay, Guillermo Izquierdo and Miguel Lucas

Abstract

Background: Recent findings have shown a correlation between the intrathecal IgG index and variants at the immunoglobulin heavy chain (IGHC) locus in patients with multiple sclerosis (MS).

Objectives: The objective of this paper is to analyse the association of the locus with MS susceptibility and its relationship with intrathecal immunoglobulin (Ig) parameters.

Methods: We genotyped the rs11621145 variant, located at the *IGHC* locus, in 2726 patients with MS and 2133 healthy controls. Associations of intrathecal IgG and IgM indexes with rs11621145 were analysed by linear regression analysis in 538 MS patients.

Results: We found that rs11621145 showed statistically significant evidence for association with susceptibility to MS (odds ratio = 0.69, $p = 1.053\text{E-}09$), though validation of this result in additional cohorts would be desirable. We confirmed the association between the IgG index and the rs11621145 ($p = 6.85\text{E-}07$, Beta = 0.207). Furthermore, rs11621145 was inversely correlated with IgM index ($p = 7.24\text{E-}04$, Beta = -0.277), and therefore marks a decreased likelihood of presenting IgM oligoclonal bands (odds ratio = 0.38, $p = 2.35\text{E-}06$).

Conclusions: Our results suggest that the polymorphism of the *IGHC* locus could be altering the switching of the Ig isotype in B cells and it may be interfering with T-dependent and T-independent antibody responses.

Keywords: Multiple sclerosis (MS), intrathecal IgG/IgM index, oligoclonal bands (OCBs), immunoglobulin heavy chain (IGHC) locus, association, SNP variant.

Date received: 1 April 2014; revised: 14 July 2014; 8 September 2014; accepted: 28 September 2014

Introduction

Multiple sclerosis (MS) is the most frequent demyelinating disease of the central nervous system (CNS) with an active inflammatory and immune component.^{1,2} MS is widely accepted as an autoimmune disease, but the aetiology and pathogenic mechanisms remain unclear. Current knowledge supports that MS develops in genetically susceptible

individuals as a result of an autoimmune attack to the myelin sheath activated through environmental factors.³ In MS, activated CD4+ T cells enter the CNS and trigger an inflammatory cascade that leads to recruitment of other immune cells such as B, CD8+ T cells and macrophages and induces the production of cytokines, antibodies and complement factors.

Multiple Sclerosis Journal

1–8

DOI: 10.1177/
1352458514556302

© The Author(s), 2014.
Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Correspondence to:

Fuencisla Matesanz
Departamento de Biología
Celular e Inmunología,
Instituto de Parasitología y
Biomedicina López Neyra
(IPBLN), Consejo Superior
de Investigaciones Científicas
(CSIC), 18016 Granada,
Spain
lindo@ipb.csic.es

María Isabel García-Sánchez
Juan Luis Ruiz-Peña
Unidad de Esclerosis
Múltiple, Hospital
Universitario Virgen
Macarena, Sevilla, Spain

Mercedes Delgado-García
Guillermo Izquierdo
Unidad de Esclerosis
Múltiple, Hospital
Universitario Virgen
Macarena, Sevilla, Spain/
These authors contributed
equally to this work

Antonio Alcina
María Fedetz
Departamento de Biología
Celular e Inmunología,
Instituto de Parasitología y
Biomedicina López Neyra
(IPBLN), Consejo Superior
de Investigaciones Científicas
CSIC, 18016 Granada, Spain

Fuencisla Matesanz
Departamento de Biología
Celular e Inmunología,
Instituto de Parasitología y
Biomedicina López Neyra
(IPBLN), Consejo Superior
de Investigaciones Científicas
CSIC, 18016 Granada, Spain/
These authors contributed
equally to this work

Óscar Fernández
María Jesús Pinto Medel
Laura Leyva
Unidad de Gestión Clínica
de Neurociencias, Instituto
de Biomedicina de Málaga
(IBIMA), Hospital Regional
Universitario de Málaga,
Málaga, Spain

Carmen Arnal

Servicio de Neurología,
Hospital Virgen de las
Nieves, Granada, Spain

Concepción Delgado

Centro Regional de
Transfusión Sanguínea
Granada-Almería, Granada,
Spain

**José Antonio López
Guerrero**

**Centro de Biología Molecular
Severo Ochoa, CSIC-UAM,**
Madrid, Spain

**Antonio González-Pérez
María E Sáez**

Centro Andaluz de Estudios
Bioinformáticos (CAEBi),
Sevilla, Spain

Luisa María Villar

**José Carlos Álvarez-
Cermeno**

Carmen Picón
Hospital Ramon y Cajal,
Departments of Immunology
and Neurology, MS Unit,
(IRYCIS), Madrid, Spain

Rafael Arroyo

Multiple Sclerosis Unit,
Hospital Clínico San Carlos,
Instituto de Investigación
Sanitaria del Hospital Clínico
San Carlos (IdISSC), Madrid,
Spain

Jezabel Varadé

Elena Urcelay
Immunology Department,
Hospital Clínico San Carlos,
Instituto de Investigación
Sanitaria del Hospital Clínico
S. Carlos (IdISSC), Madrid,
Spain

Miguel Lucas

Servicio de Biología
Molecular, Hospital
Universitario Virgen
Macarena, Facultad de
Medicina, Sevilla, Spain/
These authors contributed
equally to this work

The role of cerebrospinal fluid (CSF)-resident B cells seems to be important in CNS inflammation, but their contribution to MS evolution and progression is less clear.⁴ Nonetheless, antigen-directed affinity maturation and terminal differentiation of B cells contributing to IgG secretion and IgG oligoclonal bands (OCBs) occurs within perivascular infiltrates and meningeal lymphoid-like follicles in the CNS.^{5,6}

An elevation of intrathecal IgG synthesis and the presence of OCBs, characteristics of MS,⁷ further suggest the autoimmune origin of lesions in MS. In fact, about 95% of patients with MS show IgG-OCBs, and close to 40% show IgM OCBs in CSF.^{8,9} Moreover, it has been reported that intrathecal IgM synthesis predicts a worse disease course in the first stages of MS.¹⁰ Analysis of the IgG index, which is elevated in about 70% of patients, as well as the presence of OCBs^{11,12} have been proposed as complementary data for the diagnosis of MS.¹³

Buck et al.¹⁴ have recently reported that genetic variants in the immunoglobulin heavy chain (*IGHC*) locus are associated with the IgG index in MS, and that specific haplotypes might affect intrathecal IgG synthesis.

We aimed to study the genetic association as well as exploring the immunoglobulin (Ig) profile and OCBs in a cohort of Spanish patients with MS. Part of this cohort was included in a recent genome-wide association (GWA) study.¹⁵ Here, we recruited additional patients from the same study site and from two other Spanish centres, attaining 2726 MS patients and 2133 sex- and age-matched controls. In this work, we investigate the association with MS of single nucleotide polymorphisms (SNPs) within the *IGHC* locus and their relationship with the intrathecal IgG and IgM synthesis, as well as with the presence of OCBs for both Igs. In addition to confirming the findings of Buck et al.¹⁴ regarding the association of the *IGHC* locus with the IgG index, we observed that the selected variant was associated with both MS susceptibility and IgM status.

Patients and methods

Patients

Patients with suspicion of MS undergo lumbar puncture as part of the diagnostic procedure. Diagnosis was made by a neurologists based on McDonald criteria.¹⁶ After diagnosis patients were classified as clinically isolated syndrome (CIS), clinically definite MS, or non-MS patients. The study cohort from Hospital

Universitario Virgen Macarena (Sevilla) consisted of 362 patients with MS, and 414 patients free of MS or any other inflammatory disease (i.e. suffering mainly from headaches, hydrocephalus, Alzheimer's disease, Parkinson, behavioural changes, and stroke). The cohort from Hospital Ramón y Cajal (Madrid) consisted of 179 MS patients. Main features of the patients are shown in Table 1. Patients provided informed consent for lumbar puncture.

The entire cohort for rs11621145 genotyping study comprised 2726 patients with MS meeting established diagnostic criteria¹⁶ and 2133 healthy controls (mostly blood donors and staff). All patients and healthy controls had Caucasian origin. Patients and controls were recruited from six different Spanish hospitals: Sevilla cohort consisting of 988 cases and 1070 controls from Hospital Virgen Macarena, Hospital Virgen de la Nieves and Blood Bank of Andalucía; Málaga cohort consisting of 647 cases and 402 controls from Hospital Carlos Haya; Madrid cohort consisting of 1091 cases and 661 controls from Hospital Clínico San Carlos and Hospital Ramón y Cajal. Patients provided informed consent for blood extraction. The institutional ethics committees of these centres approved the study.

Samples and OCB studies

CSF samples were obtained by lumbar puncture from patients under suspicion of MS. The study of OCBs was performed in laboratories from Hospital Universitario Virgen Macarena (Seville) and Ramón y Cajal (Madrid). The procedure included the two-step detection of OCBs by isoelectric-focusing and incubation with antibodies to either IgG or IgM.^{17,18} Ig and albumin concentration in CSF and serum were measured by nephelometry. We quantified intrathecal IgG and IgM synthesis as the Ig index (CSF/serum Ig ratio divided by the CSF/serum albumin ratio).¹¹ These determinations are included in the protocol of routine diagnosis for MS in the above-indicated hospitals.

SNP selection and genotyping

SNP selection was based on the SNPs described by Buck et al.¹⁴ located at chromosome 14, and associated with IgG index. Linkage disequilibrium and Hardy-Weinberg (HW) equilibrium was performed with the 1000 Genome Project data for Caucasian populations.¹⁹ DNA was extracted from peripheral blood cells by standard methods using either the Maxwell Kit (Promega) or a saline precipitation procedure. The SNP rs11621145 was genotyped using Taqman® SNP Genotyping Assays (Applied Biosystems) according

Table 1. Demographic, clinical characteristics and cerebrospinal fluid parameters of the studied cohorts.

Cohorts	Madrid		Málaga		Sevilla	
	MS	Control	MS	Control	MS	Control
MS association study						
<i>N</i>	1091	661	647	402	988	1070
Age, mean year	45	48	45	42	42	41
Female (%)	54	66	67.7	67	67	67
PP (%)	4.2		1.4		6.3	
Ig Index study						
<i>N</i>	179				362	
Age, mean year at puncture (range)	37 (13–69)				35 (12–72)	
Females (%)	70				70	
IgG index, geometric mean (range)	1 (0.28–3.77)				0.96 (0.41–6.63)	
IgM index, geometric mean (range)	0.19 (0.02–2.86)				0.17 (0.01–5.53)	
IgG OCB total / + (%)	151/129 (85)				491/456 (93)	
IgM OCB total / + (%)	182/77 (42)				257/112 (44)	
<i>N</i> : number of patients; PP: primary progressive MS; IgG OCB: oligoclonal IgG bands; IgM OCB: oligoclonal IgM bands.						

to the manufacturer's specifications. Genotyping call rates were >95% in each cohort analysed and 10% of the samples were genotyped in duplicate with >99% concordance.

Statistical analysis

Departure from HW equilibrium was tested using an exact test. Genotype data were analysed to obtain allele and genotype counts using PLINK V.1.07.²⁰ For each available cohort odds ratios (OR), 95% Confidence Intervals (CI) and *p*-value were estimated by fitting sex-adjusted logistic regression models with PLINK in order to assess the associations between rs11621145 and: (a) the MS risk; (b) the presence of IgM OCB; and (c) the presence of IgG OCB. Both additive and dominant models were explored. In order to assess the association between Ig indexes and rs11621145, both indexes were transformed using natural logarithms as previously done by Buck *et al.*¹⁴ and, as happened in their study, untransformed data did not follow a Gaussian distribution, but was right-skewed. Once transformed, we analysed both IgG and IgM indexes by using the “-linear” command in PLINK that fits a linear regression. Thus, for each available cohort we obtained sex and age of puncture-adjusted beta coefficients, 95% CI, and *p*-values using both additive and dominant models. Estimates across cohorts were obtained by running meta-analyses for each endpoint using the “metan” command in Stata 12 (College Station, TX, USA). Both random and fixed-effects models were used and heterogeneity was explored by estimating *I*² index and *Q* statistic.

Results

IgG index associated variant is a risk factor for MS

Buck *et al.*¹⁴ reported five SNPs at the Ig locus associated with IgG index in MS patients by GWA analysis. Among these variants, the most associated was rs10136766. These polymorphisms present high linkage disequilibrium (LD) as determined in 379 Caucasian individuals included in the 1000 Genomes Project ($r^2 > 0.49$ –0.95). However, this polymorphism is not in HW equilibrium according to the 1000 Genomes Project data. In fact, only the second-most associated variant, rs11621145, was found in HW equilibrium, and therefore we selected this SNP as the best candidate to assess the association of this locus with IgG index and MS susceptibility.

As shown in Table 2, rs11621145 was genotyped in three Spanish cohorts whose controls were in HW equilibrium. After logistic regression analyses, we found that the dominant model was the one that best fitted the data. The rs11621145 was associated to MS in all three cohorts, showing a protective effect among G allele carriers. The meta-analysis of these results along with those from Buck *et al.*¹⁴ for rs11621145 is also indicated (Figure 1).

This locus has not been previously associated with MS in any of the GWA studies performed so far, but the coverage of this locus in the chips used is very low.

Table 2. Association analysis of rs11621145 in three Spanish MS cohorts.

Genotypes	MS N (%)		Controls N (%)				Dominant model		p
	GG	AG	AA	AG	AA	GG	OR	OR	
Madrid	126 (11.5)	405 (37.1)	560 (51.4)	72 (11)	307 (46.4)	0.8114 (0.68-0.99)	OR ¹	0.8114 (0.68-0.99)	0.036
Málaga	51 (8)	253 (39)	343 (53)	36 (9)	178 (44.2)	0.6919 (0.54-0.89)	OR ²	0.6919 (0.54-0.89)	0.004
Sevilla	88 (8.9)	356 (36)	544 (55.1)	133 (12.4)	462 (43.2)	0.6218 (0.52-0.74)	OR ³	0.6218 (0.52-0.74)	9.205E-08
Combined	265 (9.7)	1014 (37.2)	1447 (53.1)	241 (11.3)	947 (44.4)	0.6982	OR ⁴	0.6982	1.053E-09
OR: Odds ratio; OR ¹ : Fixed-effects OR estimate; OR(R) ² : Random-effects meta-analysis p-value; P ³ : Fixed-effects meta-analysis p-value; P(R) ⁴ : Random-effects meta-analysis p-value; Q ⁵ : p-value for Cochran's Q statistic; I ² : heterogeneity index (0–100).									3.046E-05
									0.14
									49.82

Figure 1. Forest plots of effect size, additive model, for rs11621145 associated with MS risk in the three cohorts analysed in the present work and in the cohort from Buck et al.¹⁴ We estimated the overall effect size using meta-analysis. The black squares represent the effect estimates for the individual cohorts and the horizontal lines represent the 95% CIs of the estimates. The diamond represents the overall estimate under fixed-effect model.

The G allele frequency of rs11621145 was 0.33 in the Spanish population, which is similar to the one found among Caucasians in the 1000 Genomes Project (0.39) but largely different when compared with other human populations (i.e. 0.97 in the African population).

Correlations of rs11621145 with IgG and IgM index

We analysed the association between rs11621145* G and the natural logarithm of the IgG index and IgM index by means of linear regression in the two MS cohorts, adjusting for sex and age at puncture (Table 3). The IgG index was associated with rs11621145 with a beta coefficient somewhat smaller than the one reported by Buck et al.¹⁴ but still statistically significant. The dominant model, as shown for MS susceptibility, seemed to fit better the data. We found that G carriers have higher values of IgG index. Interestingly, we found the exact opposite effect on the IgM index: the G carriers have a lower IgM index. Thus, G allele carriers presented lower MS susceptibility, higher intrathecal IgG index, and lower IgM index values.

In order to determine whether the associations found between IgG and IgM indexes and the rs11621145 genotypes were exclusive to MS patients, we performed correlation analysis using 414 CSF samples from individuals with non-inflammatory diseases. No association was found in these samples (Table 3).

Correlations of rs11621145 with oligoclonal IgM bands

Most MS patients enrolled in the present study are IgG OCB positives (93% and 85% in Sevilla and Madrid cohorts, respectively). We did not observe an

Table 3. Association study of the IgG and IgM index from MS and No Inflammatory Disease patients with the rs11621145 SNP by linear regression analysis.

			Sevilla			Madrid			Meta-analysis		
			<i>N</i>	BETA	<i>p</i>	<i>N</i>	BETA	<i>p</i>	<i>N</i>	BETA	<i>p</i>
MS	IgG	ADD	362	0.147	1.42E-04	176	0.101	0.066	538	0.132	2.52E-05
		DOM	362	0.215	2.6E-05	176	0.187	0.014	538	0.207	6.85E-07
	IgM	ADD	292	-0.219	0.007	173	-0.227	0.013	465	-0.223	2.2E-04
		DOM	292	-0.295	0.0063	173	-0.252	0.048	465	-0.277	7.24E-04
No MS	IgG	ADD	414	0.003	0.73						
		DOM	414	0.007	0.57						
	IgM	ADD	414	0.003	0.73						
		DOM	414	0.0064	0.92						

Sevilla corresponds to samples from the Hospital Virgen Macarena; Madrid corresponds to samples from Hospital Ramón y Cajal of Madrid
N: Number of samples; MS: multiple sclerosis samples; No MS: samples from no inflammatory diseases; BETA: Regression coefficient; *p*: Asymptotic *p*-value for *t*-statistic; ADD: additive model; DOM: dominant model.

Table 4. Association between the rs11621145 SNP and the IgM OCB.

	Allele	<i>N</i>	Dominant model	<i>p</i>	Additive model	<i>p</i>
			OR (95% CI)		OR (95% CI)	
Sevilla	G	255	0.43 (0.26–0.72)	0.0014	0.4937 (0.33–0.75)	0.0008
Madrid	G	179	0.32 (0.17–0.6)	0.00038	0.4102 (0.25–0.67)	0.00033
Meta-Analysis			0.3827	2.35E-06	0.4569	1.08E-06

Sevilla corresponds to samples from Hospital Virgen Macarena; Madrid corresponds to samples from Hospital Ramón y Cajal of Madrid.
N: Number of samples; OR: Odds ratio; *p*: Asymptotic *p*-value for *t*-statistic.

association between rs11621145 genotypes and IgG OCB. However, the power of this analysis was limited due to the low number of IgG OCB-negative patients, thus we cannot discard this association. On the other hand, we did observe an association between the presence of IgM OCB and the rs11621145 variant in both cohorts available (Sevilla and Madrid) (Table 4). In this case the additive model seemed to fit better the results. The G allele was associated with a lower likelihood of presenting IgM OCBs. The G allele was associated with protection towards MS, higher IgG index, lower IgM index and absence of IgM OCBs.

Discussion

In this work we confirm the previously reported association of the *IGHC* locus with the intrathecal IgG index in MS.¹⁴ In addition, we have found that this association does not only involve the IgG but also the IgM index and IgM OCB. The opposite effects observed in the associations between the IgG and IgM indexes with the SNP under study, along with the

localization of this polymorphism in the *IGHC* gene locus, point to an alteration of the equilibrium between innate and T-associated immune responses as the functional cause of the association.

In the T-cell-mediated immune response, IgM-positive B lymphocytes switch to IgG, IgA, or IgE upon immunization or infection.²² This process occurs by a recombination within S regions located upstream of each one of the heavy chain constant region (*C_H*) genes resulting in a change from IgM expression to one of the downstream isotypes. The isotype switching is cytokine-directed by inducing transcription from promoters located upstream to each acceptor S region. *Cis*-regulatory elements have been described within the *IGHC* gene locus and participate in the regulation of Ig *C_H* gene expression and isotype switch. Different knockout experiments have demonstrated the importance of *C_H* gene transcription for efficient isotype switch.^{23–25} In fact, a human polymorphism in the regulatory region located at the 3' of the *IGHC* gene locus, 3'RR enhancer, regulates *C_H* gene transcription.²⁶ This

polymorphism associates with selective IgA deficiency (IGAD), a common primary immunodeficiency, and with IgM concentration.²⁷ Variants in the same 3'RR enhancer have been also associated with systemic lupus erythematosus.²⁸ In the innate antibody response, responsible for the production of more than 80% of serum IgM, antibodies are produced against phylogenetically conserved antigens predominantly of non-protein nature.²⁹ This antibody response is mainly produced by CD5+ B cells and is responsible for intrathecal IgM synthesis in MS.³⁰

The polymorphisms associated with IgG index reported by Buck et al.¹⁴ and validated in this work are located in the immunoglobulin gamma heavy chain genes (*IGHG*) region, where four *IGHG* are encoded. However, since we do not know the ultimate causal variant of the Ig index alteration, it is not possible to infer whether the studied variant may be affecting *cis*-regulatory elements located upstream of the heavy chain gene modifying expression or whether it may be interfering with other process during class switching; further study in this area must be performed to clarify the underlying mechanism. It is noteworthy that the association between Ig index and rs11621145 occurs exclusively among MS patients and not in patients with non-inflammatory diseases. However, in these control patients, the intrathecal IgG would originate from the periphery. It would be important to analyse the IgG levels in a group of patients with diseases where intrathecal IgG synthesis occurs, to determine if the observed effect of the polymorphism on the Ig levels is specific to MS. The inflammatory environment of the brain in patients with MS could be the circumstance that triggers the observed effect. Cytokines produced by T helper and antigen-presenting cells determine the isotype to which B cells will switch, by inducing transcription from promoters located upstream to each acceptor S region. The promoters for the four *IGHG* are differentially regulated. They do not respond equally to regulatory stimulus, as has been shown by their differential response to the deletion of the Ikaros transcription factor in an animal model.³¹ In fact, there are several works pointing to the production of specific IgG isotypes in the CSF of patients MS.^{32,33} The specific cytokine spectrum produced in the brain of the MS patient could induce the switch to mainly one of the *IGHG* and this process could be altered by the polymorphism.

In this work we showed an association of the Ig locus with MS. However, our results were contradictory to those previously reported by Buck et al.,¹⁴ which reported a negative association of the locus with MS.

Even though in the present study we reached a large number of patients and controls from the meta-analysis between Spanish and UK cohorts, we should not forget that the small effects of non-HLA loci associated with MS require very large cohorts to demonstrate an association with the disease. On the other hand, even though all patients and controls were of Caucasian origin from Spain, we cannot totally rule out that residual population stratification could be partially responsible for the disease association observed here. For these reasons, it would be desirable for other groups to replicate this association, to reach numbers as reported in studies such as the Immunochip Project,³⁴ to completely overcome any doubts about its involvement in MS susceptibility.

The differences of association observed between cohorts were not unexpected, since previous meta-analyses have reported similar situations.³⁴ However, in this case we found notable differences between Sevilla and the rest of the cohorts. One clear difference of this cohort is that almost all patients were IgG OCB positive. We believe this could be one of the possible reasons for the observed differences in MS association between cohorts.

The homozygote individuals for the major allele (A) show an increased MS risk. The A allele is also associated with higher IgM index, higher IgM OCB and lower IgG index. One may think that since A is the major allele, the basal function should be associated with this allele, and G carriers might have acquired a higher ratio of class switching. However, when we analysed the allele frequency of rs11621145 in human populations other than Europeans, we observed that the A allele is almost non-existent in African populations. Since we do not know the ultimate causal variant of the effect, we cannot establish whether among African populations the rs11621145 polymorphism would not capture the functional variant, or that the polymorphism causing the effect in the Ig index was acquired later in human evolution and conserved in non-African populations.

The ultimate reason why these effects on the intrathecal Ig index and IgM bands can contribute to MS risk deserves further investigation. However, antibody isotype is important for determining the effector antibodies. The C_H region is bound by the Fc receptors on many cell types. Different C_H regions have different affinities for these proteins, thus greatly influencing antibody function and determining whether antibody-antigen complexes will activate cells. Pathological and clinical studies implicate antibody-dependent mechanisms in the immunopathogenesis of MS.

Therefore, any effect on the antibody isotype can affect MS susceptibility.

Acknowledgements

We thank the patients with multiple sclerosis and control subjects for making this study feasible. We also acknowledge the University Hospital Virgen Macarena Biobank (Andalusian Public Health System Biobank), integrated in the Spanish Biobanks Network (Carlos III Health Institute, Grant nº RD09/0076/00080), for its help and support in the gifts of clinical samples, used in this work.

Conflicts of interest

No conflicts of interest

Funding

Financial support for the study was provided by Fondo de Investigación Sanitaria (FIS)-Fondos Europeos de Desarrollo Regional (FEDER) [grant numbers RETICS-REEM-RD12/0032/0015-RD12/0032/0005-RD12/0032/0006-RD12/0032/0009, P12/00555, PI10/01616, PI13/01527, PI13/01466, CP10/00526, PI12-00239]; Junta de Andalucía- Fondos Europeos de Desarrollo Regional (FEDER) [grant numbers P07-CVI-02551, P09-CTS-5218]. MDG is holder of a Neuroinvest fellowship.

References

- McFarland HF and Martin R. Multiple sclerosis: A complicated picture of autoimmunity. *Nat Immunol* 2007; 8: 913–919.
- Zettl UK, Stüve O and Patejdl R. Immune-mediated CNS diseases: A review on nosological classification and clinical features. *Autoimmun Rev* 2012; 11: 167–173.
- Sospedra M and Martin R. Immunology of multiple sclerosis. *Annu Rev Immunol* 2005; 23: 683–747.
- Kuenz B, Lutterotti A, Ehling R, et al. Cerebrospinal fluid B cells correlate with early brain inflammation in multiple sclerosis. *PLoS ONE* 2008; 3: e2559.
- Prineas JW. Multiple sclerosis: Presence of lymphatic capillaries and lymphoid tissue in the brain and spinal cord. *Science* 1979; 203: 1123–1125.
- Serafini B, Rosicarelli B, Magliozzi R, et al. Detection of ectopic B-cell follicles with germinal centers in the meninges of patients with secondary progressive multiple sclerosis. *Brain Pathol* 2004; 14: 164–174.
- Disanto G, Morahan JM, Barnett MH, et al. The evidence for a role of B cells in multiple sclerosis. *Neurology* 2012; 78: 823–832.
- Kostulas VK, Link H and Lefvert AK. Oligoclonal IgG bands in cerebrospinal fluid. Principles for demonstration and interpretation based on findings in 1114 neurological patients. *Arch Neurol* 1987; 44: 1041–1044.
- Villar LM, Masjuan J, González-Porqué P, et al. Intrathecal IgM synthesis in neurologic diseases: Relationship with disability in MS. *Neurology* 2002; 58: 824–826.
- Villar LM, Masjuan J, González-Porqué P, et al. Intrathecal IgM synthesis is a prognostic factor in multiple sclerosis. *Ann Neurol* 2003; 53: 222–226.
- Lefvert AK and Link H. IgG production within the central nervous system: A critical review of proposed formulae. *Ann Neurol* 1985; 17: 13–20.
- Cowdrey GN, Tasker PJ, Gould BJ, et al. Isoelectric focusing in an immobilized pH gradient for the detection of intrathecal IgG in cerebrospinal fluid: sensitivity and specificity for the diagnosis of multiple sclerosis. *Ann Clin Biochem* 1993; 30: 463–468.
- Petereit HF and Heiss WD. New diagnostic criteria for multiple sclerosis. *Ann Neurol* 2002; 51: 533–534.
- Buck D, Albrecht E, Aslam M, et al. Genetic variants in the immunoglobulin heavy chain locus are associated with the IgG index in multiple sclerosis. *Ann Neurol* 2013; 73: 86–94.
- Matesanz F, González-Pérez A, Lucas M, et al. Genome-wide association study of multiple sclerosis confirms a novel locus at 5p13.1. *PLoS One* 2012; 7: e36140.
- Polman CH, Reingold SC, Banwell B, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald Criteria. *Ann Neurol* 2011; 69: 292–302.
- Sádaba MC, González Porqué P, Masjuan J, et al. An ultrasensitive method for the detection of oligoclonal IgG bands. *J Immunol Methods* 2004; 284: 141–145.
- Villar LM, González-Porqué P, Masjuan J, et al. A sensitive and reproducible method for the detection of oligoclonal IgM bands. *J Immunol Methods* 2002; 258: 151–155.
- 1000 Genomes Project Consortium, Abecasis GR, Altshuler D, et al. A map of human genome variation from population-scale sequencing. *Nature* 2010; 467: 1061–1073.
- Purcell S, Neale B, Todd-Brown K, et al. PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007; 81: 559–575.
- Sawcer S, Hellenthal G, Pirinen M, et al. Genetic risk and a primary role for cell-mediated immune

- mechanisms in multiple sclerosis. *Nature* 2011; 476: 214–219.
22. Stavnezer J, Guikema JE and Schrader CE. Mechanism and regulation of class switch recombination. *Annu Rev Immunol* 2008; 26: 261–292.
 23. Jung S, Rajewsky K and Radbruch A. Shutdown of class switch recombination by deletion of a switch region control element. *Science* 1993; 259: 984–997.
 24. Zhang J, Bottaro A, Li S, et al. A selective defect in IgG2b switching as a result of targeted mutation of the I gamma 2b promoter and exon. *EMBO J* 1993; 12: 3529–3537.
 25. Oruc Z, Boumédiène A, Le Bert M, et al. Replacement of Igamma3 germ-line promoter by Igamma1 inhibits class-switch recombination to IgG3. *Proc Natl Acad Sci U S A* 2007; 104: 20484–20489.
 26. Hu Y, Pan Q, Pardali E, et al. Regulation of germline promoters by the two human Ig heavy chain 3' alpha enhancers. *J Immunol* 2000; 164: 6380–6386.
 27. Giambra V, Cianci R, Lolli S, et al. Allele *1 of HS1.2 enhancer associates with selective IgA deficiency and IgM concentration. *J Immunol* 2009; 183: 8280–8285.
 28. Frezza D, Tolusso B, Giambra V, et al. Polymorphisms of the IgH enhancer HS1.2 and risk of systemic lupus erythematosus. *Ann Rheum Dis* 2012; 71: 1309–1315.
 29. Boes M, Prodeus AP, Schmidt T, et al. A critical role of natural immunoglobulin M in immediate defense against systemic bacterial infection. *J Exp Med* 1998; 188: 2381–2386.
 30. Villar LM, Sádaba MC, Roldán E, et al. Intrathecal synthesis of oligoclonal IgM against myelin lipids predicts an aggressive disease course in MS. *J Clin Invest* 2005; 115: 187–194.
 31. Sellars M, Reina-San-Martin B, Kastner P, et al. Ikaros controls isotype selection during immunoglobulin class switch recombination. *J Exp Med* 2009; 206: 1073–1087.
 32. Salier JP, Goust JM, Pandey JP, et al. Preferential synthesis of the G1m(1) allotype of IgG1 in the central nervous system of multiple sclerosis patients. *Science* 1981; 213: 1400–1402.
 33. Goust JM and Salier JP. Imbalance in recruitment of IgG (Gm) allotype-producing B-cell subsets from blood to brain in multiple sclerosis. *Cell Immunol* 1984; 88: 551–557.
 34. International Multiple Sclerosis Genetics Consortium (IMSGC), Beecham AH, Patsopoulos NA, et al. Analysis of immune-related loci identifies 48 new susceptibility variants for multiple sclerosis. *Nat Genet* 2013; 45: 1353–1360.