**THE Rho/Rac EXCHANGE FACTORS Vav2 AND Vav3 CONTROL A
LUNG-SPECIFIC PRO-METASTATIC TRANSCRIPTOMAL
PROGRAM IN BREAST CANCER CELLS**

by

**Carmen Citterio^{1,2,3}, Mauricio Menacho-Márquez^{1,2,3}, Ramón García-Escudero⁴,
Romain M. Larive^{1,2}, Olga Barreiro⁵, Francisco Sánchez-Madrid⁵, Jesús M. Paramio⁴,
and Xosé R. Bustelo^{1,2,*}**

¹Centro de Investigación del Cáncer and ²Instituto de Biología Molecular y Celular del
Cáncer, CSIC-University of Salamanca, Salamanca, Spain. ⁴CIEMAT Molecular Oncology
Unit, Madrid, Spain. ⁵Centro Nacional de Investigaciones Cardiovasculares, Madrid, Spain.

³These authors contributed equally to this work. The order was set alphabetically.

Running title: Pleiotropic roles of Vav exchange factors in breast cancer

Key words: Rho/Rac GTPases, Rho/Rac GEFs, breast cancer, Ilk, Tacstd2, inhibin, Cox2

*To whom correspondence should be addressed. E-mail: xbustelo@usal.es

ABSTRACT

Although the role of Rho/Rac exchange factors in signal transduction is well established, their therapeutic value in cancer remains obscure. Here, we report that two of those exchange factors, Vav2 and Vav3, play synergistic roles in breast cancer. They do so by sustaining tumor growth, neoangiogenesis and many of the steps involved in lung-specific metastasis. This is a proactive role, since their overexpression confers metastatic properties to otherwise non-metastatic breast cancer cells. The implication of Vav proteins in those routes could not be correlated with their expected roles on Rac1 activation and cell migration, suggesting the implication of additional signaling elements. Using microarray analyses, we found that these proteins regulate a vast transcriptional program in breast cancer cells in a pan-specific, synergistic and isoform-specific manner. Furthermore, the analysis of Vav family-regulated transcripts indicates that their abundance is controlled by Rac1-dependent and independent routes. Importantly, the Vav family-dependent transcriptome encodes therapeutically targetable proteins (Ilk, Inhibin β A, cyclooxygenase 2 and Tacstd2) that play non-redundant roles in primary tumorigenesis and lung-specific metastasis. It also contains gene signatures that can forecast disease outcome and metastasis development in breast cancer patients.

INTRODUCTION

Rho/Rac GTPases and their proximal effectors are frequently overexpressed in tumors, where they regulate biological functions that favor the proliferation, apoptosis, invasion, metastasis, and/or resistance of cancer cells to anti-cancer treatments. This has led to the idea that they could be potential anti-tumoral and anti-metastatic targets (1). Since Rho/Rac proteins are not mutated in tumors (1), one of the possible avenues to inhibit their signal transduction pathways is to block the upstream guanosine nucleotide exchange factors (GEF). These enzymes catalyze the exchange of GDP by GTP on Rho/Rac proteins, thereby favoring their rapid transition from their inactive (GDP-bound) to their active (GTP-bound) states during cell signaling (2, 3). Recent evidence supports the potential therapeutic value of these molecules in cancer. Thus, the use of knockout mice and knockdown cells has demonstrated that the elimination of individual Rho/Rac GEFs can impair the growth of skin cancer (T-cell lymphoma invasion and metastasis-inducing protein 1 [Tiam]) (4), chronic myelogenous leukemia subtypes (Vav3) (5), colorectal cancer (Tiam1, adenomatous polyposis coli-stimulated exchange factor 1[Asef1, also known as ArhGEF4], Asef2) (6, 7) and breast cancer (phosphatidylinositol 3,4,5-triphosphate-dependent Rac exchanger 1 [P-Rex1], P-Rex2) (8, 9). In addition, it has been demonstrated that inhibitors that block the interaction of a subset of GEFs with Rac1 can block chronic myelogenous leukemia in animal models (10). The role of Rho/Rac GEFs has been less characterized in the case of metastasis, although current evidence indicates that some Rac1 GEFs play either anti- (Tiam1) (11) or pro-metastatic (P-Rex1) (12) roles in different tumor types.

Despite the above evidence, the identification of the good GEF targets in different cancer/metastatic settings is complicated by a number of hurdles. One of them is the fact that Rho/Rac GEFs constitute a large enzyme family that, in humans, is composed of 65 members (2, 3). Although they all share either catalytic Dbl-homology (DH, 54 members) or Dedicator of Cytokinesis (Dock)-homology region 2 (DHR2, 11 members) domains, they differ widely in terms of ancillary domains and tissue distribution patterns. Due to this, even GEFs with the same catalytic specificity can only stimulate the downstream GTPase in specific cell types, stimulation conditions or subcellular localizations (2, 3). In addition, it has been shown that their non-catalytic domains can contribute to limit the spectrum of effectors engaged by the stimulated GTPases (13-15) and to activate GTPase-independent routes (9, 16-18). The type of oncogenic events present in tumors adds another layer of complexity, since different oncoproteins may restrict the spectrum of Rho/Rac GEFs that could be preferentially activated in a particular tumor (8, 9). For example, it is expected that tumors containing constitutive activity of the phosphatidylinositol-3 kinase would activate preferentially phosphatidylinositol 3,4,5-triphosphate-dependent Rho/Rac GEFs such as P-Rex1 and P-Rex2. All these observations indicate that the identification of therapeutically relevant Rho/Rac GEFs requires the identification of those specifically enriched/activated in tumors, the determination of the functional redundancies existing among them, and the dissection of the GEF-dependent and independent biological processes that affect cancer growth and/or metastasis *in vivo*.

In this work, we have aimed at tackling those issues in the case of Vav proteins, a specific regulatory subgroup of the DH family (19). These proteins epitomize the

complexity existing in the Rho/Rac GEF world. Thus, vertebrate species have three members (Vav1, Vav2, Vav3) with overlapping, but not identical cellular and tissues distributions. In addition, these proteins contain highly similar and complex structures that encompass regulatory, catalytic, catalytic-associated, and adaptor domains (19). As a consequence, they can induce cellular responses in both GTPase-dependent and independent manners (16, 18, 19). Finally, Vav proteins constitute a quite idiosyncratic subclass within the Rho/Rac GEF family, since they are the only GEFs positively regulated by direct tyrosine phosphorylation during signal transduction (19-22). In this work, we have used both *in silico* and animal model systems to: **(i)** Compare their abundance in relation to the rest of 51 DH family GEFs in human tumors. **(ii)** Determine whether they played specific and/or redundant roles in both primary tumorigenesis and the main metastatic steps. **(iii)** Identify non-conventional downstream targets that could mediate the Vav-dependent effects on the biology of cancer cells. This approach revealed that Vav2 and Vav3 play critical and non-redundant roles during both primary breast tumors and lung-specific metastasis. We also found a Vav2/Vav3-dependent transcriptional program that helps explaining the synergistic effect of those two proteins in primary tumorigenesis and lung-specific metastasis. These results provide, for the first time, a holistic view of the role of a Rho/Rac GEF subfamily in the main steps involved in the metastatic dissemination of a very frequent tumor, reveal a new biological program that conditions the specific tropism of metastatic breast cancer cells to the lung and, in addition, provide information about prognostic gene signatures that could be potentially relevant in this clinical setting in a near future.

RESULTS

Human breast tumors and cancer cell lines show increased abundance of Vav family transcripts

To identify tumors with enriched amounts of Vav proteins and determine the possible functional overlap with other GEFs, we analyzed the abundance in tumors of the transcripts for the 54 human DH family GEFs using metagenomic analyses. We considered that an mRNA was more abundant in a tumor when: **(i)** It showed a ≥ 2 -fold average enrichment in its abundance in that tumor when compared to other cancer types. **(ii)** Such variation had a statistical P value $\leq 1 \times 10^{-4}$ (see Materials and Methods). Using those cut-off parameters, we found that the *VAV3* mRNA was consistently enriched in breast cancer samples (**Fig. S1A**) and breast cancer cell lines (**Fig. S1B**). In this tumor type, the *VAV3* transcript was more abundant in estrogen- and progesterone receptor-positive tumors (**Table S1**). In addition, *VAV3* was also more abundant in breast luminal A/B tumors when compared to both normal breast and other breast cancer subtypes (**Table S2**). As expected from previous results from Kazanietz's lab (8), the metagenomic analyses also revealed that the *P-REX1* mRNA was enriched in breast cancer relative to other tumor types. The enrichment in abundance of *VAV3* and *P-REX1* transcripts in breast cancer was comparable, although not fully overlapping to that exhibited by *ERBB2* (v-erb-b2 erythroblastic leukemia viral oncogene homolog 2) (**Fig. S1C,D**), a gene frequently overexpressed in breast cancer (23). The *VAV1* mRNA was only enriched in abundance in a statistically significant manner in lymphoma-derived samples (**Fig. S1A,C-E**). This is probably not connected to tumor-linked requirements, since it is known that the *VAV1* gene

is preferentially expressed in healthy hematopoietic cells (**Fig. S1F**) (19). The VAV2 mRNA was detected in a number of tumor and cancer cell samples, although it did not display any statistically significant enrichment in any specific tumor type (**Fig. S1C,D**). It was, however, slightly enriched in abundance in breast luminal A/B tumors when compared to normal breast or other breast tumor subtypes, although such variation show a statistical significance ($P = 4.9 \times 10^{-3}$) below our aforementioned cut-off value (**Table S2**). The second member of the P-REX subfamily, *P-REX2*, showed no significant enrichment in the samples surveyed (**Fig. S1C-E**).

Vav proteins play critical roles in breast cancer

The above results indicated that VAV3 could play a role in breast cancer. However, the presence of other Rho/Rac GEFs that were either enriched in abundance (i.e., P-REX1) or basally expressed (i.e., VAV2) in this tumor type called into question whether such role could be meaningful or not. This question was particular important in the case of VAV2, a protein that shared similar regulatory properties with VAV3 (19). To address this issue, we first analyzed the effect of their specific knockdown in the 4T1 breast cancer cell line (24). These cells, originally isolated from a spontaneous mammary tumor developed in a wild type Balb/cJ mouse, are used extensively in the breast cancer field because: **(i)** They can grow in immunocompetent mice and, therefore, they are still subjected to the anti- and pro-tumorigenic effects of the immune system of the host. **(ii)** They can disseminate away from the primary tumor and metastasize in different organs of the recipient mouse. **(iii)** They have non-metastatic counterparts that can be used to evaluate whether a gain-of-function in

a particular signaling route can rescue specific stages of the metastatic process (24). In our particular case, these cells were also useful because they express both Vav2 and Vav3 (**Fig. S2A,B**), thereby allowing the investigation of functional redundancy/specificity issues. To clearly discern the level of functional overlap between Vav2 and Vav3, we generated knockdown cell clones lacking Vav2, Vav3, or both proteins at the same time (**Table S3**). In addition, we used one of the *Vav2;Vav3* knockdown cell clones to generate “rescued” cell line derivatives stably overexpressing Vav2, Vav3, Vav2 plus Vav3, constitutively active Vav2^{Y172F} or catalytically inactive Vav2^{R373A} (**Table S3**). The proper depletion of the indicated Vav family protein was confirmed by quantitative reverse transcription-polymerase chain reaction (qRT-PCR) (**Fig. S2A, B**). The re-expression of the appropriate Vav family member in the knockdown cells was confirmed using both qRT-PCR (**Fig. S2C,D**) and Western blot (**Fig. S2E,F**) techniques. Using a GTPase-linked immunosorbent assay (G-LISA) to determine total Rac1 activity, we observed that the simultaneous inactivation of both Vav2 and Vav3 was required to reduce basal Rac1 activity in 4T1 cells (**Fig. 1A**). By contrast, the overexpression of a single Vav family protein was enough to restore normal Rac1 activity in the *Vav2;Vav3* knockdown cell line (**Fig. 1A**). As control, the catalytically inactive Vav2^{R373A} mutant could not restore Rac1 activity when expressed in Vav2/Vav3-deficient 4T1 cells (**Fig. 1A**). Similar results were found when Rac1 activity was determined using standard pull-down experiments with a GST fusion protein containing the Rac1 binding domain of Pak1 (**Fig. 1B**). The elimination of Vav proteins in 4T1 did not have any effect on basal RhoA activity (**Fig. 1A**).

When implanted in the mammary fat pads of recipient mice, we observed that all *Vav* knockdown cell lines developed tumors with low growth kinetics (**Fig. 1C-F**). The examination of tumor-bearing animals at the end of the tumorigenesis assays also indicated that the metastasis of cancer cells to the lung was severely impaired upon the depletion of either *Vav2* or *Vav3* and totally abolished upon the simultaneous elimination of both *Vav* proteins (**Fig. 1G-I, Table S4**). This defect was tissue-specific, because all *Vav* protein deficient tumors metastasized well to other organs (**Table S4**). All the above defects were observed using two independent knockdown cell clones, indicating that they represent a bona-fide representation of the biological properties of *Vav* proteins (**Fig. 1C,D,G; Table S4**). Full rescue of control-like primary tumorigenesis (**Fig. 1E**) and lung metastasis (**Fig. 1H; Table S4**) could be achieved upon the ectopic co-expression of both *Vav* family proteins in *Vav2;Vav3* knockdown cells. By contrast, the re-expression of any of the two single *Vav* proteins (**Fig. 1E,H; Table S4**) or the constitutively active *Vav2*^{Y172F} mutant (**Fig. 1F; Table S4**) failed in restoring the tumorigenic properties of *Vav2;Vav3* knockdown cells. Since basal Rac1 activity does not change significantly between *Vav2*-, *Vav2*^{Y172F}, *Vav3*-, and *Vav2/Vav3*-reconstituted *Vav2;Vav3* knockdown cells (**Fig. 1A,B**), this result suggests that *Vav2* and *Vav3* act non-redundantly in those two processes probably by engaging both Rac1-dependent and independent routes. This idea is also consistent with the observation that the single knockdown *Vav2* and *Vav3* 4T1 cells show defects in both primary tumorigenesis (**Fig. 1C**) and metastasis (**Fig. 1G**) despite showing basal Rac1 activity comparable to that found in control cells (**Fig. 1A**).

Vav proteins have pleiotropic functions in breast cancer

To further evaluate the effect and functional redundancy of Vav proteins on primary tumor growth, we studied the implication of each of those proteins in a number of cancer-linked processes. Using immunohistochemical techniques, we observed that the tumors derived from *Vav2;Vav3* knockdown cells had reduced levels of proliferation (**Fig. 2A,B**) and neoangiogenesis (**Fig. 2C,D**). The vessels generated inside those tumors were also less leaky than those found in control samples, as assessed by in vivo permeability experiments using intravenously injected rhodamine-labeled dextrane (**Fig. 2E,F**). The rescue of all those functions required the simultaneous re-introduction of Vav2 and Vav3 in *Vav2;Vav3* knockdown cells (**Fig. 2A-F**). The absence of both Vav proteins also increased the number of apoptotic, terminal deoxynucleotidyl transferase dUTP nick end labeling (Tunel) positive cancer cells, suggesting a role for those proteins in cancer cell survival (**Fig. 2G,H**). However, this defect could be corrected by re-expressing any Vav protein in *Vav2;Vav3* knockdown cells (**Fig. 2G,H**), indicating that Vav2 and Vav3 regulate this latter route redundantly.

To analyze the role in metastasis, we analyzed the influence of Vav2 and Vav3 in the intravasation, extravasation, and lung colonization properties of breast cancer cells. By quantifying the number of circulating cancer cells present in tumor-bearing animals, we found that *Vav2;Vav3* knockdown cells intravasated 3-fold less efficiently than control cells (**Fig. 3A**). However, this defect cannot explain the lack of lung metastasis exhibited by those tumors, because the *Vav2;Vav3* knockdown cells can develop metastases in other secondary tissues (**Table S4**). Consistent with this, we observed that *Vav2;Vav3*

knockdown cells could not metastasize to the lung even when introduced intravenously in recipient mice (**Fig. 3B-E**), suggesting that the Vav protein deficiency impacts on post-intravasation metastatic steps. To investigate this possibility, we injected intravenously cancer cells that had been labeled with a cell-permeable chromophore and, 48 hours later, evaluated their extravasation to the lungs of recipient mice by confocal microscopy. These experiments indicated that *Vav2;Vav3* knockdown cells could not colonize the lung (**Fig. 3F-G**). In fact, when detected, they were usually visualized as single cells trapped inside the lung capillaries (**Fig. 3F-G**). The simultaneous re-introduction of Vav2 and Vav3 was required to restore all these intravasation (**Fig. 3A**) and post-intravasation metastatic steps (**Fig. 3E-G**) in *Vav2;Vav3* knockdown 4T1 cells.

Vav proteins play pro-tumorigenic and pro-metastatic roles in both primary breast epithelial cells and non-metastatic breast cancer cell lines

To make sure that the implication of Vav proteins in breast cancer was not limited to the 4T1 cell model, we analyzed the implication of those GEFs in this tumor type using independent experimental systems. As a first approach, we investigated whether the absence of Vav2 and Vav3 had an impact on the transformation of the mammary gland induced by the polyomavirus middle T antigen (PyMT). To this end, we isolated primary breast epithelial cells from wild type and *Vav2^{-/-};Vav3^{-/-}* mice and, after transducing them with a lentivirus containing the *PyMT* oncogene, re-implanted them in the mammary fat pads of female mice to follow the kinetics of mammary tumor formation in vivo. Unlike

the case of wild type cells, the PyMT-transduced *Vav2*^{-/-};*Vav3*^{-/-} cells could not develop tumors or metastases when implanted in recipient mice (**Fig. 4A,B**).

As a second approach, we investigated whether the overexpression of Vav2 and Vav3 could confer metastatic properties to 168FARN cells. Despite being isolated from the same mouse mammary carcinoma as 4T1 cells, 168FARN cells are not metastatic due to severe defects in both intravasation and post-intravasation steps (24). Despite the fact that the overexpression of the two Vav proteins in 168FARN cells (**Fig. 4C**) did not lead to increased Rac1 activity in vitro (**Fig. 4D**), it promoted faster tumor growth kinetics (**Fig. 4E**) as well as lung metastasis (**Fig. 4F**) in vivo. Interestingly, these cells could also infiltrate efficiently the spleen (**Fig. 4F**), indicating that the overexpression of Vav2 and Vav3 eliminates the defective intravasation step of 168FARN cells. Vav2/Vav3-overexpressing 168FARN cells, but not the control counterparts, also formed lung metastasis when intravenously injected in mice (**Fig. 4G**). Taken together, these data indicate that Vav proteins play proactive roles in both breast cancer tumorigenesis as well as in the intravasation and extravasation steps of metastatic breast cancer cells.

Vav2 and Vav3 regulate the transcriptome of breast cancer cells using common, synergistic, and isoform-specific pathways

To investigate whether the need of both Vav family proteins in breast tumorigenesis and metastasis was a reflection of their contribution to the intrinsic proliferative and migratory properties of breast cancer cells, we evaluated these two parameters in our collection of 4T1 cells in vitro. We observed that *Vav2*;*Vav3* knockdown cells showed significantly

lower proliferation (**Fig. 4H**) and invasion (**Fig. 4I**) rates than control cells. However, unlike the results obtained in orthotopically transplanted cells, the rescue of those two in vitro defects could be efficiently achieved by overexpressing Vav2 in *Vav2;Vav3* knockdown cells (**Fig. 4H,I**). These results, together with the observations made in **Figs. 1-3**, suggested that the synergistic functional interaction of Vav2 and Vav3 in breast cancer cells cannot be attributed monodimensionally to their expected action on overall Rac1 activity, migration, and/or proliferation routes.

These results led us to carry out Affymetrix microarray experiments to identifying distal effectors that could explain the cooperative action of Vav2 and Vav3 in breast cancer. We identified 2,411 transcripts whose abundance changed upon depletion of both Vav proteins in 4T1 cells (**Fig. S3A, Table S5**). Using a number of both experimental- (see Materials and Methods) and therapeutic-oriented criteria (i.e., mRNAs with reduced abundance in *Vav2;Vav3* knockdown cells that encoded therapeutically amenable proteins), we selected a final list of six transcripts whose abundance was reduced in *Vav2;Vav3* knockdown 4T1 cells for functional characterization (*Itgb6*, *Itga8*, *Ilk*, *Tacstd2*, *Inhba*, *Ptgs2*). *Itgb6* and *Itga8* encode integrin subunits that regulate cell adhesion to laminins and fibronectin, respectively (25). *Ilk* (integrin-linked kinase) expresses a pseudokinase working as an adaptor protein downstream of integrins (26). *Tacstd2* (Tumor-associated calcium signal transducer 2) encodes an epithelial cell adhesion molecule (EpCAM) family member with ascribed roles in cell adhesion and cell signaling (27). *Inhba* encodes inhibin β A (Inh β A), a ligand of the TGF β family that can form homodimers or heterodimers with Inh α and Inh β B subunits (28). *Ptgs2* encodes cyclo-oxygenase 2 (Cox2), an endoplasmic

reticulum-localized enzyme involved in prostaglandin production (29). This latter protein was chosen as a positive control, since it had been previously linked to metastatic processes (30).

We confirmed by qRT-PCR that the selected Vav-dependent transcripts were less abundant in cultured *Vav2;Vav3* knockdown cells (**Fig. S3B**) and in the primary tumors derived from them (**Fig. S3C**) than in parallel samples obtained from control cells. With the exception of the *Ptgs2* mRNA, these transcripts were also less abundant in the non-metastatic 168FARN than in the metastatic 4T1 cells (**Fig. S3D**). The overexpression of Vav2 and Vav3 in 168FARN cells, which led to the acquisition of metastatic properties in those cells (see above, **Fig. 4F,G**), increased the abundance of most of those transcripts to values either similar to (*Itga8*, *Ilk*) or higher than (*Inhba*, *Ptgs2*) those found in 4T1 cells (**Fig. S3D**). The only exceptions were the *Itgb6* and *Tacstd2* mRNA, whose abundance did not increase (*Itgb6*) or was elevated only 2-fold (*Tacstd2*) in Vav2/Vav3-overexpressing 168FARN cells (**Fig. S3D**). Given the cooperative effect of Vav2 and Vav3 in tumor growth and metastasis, we next investigated the abundance of the selected Vav family targets in single Vav2-, single Vav3-, and double Vav2/Vav3-reconstituted *Vav2;Vav3* knockdown cells (**Fig. S4A**). These experiments indicated that the rescue of the normal abundance levels of *Itga8*, *Tacstd2* and *Ptgs2* mRNAs only required the re-introduction of any single Vav family protein in the *Vav2;Vav3* knockdown cells. By contrast, the abundance of *Ilk* and *Inhba* transcripts required the simultaneous re-introduction of both Vav proteins in those cells. Normal contents of the *Itgb6* mRNA were achieved by overexpressing Vav3, but not Vav2, in *Vav2;Vav3* knockdown cells (**Fig. S4A**).

To address the issue of the involvement of Rac1 in the regulation of the Vav2/Vav3-dependent transcriptome, we stably expressed a constitutively active, fast cycling Rac1 mutant (F28L) in both control and *Vav2;Vav3* knockdown cells (**Fig. S4B,C**). G-LISA experiments indicated that the presence of Rac1^{F28L} bypassed the deficient activation of endogenous Rac1 induced by the compound *Vav2;Vav3* knockdown in 4T1 cells (**Fig. S4D**). In fact, Rac1^{F28L}-expressing cells displayed higher Rac1 activity than normal 4T1 cells independently of the abundance of endogenous Vav proteins in them (**Fig. S4D**). When expressed in *Vav2;Vav3* knockdown cells, Rac1^{F28L} could not rescue the abundance of the *Ilk* transcript, thus confirming the idea that some of the downstream effects of Vav proteins were Rac1-independent (**Fig. S4E**). In fact, when expressed in normal 4T1 cells, we observed that Rac1^{F28L} reduced the abundance of the *Ilk* mRNA (**Fig. S4E**). By contrast, the Rac1 mutant did restore the abundance of the other synergistic target (*Inhba*) and the two mRNAs commonly regulated by Vav2 and Vav3 (*Tacstd2*, *Ptgs2*) in *Vav2;Vav3* knockdown cells (**Fig. S4E**). In wild type 4T1 cells, Rac1^{F28L} promoted an increase in the abundance of both the *Inhba* and *Ptgs2* mRNAs that was significantly higher than that observed in Rac1^{F28L}-expressing *Vav2;Vav3* knockdown cells (**Fig. S4E**). Since these two cell lines display similar Rac1^{F28L} abundance (**Fig. S4B,C**) and activity (**Fig. S4D**), this result suggests that Rac1-independent routes triggered by either Vav2 or Vav3 may be required for the optimal regulation of those distal downstream targets.

To further verify the implication of GEF-independent routes in the engagement of the distal Vav family-dependent transcriptomal program, we investigated whether the synergistic response previously observed between Vav2 and Vav3 in the regulation of *Ilk*

and *Inhba* mRNA abundance (**Fig. S4A**) could be reproduced using *Vav2;Vav3* knockdown cells reconstituted with wild type *Vav3* and catalytically inactive *Vav2*^{R373A}. These experiments demonstrated that the contribution of *Vav2* to the regulation of *Ilk* gene expression was totally independent of its enzyme activity (**Fig. S4G**). By contrast, *Vav2* required GEF activity to synergize with wild type *Vav3* in the regulation of the abundance of the *Inhba* mRNA (**Fig. S4G**). The latter result suggests that the synergism observed between *Vav* proteins in the regulation of this gene is probably due to the stimulation of different spectra of Rho/Rac GTPases by *Vav2* and *Vav3* or, alternatively, by the *Vav* family member-specific activation of Rac1 in different subcellular localizations. The restoration of all *Vav* family GEF- and GEF-independent routes seems to be crucial for the metastatic efficiency of cancer cells towards the lung, because we observed that *Rac1*^{F28L}- and *Vav3/Vav2*^{R373A}-expressing *Vav2;Vav3* knockdown cells did not metastasize efficiently to the lung when injected intravenously in mice (**Fig. S4H**). Taken together, these results indicate that: (i) *Vav* proteins control gene expression using common, synergistic, and *Vav* family member-specific routes. (ii) The targets indistinctly regulated by *Vav2* and *Vav3* require exclusively Rac1-dependent (*Tacstd2*) or Rac1-dependent plus Rac1-independent (*Ptgs2*) routes. (iii) The synergistic targets can be regulated using either GTPase-independent (*Ilk*) or dependent (*Inhba*) routes (see summary in **Fig. S4I**).

To get a genome-wide estimation of the *Vav* family “common” (mRNAs whose abundance is indistinctly regulated by any of the two *Vav* proteins), “synergistic” (mRNAs whose abundance requires simultaneous signaling inputs from *Vav2* and *Vav3*), and “isoform-specific” (transcripts whose abundance is regulated exclusively by one *Vav*

family member) transcriptomal subsets, we compared the transcriptome of *Vav2;Vav3* knockdown cells, *Vav2*-, *Vav3*-, and *Vav2/Vav3*-reconstituted *Vav2;Vav3* knockdown cells using microarray analyses. The bioinformatics analysis of the data obtained revealed that 36.4%, 23.5% and 35.5% of the Vav family-dependent transcriptome could be assigned to the common-, synergistic-, and *Vav3*-dependent subclasses, respectively (**Fig. S5**). By contrast, the percentage of *Vav2*-specific genes was significantly lower in the overall Vav family-dependent transcriptome (4.6%, **Fig. S5**).

Vav-dependent targets have overlapping, but not identical roles in breast cancer

We next knocked down each of the six selected Vav target genes in 4T1 cells to address their roles in breast cancer (**Fig. S6, Table S3**). Similar to Vav proteins, we found that *Ilk*, *Tacstd2*, *InhβA* and *Cox2* were important for optimal tumor growth (**Fig. 5A**) and lung-specific metastasis (**Fig. 5B**). These data were obtained using two knockdown cell clones generated with two independent shRNAs for each target (**Fig. 5A,B**), thus ruling out the possibility of spurious, off-target effects. The analysis of tumors derived from those knockdown cell lines indicated that *Ilk*, *Tacstd2*, *InhβA* and *Cox2* were critical for cancer cell proliferation (**Fig. 5C,D**) and intratumoral angiogenesis (**Fig. 5E,F**). By contrast, *InhβA* and *Ilk* were the only selected targets with critical roles in vascular permeability and cancer cell survival, respectively (**Fig. 5G-J**). The elimination of *Itgβ6* or *Itgα8* did not show any deleterious effect in any of those biological parameters (**Fig. 5**).

The evaluation of the number of circulating cancer cells in tumor-bearing animals revealed that *InhβA* was the only target with proactive roles on cancer cell intravasation

(**Fig. 6A**). By contrast, the extravasation step was totally dependent on *Ilk*, *Tacstd2* and, to a much lower extent, on *InhβA* (**Fig. 6B,C**). Consistent with those data, long-term metastasis experiments using intravenously injected cancer cells indicated that both *Ilk* and *Tacstd2* were critical for the formation of metastatic nodules in the lung (**Fig. 6D-F**). These experiments also revealed that *InhβA* and *Cox2* were key for the formation of metastatic nodules in that tissue (**Fig. 6D-F**). Since those two proteins are not important for extravasation (see **Fig. 6B,C**), this result suggests that they play roles in the fitness of cancer cells once they reach the lung parenchyma. *Itgb6* and *Itga8* knockdown cells showed no defects in any of the above metastatic stages (**Fig. 6**).

The implication of *InhβA* and *Cox2* in post-extravasation stages led us to investigate whether Vav proteins and the other Vav targets could also regulate the fitness of cancer cells inside the lung parenchyma. Since the severe extravasation defect of *Vav2;Vav3*, *Ilk*, and *Tacstd2* knockdown cells precluded the direct examination of that issue, we artificially facilitated their extravasation in vivo by treating the mice with monocrotaline. This toxin increases the permeability of the lung vascular endothelium (31), so we speculated that it could facilitate the movement across the endothelial layer of the knockdown cells under study. Consistent with this hypothesis, we observed that monocrotaline favored the efficient extravasation of all the knockdown 4T1 cells tested, although it did not improve the already high extravasation rates of control cells (**Fig. 7A,B**). Despite this, *Vav2;Vav3*- and *Tacstd2*-deficient cells could not form lung metastases in monocrotaline-treated mice indicating that, similarly to *InhβA* and *Cox2*, those proteins participated in the intraparenchymal fitness of cancer cells (**Figs. 7C-E**). By contrast, *Ilk*

knockdown cells did form metastatic nodules in a monocrotaline-dependent manner (**Figs. 7C-E**), demonstrating that the function of *Ilk* is circumscribed just to the extravasation step. Taken together, these results indicate that *Vav2* and *Vav3* regulate many of the steps involved in the colonization of the lung by metastatic breast cancer cells and, in addition, that they do so by controlling the abundance of pro-metastatic distal targets using both common and synergistic routes (**Fig. 7F**).

The *Vav2/Vav3*-dependent transcriptome harbors prognostic gene signatures for breast cancer

To further investigate the potential clinical interest of the *Vav* route in breast cancer, we finally determined the prognostic value of the *Vav2/Vav3*-dependent transcriptome. The initial use of a small gene signature composed of *Vav2*, *Vav3*, *Ilk*, *Inhba*, *Tacstd2* and *Ptgs2* in human breast cancer datasets was not informative in terms of disease outcome prediction. Since the dual implication of their encoded products in primary tumorigenesis and metastasis could interfere with patient stratification into different prognostic categories, we decided to interrogate the entire *Vav2/Vav3*-dependent transcriptome to obtain clinically informative gene signatures. To this end, we used all the *Vav*-dependent transcripts that, according to the microarray experiments shown in **Figs. S4A** and **S5**, displayed consistent changes in abundance between the metastatic (control and *Vav2/Vav3*-reconstituted *Vav2;Vav3* knockdown cells) and non-metastatic cell lines (*Vav2;Vav3* knockdown cells) used in our study. Such strategy allowed us to eliminate transcripts whose abundance was not rescued by the re-introduction of *Vav2* and *Vav3* in the

Vav2;*Vav3* knockdown cell lines. In addition, it sieved out transcripts whose abundance were only modified under conditions of Vav protein overexpression. This approach led to the identification of 1,546 *Vav2*/*Vav3*-dependent mRNAs whose abundance correlated with the metastatic potential of our cell lines (**Fig. S7A**, **Table S6**). After mapping them to their human counterparts, we investigated whether their abundance was associated with disease outcome using human breast tumor mRNA microarray datasets. We found that patients with significant overlaps with the *Vav2*/*Vav3*-dependent mRNA signature had worse prognosis in terms of overall survival (**Table S7**), tumor recurrence (**Table S7**) and/or metastatic events (**Fig. S7B,C**; **Table S7**). To further confirm that *Vav2*/*Vav3*-regulated genes served as lung metastatic predictors, we next generated a minimal *Vav2*/*Vav3* prognostic lung metastatic signature (V2/V3LMS). By training our 1,546 mRNA signature with the Wang breast cancer dataset (32), we established a classifier group of 102 mRNAs capable of predicting lung metastasis in breast cancer (**Table S8**). When used to stratify patients, we observed that the V2/V3LMS could predict lung-metastasis-free survival in breast cancer patients in both the NKI (**Fig. S7D**) and MSKCC (**Fig. S7E**) cohorts (33, 34). By contrast, and in agreement with our biological data, it did not have any prognostic value in the case of bone metastasis-free survival (**Fig. S7F**). Interestingly, the V2/V3 LMS behaves similarly, or even better, than previously reported gene signatures with diagnostic value to forecast lung metastasis development in human patients (33, 34) (**Fig. S7G**).

DISCUSSION

Here, we have demonstrated that Vav2 and Vav3 regulate biological parameters critical for the growth of breast tumors and their metastatic dissemination to the lung (**Fig. 7F**). The effect of those proteins in primary tumorigenesis is not fully penetrant, because the depletion of both GEFs severely delays, but it does not abate, the breast tumor growth in the 4T1 model. By contrast, they are essential for lung metastasis, as assessed by the total absence of metastatic nodules or even micrometastases in animals orthotopically transplanted with *Vav2;Vav3* deficient breast cancer cells. Indeed, lung metastasis cannot develop even under very permissive experimental conditions such as the intravenous injection of large numbers of cancer cells in recipient mice. This is a tissue-specific function, because Vav-deficient cells do metastasize to other peripheral tissues. Such selectivity can be explained by the direct implication of Vav proteins in the extravasation of breast cancer cells through the lung vascular endothelium, a tight physical barrier not found in the highly fenestrated vessels of bone or liver (35). In addition, they also contribute to the fitness of breast cancer cells inside the lung parenchyma, a niche that poses biological challenges of its own for the ectopic proliferation and survival of cancer cells (35). The former function is perhaps the most relevant in this pathological context, since *Vav2;Vav3* deficient breast cancer cells only extravasate when the permeability of the endothelial barrier is artificially induced by pharmacological means (i.e., monocrotaline injections). We have also detected a role of Vav proteins in the intravasation step of metastatic breast cancer cells. Although such function is obviously not essential in our experimental setting, it may play more important roles in breast tumors with sizes more similar to those found

usually in the clinic. Favoring this idea, we have seen that the overexpression of Vav2 and Vav3 eliminates the defective intravasation properties of the non-metastatic 168FARN cancer breast cell line.

Our study has also revealed that Vav proteins control a large subset of the cancer cell transcriptome, including mRNAs encoding proteins that play critical roles in both primary tumorigenesis and lung-specific metastasis (**Fig. 7F**). This biological program is quite unique, because it does not have significant overlaps with previously described gene signatures associated to lung metastasis (33, 34). In addition, to our knowledge, no previous data have shown such a selective implication of other Rho/Rac GEFs in the tropism of any type of cancer cell to the lung. Vav2 and Vav3 control the aforementioned cancer-related processes using a combination of shared and synergistic routes that, in turn, rely on both Rac1-dependent and independent signals to become fully engaged (**Fig. 7C**). The analysis of the regulation of this transcript shows an unexpected complexity. Thus, some of the common (*Ptgs2*) and synergistic (*Inhba*) transcripts regulated by Vav proteins seem to rely on the engagement of both Rac1-dependent and independent signals for proper abundance regulation (**Fig. 7C**). Congruous with this view, we have seen that the overall positive effect of the constitutively active Rac1^{F28L} in the abundance of those two transcripts is highly dependent on the Vav2/Vav3 abundance status of breast cancer cells. Likewise, these two transcripts become more abundant in 168FARN cells upon the overexpression of Vav2 and Vav3 under conditions in which no detectable variations in overall Rac1 activity are seen. By contrast, the abundance of other Vav family common targets (*Tacstd2*) seem to rely exclusively on Rac1 signals in 4T1 cells (**Fig. 7C**). Finally, the abundance of other

transcripts that rely on synergistic signals from the two Vav proteins (*Ilk*) is controlled by Vav2 in a catalytic- and GTPase-independent manner (**Fig. 7C**). To add more complexity to this regulatory mechanism, we have observed that the rescue of the abundance of the synergistic *Inhba* transcript in *Vav2;Vav3* knockdown cells requires, in addition to wild type Vav3-derived signals, signaling contributions from the catalytic domain of Vav2. Since the overexpression of Vav3 is sufficient to rescue Rac1 activity in those cells, these results suggest that Vav3 and Vav2 may favor the specific activation of Rac1 in different subcellular localizations or, alternatively, that Vav2 may engage additional subsets of Rho/Rac proteins via its catalytic domain. These observations are consistent with other observations indicating that the knockdown of Vav2, but not Vav3, affects the proliferation and invasion of 4T1 cells in a GEF-dependent manner despite the fact that the overall Rac1 activity does not change in control, single *Vav2* knockdown, single *Vav3* knockdown, and *Vav2*^{R373A}-reconstituted *Vav2;Vav3* knockdown cells. Although all the targets identified in this work belong to the “common” and “synergistic” transcriptomal subclasses, it is quite possible that proteins whose abundance is modulated by either Vav2 or Vav3 using catalytic-dependent or independent routes could also participate in this process. The study in the near future of additional targets present in the large Vav2/Vav3-dependent transcriptomal signature of breast cancer cells will be required to address this possibility.

Consistent with the role of the Vav family in breast cancer cell biology, we have observed using metagenomic analysis that the abundance of the *VAV3* mRNA increases in samples from both specific breast cancer subtypes and immortalized breast cancer cell lines. Although the reason for the specific enrichment of the abundance of this Vav family

transcript remains to be addressed experimentally, we can speculate that it can be due to two different reasons. Firstly, our microarray analyses have shown that *Vav3* contributes to a much larger extent than *Vav2* to the regulation of the breast cancer *Vav* family-dependent transcriptome. This may indicate that, in addition to enhance the regulation of common and synergistic targets, the overexpression of *Vav3* could lead to a larger effect on the breast cancer cell transcriptome than *Vav2*. Secondly, the increase in abundance of the human *VAV3* mRNA in breast cancer cells can be a mere reflection of the specific inducible nature of its gene promoter. Consistent with this idea, we have reported previously that the expression of the *Vav3* gene, but not that of other *Vav* family genes, is highly dependent on the abundance of the aryl hydrocarbon receptor (Ahr) transcriptional factor in both mouse tissues and primary cells (36, 37). Consistent with this, *Vav3*^{-/-} and *Ahr*^{-/-} mice display very similar nervous system and cardiovascular defects (37). Additional experiments will be required to address this issue.

The present observations have also given indirect hints about the mechanistics of some of the metastatic steps of breast cancer cells. Thus, we have observed that the best biological parameter of the primary tumor that correlates with intravasation efficiency is the permeability of the intratumor blood vessels. Consistent with this view, we have seen that the two knockdown cells with more drastic defects on vascular leakiness (*Vav2*;*Vav3*- and *Inhba*-deficient 4T1 cells) are also the ones with the most severe problems in intravasation (**Fig. 7F**). In addition, unlike other unrelated pro-metastatic biological programs (30), we have seen that the intravasation and extravasation steps do not require the same exact subsets of *Vav*-dependent target genes (**Fig. 7F**). This result suggests that the extravasation

step is much more stringent functionally than the intravasation step, probably because the metastatic cells have to deal with a more naïve and tight endothelium in the lung than in the intratumoral context. Alternatively, this functional specificities may be related with the fact that the adhesion forces required to attach to the lung endothelium must be stronger, given that this adhesion step is counterbalanced by the shear forces induced by the blood flow inside the lung arterioles. Finally, we could not detect any obvious correlation between the defects found in the primary tumor and the long-term fitness of the metastatic cells in the lung parenchyma (**Fig. 7F**). This suggests that, despite the fact that both processes require the engagement of common biological processes such as angiogenesis and proliferation (38, 39), they are still controlled by different signaling mechanisms.

The circumscribed effect of Vav proteins to lung metastasis limits the potential interest of these GEFs and its distal downstream targets as general anti-metastatic drug targets, since it is clear that their inhibition will not limit the dissemination and growth of cancer cells in other peripheral tissues. Despite this, the observation that the elimination of Vav proteins, *InhβA*, *Tacstd2* and *Cox2* compromise the fitness of cancer cells in the lung parenchyma indicates that their inhibition could have some interest in the case of patients with metastasis circumscribed to the lung. In any case, the most obvious application of the present work is the observation that the Vav-dependent transcriptome harbors mRNA subsignatures capable of forecasting both disease outcome and lung metastasis in breast cancer patients. Prediction of this latter parameter is of obvious clinical interest, because that tissue is one of the main landing sites for metastatic breast cancer cells. In addition, it has a direct correlation with tumor recurrence and overall patient survival (40). The Vav-

dependent gene signature identified here has no prognostic value for bone metastasis, a result that further underscores the tight functional link between the Vav route and the tropism of breast cancer cells to the lung.

The specific effect of Vav proteins on the late stages of lung metastasis raises additional questions regarding the participation of Rho/Rac GEFs in breast cancer tumorigenesis. For example, it will be interesting to verify in the near future whether the tumorigenic and metastatic steps that are less Vav2;Vav3-dependent are so because other Rho/Rac GEFs are in charge of them. This is quite possible in the case of primary tumorigenesis, since a previous report has shown that P-Rex1, a phosphatidylinositol 3-kinase-dependent Rac1 GEF (2, 3) highly expressed in breast cancer ((8) and this work, **Fig. S1A-E**), is important for the growth of breast cancer cell lines in the primary tumor site (8). Thus, it is possible that P-Rex1, Vav2 and Vav3 would work in a combined fashion in that specific functional niche. At this point, however, there is no information about the role of P-Rex1 or other Rho/Rac GEFs in all the metastatic stages of breast cancer cells *in vivo*. It will be interesting therefore to conduct in the near future a functional screening with all the Rho/Rac GEFs that are either enriched or basally expressed in breast cancer (**Fig. S1A,B**) to dissect the specific contribution of each of them to all the biological parameters tested in the present work. That would reveal the Rho/Rac GEF-dependent stages of the metastatic process and, in addition, the specific functional tasks that breast cancer-expressed GEFs play in that tumor. Finally, another question that remains to be addressed is whether the roles of Vav2 and Vav3 unveiled in this work are specific just for breast cancer cells or, alternatively, whether they are common to other metastatic tumors

with tropism towards the lung. Future work will be needed to tackle all those lingering questions.

MATERIALS AND METHODS

Metagenomic analysis

The 54 Rho/Rac GEF family members were used together to interrogate the Oncomine™/Compendia Bioscience database (<https://www.oncomine.org/resource/login.html>) (41) in order to identify those with statistically significant enrichment in their abundance in specific types of human tumors, cancer cell lines or healthy tissues. Score criteria for significant enrichment included: *analysis type*, differential analysis; *P value* $\leq 1 \times 10^{-4}$; *fold change* ≥ 2 ; *gene rank*, top 1%; *tumor samples in microarray analysis* ≥ 151 . Visualization of mRNA profiles was done using the tools present in the Oncomine™/Compendia Bioscience. In all cases, the 20 top hitters were graphically shown regardless of the statistical significance of their variation among samples. In the case of breast cancer subtypes (**Table S1**), the screening criteria were the same as above, including the cancer type (breast cancer) as additional parameter. The analysis of the abundance of VAV2 and VAV3 mRNAs in luminal breast tumors (**Table S2**) was done using a microarray breast cancer dataset containing 508 samples (42). Intrinsic subtype classification and normalized microarray data were downloaded from the Gene Expression Omnibus (GEO) website (GSE25066). Z-values were calculated from log2-based microarray abundance data for both mRNAs and the mean and standard errors calculated using ANOVA.

Plasmid construction

To generate the lentiviral vector encoding HA-tagged Vav2 (pCCM33), a cDNA fragment encoding mouse wild type Vav2 was amplified by PCR using as template the pCMV-Vav2

plasmid (43). Primers were modified to introduce a *SpeI* site (underlined)/HA epitope (bold face) at the 5' end and a *NotI* site (underlined) at the 3' end in the amplified product. The sequence of the forward and reverse primers were 5'-AAT AAC TAG TGC CAC CAT **GTA CCC ATA CGA CGT CCC AGA CTA CGC** TGA GCA GTG GCG GCA ATG CGG C-3' and 5'-ATA GAC CGC GGC CGC TCA CTG GAT GCC CTC CTC TTC TAC GTA-3', respectively. The amplified PCR cDNA fragment was digested with the indicated enzymes and cloned into the *SpeI/NotI*-linearized lentiviral vector (pLVX-IRES-Hyg, Clontech). The pCCM33 vector was subsequently used to generate plasmids encoding catalytically inactive (R373A mutant, pCCM36 vector) and catalytically active (Y172F mutant, pCCM35 vector) versions of HA-tagged Vav2 using site-directed mutagenesis (Quickchange kit, Stratagene). Primers used to introduce those mutations were 5'-GTA CAT CAA TGA AGT GAA GGC GGA CAA GGA GAC CTT GAA G-3' (forward, for pCCM36), 5'-CTT CAA GGT CTC CTT GTC CGC CTT CAC TTC ATT GAT GTA C-3' (reverse, for pCCM36), 5'-GAT GAA GGA GAT GAC ATT TTT GAG GAC ATC ATC AGG GTG-3' (forward, for pCCM35) and 5'-CAC CCT GAT GAT GTC CTC AAA AAT GTC ATC TCC TTC ATC-3' (reverse, for pCCM35). To generate the lentiviral vector encoding Myc-tagged Vav3 (pCCM31), a cDNA fragment encoding mouse wild type Vav3 was amplified by PCR using the pC.HA-Vav3 plasmid (Addgene) as template. Primers used in the amplification step were modified to introduce a *XhoI* site (underlined) and a Myc-tag (bold face) at the 5' end as well as a *NotI* site (underlined) at the 3' end of the final cDNA product. The sequence of the forward and reverse primers were 5'-ATA CTC GAG GCC ACC ATG **GAA CAA AAA CTT ATT TCT GAA GAA GAT CTG** GAG CCG

TGG AAG CAG TGC GCT-3' and 5'-ATA GAC CGC GGC CGC TTA TTC ATC CTC TTC CAC ATA TGT GG-3', respectively. Upon digestion with *XhoI* and *NotI*, the *Vav3* cDNA was ligated into *XhoI/NotI*-linearized pLVX-IRES-Hyg. A bicistronic lentiviral vector encoding human wild type *Vav3* and GFP (pCQS1) has been described before (44). To generate a lentivirus encoding PyMT (pRL22), the *PyMT* cDNA was amplified from the pBabe-Hygro-PyMT (Addgene) and cloned into *EcoRI/BamHI*-linearized pHIV-ZsGreen (Addgene). The forward and reverse oligonucleotides for the amplification step were 5'-GCG CGC GAA TTC ATG GAT AGA GTT CTG AGC-3' and 5'-GCG CGC GGA TCC CTA GAA ATG CCG GGA AC-3', respectively (restriction sites are underlined). To generate the lentiviral vector encoding a fast cycling form of Rac1 (pMMM1), the mouse cDNA fragment encoding the mutated Rac1 sequence was amplified by PCR using as template the previously described pMIEG3-Rac1^{F28L} plasmid (45). Primers were modified to introduce a *XhoI* site (underlined) at the 5' end and a *BamHI* site (underlined) at the 3' end in the amplified product. The sequence of the forward and reverse primers were 5'-AGT CCT CGA GAT GCA GGC CAT CAA GTG TG-3' and 5'-AGT CGG ATC CGC ATT TTC TCT TCC TCT TCT TGA C-3', respectively. The amplified PCR cDNA fragment was digested with the indicated enzymes and cloned into the *XhoI/BamHI*-linearized pLVX-IRES-Hyg.

Cell culture

All cells were incubated at 37°C with 5% CO₂. 4T1 cells were cultured in Roswell Park Memorial Institute (RPMI) media supplemented with 10% fetal calf serum (FCS),

penicillin (10 µg/ml), and streptomycin (100 µg/ml). The non-metastatic 67NR, 4OT7 and 168FARN cell lines were gently provided by Dr. F. Miller (Wayne State University School of Medicine, Detroit, MI) (24) and maintained in culture in the above conditions. Cultures of primary mouse mammary epithelial cells were generated from 8- to 12-week-old wild-type or *Vav2*^{-/-};*Vav3*^{-/-} female mice (FVB background) (46), cultured in mammary epithelial cell primary growth medium (DMEM:F12, 5 µg/ml insulin, 5 ng/ml EGF, 10% FBS, 100 µg/ml streptomycin, 100 U/ml penicillin and 50 µg/ml gentamycin), and transduced in suspension with the pRL22 lentivirus as indicated (47). The Lenti-X 293T lentiviral packaging cell line (Clontech) was maintained in DMEM supplemented with 10% tetracycline-free FCS, penicillin (10 µg/ml) and streptomycin (100 µg/ml). To produce lentiviral particles, the appropriate vector was transfected into Lenti-X 293T cells using the Lenti-X HT packaging mix (Clontech). Viral particles were collected 48 hours after transfection and concentrated using the Lenti-X concentrator kit (Clontech). Viral titers were determined with the Lenti-X qRT-PCR Titration Kit (Clontech) according to the manufacturer's instructions.

Generation of knockdown cell lines

The shRNA-mediated knockdown of transcripts for *Vav2*, *Vav3* and the selected *Vav* family downstream targets was carried out using already packaged Mission TRC lentiviral particles (Sigma-Aldrich) according to the manufacturer's protocol. The catalogue numbers and shRNA sequences yielding the greatest knockdown were TRC0000097094 (5'-CCG GGC CTG CAT CTC TGG TTT AGA TCT CGA GAT CTA AAC CAG AGA TGC AGG

CTT TTT G-3') for the mouse *Vav2* mRNA; TRC0000097124 (5'-CCG GCC AGC ATT TCT CGT CTT AAA TCT CGA GAT TTA AGA CGA GAA ATG CTG GTT TTT G-3') for the mouse *Vav3* mRNA; TRCN0000067219 (5'-CCG GGC CGA GAG ATT CAA TGA AAT TCT CGA GAA TTT CAT TGA ATC TCT CGG CTT TTT G-3') for the mouse *Itgb6* mRNA; TRCN0000066240 (5'- CCG GCG GGA TGA GTT TCT CCT CTA TCT CGA GAT AGA GGA GAA ACT CAT CCC GTT TTT G-3') for the mouse *Itga8* mRNA; TRCN0000022516 (5'-CCG GGC TGA CAC CAA TGC AGT GAA TCT CGA GAT TCA CTG CAT TGG TGT CAG CTT TTT-3') and TRCN0000022515 (5'- CCG GGC ACG GAT TAA TGT GAT GAA TCT CGA GAT TCA TCA CAT TAA TCC GTG CTT TTT -3') for the mouse *Ilk* mRNA (A and B cell clones, respectively); TRCN0000112701 (5'-CCG GGC TCT TCC AAG AAC GCT ACA ACT CGA GTT GTA GCG TTC TTG GAA GAG CTT TTT G-3') and TRCN0000112700 (5'- CCG GGC TAG ATA GCT TGG TTA AGA ACT CGA GTT CTT AAC CAA GCT ATC TAG CTT TTT G -3') for mouse *Tacstd2* mRNA (A and B cell clones, respectively); TRCN0000067741 (5'-CCG GGC AAG GTC AAC ATT TGC TGT ACT CGA GTA CAG CAA ATG TTG ACC TTG CTT TTT G-3') and TRCN0000067738 (5'- CCG GGC TGT CAA GAA GCA CAT CTT ACT CGA GTA AGA TGT GCT TCT TGA CAG CTT TTT G-3') for mouse *Inhba* mRNA (A and B cell clones, respectively); and TRCN0000067938 (5'-CCG GGC ACA GGA TTT GAC CAG TAT ACT CGA GTA TAC TGG TCA AAT CCT GTG CTT TTT G-3') and TRCN0000067942 (5'- CCG GCC GTA CAC ATC ATT TGA AGA ACT CGA GTT CTT CAA ATG ATG TGT ACG GTT TTT G-3') for the mouse *Ptgs2* mRNA (A and B cell clones, respectively). Lentiviral particles containing the empty pLOK.1puro

vector (Sigma-Aldrich) were used to generate the “Control” 4T1 cells used in these experiments. Cells were incubated with lentiviral particles in the presence of 8 µg/ml polybrene (Sigma), selected with puromycin (3 µg/ml; Sigma). The designation and characteristics of the control and knockdown cell lines used in this study is summarized in **Table S3**.

Re-expression of Vav proteins in knockdown 4T1 cell lines

To generate single Vav2 and Vav3 rescued KD_{2/3(A)} 4T1 cells (see **Table S3**), this knockdown cell clone was infected with lentiviral particles encoding the indicated proteins in the presence of 8 µg/ml polybrene and then selected using 200 µg/ml hygromycin (Roche). As control, we carried out the same selection procedure with KD_{2/3(A)} 4T1 cells that were infected with lentivirus containing the empty pLVX-IRES-Hyg vector. To generate the rescued KD_{2/3(A)} cell line expressing either wild type Vav2 plus wild type Vav3 or Vav2^{R373A} plus wild type Vav3, Vav2;Vav3 knockdown cells were first infected with lentiviral particles produced from either the pCCM33 (encoding wild type HA-tagged Vav2) or the pCCM36 (encoding HA-tagged Vav2^{R373A}) vectors and then with lentiviral particles containing the pCQS1 vector (encoding wild type, HA-tagged Vav3). Cells were then selected with hygromycin (present in the Vav2-encoding vectors) and subsequently sorted by flow cytometry to isolate GFP positive cells (which was expressed bicistronically from the same Vav3-encoding pCQS1 vector (44)). The generation of 168FARN cells overexpressing wild type Vav2 and Vav3 was done following the above protocol. The

designation and features of the control and knockdown cell lines used in this study is summarized in **Table S3**.

Analysis of mRNA abundance

Total RNA was extracted from the indicated cells using Trizol (Sigma) and quantitative RT-PCR performed using the QuantiTect SYBR Green RT-PCR kit (Qiagen) and the iCycler machine (Bio-Rad) or, alternatively, the Script One-Step RT-PCR kit (BioRad) and the StepOnePlus Real-Time PCR System (Applied Biosystems). Raw data were then analyzed using either the iCycler iQ Optical System software (Bio-Rad) or the StepOne software v2.1 (Applied Biosystems). As normalization control, we used the abundance of the endogenous *P36b4* transcript. Primers used for transcript quantitation included 5'-AAG CCT GTG TTG ACC TTC CAG-3' (forward for mouse *Vav2*) and 5'-GTG TAA TCG ATC TCC CGG GAT-3' (reverse for mouse *Vav2*), 5'-AAT AGA TCT CCA GCA GTA C-3' (forward for mouse *Vav3*) and 5'-TGG TGT TGA ATG GCC CTT G-3' (reverse for mouse *Vav3*), 5'-TCA CGG CTT CCA GCT TTG GTC T-3' (forward for mouse *Itgb6*) and 5'-CTG CAC CAC CCC AGG CAC AG-3' (reverse for mouse *Itgb6*), 5'-TCC TCT GGG CTC CAC GGC TC-3' (forward for mouse *Itga8*) and 5'-ACG CAA CAG AGA CGC GCG AA-3' (reverse for mouse *Itga8*), 5'-AGG TCG GAA GGG AGG GAC CG-3' (forward for mouse *Ilk*) and 5'-GTC CAT AGC AGC GTC CCG GC-3' (reverse for mouse *Ilk*), 5'-CAC CAT GGC GAG GGG CTT GG-3' (forward for mouse *Tacstd2*) and 5'-CGC CTT GAG CAG CAG GCA CT-3' (reverse for mouse *Tacstd2*), 5'-GCC GAG TCA GGC ACA GCC AG-3' (forward for mouse *Inhba*) and 5'-TTC CTC GGC CTC ATC CCC CG-

3' (reverse for mouse *Inhba*), 5'- GGG CCC TTC CTC CCG TAG CA-3' (forward for mouse *Ptgs2*) and 5'-CCA TGG CCC AGT CCT CGG GT-3' (reverse for mouse *Ptgs2*), 5'- TAT GGG ACA CAG CTG GAC AA-3' (forward for mouse *Rac1*) and 5'-ACA GTG GTG TCG CAC TTC AG-3' (reverse for mouse *Rac1*), 5'-TTG ATG ATG GAG TGT GGC ACC-3' (forward for mouse *P36b4*) and 5'-GTG TTT GAC AAC GGC AGC ATT-3' (reverse for mouse *P36b4*).

Immunoblotting and GTPase activity assays

Immunoblots and GST-Pak1 pull-down experiments to determine Rac1 activity in cells were performed as indicated (48). Rac1 and RhoA activities were also quantified using the appropriate G-LISA activation kit (Cytoskeleton) according to the manufacturer's specifications (Cytoskeleton).

Animal studies with 4T1 and 168FARN cells

All animal work was done following protocols approved by the Bioethics committees of both the University of Salamanca and CSIC. 6-8 wk-old female Balb/c mice (Charles River) were used for the in vivo studies.

To carry out primary breast cancer tumorigenesis analysis, 5×10^3 viable cells were resuspended in 100 μ l of phosphate buffered saline solution (PBS) and injected orthotopically into the fourth right mammary gland of the recipient mouse. Primary tumor growth was analyzed by measuring tumor length, width and depth, and by calculating

tumor volume. Metastatic dissemination of tumor cells was quantified both de visu and histologically after sacrificing the mice 35 days after orthotopic implantation.

To carry out direct, long-term lung colonization assays, cells were resuspended in PBS as above and injected into the lateral tail vein (unless otherwise stated, 5×10^4 cells were injected in the experiments). After 2 weeks, mice were euthanized, the chest cavity exposed through a midline chest incision, the trachea cannulated with a 20-gauge catheter, and lungs slowly inflated using 1 ml of India ink (Parker, 1:16 dilution in PBS). Lungs were extracted, immersed in Fekete's solution (100 ml 70% ethanol (Sigma), 10 ml 4% formaldehyde (Leica), and 5 ml 100% glacial acetic acid (Sigma)) to destain and metastatic nodules counted de visu. Alternatively lungs were extracted, fixed in 4% formaldehyde, and embedded in paraffin for hematoxylin/eosin staining. In the case of studies conducted with monocrotaline-treated mice, animals were injected twice intravenously with monocrotaline (1.5 mg/ml, Sigma) the week before the introduction of cancer cells through the tail vein. After 2 weeks, animals were sacrificed and lungs extracted as indicated above.

Vascular permeability was evaluated indirectly by measuring the effusion in vivo of a fluorescent-labeled dextran from the blood vessels into the primary tumor. For that purpose, a rhodamine-conjugated 70 kDa dextran (Invitrogen) was injected intravenously (2 mg per 20 g body weight) in tumor-bearing mice. 1 hour later, mice were euthanized, the primary breast tumors extracted, cut in 30 μ m-thick sections, and dextran signals monitored by confocal microscopy (TCS SP5, Leica). Images were collected using the

LAS AF software (Leica) and fluorescence intensities quantified using the ImageJ software (NIH).

To measure the intravasation of cancer cells, we collected ≈ 1 ml of blood from the heart atrium of tumor-bearing mice and cells pelleted by centrifugation. After the elimination of red cells by two lysis cycles in ACK buffer (Gibco), we extracted total RNA from the remaining blood cells using Trizol (Sigma) to be used in qRT-PCR experiments to quantify the presence of circulating tumor cells. This was done by calculating the abundance of the mRNA for the puromycin resistance gene (present exclusively in all the 4T1 cells used in this study) in relation to that of an endogenous mouse mRNA (*B2m*). Primers used in these experiments were 5'-GTGTGGGCCCTGTTCTGCC-3' (forward viral *Puro^R* transcript); 5'-GCGCACCGTGGGCTTGACT-3' (reverse for viral *Puro^R* transcript); 5'-ATGGCTCGCTCGGTGACCCT-3' (forward for *B2m* mRNA) and 5'-CCATTCTCCGGTGGGTGGCG-3' (reverse for *B2m* mRNA). Values were corrected according to the abundance values of the viral gene in each cell line when maintained in tissue culture.

To measure the extravasation of breast cancer cells, the indicated cell lines were labeled with 5 μ M of a green cell tracker (CellTracker green CMFDA, Invitrogen) for 1 hour and injected intravenously into mice (1×10^6 cells/mice). After 48 hours, mice were injected intravenously with a rhodamine-conjugated lectin (Vector Laboratories) to stain lung capillaries and killed 60 min later. Lungs were then extracted, fixed in paraformaldehyde, and examined by confocal fluorescence microscopy to visualize the number of cancer cells (green color) present in the lung.

Animal studies with primary breast epithelial cells

Normal, primary breast epithelial cells were obtained from either wild type or *Vav2^{-/-};Vav3^{-/-}* female mice (3 to 12 week old; FVB background) as indicated above and, after transduction with the PyMT-encoding pRL22 lentivirus, were reimplanted into the cleared inguinal mammary fat pads of wild type female mice as described (47).

Immunostaining techniques

All immunohistochemical procedures presented in this work were carried out by the personnel of the Molecular Pathology Unit of our Institution. Tissues were extracted, fixed in 4% paraformaldehyde (Sigma), cut in 2-3 μ m thick sections, and stained with hematoxylin/eosin. For immunohistochemical staining, paraffin-embedded sections were incubated with rabbit polyclonal antibodies to either Ki67 (Novocastra, 1:100 dilution) or CD31 (Abcam, 1:50 dilution). After an overnight incubation, tissue slides were rinsed with PBS, incubated for 1 hour at room temperature in a milk/PBS solution containing a goat anti-rabbit IgG-horseradish peroxidase (GE Healthcare), rinsed with PBS, and developed with diaminobenzidine (Dako). Quantification of signals was done blindly using the Metamorph-Metaview software (Universal Imaging). Apoptotic cells were detected using the TUNEL-based In Situ Cell Detection kit (Roche). To this end, sections were deparaffinized, hydrated, digested with proteinase K (Dako) for 30 min at 37°C, subjected to the TUNEL reaction according to the manufacturer's instructions, and TUNEL positive cells scored de visu using a standard immunofluorescence microscope (CTR600, Leica).

Determination of in vitro proliferation rates

We used the 3-(4,5-dimethylthiazol-2-yl) 2,5-diphenyltetrazolium bromide (MTT) assay (Sigma) to determine the growth properties of cell lines under study. To this end, cells were plated onto 24-well dishes (7,000 cells/well) and cultured in RPMI supplemented with 10% calf serum for the indicated periods of time. At the indicated time-point, the culture medium of each well was discarded and replaced by 250 µl of an MTT solution (0.5 mg/ml) in PBS. After 1 hour at 37°C in a 5% CO₂ atmosphere, 500 µl of dimethyl sulfoxide (Sigma)/well were added to dissolve the formazan crystals formed and the absorbance at the 570 nm wavelength measured 15 min later using a microplate reader (Ultraevolution, Tecan).

Cell invasion assays

The ability of the cells to invade was assessed using CytoSelect 24-Well Cell Invasion assay (Cell Biolabs) according to the manufacturer's protocol. Briefly, 3×10^5 cells in serum-free medium were plated onto the basement membrane layer of the cell culture insert of the invasion chamber. After incubation for 48 hours at 37°C in 5% CO₂, the inserts were stained with Cell Stain solution (Cell Biolabs), incubated with Extraction Solution (Cell Biolabs) and the absorbance at the 560 nm wavelength measured 10 min later using the Ultraevolution microplate reader.

Microarray analyses

All microarray experiments were performed by the personnel of the Genomics and Proteomics Unit of our Institution. Transcriptomal changes were determined using the Mouse Gene 1.0 ST arrays (Affymetrix). Two independent experiments were performed to identify the Vav2/Vav3-dependent transcriptome. In the first one, we compared the transcriptomes of Control, KD_{2/3(A)} and KD_{2/3(B)} (**Table S3**) to identify Vav family-dependent genes. In the second one, we compared the transcriptomes of KD_{2/3(A)}, KD_{2/3+}V₂, KD_{2/3+}V₃ and KD_{2/3+}V₂+V₃ cells (**Table S3**). In all cases, total cellular RNA was extracted from three independent exponential cultures of the appropriate cell lines using the RNeasy kit (Qiagen), quantified using 6000 Nano Chips (Agilent Technologies), and used (2.3 µg/sample) to generate labeled cRNA probes according to the manufacturer's instructions (Affymetrix). Upon microarray hybridization, raw data were normalized, filtered and analyzed with the Bioconductor software (www.bioconductor.com) using the *Affy* and *Siggenes* applications. For the graphical presentation of microarray data, we performed hierarchical clustering analysis using the Bioconductor HCLUST tool. Details about those procedures can be found elsewhere (49, 50).

Criteria for selecting Vav-dependent target genes for functional characterization

We used a number of sieving criteria to select the Vav targets to be knocked down in parental 4T1 cells using shRNA-based strategies. Firstly, we decided to only focus on mRNAs whose abundance was reduced in Vav2;Vav3-deficient cells, since we reasoned that it would be easier to cripple the prometastatic program of 4T1 cells by eliminating

single proteins with positive roles in this process rather than overexpressing single negative metastatic regulators. Likewise, from a clinical and therapeutic point of view, it is easier to treat patients with drug inhibitors rather than by gain-of-function approaches. Keeping in mind this clinical-oriented view, we also decided to select mRNAs encoding products potentially amenable to either chemical (i.e., enzymes) or antibody-based (i.e., surface molecules) inhibitory strategies. Furthermore, we favored genes encoding proteins whose inhibition could have a major impact in the biology of the metastatic cancer cell due to their enzymatic, adaptor and/or signaling properties. We also took into consideration additional empirical criteria for selecting those targets. Thus, we were interested on selecting targets displaying: **(i)** High fold change variations between the control and double *Vav2*;*Vav3*-deficient 4T1 cells. **(ii)** Homogenous behavior in the two independent clones of *Vav2*;*Vav3*-deficient 4T1 cells. **(iii)** Similar variations in vivo when using microarray-independent techniques (i.e., qRT-PCR). **(iv)** Similar abundance changes in cultured cells and in the tumors derived from them in recipient mice. Some of the genes initially selected for analysis did not fulfill some of the latter criteria. For example, *Igfr2*, *PIP5K1a*, *Plau*, or *Mns* did not passed criteria # iii. Likewise, *Adamts1*, *Gadd45b* and *Bgalt6* did not fulfill criteria # iv (they were deregulated in cultured *Vav2*;*Vav3*-deficient cells but not in primary tumors derived from them). After all those verification steps, we selected 6 putative target genes (*Itgb6*, *Itga8*, *Ilk*, *Tacstd2*, *Inhba*, *Ptgs2*) for further characterization in the present work.

Calculation of the percentage of common, synergistic, and Vav family member-specific transcripts in the Vav family-dependent transcriptome of breast cancer cells

These transcriptomal subsets were calculated using Pavlidis Template Matching (PTM) analysis from the microarray data obtained from $KD_{2/3(A)}$, $KD_{2/3(B)}$, $KD_{2/3+V_2}$, $KD_{2/3+V_3}$ and $KD_{2/3+V_2+V_3}$ cells, using a Q threshold significance cut-off value ≤ 0.0025 . Control samples were included in the heatmaps of each cluster after that statistical analysis.

Overlapping analysis with human cancer microarray studies

The OncomineTM/ Compendia Bioscience database (41) was used to search for general overlaps between the Vav2/Vav3-dependent transcriptome and clinical data from human breast cancer datasets. To this end, the microarray data obtained in our two independent experiments (control vs $KD_{2/3(A)}$ and $KD_{2/3(B)}$; $KD_{2/3(A)}$ vs $KD_{2/3+V_2/V_3}$) were analyzed side by side bioinformatically to identify transcripts showing consistent and specular changes between those two analyses and that, therefore, could be directly linked to the known metastatic properties of each of those cell lines. Briefly, the abundance of each mRNA was transformed to z-scores (mean value = 0, standard deviation = 1) in each independent 4T1 cell line experiment dataset to allow differential abundance analysis using a unique dataset file. We performed PTM analysis (51), in which the Pearson's correlation coefficient is computed among the intensities measured for each transcript and the values of an independent variable. The threshold significance values used in those analyses where: Q value ≤ 0.0025 ; and Pearson coefficient either ≥ 0.82 (in the case of transcripts whose abundance was increased in reconstituted $KD_{2/3+V_2/V_3}$ and control cells) or ≤ -0.82 (in the

case of transcripts was abundance was reduced in reconstituted KD_{2/3+V2/V3} and control cells).

The mouse Probe IDs for the identify transcripts in the above analyses were then humanized using the Ailun web tool (<http://ailun.stanford.edu/>) (52) and loaded onto the OncomineTM/Compendia database. Association of the mapped Vav2/Vav3-signature with clinical parameters was tested using Fisher's exact test and it was considered significant for Odds Ratio ≥ 1.5 and $P \leq 0.0001$. Overlapping mRNAs with the Loi (53) and NKI (33) datasets were used to stratify the patients. To this end, the sum of the abundance values of mRNAs that were induced in both the Vav2/Vav3-signature and in metastatic human breast tumors was obtained and subtracted with the sum of the abundance values of mRNA repressed in the Vav2/Vav3-signature and metastatic human breast tumors. The value obtained (Vav/Vav3-dependent signature score) was used to divide the patients into high and low score groups and survival probabilities were obtained for each patient group.

Development of a prognostic Vav2/Vav3 gene signature for lung-specific metastasis in breast cancer patients

Raw microarray data from the Wang dataset (32) (GEO identifier GSE2034; $n = 286$ patients) was obtained and signal intensity values generated using the RMAExpress software (<http://rmaexpress.bmbolstad.com>). The censored lung metastasis at 5 years as endpoint from that dataset was then used to perform a Cox regression proportional hazard analysis with the humanized Vav2/Vav3 mRNA signature described above. This analysis yields a Cox coefficient (which gives values above or below 0 if the abundance of an

mRNA is increased or decreased in metastatic tissue, respectively) and, in addition, performs a Wald test (54, 55) to check the null hypothesis of the Cox coefficient being 0. This analysis resulted in the identification of a minimal Vav2/Vav3 lung metastasis signature (referred to hereafter as V2/V3LMS) composed of 120 probesets representing 102 independent mRNAs with significant Cox values ($P \leq 0.05$). 76 (with increase abundance in $KD_{2/3+V2/V3}$ cells) and 26 (in decreased abundance in $KD_{2/3+V2/V3}$ cells) of those probesets displayed Cox coefficients above and below 0, respectively (**Table S7**).

To test the prognostic value of the V2/V3LMS, the V2/V3LMS Risk Score was calculated using data from each of the patients present in the NKI ($n = 295$) (33) and MSKCC ($n = 83$) (34) datasets. In the case of the MSKCC dataset (generated using Affymetrix HG-U133A arrays), this step was done using the gene $\log_2$ expression values (x_i) and Wald statistics (s_i) using the following formula:

$$\text{V2/V3LMS Risk Score} = \sum_{i=1}^{120} s_i \cdot x_i$$

In the case of the NKI dataset (generated using the Agilent array platform), the mean for the s_i values associated with different Affymetrix probesets of each mRNA was first determined so that only a single s_i value was assigned for transcripts in the Agilent array that had more than one probeset in the Affymetrix Mouse Gene 1.0 ST arrays. The Risk Score for each patient was then calculated based on the abundance values of V2/V3LMS mRNAs (with the exception of six of them which were not present in the Agilent arrays: *AQP1*, *C14orf159*, *CCDC21*, *ELOVL1*, *RBM34* and *TINAGLI*). Risk values were used to subdivide patients into high (percentile 33) and low risk (percentiles 66 and

100) subsets and, subsequently, a Kaplan-Meier analysis was performed using lung-specific (in the case of NKI's and MSKCC's datasets) or bone-specific metastasis (in the case of the MSKCC's dataset) as final endpoints.

The comparison of the predictive value of the V2/V3 LMS with previously described mRNA signatures (33, 34) was done as indicated above.

Statistics. Depending on the type of experiment, data obtained in wet lab experiments were analyzed using either the Students's *t* or the Mann-Whitney test. Values are given as mean $\pm$ sem. *P* values ≤ 0.05 , 0.01 and 0.001 were indicated by *, ** and *** in all figures, respectively. Unless otherwise stated, *n* refers to independent experiments, each performed in triplicate.

SUPPLEMENTARY MATERIALS

Fig. S1. Rho/Rac GEF transcripts have difference abundance in human tumors.

Fig. S2. Abundance of Vav family mRNAs and proteins in cell lines used in this study.

Fig. S3. Vav proteins control a large transcriptomal program in breast cancer cells.

Fig. S4. Mechanistics of the regulation of the breast cancer cell transcriptome by Vav proteins.

Fig. S5. Vav proteins control the breast cancer cell transcriptome using common, synergistic, and Vav family member-specific routes.

Fig. S6. Generation of knockdown cell lines for Vav family-dependent targets.

Fig. S7. Vav-dependent mRNA signatures have prognostic value in breast cancer.

Table S1. Differential abundance of Rho/Rac GEF in human breast cancer subtypes.

Table S2. Abundance of *Vav2* and *Vav3* mRNAs in the breast cancer luminal A and B subtypes.

Table S3. Cell lines used in this study.

Table S4. Metastasis of indicated 4T1 knockdown cell clones in recipient mice.

Table S5. Transcriptomal changes in 4T1 cells lacking Vav2 and Vav3.

Table S6. The Vav2/Vav3-dependent transcriptome associated to the metastatic potential of the 4T1 cells used in this study.

Table S7. Overexpressed and repressed genes from **Table S5** associated with human cancer clinical outcome.

Table S8. Genes belonging to the Vav2/Vav3-lung metastasis signature.

REFERENCES AND NOTES

1. D. Vigil, J. Cherfils, K. L. Rossman, C. J. Der, Ras superfamily GEFs and GAPs: Validated and tractable targets for cancer therapy? *Nat Rev Cancer* 10, 842-857 (2010).
2. J. L. Bos, H. Rehmann, A. Wittinghofer, GEFs and GAPs: Critical elements in the control of small g proteins. *Cell* 129, 865-877 (2007).
3. K. L. Rossman, C. J. Der, J. Sondek, GEF means go: Turning on rho gtpases with guanine nucleotide-exchange factors. *Nat Rev Mol Cell Biol* 6, 167-180 (2005).
4. A. Malliri, R. A. van der Kammen, K. Clark, M. van der Valk, F. Michiels, J. G. Collard, Mice deficient in the Rac activator Tiam1 are resistant to Ras-induced skin tumours. *Nature* 417, 867-871 (2002).
5. K. H. Chang, A. Sanchez-Aguilera, S. Shen, A. Sengupta, M. N. Madhu, A. M. Ficker, S. K. Dunn, A. M. Kuenzi, J. L. Arnett, R. A. Santho, X. Agirre, J. P. Perentesis, M. W. Deininger, Y. Zheng, X. R. Bustelo, D. A. Williams, J. A. Cancelas, Vav3 collaborates with p190-bcr-abl in lymphoid progenitor leukemogenesis, proliferation and survival. *Blood*, in press (2012).
6. A. Malliri, T. P. Rygiel, R. A. van der Kammen, J. Y. Song, R. Engers, A. F. Hurlstone, H. Clevers, J. G. Collard, The Rac activator Tiam1 is a Wnt-responsive gene that modifies intestinal tumor development. *J Biol Chem* 281, 543-548 (2006).
7. Y. Kawasaki, S. Tsuji, K. Muroya, S. Furukawa, Y. Shibata, M. Okuno, S. Ohwada, T. Akiyama, The adenomatous polyposis coli-associated exchange factors Asef and Asef2 are required for adenoma formation in APC(min/+) mice. *EMBO Rep* 10, 1355-1362 (2009).

8. M. S. Sosa, C. Lopez-Haber, C. Yang, H. Wang, M. A. Lemmon, J. M. Busillo, J. Luo, J. L. Benovic, A. Klein-Szanto, H. Yagi, J. S. Gutkind, R. E. Parsons, M. G. Kazanietz, Identification of the Rac-GEF P-Rex1 as an essential mediator of Erbb signaling in breast cancer. *Mol Cell* 40, 877-892 (2010).
9. B. Fine, C. Hodakoski, S. Koujak, T. Su, L. H. Saal, M. Maurer, B. Hopkins, M. Keniry, M. L. Sulis, S. Mense, H. Hibshoosh, R. Parsons, Activation of the PI3K pathway in cancer through inhibition of pten by exchange factor P-Rex2a. *Science* 325, 1261-1265 (2009).
10. E. K. Thomas, J. A. Cancelas, Y. Zheng, D. A. Williams, Rac GTPases as key regulators of p210-Bcr-Abl-dependent leukemogenesis. *Leukemia* 22, 898-904 (2008).
11. K. Uhlenbrock, A. Eberth, U. Herbrand, N. Daryab, P. Stege, F. Meier, P. Friedl, J. G. Collard, M. R. Ahmadian, The RacGEF Tiam1 inhibits migration and invasion of metastatic melanoma via a novel adhesive mechanism. *J Cell Sci* 117, 4863-4871 (2004).
12. C. R. Lindsay, S. Lawn, A. D. Campbell, W. J. Faller, F. Rambow, R. L. Mort, P. Timpson, A. Li, P. Cammareri, R. A. Ridgway, J. P. Morton, B. Doyle, S. Hegarty, M. Rafferty, I. G. Murphy, E. W. McDermott, K. Sheahan, K. Pedone, A. J. Finn, P. A. Groben, N. E. Thomas, H. Hao, C. Carson, J. C. Norman, L. M. Machesky, W. M. Gallagher, I. J. Jackson, L. Van Kempen, F. Beermann, C. Der, L. Larue, H. C. Welch, B. W. Ozanne, O. J. Sansom, P-Rex1 is required for efficient melanoblast migration and melanoma metastasis. *Nat Commun* 2, 555 (2011).

13. K. Zhou, Y. Wang, J. L. Gorski, N. Nomura, J. Collard, G. M. Bokoch, Guanine nucleotide exchange factors regulate specificity of downstream signaling from Rac and Cdc42. *J Biol Chem* 273, 16782-16786 (1998).
14. R. J. Buchsbaum, B. A. Connolly, L. A. Feig, Interaction of Rac exchange factors Tiam1 and Ras-GEF1 with a scaffold for the p38 mitogen-activated protein kinase cascade. *Mol Cell Biol* 22, 4073-4085 (2002).
15. R. J. Buchsbaum, B. A. Connolly, L. A. Feig, Regulation of p70 S6 kinase by complex formation between the Rac guanine nucleotide exchange factor (Rac-GEF) Tiam1 and the scaffold spinophilin. *J Biol Chem* 278, 18833-18841 (2003).
16. A. Saveliev, L. Vanes, O. Ksionda, J. Rapley, S. J. Smerdon, K. Rittinger, V. L. Tybulewicz, Function of the nucleotide exchange activity of Vav1 in T cell development and activation. *Sci Sig* 2, ra83 (2009).
17. M. R. Schiller, A. Blangy, J. Huang, R. E. Mains, B. A. Eipper, Induction of lamellipodia by kalirin does not require its guanine nucleotide exchange factor activity. *Exp Cell Res* 307, 402-417 (2005).
18. M. R. Kuhne, G. Ku, A. Weiss, A guanine nucleotide exchange factor-independent function of Vav1 in transcriptional activation. *J Biol Chem* 275, 2185-2190 (2000).
19. X. R. Bustelo, Regulatory and signaling properties of the Vav family. *Mol Cell Biol* 20, 1461-1477 (2000).
20. P. Crespo, K. E. Schuebel, A. A. Ostrom, J. S. Gutkind, X. R. Bustelo, Phosphotyrosine-dependent activation of Rac-1 GDP/GTP exchange by the Vav proto-oncogene product. *Nature* 385, 169-172 (1997).

21. K. E. Schuebel, N. Movilla, J. L. Rosa, X. R. Bustelo, Phosphorylation-dependent and constitutive activation of rho proteins by wild-type and oncogenic Vav-2. *EMBO J* 17, 6608-6621 (1998).
22. N. Movilla, X. R. Bustelo, Biological and regulatory properties of Vav-3, a new member of the vav family of oncoproteins. *Mol Cell Biol* 19, 7870-7885 (1999).
23. J. Baselga, S. M. Swain, Novel anticancer targets: Revisiting ErbB2 and discovering ErbB3. *Nat Rev Cancer* 9, 463-475 (2009).
24. C. J. Aslakson, F. R. Miller, Selective events in the metastatic process defined by analysis of the sequential dissemination of subpopulations of a mouse mammary tumor. *Cancer Res* 52, 1399-1405 (1992).
25. R. O. Hynes, Integrins: Bidirectional, allosteric signaling machines. *Cell* 110, 673-687 (2002).
26. S. A. Wickstrom, A. Lange, E. Montanez, R. Fassler, The ilk/pinch/parvin complex: The kinase is dead, long live the pseudokinase! *EMBO J* 29, 281-291 (2010).
27. R. Cubas, M. Li, C. Chen, Q. Yao, Trop2: A possible therapeutic target for late stage epithelial carcinomas. *Biochim Biophys Acta* 1796, 309-314 (2009).
28. D. J. Bernard, S. C. Chapman, T. K. Woodruff, Mechanisms of inhibin signal transduction. *Recent Prog Horm Res* 56, 417-450 (2001).
29. R. J. Flower, The development of Cox2 inhibitors. *Nat Rev Drug Discov* 2, 179-191 (2003).

30. G. P. Gupta, D. X. Nguyen, A. C. Chiang, P. D. Bos, J. Y. Kim, C. Nadal, R. R. Gomis, K. Manova-Todorova, J. Massague, Mediators of vascular remodelling co-opted for sequential steps in lung metastasis. *Nature* 446, 765-770 (2007).
31. M. Kido, T. Hirose, K. Tanaka, T. Kurozumi, Y. Shoyama, Increased alveolar-capillary membrane permeability by monocrotaline. *Jpn J Med* 20, 170-179 (1981).
32. Y. Wang, J. G. Klijn, Y. Zhang, A. M. Sieuwerts, M. P. Look, F. Yang, D. Talantov, M. Timmermans, M. E. Meijer-van Gelder, J. Yu, T. Jatkoa, E. M. Berns, D. Atkins, J. A. Foekens, Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet* 365, 671-679 (2005).
33. M. J. van de Vijver, Y. D. He, L. J. van't Veer, H. Dai, A. A. Hart, D. W. Voskuil, G. J. Schreiber, J. L. Peterse, C. Roberts, M. J. Marton, M. Parrish, D. Atsma, A. Witteveen, A. Glas, L. Delahaye, T. van der Velde, H. Bartelink, S. Rodenhuis, E. T. Rutgers, S. H. Friend, R. Bernards, A gene-expression signature as a predictor of survival in breast cancer. *New Eng J Med* 347, 1999-2009 (2002).
34. A. J. Minn, G. P. Gupta, P. M. Siegel, P. D. Bos, W. Shu, D. D. Giri, A. Viale, A. B. Olshen, W. L. Gerald, J. Massague, Genes that mediate breast cancer metastasis to lung. *Nature* 436, 518-524 (2005).
35. D. X. Nguyen, P. D. Bos, J. Massague, Metastasis: From dissemination to organ-specific colonization. *Nat Rev Cancer* 9, 274-284 (2009).
36. J. M. Carvajal-Gonzalez, S. Mulero-Navarro, A. C. Roman, V. Sauzeau, J. M. Merino, X. R. Bustelo, P. M. Fernandez-Salguero, The dioxin receptor regulates the constitutive

- expression of the Vav3 proto-oncogene and modulates cell shape and adhesion. *Mol Biol Cell* 20, 1715-1727 (2009).
37. V. Sauzeau, J. M. Carvajal-Gonzalez, A. S. Riolobos, M. A. Sevilla, M. Menacho-Marquez, A. C. Roman, A. Abad, M. J. Montero, P. Fernandez-Salguero, X. R. Bustelo, Transcriptional factor aryl hydrocarbon receptor (Ahr) controls cardiovascular and respiratory functions by regulating the expression of the Vav3 proto-oncogene. *J Biol Chem* 286, 2896-2909 (2011).
 38. G. P. Gupta, J. Massague, Cancer metastasis: Building a framework. *Cell* 127, 679-695 (2006).
 39. S. Valastyan, R. A. Weinberg, Tumor metastasis: Molecular insights and evolving paradigms. *Cell* 147, 275-292 (2011).
 40. E. F. Solomayer, I. J. Diel, G. C. Meyberg, C. Gollan, G. Bastert, Metastatic breast cancer: Clinical course, prognosis and therapy related to the first site of metastasis. *Breast Cancer Res Treat* 59, 271-278 (2000).
 41. D. R. Rhodes, S. Kalyana-Sundaram, V. Mahavisno, R. Varambally, J. Yu, B. B. Briggs, T. R. Barrette, M. J. Anstet, C. Kincead-Beal, P. Kulkarni, S. Varambally, D. Ghosh, A. M. Chinnaiyan, Oncomine 3.0: Genes, pathways, and networks in a collection of 18,000 cancer gene expression profiles. *Neoplasia* 9, 166-180 (2007).
 42. C. Hatzis, L. Pusztai, V. Valero, D. J. Booser, L. Esserman, A. Lluch, T. Vidaurre, F. Holmes, E. Souchon, H. Wang, M. Martin, J. Cotrina, H. Gomez, R. Hubbard, J. I. Chacon, J. Ferrer-Lozano, R. Dyer, M. Buxton, Y. Gong, Y. Wu, N. Ibrahim, E. Andreopoulou, N. T. Ueno, K. Hunt, W. Yang, A. Nazario, A. DeMichele, J.

- O'Shaughnessy, G. N. Hortobagyi, W. F. Symmans, A genomic predictor of response and survival following taxane-anthracycline chemotherapy for invasive breast cancer. *JAMA* 305, 1873-1881 (2011).
43. L. Rodrigues, M. Pires de Miranda, M. J. Caloca, X. R. Bustelo, J. P. Simas, Activation of Vav by the gammaherpesvirus M2 protein contributes to the establishment of viral latency in b lymphocytes. *J Virol* 80, 6123-6135 (2006).
44. C. Quevedo, V. Sauzeau, M. Menacho-Marquez, A. Castro-Castro, X. R. Bustelo, Vav3-deficient mice exhibit a transient delay in cerebellar development. *Mol Biol Cell* 21, 1125-1139 (2010).
45. F. Guo, Y. Zheng, Rho family GTPases cooperate with p53 deletion to promote primary mouse embryonic fibroblast cell invasion. *Oncogene* 23, 5577-5585 (2004).
46. V. Sauzeau, M. Jerkic, J. M. Lopez-Novoa, X. R. Bustelo, Loss of Vav2 proto-oncogene causes tachycardia and cardiovascular disease in mice. *Mol Biol Cell* 18, 943-952 (2007).
47. R. M. Larive, A. Abad, C. M. Cardaba, T. Hernandez, M. Canamero, E. de Alava, E. Santos, B. Alarcon, X. R. Bustelo, The ras-like protein R-Ras2/TC21 is important for proper mammary gland development. *Mol Biol Cell* 23, 2373-2387 (2012).
48. A. Castro-Castro, V. Ojeda, M. Barreira, V. Sauzeau, I. Navarro-Lerida, O. Muriel, J. R. Couceiro, F. X. Pimentel-Muinos, M. A. Del Pozo, X. R. Bustelo, Coronin 1a promotes a cytoskeletal-based feedback loop that facilitates Rac1 translocation and activation. *EMBO J* 30, 3913-3927 (2011).

49. I. M. Berenjeno, F. Nunez, X. R. Bustelo, Transcriptomal profiling of the cellular transformation induced by Rho subfamily gtpases. *Oncogene* 26, 4295-4305 (2007).
50. I. M. Berenjeno, X. R. Bustelo, Identification of the Rock-dependent transcriptome in rodent fibroblasts. *Clin Transl Oncol* 10, 726-738 (2008).
51. P. Pavlidis, W. S. Noble, Analysis of strain and regional variation in gene expression in mouse brain. *Genome Biol* 2, Research0042.0041-0042.0015 (2001).
52. R. Chen, L. Li, A. J. Butte, Ailun: Reannotating gene expression data automatically. *Nat Meth* 4, 879 (2007).
53. S. Loi, B. Haibe-Kains, C. Desmedt, P. Wirapati, F. Lallemand, A. M. Tutt, C. Gillet, P. Ellis, K. Ryder, J. F. Reid, M. G. Daidone, M. A. Pierotti, E. M. Berns, M. P. Jansen, J. A. Foekens, M. Delorenzi, G. Bontempi, M. J. Piccart, C. Sotiriou, Predicting prognosis using molecular profiling in estrogen receptor-positive breast cancer treated with tamoxifen. *BMC Genomics* 9, 239 (2008).
54. S. D. Silvey, The lagrangian multiplier test. *Ann Math Stat* 30, 389-407 (1959).
55. A. Wald, Tests of statistical hypotheses concerning several parameters when the number of observations is large. *Trans Am Math Soc* 54, 426-482 (1943).
56. **Acknowledgements:** We thank the personnel of the Genomics & Proteomics, Bioinformatics, Molecular Pathology, and Microscopy units of our Center for their expert technical help, A. Abad and M. Blázquez for technical lab work, M. Dosil for comments on the manuscript, and F. Miller for providing 67NR, 168FARN and 4TO7 cells.

Funding: XRB work was supported by the Spanish Ministry of Economy and Competitiveness (SAF2009-07172 and RD06/0020/0001), the Castilla-León Autonomous

Government (GR97), the 7th Framework European Union Program (FP7-HEALTH-2007-A-201862), and the Spanish Association Against Cancer. The Spanish XRB's funding is partially supported by the European Regional Development Fund. Work in JMP laboratory is partially supported by grants from both the Spanish Ministry of Economy and Competitiveness (SAF2006-00121 and RD06/0020/0029) and Madrid Autonomous Government (S2006/BIO-0232). CC's and MM-M's salaries were partially supported by both the JAE-Doc program, the Spanish National Research Council and the RD06/0020/0001 grant and the European Regional Development Fund.

Author contributions: CC and MM-M performed the experiments shown in **Figs. 1; 3A-F; 4C-G; 5A,B; 6A,D-F; 7; S2; S3B-D; S4A,F; S6; and Table S4**; CC generated most of the cell lines, generated the raw microarray data, and carried out the experiments shown in **Figs. 2; 3G, 4H,I and 5C-J**. MM-M generated some of the reconstituted cell lines used in the work and carried out the experiments shown in **Fig. S4B-I**. RG-E and JMP carried out the metagenomic experiments shown in **Table S2** and the bioinformatics analysis presented in **Figs. S6 and S7** as well in **Tables S6-S8**. RML performed the experiments shown in **Fig. 4A,B**. OB and FS-M collaborated in the short-term extravasation experiments shown in **Fig. 3**. Artwork was generated by CM, MM-M, RG-E, RML or XRB and finally edited by XRB. XRB directed the work, carried out the metagenomic analyses shown in **Fig. S1** and wrote the manuscript.

Competing interests: None.

Data and materials availability: The microarray data generated in this work have been uploaded to the Gene Expression Omnibus database (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?token=fbefjswiwsqksfu&acc=GSE33348>).

FIGURE LEGENDS

FIGURE 1. Defective tumorigenesis and lung metastasis of Vav-deficient 4T1 cells. **(A-B)** Rac1 (A,B) and RhoA (A) activity levels in indicated cell lines calculated using G-ELISA (A) and GST-Pak-pull-down (B) experiments ($n = 3$). KD, knockdown cell line; for a description and designation of the cell lines utilized in these experiments, check **Table S3**. **(C-F)** Growth kinetics of tumors induced by the indicated cell lines. Statistics were performed at the 35 days time point. The number of independent samples in panel C were: Control ($n = 15$), KD_{2(A)} ($n = 4$), KD_{2(B)} ($n = 4$), KD_{3(A)} ($n = 4$), KD_{3(B)} ($n = 4$), KD_{2/3(A)} ($n = 8$). In panels D,E, $n = 8$. In panel F, $n = 4$. **(G,H)** Number of metastatic nodules detected in mice harboring tumors induced by the indicated cell lines ($n = 5$ animals). **(I)** Images showing the lack of macro- and micrometastasis in the lung of a mouse containing a tumor derived KD_{2/3(A)} cells. As control, we include a lung section from a mouse harboring a tumor derived from the control cell line. Scale bars, 100 (top panel of each group) and 200 μm (bottom panel of each group).

FIGURE 2. Vav family proteins have overlapping, but not identical roles in primary tumorigenesis. **(A,B)** Example (A; scale bar, 50 μm) and quantitation (B; $n = 3$ sections/tumor, 4 animals) of the proliferation present in tumors derived from the indicated cell lines. **(C,D)** Example (C; scale bar, 100 μm) and quantitation (D; $n = 3$ sections/tumor, 3 animals) of the angiogenesis present in tumors developed from the indicated cell lines. **(E,F)** Examples (E; scale bar, 100 μm) and quantification (F; $n = 2$ sections/tumor, 3 animals) of the vessel permeability present in tumors derived from indicated cell lines.

(**G,H**) Example (G; scale bar, 50 μm) and quantification (H; $n = 3$ sections/tumor, 4 animals) of the apoptosis present in tumors derived from the indicated cell lines.

FIGURE 3. Vav proteins work in several metastatic stages. (**A**) Percentage of circulating cancer cells present in mice containing tumors derived from the indicated cell lines ($n = 8$ in experiments with control, $\text{KD}_{2/3}$ (A), and $\text{KD}_{2/3}+\text{V}_{2/3}$ cells and 4 in rest of experiments). a.u., arbitrary units. (**B-E**) Number (B, E) and extension of metastasis (C,D) formed in the lung by the indicated intravenously injected cell lines ($n = 3$ lung sections/mice, 4 mice in each experimental condition). Scale bar, 100 μm . (**F**) Lungs from mice intravenously injected with the indicated chromophore-labeled cell lines were imaged by confocal microscopy to visualize cancer cells (F, green color) and the vascular endothelium (F, red color). Scale bars, 100 (left panel of each group) and 25 μm (right panel of each group). (**G**) Quantification of the number of chromophore-labeled cancer cells present in either whole lung sections (left) or the indicated lung localizations (right) ($n = 6$ confocal sections/animal, 4 mice in each condition).

FIGURE 4. Pro-tumorigenic and pro-metastatic roles of Vav proteins in independent experimental systems. (**A,B**) Primary tumorigenesis (A,B) and lung metastasis (B) induced by *PyMT*-transformed wild type and *Vav2*^{-/-};*Vav3*^{-/-} primary breast epithelial cells upon implantation in the fat pads of recipient wild type female mice ($n = 11$ animals). (**C**) Abundance of *Vav2* (left) and *Vav3* mRNAs (right) in the indicated cells ($n = 3$). (**D**) Rac1 activity in the indicated cell lines detected using G-LISA assays ($n = 3$). (**E,F**) Growth

kinetics (E) and lung metastasis (F) displayed by breast tumors induced by the indicated orthotopically transplanted cell lines ($n = 6$ animals). In F, the x/y ratio indicates the number (x) of animals that scored positive for metastasis in the indicated tissues versus the total number (y) of mice analyzed in these experiments. (G) Example (left) and quantification (right) of metastases generated by the indicated, intravenously injected cell lines ($n = 4$ animals). (H,I) In vitro proliferative (H) and invasion (I) properties of indicated cell lines ($n = 3$).

FIGURE 5. Vav family targets play overlapping, but not identical roles in biological processes linked to primary tumorigenesis. (A) Growth of tumors derived from the indicated, orthotopically transplanted cell lines ($n = 4$ animals). KD, knockdown cell. Clone A and B refer to two knockdown cell clones generated using two independent shRNA for the indicated target (see also **Table S3**). (B) Metastasis found in tissues (top) of mice bearing tumors from the indicated cell line clones. Values are given as in **Fig. 4F**. ND, not determined. (C,D) Example (C; scale bar, 50 μm) and quantitation (D; $n = 3$ sections/tumor, 4 animals) of the proliferation of tumors derived from the indicated cell lines. (E,F) Example (E; scale bar, 100 μm) and quantitation (F; $n = 3$ sections/tumor, 3 animals) of angiogenesis present in tumors derived from the indicated cell lines. (G,H) Example (G; scale bar, 100 μm) and quantitation (H; $n = 2$ sections/tumor, 3 animals) of the vessel permeability of tumors derived from indicated cell lines. (I,J) Example (I; scale bar, 50 μm) and quantitation (J; $n = 3$ sections/tumor, 3 animals) of the apoptosis found in tumors from the indicated cell lines.

FIGURE 6. Implication of Vav-dependent targets in the metastatic steps of breast cancer cells. **(A)** Percentage of circulating cancer cells present in mice harboring tumors derived from the indicated cell lines ($n = 4$ animals). The description of each cell line can be found in **Table S3**. **(B,C)** Visualization **(B)** and quantification of the extravasation rates **(C)** of the indicated chromophore-labeled cell lines after being intravenously injected in recipient mice. Scale bars in **B**, 100 (left panel of each group) and 25 μm (right panel of each group) ($n = 6$ confocal sections/animal, 4 mice in each condition). **(D-F)** Number **(D)** and extension **(E,F)** of the metastatic nodules formed by the indicated, intravenously injected cell lines ($n = 3$ independent lung sections/mice, 4 mice in each condition). Scale bar, 100 μm .

FIGURE 7. The Vav route is involved in the survival of cancer cells inside the lung. **(A,B)** Visualization **(A; green color; scale bar, 100 μm)** and quantification **(B; $n = 6$ confocal sections/animal, 4 mice in each condition)** of the extravasation of the indicated chromophore-labeled cell lines upon intravenous injection in monocrotaline-treated (+MCT) and untreated (-MCT) mice. Endothelial cells are labeled in red in panel **A**. **(C,D)** Visualization **(C)** and quantification **(D; $n = 4$ mice)** of the metastatic nodules induced by the indicated cell lines upon their intravenous injection in MCT treated (+) or untreated (-) mice. **(E)** Area of the metastatic nodules generated by the indicated cells cancer cells in the lung parenchyma ($n = 3$ sections/mice, 4 mice per experimental condition). **(F)** Summary of our results. The implication of a protein in the indicated process is proportional to the

thickness and darkness of horizontal bars. Rac1-dependent and independent routes are depicted in black and gray arrows, respectively. In the case of Ilk, we cannot rule out that the regulation of its abundance will be mediated by the Vav3-mediated stimulation of another Rho/Rac GTPase. However, Vav2 contributes to this response in a GEF-independent manner.

Figure 2
Citterio et al.

Figure 3
Citterio et al.

Figure 4
Citterio et al.

Figure 5
Citterio et al.

Figure 6
Citterio et al.

Figure 7
Citterio et al.

Supplementary Materials for

The Rho/Rac exchange factors Vav2 and Vav3 control a lung-specific pro-metastatic transcriptomal program in breast cancer cells

Carmen Citterio, Mauricio Menacho-Márquez, Ramón García-Escudero, Romain M. Larive, Olga Barreiro, Francisco Sánchez-Madrid, Jesús Paramio, and Xosé R. Bustelo*

*To whom correspondence should be addressed. E-mail: xbustelo@usal.es

This PDF file includes^a:

Figure S1. Rho/Rac GEF transcripts have difference abundance in human tumors.

Figure S2. Abundance of Vav family proteins and Rac1 activity in cell lines used in this study.

Figure S3. Vav proteins control a large transcriptomal program in breast cancer cells.

Figure S4. Mechanistics of the regulation of the breast cancer cell transcriptome by Vav proteins.

Figure S5. Vav proteins control the breast cancer cell transcriptome using pan-specific, synergistic, and Vav family member-specific routes.

Figure S6. Generation of knockdown cell lines for Vav family-dependent targets.

Figure S7. Vav-dependent mRNA signatures have prognostic value in breast cancer.

Table S1. Differential abundance of Rho/Rac GEFs in human breast cancer subtypes.

Table S2. Abundance of human VAV2 and VAV3 mRNAs in the breast cancer luminal A and B subtypes.

Table S3. Cell lines used in this study.

Table S4. Metastasis of indicated 4T1 knockdown cell clones in recipient mice.

^a**Tables S5-S8** in Excel format are located in an independent online file.

FIGURE S1. Rho/Rac GEF transcripts have difference abundance in human tumors.

(A,B) Distribution of transcripts for Rho/Rac GEFs in a variety of human tumors (A, Bittner Multicancer database) and cancer cell lines (B, Adai Cell Line dataset). The relative abundance of a specific transcript is indicated by gradients of red (high abundance) and blue (low abundance). *n*, number of independent clinical samples (A)

and cell lines (B) analyzed of each tumor subtype. Rho/Rac GEFs with statistically significant enrichment in abundance in breast tumors are indicated in red. The figure only shows the 20 top hitters of the Rho/Rac GEFs interrogated in the metagenomic analysis. **(C,D)** Relative abundance of transcripts for *VAV* and *P-REX* family members in human tumors (C, Bittner Multicancer database) and cancer cell lines (D, Adai Cell Line dataset). **(E)** Relative abundance of *VAV* and *P-REX* family transcripts in the Adai Cell Line dataset. The number of cell lines analyzed of each cancer cell type is indicated in brackets (bottom). **(F)** Distribution of Rho/Rac GEFs in human healthy tissues (Roth Normal dataset). The abundance profiles of mRNAs for human Rho/Rac GEFs in the mammary gland are boxed. The profiles of the human *VAV1*, *VAV2*, *VAV3* and *PREX1* mRNAs are indicated by arrowheads.

FIGURE S2. Abundance of Vav family mRNAs and proteins in cell lines used in this study. **(A-D)** qRT-PCR analysis showing the abundance of *Vav2* (A,C) and *Vav3* (B,D) transcripts in the indicated 4T1 cell line derivatives ($n = 3$). KD, knockdown cell. For further information about these cell lines, see **Table S3**. **(E,F)** Detection of ectopic HA-

tagged Vav2 (E) and Myc-tagged Vav3 (F) in total cellular extracts obtained from the indicated cell lines (E,F; top) using Western blot (WB) analysis with the indicated antibodies ($n = 3$).

FIGURE S3. Vav proteins control a large transcriptomal program in breast cancer cells.

(A) Hierarchical cluster diagram of mRNAs whose abundance change upon the knockdown of Vav2 and Vav3 in 4T1 cells. Two independent clones of Vav2;Vav3 knockdown cells (KD_{2/3(A)}, KD_{2/3(B)}) are shown. Information about cell lines can be found in **Table S3**. Columns and rows represent experimental replica (three/cell line) and

genes, respectively. Signal log ratio abundance levels are depicted on a dark blue (highest abundance) to dark red (lowest abundance) scale. **(B-D)** Determination by qRT-PCR of the abundance of the indicated mRNAs in 4T1 cell lines (B,D), in tumors derived from 4T1 cells (C), and 168FARN cells (D). Values (%) are given in relation to the abundance of each transcript in the appropriate control cell line (which was given in each case an arbitrary value of 100%). $n = 3$, except in the graph on the right of panel C, where $n = 1$ (with three independent tumor-bearing mice).

FIGURE S4. Mechanistics of the regulation of the breast cancer cell transcriptome by Vav proteins. **(A)** qRT-PCR showing the abundance of indicated mRNAs in 4T1 cell lines ($n=3$). Values (%) are given in relation to the abundance of each transcript in the

control cell line (which was given a 100% value) ($n = 3$). **(B,C)** Abundance of the *Rac1* mRNA (B) and its encoded protein (C) in the indicated cell lines using qRT-PCR and immunoblot analysis, respectively ($n = 3$). The reduced abundance of the tubulin control detected in the two *Rac1*^{F28L}-expressing cells (C) has been reproducibly observed in three independent experiments. **(D)** *Rac1* activity in the indicated cell lines determined using G-LISA assays ($n = 3$). **(E)** qRT-PCR determination of the abundance of *Ilk*, *Tacstd2*, *Inhba* and *Ptgs2* mRNAs in the indicated cell lines ($n = 3$). Values are given as in panel A. **(F,G)** qRT-PCR determination of the abundance of the indicated mRNAs in our cell lines ($n = 3$). **(H)** Lung metastatic nodules induced by the indicated cell lines upon intravenous injection in recipient mice ($n = 4$ animals). **(I)** Summary of the results obtained in these experiments. Inhibition steps are indicated by blunted lanes.

FIGURE S5. Vav proteins control the breast cancer cell transcriptome using pan-specific, synergistic, and Vav family member-specific routes. **(A,D)** Hierarchical cluster diagram of mRNAs belonging to the pan-specific (A), synergistic (B), Vav2-specific (C) and Vav3-specific (D) subtypes. Signal log ratio abundance levels are depicted on a dark blue (highest increase in abundance) to dark red (highest reduction in abundance) scale. Information about cell lines can be found in **Table S3**. In the pan-specific subclass, the abundance of the mRNAs is regulated in the same manner in Vav2-, Vav3- and Vav2/Vav3-reconstituted *Vav2;Vav3* knockdown cells (A). In the synergistic subclass, the mRNAs are regulated by the two co-expressed Vav proteins but not when each single Vav protein is expressed alone in *Vav2;Vav3* knockdown cells (B). In the Vav2-specific subclass, the mRNA abundance is regulated in the same manner in Vav2- and

Vav2/Vav3-reconstituted *Vav2;Vav3* knockdown cells but not in Vav3-reconstituted *Vav2;Vav3* knockdown cells (C). In the Vav3-specific subclass, the mRNA abundance is regulated similarly in Vav3- and Vav2/Vav3-reconstituted *Vav2;Vav3* knockdown cells but not in Vav2-reconstituted *Vav2;Vav3* knockdown cells (D). **(E)** Percentage (%) of each mRNA subclass in the total Vav2/Vav3-dependent transcriptome.

FIGURE S6. Generation of knockdown cell lines for Vav family-dependent targets. qRT-PCR determination of the abundance of the indicated transcripts in their respective knockdown cell lines. KD, knockdown cell. Clone A and B refer to two knockdown cell clones generated using two independent shRNA for the indicated target (see also **Table S3**). Values are given in reference to the abundance of those mRNAs in parental 4T1 cells (which was given an arbitrary value of 100%) ($n = 3$).

FIGURE S7. Vav-dependent mRNA signatures have prognostic value in breast cancer.

(A) Hierarchical cluster diagram of mRNAs whose abundance changed consistently between the metastatic and non-metastatic cell lines used in this study. (B,C) Overlap

between the Vav-dependent mRNA signature and transcripts deregulated in metastatic primary breast cancer samples present in Loi's (B) and NKI's (C) datasets. H and L, patient groups with high and low similarity with the Vav signature, respectively. Panels on the right (B,C) show the heatmaps of overlapping genes. The size of each patient group (blue, low score; red, high score) is indicated on the left. The number of mRNAs with either increase (red) or reduced (blue) abundance in the Vav signature and metastatic tumors is shown at the bottom of each panel. **(D-F)** Patients present in the NKI (D) and MSCKK (E,F) cohorts were stratified into high (H) and low (L) groups according to their similarity to the V2/V3LMS and either the lung (D,E) or bone (F) metastasis-free survival curves plotted. **(G)** The breast cancer NKI dataset was used to compare the predictive value of the V2/V3 LMS, Minn's and MammaPrint mRNA signatures. The H and L patient groups were assigned as indicated in D-F.

TABLE S1. Differential abundance of Rho/Rac GEF in human breast cancer subtypes^a

Breast cancer type	GEF mRNA	Fold change	P value	Database ^b	Number of samples ^c
ER positive^d	<i>VAV3</i>	2.37	7.86×10^{-11}	Bittner's	336
	<i>P-REX1</i>	1.89	3.17×10^{-10}		
	<i>VAV3</i>	2.74	1.07×10^{-8}	Chin's	118
	<i>DNMBP</i>	1.97	1.31×10^{-8}		
	<i>VAV3</i>	2.64	1.15×10^{-11}	Desmedt's	198
	<i>DNMBP</i>	1.59	3.87×10^{-5}		
	<i>VAV3</i>	2.27	5.42×10^{-16}	Hess'	133
	<i>MCF2L</i>	1.50	1.17×10^{-6}		
	<i>P-REX1</i>	4.20	6.65×10^{-18}	Lu's	129
	<i>VAV3</i>	3.77	2.60×10^{-15}		
	<i>VAV3</i>	2.80	2.00×10^{-17}	Wang's	286
	<i>DNMBP</i>	1.56	1.83×10^{-11}		
PR positive^e	<i>VAV3</i>	2.59	2.21×10^{-8}	Chin's	118
	<i>FARP1</i>	1.64	7.6×10^{-7}		
	<i>VAV3</i>	2.78	2.88×10^{-11}	Minn 2's	121
	<i>MCF2L</i>	1.45	3.2×10^{-4}		

^aThese values were obtained using metagenomic analyses with the OncomineTM/Compendia Bioscience tool. As a comparison, we include the second GEF showing the highest enrichment in a particular breast cancer subtype regardless of whether it fit or not the search parameters. With the exception of *P-REX1* in the Lu's database, no other Rho/Rac GEF-encoding gene gave a positive score under our search criteria.

^bName of the microarray OncomineTM/Compendia Bioscience dataset that scored positive in these analyses.

^cNumber of independent tumor samples used in the microarray experiments of the indicated databases.

^dCompared to estrogen receptor (ER) negative tumors.

^eCompared to progesterone receptor (PR) negative tumors.

TABLE S2. Abundance of *Vav2* and *Vav3* mRNAs in the breast cancer luminal A and B subtypes^a

<i>Vav2</i> mRNA (<i>P</i> value = 0.0049)					
	Normal-like	Basal	Her2⁺	Luminal A	Luminal B
Mean value	-0.04 ^b	-0.20	-0.03	0.13	0.25
Standard error	0.15	0.07	0.16	0.08	0.11
<i>n</i>	44	189	37	160	78
<i>Vav3</i> mRNA (<i>P</i> value = 9.2 x10⁻⁵⁸)					
	Normal-like	Basal	Her2⁺	Luminal A	Luminal B
Mean value	-0.05	-0.79	0.06	0.64	0.59
Standard error	0.12	0.06	0.13	0.06	0.09
<i>n</i>	44	189	37	160	78

^aThe abundance of the two transcripts was analyzed bioinformatically as described in Materials and Methods).

^bMean values (and corresponding standard errors) in intrinsic subtype group were obtained from the z-values of each gene.

TABLE S3. Cell lines used in this study (it continues in the next page)

Cell line	Features
Control	4T1 cells harboring the empty pLKO.1puro lentiviral vector
KD ₂ (A)	Clone of <i>Vav2</i> -deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD ₂ (B)	Clone of <i>Vav2</i> -deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD ₃ (A)	Clone of <i>Vav3</i> -deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD ₃ (B)	Clone of <i>Vav3</i> -deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD _{2/3} (A)	Clone of double <i>Vav2</i> ; <i>Vav3</i> -deficient 4T1 cells generated by transducing lentiviral particles each encoding shRNAs to mouse <i>Vav2</i> and <i>Vav3</i> transcripts
KD _{2/3} (B)	Clone of double <i>Vav2</i> ; <i>Vav3</i> -deficient 4T1 cells generated by transducing lentiviral particles each encoding shRNAs to mouse <i>Vav2</i> and <i>Vav3</i> transcripts
KD _{2/3} +ev	KD _{2/3(A)} cells harboring an empty pLVX-IRES-Hyg lentiviral vector
KD _{2/3} +V ₂	KD _{2/3(A)} cells in which an HA-tagged version of wild type <i>Vav2</i> was re-expressed using lentiviral delivery methods
KD _{2/3} +V ₂ (Y172F)	KD _{2/3(A)} cells in which an HA-tagged, constitutively active version of <i>Vav2</i> was re-expressed using lentiviral delivery methods
KD _{2/3} +V ₂ (R373A)	KD _{2/3(A)} cells in which an HA-tagged, catalytically inactive version of <i>Vav2</i> was re-expressed using lentiviral delivery methods
KD _{2/3} +V ₃	KD _{2/3(A)} cells in which an Myc-tagged version of wild type <i>Vav3</i> was re-expressed using lentiviral delivery methods
KD _{2/3} +V ₂ +V ₃	KD _{2/3(A)} cells in which the wild type versions of <i>Vav2</i> (HA-tagged) and <i>Vav3</i> (Myc-tagged) were re-expressed using lentiviral delivery methods
168FARN+V ₂ +V ₃	168FARN cells in which the wild type versions of <i>Vav2</i> (HA-tagged) and <i>Vav3</i> (Myc-tagged) were overexpressed using lentiviral delivery methods

TABLE S3 (cont). Cell lines used in this study

Cell line	Features
KD _{2/3} +V _{2(R373A)} +V ₃	KD _{2/3(A)} cells in which an HA-tagged, catalytically inactive version of Vav2 was re-expressed together with the Myc-tagged wild type Vav3 using lentiviral delivery methods
Control+Rac1(F28L)	4T1 cells harboring the empty pLKO.1puro lentiviral vector and expressing a fast cycle mutant (F28L) of Rac1
KD _{2/3} +Rac1(F28L)	KD _{2/3(A)} cells in which an a fast cycle mutant (F28L) of Rac1 was ectopically expressed using lentiviral delivery methods
KD-Itgb6 (A)	Clone of integrin β 6-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD-Itgb6 (B)	Clone of integrin β 6-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus. This shRNA is different from the one used for the generation of the KD-Itgb6 (A) clone
KD-Itga8(A)	Clone of integrin α 8-deficient 4T1 cells generated by transducing a shRNA-encoding lentivirus
KD-Itga8(B)	Clone of integrin α 8-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus. This shRNA is different from the one used for the generation of the KD-Itga8 (A) clone
KD-Ilk (A)	Clone of Ilk-deficient 4T1 cells generated by transducing a shRNA-encoding lentivirus
KD-Ilk (B)	Clone of Ilk-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus. This shRNA is different from the one used for the generation of the KD-Ilk (A) clone
KD-Tacstd2 (A)	Clone of Tacstd2-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD-Tacstd2 (B)	Clone of Tacstd2-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus. This shRNA is different from the one used for the generation of the KD-Tacstd2 (A) clone
KD-Inhba (A)	Clone of Inhibin β A-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus
KD-Inhba (B)	Clone of Inhibin β A-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus. This shRNA is different from the one used for the generation of the KD-Inhba (A) clone
KD-Ptgs2 (A)	Clone of cyclo-oxygenase 2-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus

TABLE S3 (cont). Cell lines used in this study

Cell line	Features
KD-Ptgs2 (B)	Clone of cyclo-oxygenase 2-deficient 4T1 cells generated by transducing an shRNA-encoding lentivirus. This shRNA is different from the one used for the generation of the KD- Ptgs2 (A) clone

TABLE S4. Metastasis of indicated 4T1 knockdown cell clones in recipient mice.

Cell line ^a	Lung	Lymph nodes	Spleen	Liver	Intestine
Control	20/20 ^b	20/20	18/20	3/20	3/20
KD_{2/3(A)}	1/20	20/20	20/20	4/20	1/20
KD_{2/3(B)}	0/15	14/15	14/15	5/15	1/15
KD_{2(A)}	2/5	4/5	4/5	0/5	0/5
KD_{2(B)}	3/5	5/5	4/5	1/5	0/5
KD_{3(A)}	3/5	4/5	4/5	0/5	0/5
KD_{3(B)}	2/5	5/5	5/5	1/5	1/5
KD_{2/3}+V₂	2/8	8/8	7/8	0/8	0/8
KD_{2/3}+V_{2(Y172F)}	0/4	4/4	2/4	0/4	0/4
KD_{2/3}+V₃	3/8	8/8	4/8	0/8	0/8
KD_{2/3}+V₂+V₃	8/8	8/8	8/8	1/8	0/8

^aFor information about each cell line, see **Table S3**.

^bThe x/y ratio indicates the number of animals that scored positive for metastasis in the indicated tissues (x) versus the total number of mice analyzed in these experiments (y).