

1 **New insights into meat by-products utilization**

2 Fidel Toldrá^{1,*}, Leticia Mora¹ and Milagro Reig²

3 *¹Instituto de Agroquímica y Tecnología de Alimentos (CSIC), Avenue Agustín Escardino*
4 *7, 46980 Paterna (Valencia), Spain and ²Instituto de Ingeniería de Alimentos para el*
5 *Desarrollo, Universidad Politécnica de Valencia, Camino de Vera s/n, 46022 Valencia,*
6 *Spain*

7
8
9
10 *Author for correspondence (F. Toldrá), e-mail: ftoldra@iata.csic.es

11
12
13
14
15
16
17
18
19
20 Running title: New insights into meat by-products utilization
21

22 **Abstract**

23 Meat industry generates large volumes of by-products like blood, bones, meat
24 trimmings, skin, fatty tissues, horns, hoofs, feet, skull and viscera among others that are
25 costly to be treated and disposed ecologically. These costs can be balanced through
26 innovation to generate added value products that increase its profitability. Rendering
27 results in feed ingredients for livestock, poultry and aquaculture as well as for pet foods.
28 Energy valorisation can be obtained through the thermochemical processing of meat and
29 bone meal or the use of waste animal fats for the production of biodiesel. More recently,
30 new applications have been reported like the production of polyhydroxyalkanoates as
31 alternative to plastics produced from petroleum. Other interesting valorisation strategies
32 are based on the hydrolysis of by-products to obtain added value products like bioactive
33 peptides with relevant physiological effects as antihypertensive, antioxidant,
34 antidiabetic, antimicrobial, etc. with promising applications in the food, pharmaceutical
35 and cosmetics industry. This paper reports and discusses the latest developments and
36 trends in the use and valorisation of meat industry by-products.

37

38 **Keywords:** animal by-products, meat by-products, offal, skin, bones, trimmings,
39 bioactive peptides, hydrolysed proteins, biodiesel

40 **1. Introduction**

41 Meat industry generates large volumes of by-products like blood, bones, meat
42 trimmings, skin, fatty tissues, horns, hoofs, feet, skull and viscera among others that are
43 costly to be treated and disposed ecologically (Ryder, Ha, El-Din Bekhit and Carne,
44 2015). These costs can be balanced through innovation to generate added value products
45 that increase its profitability. On the other hand, unappropriated treatment or handling
46 of such by-products raised relevant crisis in the past such as the spread of the
47 spongiform encephalopathies. The European Commission published the Regulation
48 (EC) 1069/2009 laying down health rules as regards animal by-products and derived
49 products not intended for human consumption and repealing Regulation (EC)
50 1774/2002. Later, the European Commission published the Regulation (EC) 142/2011
51 that was implementing the Regulation 1069/2009. Rules were also provided by the Food
52 and Drug Administration (FDA, 2004) to prevent the establishment and spread of
53 bovine spongiform encephalopathy (BSE) in the United States, including a prohibition
54 on the use of high-risk, cattle-derived materials that can carry the BSE agent which are
55 defined as specified risk material. This means that adequate disposal of by-products
56 may increase the cost to processors and makes necessary to produce new substances or
57 products capable to cover the disposal costs (Toldrá, Mora, Aristoy and Reig, 2012).
58 It must be taken into account that certain meat by-products can be considered as foods
59 of interest depending on the country and local traditions while in other places they can
60 be considered as inedible foods (Ockerman & Basu, 2004a). In fact, some by-products
61 with high nutritional value like blood, liver, lung, heart, kidney, brains, spleen and tripe
62 constitute part of the diet and culinary recipes in many countries worldwide (Nollet &
63 Toldrá, 2011). Of course, the nutritional composition depends on each particular type of
64 by-product and the animal species from which they are obtained (Honikel, 2011). Other
65 by-products like lard may be used for cooking.

66 Meat by-products may constitute a valuable resource if handled properly to produce
67 added value substances or products (Zhang, Xiao, Samaraweera, Lee & Ahn, 2010,
68 Toldrá and Reig, 2011). Efficient use of by-products may arise up to 11.4% and 7.5% of
69 the gross income of beef and pork (Jayathilakan, Sultana and Radhakrishna, 2012).
70 There is a large variety of meat by-products but, in general, most of them contain good
71 amounts of nutrients like essential amino acids, minerals and vitamins (Aristoy &
72 Toldrá, 2011, Honikel, 2011, Kim, 2011), constituting good valorization opportunity for
73 the meat industry (Valta, Damala, Orli, Papadaskalopoulou, Moustakas, Malamis and

74 Loizidou, 2015). There are numerous applications based on new or improved
75 technologies for processing meat by-products like edible food ingredients for the food,
76 feed and pet food industry (see Figure1). Meat by-products can be considered as raw
77 materials for the generation of biomolecules of interest like protein hydrolysates with
78 relevant bioactivities or enzymes (Lasekan, Abu Bakar and Hashim, 2013), extracts
79 with functional properties (Chernukha, Fedulova and Kotenkova, 2015) or bioactive
80 peptides (Mora, Reig and Toldrá, 2014; Martínez-Alvarez, Chamorro and Brenes,
81 2015).

82 Other applications are addressed towards inedible products like fertilizers, substances of
83 interest for the chemical or pharmaceutical industry or energy generation (see Figure 1).
84 Energy generation is an active area mainly focused on the biodiesel production from
85 waste animal fats (Banckovic-Ilic, Stojkovic, Stamenkovic and Veljkovic, 2014;
86 Adewale, Dumont and Ngadi, 2016) or even a second generation of bioderived diesel
87 fuel, also known as bio gas oil (Balandincz and Hancsók, 2015).
88 This manuscript reports and discusses the latest developments and trends in the use and
89 valorisation of meat industry by-products.

90

91 **2. Food applications**

92

93 *Applications as functional ingredients*

94 Bioactive peptides are sequences usually between 2 and 20 amino acids that exert a
95 biological function in one or several of the physiological systems in human being. In
96 this sense, hypocholesterolemic, antioxidant and antithrombotic peptides have been
97 described to modulate the cardiovascular system whereas mineral binding and
98 immunomodulatory peptides act in gastrointestinal and immune systems, respectively.
99 Some groups of peptides are able to participate in multiple system reactions. Thus,
100 opioid agonist and antagonists can act on nervous, gastrointestinal, and immune
101 systems, whereas antimicrobial peptides can modulate gastrointestinal and immune
102 systems (Lafarga and Hayes, 2014).

103 Bioactive peptides need to be liberated from their origin protein in order to exert the
104 biological function as they are inactive within the parent protein (Vercauteren, Van
105 Camp, and Smagghe, 2005). Some bioactive peptides are released during food
106 processing either in fermentation or curing stages, whereas others are generated during
107 gastrointestinal digestion. The main problem of naturally generated peptides is the

108 difficulty in controlling the hydrolysis conditions because many endogenous enzymes
109 are acting at the same time and a wide profile of peptides showing different sizes and
110 characteristics is generated (Mora, Gallego, Escudero, Reig, Aristoy & Toldrá, 2015).
111 For this reason, the digestion of protein extracts under controlled hydrolysis conditions
112 using known enzymes such as alcalase, pepsin, thermolysine, trypsin, etc., allows the
113 control of the generated bioactive peptides as well as the obtention of more
114 homogeneous batches.

115 The use of by-products as a source of bioactive peptides has been extensively studied
116 during the last years. In this sense, blood and collagen, very important by-products from
117 slaughterhouses and meat industry, have been the most assayed (Ryder, El-Din Bekhit,
118 McConnell and Carne, 2016).

119 Blood is a rich source of proteins where hemoglobin, an iron-containing protein, is the
120 most abundant complex (Ofori and Hsieh, 2014). It is obtained all around the world and
121 even though is used as food ingredient in Europe, Asia, and Africa, its production is
122 more copious than needed. Its value as a source of bioactive peptides has been studied
123 in both the cellular fraction (hemoglobin cells) and the plasma fraction, and their
124 hydrolysates have been described to exert antimicrobial, antioxidant, ACE-inhibitory,
125 and opioid activities (Chang, Wu and Chiang, 2007). However, antimicrobial peptides
126 derived from hemoglobin hydrolysates have been the most studied (Nedjar-Arroume et
127 al., 2004; Marya, Kouach, Briand and Guillochon, 2005; Briand and Guillochon, 2006,
128 2008). Bovine hemoglobin hydrolysate obtained with pepsin in the presence of 30%
129 ethanol resulted in the novel identification of 67-106, 73-105, 99-105, and 100-105
130 fragments of the α -chain of bovine hemoglobin. These peptides exert an antibacterial
131 activity against *Kocuria luteus* A270, *Listeria innocua*, *Escherichia coli*, and
132 *Staphylococcus aureus* with a MIC between 187.1 and 35.2 μ M as well as an ACE
133 inhibitory activity with IC₅₀ values from 42.55 to 1,095 μ M (Adje et al 2011a). On the
134 other hand, Hu et al. (2011) identified the peptide VNFKLLSHSLLVTLASHL from α -
135 chain bovine hemoglobin showing antimicrobial activity against *E. coli*, *S. aureus*, and
136 *Candida albicans* when assessed. The minimal peptide sequences necessary to show
137 antimicrobial activity after a pepsin enzyme digestion of α - and β -chain hemoglobin
138 proteins have been described to be KYR and RYH, respectively, and were studied
139 against *E. coli*, *Salmonella enteritidis*, *L. innocua*, *Micrococcus luteus*, and *S. aureus*
140 (Catiau et al 2011a, 2011b). The sequences obtained from blood protein hydrolysates in
141 recent years are shown as Table 1.

142 The generation of bioactive peptides depends to a high extent on the enzymes and
143 substrate used in the hydrolysis. In fact, the hydrolysis degree determines the extent of
144 hydrolysis whereas the digestion conditions (temperature, pH, and time) are very
145 important to obtain the bioactive peptides. On the other hand, peptide size and amino
146 acid sequences are crucial for the bioactive potential of the peptides (Yu, Hu,
147 Miyaguchi, Bai, Du and Lin, 2006). As an example, antimicrobial peptides have been
148 shown to be mostly hydrophobic as higher hydrophobicity is necessary in the affinity
149 with the outer membrane of microbials. In fact, there is an interaction with negatively
150 charged membrane phospholipids by tyrosine residues together with arginine and lysine
151 which can act as peptide anchors in membranes (Lopes, Fedorov and Castanho, 2005).
152 ACE-inhibitory peptides, also well-studied in hemoglobin hydrolysates, have been
153 described to contain proline, lysine or aromatic residues. In fact, ACE binding is
154 influenced by a proline residue at any of the three last positions of the C-terminal site.
155 Antimicrobial and ACE-inhibitory peptides derived from bovine and porcine
156 hemoglobin and plasma have been described in Table 1. Some opioid peptides with
157 potential to have an effect on nervous and gastrointestinal systems have also been
158 described from animal blood sources (Zhao et al., 1997, 1994; Kapel et al., 2003;
159 Froidevaux et al., 2008). However, there is a lack of studies about the antioxidant
160 capability of hemoglobin-derived peptides.

161 Collagen is the most abundant protein in many by-products obtained from meat
162 industry. In fact, it is the main constituent in skin, hide, bones, and cartilages. The
163 nutritional value of collagen is very low because it lacks essential amino acids but, on
164 the other hand, collagen is very useful as a source of bioactive peptides (Morimatsu,
165 2008, Dierckx and Smagghe, 2011). Despite many recent studies have been focused on
166 the bioactive properties of collagen hydrolysates, most of the published studies have
167 been focused on fisheries by-products. In collagen hydrolysates, ACE-inhibitory and
168 antioxidant activities resulted to be the most relevant when enzymes such as alcalase,
169 trypsin, chymotrypsin, neutrase, flavorenzyme, pepsin, bromelain and papain were used
170 (Saiga et al., 2008; Gómez-Guillén et al. 2011; Di Bernardini, Mullen, Bolton, Kerry,
171 O'Neill & Hayes, 2012). In this sense, Herregods et al (2011) reported that thermolysin
172 hydrolysate showed the highest *in vitro* ACE inhibitory activity as well as an important
173 *in vivo* antihypertensive effect in spontaneously hypertensive rats. Recently, a MALDI-
174 ToF mass spectrometry methodology has been used to determine the animal origin from
175 collagen trypsinated peptides in food preparations and galenic formulations. The

176 differentiation between pork and bovine gelatin was performed through the mass spectra
177 (Flaudrops et al., 2015).

178

179 *Technological applications*

180 The cellular fraction that contains red blood cells, white blood cells and platelets, can be
181 used as colour enhancer for sausages even though it has limited applications in foods
182 due to the dark colour of hemoglobin, sensory adverse effects or even hygiene (Ofori &
183 Hsieh, 2011). Better flavor can be obtained if hemoglobin is removed and used to
184 replace fat in meat products (Viana, Silva, Delvivo, Bizzotto & Silvestre, 2005).

185 A heme iron polypeptide that helps for a better iron absorption can be generated through
186 enzymatic hydrolysis of hemoglobin (Nissenson, Berns, Sakiewickz, Ghaddar, Moore &
187 Schleicher, 2003).

188 Interesting technological properties for food processing can be obtained from blood
189 proteins (Hsieh and Ofori, 2011). So, immunoglobulins, fibrinogen and serum albumin
190 contribute to gelation and emulsification (Cofrades, Guerra, Carballo, Fernández-Martín
191 & Jiménez-Colmenero, 2000) while other plasma proteins contribute to proteins cross-
192 linking (Kang & Lanier, 1999), proteins enrichment (Yousif, Cranston and Deeth, 2003)
193 or foaming (Del, Rendueles and Díaz, 2008). High antioxidant activity has been
194 reported in red blood cell fractions from sheep, pig, cattle and red deer (Bah, Bekhit,
195 Carne and McConnell, 2016). Also, antimicrobial activity against *E. coli*, *S. aureus* and
196 *P. aeruginosa* was reported in sheep white blood cells (Bah et al., 2016).

197 The enzyme thrombin and fibrinogen are used for binding of meat pieces and, for
198 instance, reconstitute meat steaks or generate meat emulsions increasing the hardness
199 and springiness. Fibrinogen is converted by thrombin into insoluble fibrin that form
200 fibers by aggregation. The final results is a three-dimensional network fibrin clot
201 (Lennon, McDonald, Moon, Ward & Kenny, 2010) with more or less strength
202 depending on the size and moisture of the pieces and the conditions of pH and
203 temperature used (Chen & Lin, 2002). Thrombin and fibrinogen are registered under the
204 trade mark Fibrimex® and commercialised as a binder for meat processing to
205 manufacture restructured meat products.

206 Gelatin is obtained from collagen through hydrolysis and is widely used in the food
207 industry because of its good gel-forming ability, but also as clarifying agent, stabiliser
208 or protective coating material (Djagny, Wang & Xu, 2001; Gómez-Guillen et al., 2011).

209 Animal rendering yields proteins that can reduce the surface tension and produce foams
210 (Bressler, 2009). Protein hydrolysates are also used as flavor ingredients; their sensory
211 properties depending on the balance and content of small peptides and free amino acids
212 (Maehashi, Matsuzaki, Yamamoto & Udaka, 1999).

213

214 **3. Feed and pet food applications**

215 Raw or rendered animal by-products have been traditionally used as ingredients in feeds
216 and pet foods. About 25 million tonnes per year of animal by-products derived from
217 meat industries in the US and 15 million tonnes in the European Union are processed by
218 rendering to produce high quality fats and proteins (Hamilton, 2016). In fact, animal by-
219 products constitute a good source of nutrients like essential amino acids, fatty acids,
220 minerals and trace elements, B vitamins and some fat-soluble vitamins (Nollet and
221 Toldrá, 2011; Honikel, 2011). Examples are protein or blood meals (Alexis & Robert,
222 2004; Pérez-Gálvez, Almécija, Espejo, Guadix and Guadix, 2011), amino acids
223 solutions obtained from blood (Giu & Giu, 2010) or meat and bone meal ashes obtained
224 after co-incineration (Goutand, Cyr, Deydier, Guilet and Clastres, 2008). Meat and bone
225 meal is also a good source of essential amino acids and group B vitamins for animal
226 feeds (Jayathilakan et al., 2012). Protein hydrolysates have been reported to be
227 successful in aquaculture (Gilbert, Wong and Webb, 2012). Excessive bitterness in
228 protein hydrolysates can be reduced by cleaving hydrophobic amino acids from peptides
229 and make the palatability more appealing in pet foods (Nchienzia, Morawicki and
230 Gadang, 2010). Rendered meat by-products are also used as ingredients for dogs pet
231 foods (Murray, Patil, Fahey, Merchen and Hughes, 1997).

232 Meat by-products protein hydrolysates represent an interesting alternative to soybean
233 meal because the absence of antinutritional factors or allergenic proteins and the
234 presence of large amounts of all essential amino acids (Martínez-Alvarez, Chamorro
235 and Brenes, 2015). Other by-products like hair, nail, feather and outer layer of skin
236 containing keratin, can be profitable after hydrolysis with the enzyme keratinase
237 (Deivasigamani & Alagappan, 2008; Lasekan, Abu Bakar and Hashim, 2015). This
238 enzyme is predominantly a serine peptidase with a broad range of neutral-alkaline pH
239 for activity, pH ranging 6.0-13.0, and able to hydrolyse keratin under reducing
240 conditions (Brandelli, Sala and Kalil, 2015).

241

242 **4. Energy generation applications**

243 In recent years, biodiesel has been produced and is now replacing progressively the
244 diesel fuel due to its advantages like being biodegradable, non-toxic and with a
245 favorable combustion emission profile that leads to reductions in carbon dioxide, carbon
246 monoxide, particulate matter and unburned hydrocarbons (Gerpen, 2005; Moreira, Dias,
247 Almeida & Alvim-Ferraz, 2010). Further, the use of biodiesel does not imply significant
248 modifications in engines.

249 Low cost animal fat by-products are used as raw materials that are transesterified with a
250 low molecular weight alcohol to yield a mixture of fatty acid methyl esters and glycerol
251 as a side product (Bhatti, Hanif, Qasim & Rheman, 2008; Moreira et al., 2010). Hydro-
252 oxygenation and hydroisomerization in tubular reactors has been proposed to increase
253 biodiesel profitability (Herskowitz, 2008), also supercritical transesterification
254 (Marulanda, Anitescu & Tavlarides, 2010). Other recent studies focus on the improved
255 production of biodiesel by using ultrasounds assisted transesterification of the animal
256 fats (Adewale et al., 2016). Animal fats have some limitations due to its protein and
257 phosphoacylglycerols content that makes a degumming process necessary, the presence
258 of water that requires of vacuum drying and the high content of saturated fatty acids that
259 need to be reduced through winterization process or additives addition (Banckovic-Ilic
260 et al., 2014).

261 The developments have continued and nowadays a new 2nd generation, so-called bio gas
262 oil is facing prompt application. Triacylglycerols are converted into a mixture of iso and
263 normal paraffin via heterogeneous catalytic hydrogenation. Raw materials like brown
264 greases have been also assayed with positive results (Baladincz and Hancsók, 2015).

265

266 **5. Medical and pharmaceutical applications**

267 Pork skin can be used as dressing for burns or skin ulcers in humans (Jayathilakan et al.,
268 2012). Glands and organs constitute edible meat by-products with good nutritive value
269 that are consumed in different regions of the world (Nollet and Toldrá, 2011) and, in
270 fact, some of them are consumed for medicinal purposes in countries like China, Japan
271 and India, or used as a source of particular pharmaceutical substances. This is the case
272 of bile from the gall bladder, melatonin from the pineal gland, heparin from the liver,
273 progesterone and oestrogen from ovaries, insulin from pancreas, etc. (Jayathilakan et al.,
274 2012). Protein hydrolysates, especially those from collagen can generate peptides to be
275 used in treatments against osteoarthritis by accumulation in the joint cartilage (Bello
276 and Oeser, 2006). Hydrolysed collagen exerts a positive effect on bones and joints. In

277 fact, these hydrolysates with added hyaluronic acid are being commercialised for better
278 performance of joints and pain relief in humans.

279 Low molecular weight ultrafiltrates (<30kDa) obtained from pig aorta extracts were
280 assayed with laboratory guinea pigs and such extracts were reported to exert substantial
281 reductions in atherogenic lipoproteins, atherogenic index and total and residual
282 cholesterol (Chernukha, Fedulova and Kotenkova, 2015).

283

284 **6. Fertilizer applications**

285 Large amounts of meat and bone meal are generated in all countries and an interesting
286 approach is the thermochemical processing including pyrolysis, combustion and
287 gasification. The most analysed are co-combustion with coal and pyrolysis. The
288 resulting ashes demonstrate a high content of phosphorus which makes them suitable as
289 fertilisers and the gas emissions are within the international regulations and contains
290 combustibles to be used for energy production (Coutand, Cyr, Deydier, Guilet and
291 Clastres, 2008; Cascarosa, Gea and Arauzo, 2012). The incineration of animal by-
292 products results in good mineral fertilisers. In addition, the use of heat recovery allows
293 for efficient energy recovery (Nujak, 2015).

294

295 **7. Chemical applications**

296 Rendered fats have many applications in cosmetic industry for products like hand and
297 body lotions, creams and bath products. Fatty acids are used in the chemical industry for
298 rubber and plastic polymerization, softeners, lubricants and plasticizers (Ockerman and
299 Basu, 2006). Collagen, gelatin and glycerin are also used in chemical industry as
300 ingredients for surfactants, paints, varnishes, adhesives, antifreeze, cleaners and polishes
301 (Pearl, 2004). New applications using rendered fats have been reported like the
302 production of polyhydroxyalkanoates with a recombinant strain of *Ralstonia eutropha*
303 (Riedel, Jahns, Koenig, Bock, Brigham, Bader and Stahl, 2015). Such polymer has the
304 advantage being biodegradable and constitutes an attractive alternative to plastics
305 produced from petroleum.

306 There are many applications for hides that traditionally have been used for leather-based
307 articles like clothes, shoes, belts, handbags and purses (Ockerman & Basu, 2004b).

308

309 **8. Conclusions**

310 There are many applications of meat by-products like feed ingredients for livestock,
311 poultry and aquaculture as well as for pet foods, energy valorisation through biodiesel
312 production, new substances as alternative to plastics and protein hydrolysates to be used
313 for technological purposes or as a source of bioactive peptides with relevant
314 physiological effects. Research efforts are going ahead to produce new substances with
315 new applications or improving those existing processes. So, the innovation is
316 continuously addressed towards adding value and finding new applications to meat by-
317 products.

318

319 **Acknowledgements**

320 Grant from AGL2014-57367-R from MINECO (Spain) and FEDER funds and Grant
321 GV/2015/138 from Generalitat Valenciana (Spain) and JAEDOC-CSIC postdoctoral
322 contract of L.M. co-funded by the European Social Found are acknowledged. Work
323 prepared within the Unidad Asociada IAD (UPV)-IATA (CSIC) framework.

324

325 **References**

326 Adewale, P., Dumont, M-J., Ngadi, M. (2016). Enzyme-catalyzed synthesis and kinetics
327 of ultrasonic assisted methanolysis of waste lard for biodiesel production. *Chemical*
328 *Engineering Journal*, 284, 158–165.

329 Adje, E. Y., Balti, R., Kouach, M., Guillochon, D., & Nedjar-Arroume, N. (2011a). α
330 67-106 of bovine hemoglobin: a new family of antimicrobial and angiotensin I-
331 converting enzyme inhibitory peptides. *European Food Research and Technology*, 232,
332 637–646.

333 Adje, E. Y., Balti, R., Kouach, M., Dhulster, P., Guillochon, D. (2011b) Obtaining
334 antimicrobial peptides by controlled peptic hydrolysis of bovine hemoglobin.
335 *International Journal of Biological Macromolecules* 49, 143–153

336 Alexis, A. & Robert, J. (2004) Animal byproduct conversion system and method. US
337 patent 7,000,333.

338 Aristoy, M.C. & Toldrá, F. (2011) Essential amino acids. In: *Handbook of Analysis of*
339 *Edible Animal By-Products* (L.M.L. Nollet and F. Toldrá, Eds.), pp 123-135, CRC Press,
340 Boca Raton FL, USA.

341 Bah, C.S.F., El-Din, A., Bekhit, A., Carne, A., McConnell, M.A. (2016) Composition
342 and biological activities of slaughterhouse blood from red deer, sheep, pig and cattle.
343 *Journal of the Science of Food & Agriculture*, 96, 79–89.

344 Bajuk, J.W. (2015) New insights into waste management-Meat industry. *Renewable*
345 *Energy*, 83, 1174-1186.

346 Baladincz, P., Hancsó, J. (2015) Fuel from waste animal fats. *Chemical Engineering*
347 *Journal*, 282, 152–160.

348 Banković-Ilić, I.B., Stojković, I.J., Stamenković, O.S., Veljković, V.B., Hung, Y-T.
349 (2014) Waste animal fats as feed stocks for biodiesel production. *Renewable and*
350 *Sustainable Energy Reviews*, 32, 238–254.

351 Bello, A. E., & Oesser, S. (2006) Collagen hydrolysate for the treatment of
352 osteoarthritis and other joint disorders: A review of the literature. *Current Medical*
353 *Research and Opinion*, 22, 2221–2232.

354 Bhatti, H.N., Hanif, M.A., Qasim, M. & Rheman, A.U. (2008) Biodiesel production
355 from waste tallow. *Fuel*, 87, 2961-2966.

356 Brandelli, A., Sala, L. & Kalil, S.J. (2015) Microbial enzymes for bioconversion of
357 poultry waste into added-value products. *Food Research International*, 73, 3–12.

358 Bressler, D. (2009) Protein based foaming agents and methods of making thereof.
359 World Patent WO2009053852.

360 Bujak, J.W. (2015) New insights into waste management - Meat industry. *Renewable*
361 *Energy*, 83, 1174-1186

362 Cascarosa, E., Gea, G., Arauzo, J. (2012) Thermochemical processing of meat and bone
363 meal: A review. *Renewable and Sustainable Energy Reviews*, 16, 942– 957.

364 Catiau, L., Traisnel, J., Delval-Dubois, V., Chihib, N.E., Guillochon, D. & Nedjar-
365 Arroume, N. (2011) Minimal antimicrobial peptidic sequence from hemoglobin alpha-
366 chain: KYR. *Peptides*, 32, 633-638.

367 Catiau, L., Traisnel, J., Chihib, N.E., Le Flem, G., Blanpain, A., Melnyk, O.,
368 Guillochon, D. & Nedjar-Arroume, N. (2011) RYH: A minimal peptidic sequence
369 obtained from beta-chain hemoglobin exhibiting an antimicrobial activity. *Peptides*, 32,
370 1463-1468.

371 Chang, C-Y., Wu, K-C. & Chiang, S.H. (2007) Antioxidant properties and protein
372 compositions of porcine haemoglobin hydrolysates. *Food Chemistry*, 100, 1537-1543.

373 Chen, M.J. & Lin, C.W. (2002) Factors affecting the water-holding capacity of
374 fibrinogen/plasma protein gels optimized by response surface methodology. *Journal of*
375 *Food Science*, 67, 2579-2582.

376 Chernukha, I.M., Fedulovaa, L.V., Kotenkovaa, E.A. (2015) Meat by-product is a
377 source of tissue-specific bioactive proteins and peptides against cardio-vascular diseases
378 *Procedia Food Science*, 5, 50 – 53.

379 Cofrades, S., Guerra, N.I.A., Carballo, J., Fernández-Martin, F. & Jiménez-Colmenero,
380 F. (2000) Plasma protein and soy fiber content effect on bologna sausage properties
381 as influenced by fat levels. *Journal of Food Science*, 65, 281-287.

382 Coutand, M., Cyr, M., Deydier, D., Guilet, R., Clastres, P. (2008) Characteristics of
383 industrial and laboratory meat and bone meal ashes and their potential applications.
384 *Journal of Hazardous Materials*, 150, 522–532.

385 Daoud, R., Dubois, V., Bors-Dodita, L., Nedjar-Arroume, N., Krier, F., Chihib, N. E.,
386 Marya, P., Kouach, M., Briand, G. & Guillochon, D. (2005) New antibacterial peptide
387 derived from bovine hemoglobin. *Peptides*, 26, 713-719.

388 Deivasigamani, B. & Alagappan, K.M. (2008) Industrial application of keratinase and
389 soluble proteins from feather keratine. *Journal of Environmental Biology*, 29, 933-936.

390 Del, P.H., Rendueles, M., Díaz, M. (2008) Effect of processing on functional properties
391 of animal blood plasma. *Meat Science*, 78, 522–528

392 Deng H., Zheng J., Zhang F., Wang Y., Kan J. 2014 Isolation of angiotensin I-
393 converting enzyme inhibitor from pepsin hydrolysate of porcine hemoglobin. *European*
394 *Food Research and Technology* 239, 933-940.

395 Di Bernardini, R., Mullen, A.M., Bolton, D., Kerry, J., O'Neill, E. & Hayes M. (2012).
396 Assessment of the angiotensin-I-converting enzyme (ACE-I) inhibitory and antioxidant
397 activities of hydrolysates of bovine brisket sarcoplasmic proteins produced by papain
398 and characterisation of associated bioactive peptidic fractions. *Meat Science*, 90, 1, 226-
399 235.

400 Djagny, K.B., Wang, Z. & Xu, S. (2001) Gelatin: A valuable protein for food and
401 pharmaceutical industries, Review. *Critical Reviews in Food Science and Nutrition* 41,
402 481-492.

403 Duarte, R. T., Carvalho Simoes, M. C. & Sgarbieri, V. C. (1999). Bovine blood
404 components: fractionation, composition, and nutritive value. *Journal of Agricultural &*
405 *Food Chemistry*, 47, 231-236.

406 Flaudrops, C., Armstrong, N., Raoult, D., and Chabrière, E.(2015). Determination of the
407 animal origin of meat and gelatin by MALDI-ToF-MS. *Journal of Food Composition*
408 *and Analysis*, 41, 104-112.

409 Fogac, A.C., Da Silva, P.I., Teresa Jr, M., Miranda, M., Bianchi, A.G. & Miranda, A.
410 (1999) Antimicrobial activity of a bovine hemoglobin fragment in the tick *Boophilus*
411 *microplus*. *Journal of Biological Chemistry*, 274, 25330–25334.

412 Froidevaux, R., Vanhoute, M., Lecouturier, D., Dhulster, P. & Guillochon, D. (2008)
413 Continuous preparation of two opioid peptides and recycling of organic solvent using
414 liquid/liquid extraction coupled with aluminium oxide column during haemoglobin
415 hydrolysis by immobilized pepsin. *Process Biochemistry*, 43, 431-437.

416 Gerpen, J.V. (2005) Biodiesel processing and production. *Food Processing Technology*,
417 86, 1097-1107.

418 Gilbert, E. R., Wong, E. A., & Webb, K. E. (2008) Board-invited review: Peptide
419 absorption and utilization: Implications for animal nutrition and health. *Journal of*
420 *Animal Science*, 86, 2135–2155.

421 Giu, H.M. & Giu, P.D. (2010) Blood waste treatment system for slaughtered animals,
422 and method for producing high quality amino acid solution using blood waste. World
423 patent WO2010KR00586 20100201.

424 Glamsta E.-L., Meyerson B., Silberring J., Terenius L. & Nyberg F. (1992) Isolation of a
425 hemoglobin-derived opioid peptide from cerebrospinal fluid of patients with
426 cerebrovascular bleedings. *Biochemical Biophysical Research Communications*, 184,
427 1060-1066.

428 Gómez-Guillén, M.C., Giménez, B., López-Caballero, M.E. & Montero, M.P. (2011)
429 Functional and bioactive properties of collagen and gelatin from alternative
430 sources: A review. *Food Hydrocolloids*, 25, 1813-1827.

431 Hamilton, C.R. (2016) Real and perceive issues involving animal proteins. In: [Protein](#)
432 [Sources for the Animal Feed Industry](#). FAO report. www.fao.org. Accessed 12 april
433 2016.

434 Hazato, T. & Kase, R. (1986) Isolation of angiotensin-converting enzyme inhibitor from
435 porcine plasma. *Biochemical Biophysical Research Communications*, 139, 52–55.

436 Herregods, G., Van Camp, J., Morel, N., Ghesquière, B., Gevaert, K., Vercruyse, L.,
437 Dierckx, E. & Smagghe, G. (2011) Angiotensin I-converting enzyme inhibitory activity
438 of gelatin hydrolysates and identification of bioactive peptides. *Journal of Agricultural*
439 *& Food Chemistry*, 59, 552-558.

440 Herskowitz, M. (2008) Reaction system for production of diesel fuel from vegetable
441 and animal oils. World Patent WO2008035155.

442 Honikel, K.O. (2011) Composition and calories. In: *Handbook of Analysis of Edible*
443 *Animal By-Products* (L.M.L. Nollet and F. Toldrá, Eds.), pp 105-121, CRC Press, Boca
444 Raton FL, USA.

445 Hsieh, Y.H. & Ofori, J.A. (2011) Food-grade proteins from animal by-products. Their
446 usage and detection methods. In: *Handbook of Analysis of Edible Animal By-Products*
447 (L.M.L. Nollet and F. Toldrá, Eds.), pp 3-11, CRC Press, Boca Raton FL, USA.

448 Hu, J., Xu, M., Hang, B., Wang, L., Wang, Q., Chen, J., et al. (2011) Isolation and
449 characterization of an antimicrobial peptide from bovine hemoglobin α -subunit. *World*
450 *Journal of Microbiology and Biotechnology*, 27, 767–771.

451 Hyun, C.K. & Shin, H.K. (1998) Utilization of bovine blood plasma obtained from a
452 slaughterhouse for economic production of probiotics. *Journal of Fermentation*
453 *Bioengineering*, 86,34–7.

454 Jayathilakan, K., Sultana, K., Radhakrishna, K., Bawa, A.S. (2012) Utilization of
455 byproducts and waste materials from meat, poultry and fish processing industries: a
456 review. *Journal of Food Science & Technology*, 49, 278–293.

457 Kang, I.S. & Lanier, T.C. (1999) Bovine plasma proteins functions in surimi gelation
458 compared with cysteine protease inhibitors. *Journal of Food Science*, 64, 842-846.

459 Kapel, R., Froidevaux, R., Nedjar-Arroume, N., Fertin-Bazus, A., Dhulster, P. &
460 Guillochon, D. (2003) Continuous production of a peptidic fraction containing the
461 intermediate opioid peptide LVV-haemorphin-7 (LVVh-7) by peptic hydrolysis of
462 bovine haemoglobin in a continuous membrane reactor. *Biotechnology and Applied*
463 *Biochemistry*, 37, 317-324.

464 Kim, Y-N. (2011) Vitamins. In: *Handbook of Analysis of Edible Animal By-Products*
465 (L.M.L. Nollet y F. Toldrá, Eds.), pp 161-182, CRC Press, Boca Raton FL, USA.

466 Kohmura, M.; Nio, N.; Kubo, K.; Minoshima, Y.; Munekata, E.; Ariyoshi, Y. (1989)
467 *Agricultural Biological Chemistry*, 53, 2107.

468 Lafarga, T. & Hayes, M. (2014). Bioactive peptides from meat muscle and by-products:
469 generation, functionality and application as functional ingredients. *Meat Science*, 98, 2,
470 227-239.

471 Lasekan, A., Abu Bakar, F., Hashim, D. (2013) Potential of chicken by-products as
472 sources of useful biological resources. *Waste Management*, 33, 552–565.

473 Lee, S.H. & Song, K.B. (2009a) Isolation of a calcium-binding peptide from
474 hydrolysates of porcine blood plasma protein. *Journal of Korean Society of Applied*
475 *Biology and Chemistry*, 52,290–594.

476 Lee, S.H. & Song, K.B. (2009b) Purification of an iron-binding nona-peptide from
477 hydrolysates of porcine blood plasma protein. *Process Biochemistry*, 44, 378–381.

478 Lennon, A.M., McDonald, K., Moon, S.S., Ward, P. & Kenny, T.A. (2010)
479 Performance of cold-set binding agents in re-formed beef steaks. *Meat Science*, 85,
480 620–624.

481 Liu, X., Song, C., Chen, R., Jiang, X., Jin, Y. & Zou, H. (2010) Identification of
482 angiotensin I-converting enzyme inhibitors in peptides mixture of hydrolyzed red deer
483 plasma with proteomic approach. *Chinese Journal of Chemistry*, 28,1665–1672.

484 Lopes, S.C., Fedorov, A. & Castanho, M.A. (2005) Lipidic membranes are potential
485 “catalysts” in the ligand activity of multifunctional pentapeptide neokyotorphin.
486 *Chemistry and Biochemistry*, 6, 697-702.

487 Maehashi, K., Matsuzaki, M., Yamamoto, Y. & Udaka, S. (1999) Isolation of peptides
488 from an enzymatic hydrolysate of food proteins and characterization of their taste
489 properties. *Bioscience, Biotechnology and Biochemistry*, 63, 555-559

490 Martínez-Alvarez, O., Chamorro, S., Brenes, A. (2015) Protein hydrolysates from
491 animal processing by-products as a source of bioactive molecules with interest in
492 animal feeding: A review. *Food Research International*, 73, 204–212.

493 Marulanda, V.F., Anitescu, G. & Tavlarides, L.L. (2010) Investigations on supercritical
494 transestrification of chicken fat for biodiesel production from low cost lipid feedstocks.
495 *Journal of Supercritical Fluids*, 54, 53-60.

496 Mito, K., Fujii, M., Kuwahara, M., Matsumura, N., Shimizu, T., Sugano, S. & Karaki,
497 H. (1996) Antihypertensive effect of angiotensin I-converting enzymeinhibitory
498 peptides derived from hemoglobin. *European Journal of Pharmacology*, 304,93–98.

499 Mora, L., Reig, M., Toldrá, F. (2014) Bioactive peptides generated from meat industry
500 by-products. *Food Research International*, 65, 344-349.

501 Mora, L., Gallego, M., Escudero, E., Reig, M., Aristoy, M-C. & Toldrá, F. (2015) Small
502 peptides hydrolysis in dry-cured meats. *International Journal of Food Microbiology*,
503 212, 9-15.

504 Moreira, A.L., Dias, J.M., Almeida, M.F. & Alvim-Ferraz, C.M. (2010) Biodiesel
505 production through transesterification of poultry fat at 30°C. *Energy Fuels*, 24, 5717-
506 5721.

507 Murray, S.M., Patil, A. R., Fahey, Jr., G.C., Merchen, N.R. & Hughes, D.M. (1997)
508 Raw and Rendered Animal By-Products as Ingredients in Dog Diets. *Journal of Animal*
509 *Science*, 75, 2497-2505.

510 Nchienzia, H. A., Morawicki, R. O., & Gadang, V. P. (2010). Enzymatic hydrolysis of
511 poultry meal with endo- and exopeptidases. *Poultry Science*, 89, 2273–2280.

512 Nedjar-Arroume, N., Dubois-Delval, V., Miloudi, K., Daoud, R., Krier, F., Kouach, M.,
513 Briand, G. & Guillochon, D. (2006) Isolation and characterization of four antibacterial
514 peptides from bovine hemoglobin. *Peptides*, 27, 2082-2089.

515 Nedjar-Arroume, N., Dubois-Delval, V., Adje E. Y., Traisnel, J., Krier, F., Kouach, M.,
516 Briand, G. & Guillochon, D. (2008) Bovine hemoglobin: an attractive source of
517 antibacterial peptides. *Peptides*, 29, 969-977.

518 Nissenson, A. R., Berns, J. S., Sakiewicz, P., Ghaddar, S., Moore, G. M., Schleicher, R.
519 B. & Seligman, P.A. (2003) Clinical evaluation of heme iron polypeptide: sustaining a
520 response to rHuEPO in hemodialysis patients. *American Journal of Kidney Disorders*,
521 42, 325-330.

522 Nollet, L.M.L. & Toldrá, F. (2011) Introduction. Offal meat: Definitions, regions,
523 cultures, generalities. In: *Handbook of Analysis of Edible Animal By-Products* (L.M.L.
524 Nollet & F. Toldrá, Eds.), pp 3-11, CRC Press, Boca Raton FL, USA.

525 Ockerman, H.W. & Basu, L. (2004a) By-products. In: *Encyclopedia of Meat Sciences*
526 (W. Jensen, C. Devine & M. Dikemann, Eds.), pp 104-112, London, UK: Elsevier Science
527 Ltd.

528 Ockerman, H.W. & Basu, L. (2004b) Hides and skins. In: *Encyclopedia of Meat*
529 *Sciences* (W. Jensen, C. Devine & M. Dikemann, Eds.), pp 125-138, London, UK:
530 Elsevier Science Ltd.

531 Ockerman, H.W. & Basu, L. (2006) Edible rendering-rendered products for human use.
532 In: *Essential rendering: All about the animal by-products industry* (D.L. Meeker, Ed.), pp
533 95-110, National Renderers Association, Arlington, VA, USA.

534 Ofori, J.A. & Hsieh, Y-H. P. (2014) Issues Related to the Use of Blood in Food and
535 Animal Feed. *Critical Reviews in Food Science & Technology*, 54, 687-697.

536 Park, E., Won, M., Lee, H-H. & Bin Song, K. (1996) Angiotensin converting enzyme
537 inhibitory pentapeptide isolated from supernatant of pig plasma treated by
538 trichloroacetic acid. *Biotechnology Techniques*, 10, 479–480.

539 Pearl, G.G. (2004) Inedible. In: *Encyclopedia of Meat Sciences* (W. Jensen, C. Devine &
540 M. Dikemann, Eds.), 112-125, London, UK: Elsevier Science Ltd.

541 Pérez-Galvez, R., Almecija, M.C., Espejo, F.J., Guadix, E.M., Guadix, A. (2011)
542 Biobjective optimisation of the enzymatic hydrolysis of porcine blood protein.
543 *Biochemical Engineering Journal*, 53, 305–310.

544 Piot, J.M., Zhao, Q.Y., Guillochon, D., Ricart, G. & Thomas, D. (1992) Isolation and
545 characterization of two opioid peptides from a bovine hemoglobin peptic hydrolysate.
546 *Biochemical and Biophysical Research Communications*, 189, 101–110.

547 Ren, Y., Wan, D.-G., Lu, X., Chen, L., Zhang, T. & Guo J. (2011) Isolation and
548 characterization of angiotensin I converting enzyme inhibitor peptides derived from
549 porcine hemoglobin. *Scientific Research and Essays*, 6, 6262-6269.

550 Riedel, S.L., Jahnsa, S., Koeniga, S., Bocka, M.C.E., Brigham, C.J., Bader, J. &
551 Stahla, U. (2015) Polyhydroxyalkanoates production with *Ralstonia eutropha* from
552 lowquality waste animal fats, *Journal of Biotechnology*, 214, 119–127.

553 Ryder, K., Ha, M., El-Din Bekhit, A., & Carne, A. (2015). Characterisation of novel
554 fungal and bacterial protease preparations and evaluation of their ability to hydrolyse
555 meat myofibrillar and connective tissue proteins. *Food Chemistry*, 172, 197–206.

556 Ryder, K., El-Din Bekhit, A., McConnell, M. & Carne, A. (2016). Towards generation
557 of bioactive peptides from meat industry waste proteins: Generation of peptides using
558 commercial microbial proteases. *Food Chemistry*, 208, 42–50

559 Saiga, A., Iwai, K., Hayakawa, T., Takahata, Y., Kitamura, S., Nishimura, T. &
560 Morimatsu, F. (2008). Angiotensin I-converting enzyme-inhibitory peptides obtained
561 from chicken collagen hydrolysate. *Journal of Agricultural and Food Chemistry* 56,
562 9586-9591.

563 Toldrá, F. & Reig, M. (2011) Innovations for healthier processed meats. *Trends in Food*
564 *Science & Technology*, 22, 517-522.

565 Toldrá, F., Aristoy, M-C., Mora, L., Reig, M. (2012) Innovations in value-addition of
566 edible meat byproducts. *Meat Science*, 92, 290-296.

567 Valta, K., Damala, P., Orli, E., Papadaskalopoulou, C., Moustakas, K., Malamis, D. &
568 Loizidou, M. (2015) Valorisation Opportunities Related to Wastewater and Animal
569 By-Products Exploitation by the Greek Slaughtering Industry: Current Status and Future
570 Potentials. *Waste Biomass Valor*, 6, 927–945.

571 Vercruysse, L., Van Camp, J. & Smaghe, G. (2005) ACE inhibitory peptides derived
572 from enzymatic hydrolysate of animal protein: A review. *Journal of Agricultural &*
573 *Food Chemistry*, 53, 8106-8115.

574 Viana, F.R., Silva, V.D.M., Delvivo, F.M., Bizzotto, C.S. & Silvestre, M.P.C. (2005).
575 Quality of ham pâté containing bovine globin and plasma as fat replacers. *Meat Science*,
576 70, 153-160.

577 Yousif, A.M., Cranston, P. & Deeth, H.C. (2003) Incorporation of bovine dry blood
578 plasma into biscuit flour for the production of pasta. *LWT Food Science and*
579 *Technology*, 36, 295-302.

580 Yu, Y., Hu, J., Miyaguchi, Y., Bai, X., Du, Y. & Lin, B. (2006) Isolation and
581 characterization of angiotensin I-converting enzyme inhibitory peptides derived from
582 porcine hemoglobin. *Peptides*, 27, 2950–2956.

583 Zhang, W., Xiao, S., Samaraweera, H., Lee, E.J. & Ahn, D.U. (2010). Improving
584 functional value of meat products. *Meat Science*, 86, 15-31.

585 Zhao, Q., Sannier, F., Garreau, I., Guillochon, D. & Piot, J.M. (1994) Inhibition and
586 inhibition kinetics of angiotensin cinverting enyzme activity by hemorphins isolated
587 from peptic bovine hydrolysate. *Biochemical and Biophysical Research*
588 *Communications*, 204, 216-223.

589 Zhao, Q., Garreau, I., Sanier, F. & Piot, J.M. (1997) Opioid peptides derived from
590 hemoglobin: hemorphins. *Biopolymers*, 43, 75-98.

591

592

593 **Legends for the figures**

594

595 Figure 1.- Flow diagram of main routes of applications for meat by-products

596

597