Running title: Genetic basis of the hoopoe microbiome

Title: **The microbiome of the uropygial secretion in hoopoes is shaped along the nesting phase**

*Martínez-García, Ángela¹, Martín-Vivaldi, Manuel³, Ruíz-Rodríguez Magdalena¹,
Martínez-Bueno, Manuel²; Arco, Laura³, Rodríguez-Ruano, Sonia M.², Peralta-
Sánchez, Juan Manuel³, Soler, Juan José^{1*}*

¹ Estación Experimental de Zonas Áridas (CSIC) E-04120 Almería, Spain

² Departamento de Microbiología Universidad de Granada, E-18071 Granada, Spain

³ Departamento de Zoología Universidad de Granada, E-18071 Granada, Spain

*corresponding author;

Email: jsoler@eeza.csic.es

Tel: 34 950 281045

Fax: 34 950 277100

Word count: 6630

Table count: 1

Figure count: 3

Abstract

Microbial symbiont acquisition by hosts may determine the effectiveness of the mutualistic relationships. A mix of vertical and horizontal transmission may be advantageous for hosts by allowing plastic changes of microbial communities depending on environmental conditions. Plasticity is well known for gut microbiota, but is poorly understood for other symbionts of wild animals. We here explore the importance of environmental conditions experienced by nestling hoopoes (*Upupa epops*) during the late nesting phase determining microbiota in their uropygial gland. In cross-fostering experiments of 8 days old nestlings, “sibling-sibling” and “mother-offspring” comparisons were used to explore whether the bacterial community naturally established in the uropygial gland of nestlings could change depending on experimental environmental conditions (i.e. new nest environment). We found that the final microbiome of nestlings was mainly explained by nest of origin. Moreover, cross-fostered nestlings were more similar to their siblings and mothers than to their stepsiblings and stepmothers. We also detected a significant effect of nest of rearing, suggesting that nestling hoopoes acquire most bacterial symbionts during the first days of life, but that the microbiome is dynamic and can be modified along the nestling period depending on environmental conditions. Estimated effects of nest of rearing, but also most of those of nest of origin are associated to environmental characteristics of nests, which are extended phenotypes of parents. Thus, natural selection may favor the acquisition of appropriated microbial symbionts for particular environmental conditions found in nests.

Key-words: Cross-fostering experiment, Horizontal transmission, Microbial symbiont, Microbial transmission, Parent-Offspring comparisons, Plasticity, *Upupa epops*, Vertical transmission

Introduction

Hosts may acquire symbionts directly by vertical transmission from parents to offspring [1, 2], or by horizontal transmission from the environment [3]. Although the vast majority of symbioses described in eukaryotes involve bacteria [1, 4], studies on mechanisms of bacterial transmission are limited to a handful of model systems [5, 6]. Horizontally transmitted bacteria are known for squids [3], tubeworms [7] and mussels [8], while mechanisms of vertical transmission have been detected for instance in ascidians [9], bryozoans [10] and earthworms [11]. For some other model systems, microbial symbionts are acquired both vertically and horizontally, as it is the case for beneficial gastrointestinal microbiomes of animals [12] or for enterococci of the uropygial gland of hoopoes (*Upupa epops*) [13].

Modes of bacterial acquisition may determine the effectiveness of the mutualistic relationship [14]. On the one hand, fitness of vertically transmitted symbionts is closely related to that of their hosts and, thus, enhancing reproductive success of hosts will directly benefit their own performance [14-17]. On the other hand, different symbionts may provide hosts with characteristics that are more appropriate for particular environments and/or symbiont effectiveness may vary with environmental conditions. Although vertical transmission is identified as a key route to host specialization on effective symbionts [14, 18], hosts with horizontally transmitted symbionts have the opportunity to adjust the community of symbionts to environmental characteristics [19]. In situations of variable environmental conditions with unpredictable selection pressures, a mix of vertical and horizontal transmission shaping symbiont bacterial communities might be of selective advantage for hosts because it would guarantee the simultaneous presence of potential beneficial microorganisms from

an for different environments [5, 14, 20]. In this mixed mode of symbiont acquisition the symbiotic community that hosts acquire from mothers and/or environment during the first days of life could change in relation to variable environmental conditions experienced later, in subsequent phases of life. Symbiotic community changes (i.e., plasticity) of hosts in relation to environmental changes are well known for gut microbiota [21, 22] but are poorly understood for other symbionts of wild animals.

We experimentally explore whether microbiome of the uropygial gland of nestling hoopoes change along the nesting period in relation to environmental conditions hosted in their uropygial gland. Symbiotic bacteria have only been detected by traditional culture methods in incubating females and nestlings [23, 24]. Nowadays we know that males and non-reproducing females also harbor bacteria in their gland, but at very low density [25]. Interestingly, we also know that antibiotic producing enterococci are transmitted from mother hoopoes to offspring soon after hatching (vertical transmission), and that hatchlings (i.e. before uropygial gland functions) are able to incorporate new enterococci symbionts from new environments after cross-fostering experiments (horizontal transmission) [13]. The question that we try to answer here is whether the bacterial community, once it is established in the uropygial gland of nestlings (i.e. functioning), could change depending on experimental environmental conditions (i.e. new nest environment). We approached this aim by cross-fostering nestlings of intermediate age (i.e. with an established bacterial community) from different nests and characterizing bacterial communities of fledglings by mean of ARISA (Automatic Ribosomal Intergenic Spacer Analysis). We first describe the bacterial communities in uropygial secretions of nestlings and females in terms of prevalence, richness, and Operational Taxonomic Units (OTUs) composition. The relative contribution of factors acting during early (i.e. genetic plus early environment

including any pre-manipulation maternal effects) and late (experimental nest environment) nesting phase were determined by (i) comparing the proportion of variance explained by nest of origin and nest of rearing respectively. Relative contribution of early and later environments explaining microbiota of the uropygial secretion was also explored by comparing (ii) levels of similarity between siblings reared in different nests and between stepsiblings reared in the same nest, and (iii) between cross-fostered nestlings and biological and stepmothers.

Materials and methods

Study species, study area and general methods

The hoopoe is distributed throughout Europe, Asia and Africa, inhabiting open woods or open areas as steppes, grasslands, pastures, semi-deserts, or crops whenever they have scattered trees, walls or buildings providing holes for nesting and soil without tall vegetation for feeding [26-28]. Females lay one or two clutches of 6-8 eggs along the breeding season, between February and July [29]. Incubation lasts 17 days and starts with the first or second egg, which results in eggs hatching asynchronously at 24 h or even greater intervals [30].

The fieldwork was performed during the breeding seasons 2010-2011 in a wild population located in the Hoya de Guadix (37°18'N, 38°11'W), southern Spain, where hoopoes breed in crops, forests and gullies within nest-boxes placed in trees or buildings. In 2011, hoopoes were also sampled in a captive population descendant from our wild population and breeding in captivity since 2008. The captive pairs were distributed in two different subpopulations located in South-eastern Spain, one of them in installations of the University of Granada in Hoya of Guadix (Granada), and the other one in facilities of the Estación Experimental de Zonas Áridas (CSIC) at the Finca

Experimental La Hoya in Almería (36°50'N, 2°28'W). All females and nestlings were ringed with numbered rings and females also with color rings for individual recognition.

Nestboxes in the wild were visited twice per week, from mid-February to the end of July to record laying date, clutch size and hatching date. Pairs of hoopoes breeding in captivity were housed in independent cages at least 3m x 2m x 2m installed in the open, scattered and isolated to avoid interactions between pairs and ensure successful breeding. Cages had access to soil and were provided with live food (crickets, vitamin-enriched fly larvae and meat (beef heart)) *ad libitum* and were visited daily.

Experimental design and sampling

The cross-fostering experimental design consisted in the exchange of two experimental nestlings among pairs of nests of similar (± 1 day) hatching date and similar brood size. The exchange was carried out when the oldest nestling in each nest was 8 days old (i.e., when nestlings start to produce secretion containing bacteria). Two of the four heaviest nestlings in each nest were randomly selected and exchanged with those from another nest (i.e. with the same age and similar weight). Comparisons were later performed with all nestling in the experimental nests. Nestlings were individually marked by painting their tarsus with permanent innocuous markers. Cross-fostering experiments were performed between wild nests in 2010 and in 2011 between one nest in captivity and the other in wild conditions. When this was not possible, experimental nestlings were exchanged between two captivity nests, or between two wild nests. This was done so to increase phenotypic variance among cross-fostered nests that allow a more realistic estimation of the effects of nest of origin and of nest of rearing (Falconer 1989).

Transport of nestlings between nests lasted approximately 1 h and was done in a portable incubator at 37 °C to reduce stress due to temperature change.

Uropygial secretions of females were sampled before hatching date (i.e. 14 days after laying the first egg), whereas those of nestlings were sampled 10 days after nest exchange (i.e. oldest nestlings had 18 days old). Incubating females were captured within the nest-box by hand, quickly sampled, and released again within the nest to reduce disturbance. For each capture, we wore new sterile latex gloves cleaned with 70% ethanol for the whole process to limit external bacterial contamination. Before collecting samples from uropygial gland, we softly washed the circlet of feathers and surrounding skin with a cotton swab slightly soaked in ethanol to reduce the risk of contamination with external bacteria. After evaporation of the alcohol, a sterile micropipette tip (1-10 µl micropipette (Finpipette)) was introduced in the gland papilla. The papilla was pressed softly with a finger and the uropygial secretion entirely collected was transferred to a sterile microfuge tube. Afterwards, 5µl were separated in a different sterile microfuge tube for the molecular analyses. Nestling hoopoes were sampled with identical protocol than adult females were. For further molecular analyses, all samples were individually stored in 1.2 ml sterile microfuge tubes in a portable cooler (1-3 °C) until being stored in the lab at -20° C the same day of sampling.

We sampled 44 nests and got information for the 44 breeding females and for 165 nestlings; 93 of them did grow in the same nests where they hatched, whereas 72 were moved to foreign nests. However, final sample sizes were reduced due to predation of wild nests or failures with ARISA. We obtained complete information of siblings that were reared in the same nests of hatching ($N = 57$) or moved to another nests ($N = 44$) for 28 nests. Only for 21 of these nests, we got the necessary information to compare the bacterial community of experimental nestlings with that of their foster

and genetic siblings on the one hand, and with the bacterial community of their mother and stepmother on the other hand.

Laboratory work

Bacterial genomic DNA for the uropygial secretion samples was extracted with a commercial kit (The FavorPrep™ Blood Genomic DNA Extraction Kit, Favorgen). ARISA (Fisher and Triplett 1999), which amplifies an intergenic transcribed spacer (ITS) region between the prokaryotic 16S and 23S rDNA, was used to characterize the composition of bacterial communities. This region is highly variable both in size and sequence between species and, thus, offers an appropriate taxonomic resolution for microbiota characterization (Danovaro et al. 2006). The ITS was amplified using the primer pair ITSF (5'-GTCGTAACAAGGTAGCCGTA-3') and ITSReub (5'-GCCAAGGCATCCACC-3') [31]. The primer ITSReub was labeled fluorescently with 6-FAM. Amplifications were performed in 50 µL reaction volumes containing ultrapure H₂O, 20 µL of 5 PRIME MasterMix (2.5x) including 1.5mM Mg(OAC)₂, 200 µM dNTPs, 1.25 U Taq DNA polymerase 0.2 µM of primers and 5µL of diluted DNA 1:10. PCRs were carried out in Eppendorf Mastercycler Nexus. Fragments were amplified under the following conditions: initial denaturation at 94 °C 2 min, followed by 30 cycles with denaturation at 94 °C 45 s, annealing at 52 °C 45 s and extension at 72 °C 1 min, with a final extension at 72 °C 5 min. Amplified PCR products were diluted 1:10 and denatured by heating in formamide. Fragment lengths were determined by mean of automated fluorescent capillary electrophoresis on 3130 Genetic Analyzer. Electropherogram peak values were calculated after interpolation with an internal size standard named GeneScan™ 1200 LIZ dye Size Standard (both Applied Biosystems).

These analyses were performed at the Scientific Instrumentation Center of the University of Granada.

Statistical analysis

Peak Scanner 1.0 (Applied Biosystems) was used to determine fragment length (i.e. OTUs) in terms of base pairs (bp). For binning DNA fragment lengths from different samples we used available scripts in R-environment [<http://cran.r-project.org/>] at www.ecology-research.com [32] with a window size of 4 base pairs (bp) and a distance of two consecutive binning frames (i.e. shift) of 0.1. Peaks with RFI values of $< 0.09\%$ were considered as background peaks. Only fragments above a threshold of 50 fluorescence units and ranging between 100 and 1,000 bp [32] were used for constructing and analyze the presence-absence matrices depicting bacterial communities.

We described the bacterial community harbored in uropygial secretion of females and nestlings with information obtained from ARISA for all individuals (44 females and 165 nestlings). To explore the differences in bacterial richness (number of OTUs per sample) between adult females and nestlings, we performed ANOVAs with one fixed factor (adult females *vs.* nestlings). Moreover, we explored differences in prevalence of OTUs detected in uropygial secretions of adult females and nestlings, but considering the most frequent OTUs; i.e., those that were detected in more than 30% of females or nestlings uropygial secretion sampled. Pearson correlation coefficients were used to explore whether OTU's prevalence in females and nestling samples were related. We did this analysis with all detected OTUs and also including only those that were present in more than 30% of females or nestlings sampled. Furthermore, we analyzed differences in the composition of bacterial communities hosted in uropygial

secretions of females and nestlings by one-way PERMANOVAs analysis (Jaccard's distance), taking into consideration all females and only non-moved nestlings. Trying to reduce probability of detecting significant differences among females and nestlings due to rare OTUs, we only considered those that appeared in more than 3 samples of females or nestlings. We used classical multidimensional scaling analysis, Principal Coordinates Analysis (PCoA) to graphically show variation in bacterial communities of uropygial secretions of females and nestlings. This technique represents the objects (communities) on a plot with canonical axes, where the distance between the objects shows their underlying similarity [33].

Cross-fostering experiments are a well-established approach for partitioning phenotypic variance in its genetic and environmental components in mixed statistical models that include the identity of nest of origin and rearing (nested within nest of origin) as random factors [34]. This experimental approach has been previously used to explore genetic and environmental influences determining cloacal bacterial assemblages in great tit (*Parus major*) [35], and the enterococci community of the uropygial gland of hoopoes [13]. Here, we performed cross-fostering experiments of nestlings of intermediate age to estimate the relative importance of early and late nestling periods explaining the whole microbiota of the uropygial gland of hoopoes. The effects of early nestling phase, which include genetic component and any pre-manipulation maternal and environmental effects, were estimated by the proportion of variance of microbiome composition explained by nest of origin and by mother-offspring (i.e. vertical transmission) and sibling-sibling comparisons. The effects of late nesting phase, which would only include environmental components, were estimated by the proportion of variance associated to the nest of rearing and by stepmother-offspring and sibling-stepsibling comparisons.

The similarity matrix among all bacterial communities of the sampled individuals were based in Jaccard's distance [36]. The similarity values were used as the dependent variables of PERMANOVA model using type III estimation of mean squares. This model try to explain similarity among nestlings including two random factors: nest of origin and nest of rearing (nested within nest of origin). For this model we used only the 28 nests for which we have information for moved and non-moved nestlings from the same nest of origin. Finally, for the 21 experiments with all the information (see above), we estimated mean values of similarities among bacterial communities of experimental nestlings and those of their genetic (reared in different nests but genetically related) or foster (reared in the same environment but genetically unrelated) siblings. We estimated for the same nests mean values of similarities among bacterial communities of experimental nestlings and their genetic or foster mothers.

All multivariate analyses and figures trying to explain similarity matrices (PERMANOVAs) were performed with PRIMER v7 (PRIMER-E) software (Anderson et al. 2008). Statistical inferences (e.g., P-values) of all PERMANOVAs were based on 9999 permutations. Statistical tests trying to explain variation in bacterial richness and prevalence of different bacterial strains, as well as those comparing mean values of similarities estimated for genetically related and unrelated individuals, were performed with STATISTICA10 software [37].

Results

Description of bacterial communities in uropygial secretions: prevalence, richness, and composition

We detected 143 different OTUs (length of the ITS fragment varying between 100 bp and 847 bp) in the bacterial community of the uropygial secretion of female and nestling

hoopoes, 141 of which were present in nestlings, and 116 in females. All except two OTUs that were detected in females at very low prevalence (143 bp and 603 bp, 2.22% and 4.44%, respectively) were also present in nestling samples. Prevalence of detected OTUs ranged from 0.61% (OTU with 847 bp) to 84.44% (OTU with 183 bp), and were similar for females and nestlings as shown by the strong positive relationships among their values (Appendix 1, $R^2 = 0.89$, $N = 143$, $t = 34.2$, $p < 0.0001$). This relationship was evident even when only considering the 28 OTUs that were present in more than 30% of samples from females or nestlings (Fig. 1, $R^2 = 0.73$, $N = 28$, $t = 8.44$, $p < 0.0001$). Richness of bacterial community of the uropygial secretions of nestlings (mean (SE) = 22.64 (0.66)) was also similar to that of females (mean (SE) = 21.78 (1.37)) ($F = 0.34$, $df = 1$, 207, $p = 0.55$). Additionally, composition of bacterial communities of nestlings and females did not significantly differ (one-way PERMANOVA, $F = 1.53$, $df = 1$, 135, $p = 0.0572$, Fig. 2).

Effects of early and late nesting phase on the bacterial community

The similarity matrix among bacterial communities of the uropygial gland of experimental nestlings was significantly explained by nest of origin and nest of rearing (Table 1). The proportion of variance explained by the nest of origin was relatively larger than that explained by nest of rearing (Table 1). This result suggests that the influence of the early nesting phase (i.e., genetic factors and/or pre-manipulation maternal and environmental effects) explaining uropygial microbiome of hoopoes was relatively larger than that of late nesting period (i.e., environmental influence and maternal effects after the experiment). This inference was further confirmed by the significantly larger similarity values of comparisons of siblings reared in different nests than those for comparisons of stepsiblings reared in the same nest (GLM, $F = 19.33$, $df =$

1, 20, $p = 0.0002$; Fig. 3a). Results from comparisons of similarities between bacterial communities of cross-fostered nestlings and those of their foster and genetic mothers; (Fig.3b) were also in accordance with a relatively larger influence of the early nesting phases determining the bacterial community of the uropygial secretion of hoopoe nestlings (GLM, $F = 20.42$, $df = 1, 20$, $p = 0.0002$).

Importantly, we found statistically significant effects of nest of rearing explaining bacterial community of nestling hoopoes (Table 1), which suggest that environmental conditions experienced by hoopoes during the late nesting phase also contribute to mould microbial community.

Discussion

Our main results are twofold. The first one is that bacterial communities of the uropygial secretion of hoopoe nestlings did not differ significantly from those of their mothers. The second group of results pointed out strong pre-manipulation effects explaining the composition of bacterial communities in experimental cross-fostered nestlings. We also detected a significant effect of nest of rearing, suggesting that environmental characteristics experienced after experimental treatment contributes to the microbiome of the uropygial secretion of hoopoes. Below, we discuss the importance of these findings for understanding the mechanisms of acquisition of bacterial symbionts by nestling hoopoes, and the implication for coevolutionary relationships between hoopoes and bacteria of their uropygial secretion.

Previous work has shown that the prevalence of cultivable bacterial strains (i.e., enterococci) in the uropygial secretion of females and nestlings differs [13]. These differences were at least partially due to the effect of few enterococci strains that appeared at a higher prevalence in samples of females or nestlings [13]. However, when

considering the bacterial community as a whole, differences between females and nestlings did not reach statistical significance, and prevalence of different OTUs in samples from females and from nestlings correlated positively. In terms of bacterial diversity, even when considering the group of enterococci, estimates for females and nestlings did not differ significantly [13]. Thus, although prevalence of some OTUs in communities of females and nestlings may differ, the microbiome of the uropygial secretion of females and nesting hoopoes is quite similar.

Results from previous cross-fostering experiments performed by Ruiz-Rodríguez et al. [13] with recently hatched nestlings that were exchanged before they started to produce secretion (4 days old) strongly suggested that enterococci from the new environment are incorporated into the community of the uropygial secretion of nestlings, although some strains came from the uropygial secretion of mothers [13]. We here considered the whole microbiome of the uropygial gland of hoopoes, and performed the experiment with nestlings of intermediate age, once they started to produce uropygial secretion. Even with these nestlings, we also found a significant effect of nest of rearing and of nest of origin explaining variation in microbial community. Thus, the effect of nest of origin detected here also included environmental effects before the experiment and, accordingly, was stronger than that detected for enterococci community in cross-fostering experiments performed with 4-days-old nestlings. Moreover, the detected effect of nest of rearing of 8 days old cross-fostered nestlings confirms that experimental nestlings incorporate new bacteria to their uropygial microbiome once the uropygial gland is functioning. Consequently, results from these two experiments considered together suggest that nestling hoopoes acquire most bacterial symbionts during the first days of life, but that the microbiome of

hoopoes is dynamic and can be modified along the nestling period depending on environmental conditions.

The effect of either nest of origin or nest of rearing included possible maternal effects that respectively occur before (from genetic mother) and after (from stepmother) the experimental translocation of nestlings [34, 38]. Mechanisms of vertical transmission of symbionts is by definition a maternal effect that contributes to offspring phenotype, but that is genetically determined in mothers (i.e. indirect genetic effect, [39-41]). Thus, the detected effects of nest of origin on the microbial community of nestling secretions, as well as the relatively high similarities between related nestlings, and between nestlings and mothers, may be explained by direct vertical transmission of symbionts from mother to offspring. However, previous results suggested that direct contacts with mother or nest material are not necessary for hatchling hoopoes to develop normal uropygial glands and acquire enterococci symbionts [13]. Thus, it is unlikely that the strong influence of nest of origin, and the similarity among bacterial communities of related individuals detected here, were exclusively explained by vertical transmission. An additional interpretation of these results is that the first microbes to colonize the hatchling glands will exclude subsequent colonists. So an empty gland acquires any microbe, but this is biased in nature because the natal nest is dominated by microbes from the mother, and later from the hatchling itself after it has been colonized. Thus, the transmission would be technically horizontal (because it has an external phase) in this scenario, but the external phase is determined by the mother similar to vertical transmission.

An alternative explanation of the strong effects of nest of origin is that related hoopoes share characteristics of their uropygial gland and/or secretion (i.e., chemical properties) that influence the composition of the bacterial community established.

Bacteria from the environment that were compatible with characteristics of the uropygial gland and secretion of hoopoes would colonize hosts. Within the uropygial gland, competitive ability of different bacterial strains would depend on the particular environment (i.e. chemicals, resources, etc.) provided by hoopoes, which would determine the stabilized microbiome of the uropygial secretion [see 19]. Even if this was the explanation of the large detected microbiome similarities among relatives, the detected effects of nest of rearing suggest a plastic microbiome response to environmental changes after reaching certain stability level (i.e. uropygial gland functioning).

Nests of birds are considered as extended phenotypes of parents because nest site selection, nest building behavior, nest sanitation behavior, etc. are characters with strong genetic components [42, 43]. Thus, nests are indirect genetic (i.e., parental) effects for nestlings where natural selection would work [39]. Estimated effects of nest of rearing, but also most of those of nest of origin are likely associated to environmental characteristics of nests (included bacterial communities and characteristics of mothers). Thus, independently of the relative importance of genetic, environmental and maternal effects, the factors determining the bacterial community of the uropygial secretion of nestling hoopoes have a considerable genetic background to be modulated by natural selection.

Detecting mechanisms explaining how beneficial microbiomes are established and maintained within their hosts is a major question in evolutionary biology [5, 44]. Here, we found a strong effect of nest of origin that likely included indirect genetic effects, but also evidence of an influence of the environment. This effect suggests the composition of the microbial community in the uropygial secretion of hoopoes for which evidence of beneficial effects for hosts are accumulating [23, 45-47] is plastic.

We expect that these results will encourage further research directed to detect factors driving phenotypic plasticity of the symbiotic microbiome of hoopoe uropygial gland, including physiological and morphological characteristics of the gland as well as characteristics of the microbiome of mothers, siblings and nests.

Acknowledgements

We thank Estefanía López Hernández and Olga Corona Forero for help in laboratory work; Ana Belén García, Jonathan Romero Masegosa, Manuel Soto Cárdenas, Natalia Juárez García Pelayo and Jorge Doña Reguera for help in caring of captive hoopoes. Support by funding was provided by Spanish Ministerio de Economía y Competitividad, European funds (FEDER) (CGL2013-48193-C3-1-P, CGL2013-48193-C3-2-P) and Junta de Andalucía (P09-RNM-4557). AM-G had a predoctoral grant from the Junta de Andalucía (P09-RNM-4557).

Authorship statement

JJS and MM-V designed the study, JJS and AM-G carried out statistical analyses. AM-G together with SRR performed molecular analyses. AM-G, LA and JMPS performed most of the field work. AM-G wrote the first version of the manuscript with substantial contribution from MRR and JJS. All authors contributed to the final version of the article.

Reference List

1. Moran NA, Wernegreen JJ (2000) Lifestyle evolution in symbiotic bacteria: insights from genomics. *Trends Ecol Evol* 15:321-326
2. Darby AC, Douglas AE (2003) Elucidation of the transmission patterns of an insect-borne Bacterium. *Appl Environ Microbiol* 69:4403-4407

3. Nyholm SV, McFall-Ngai M (2004) The winnowing: establishing the squid-vibrio symbiosis. *Nat Rev Microbiol* 2:632-642. doi: 10.1038/nrmicro957
4. Lindquist N, Barber PH, and Weisz JB (2005) Episymbiotic microbes as food and defence for marine isopods: unique symbioses in a hostile environment. *Proc R Soc Lond B* 272:1209-1216
5. Chaston J, Goodrich-Blair H (2010) Common trends in mutualism revealed by model associations between invertebrates and bacteria. *FEMS Microbiol Rev* 34:41-58. doi: 10.1111/j.1574-6976.2009.00193.x
6. Salem H, Florez L, Gerardo N, and Kaltenpoth M (2015) An out-of-body experience: the extracellular dimension for the transmission of mutualistic bacteria in insects. *Proc R Soc Lond B* 282. doi: 10.1098/rspb.2014.2957
7. Nussbaumer AD, Fisher CR, and Bright M (2006) Horizontal endosymbiont transmission in hydrothermal vent tubeworms. *Nature* 441:345-348. doi: 10.1038/nature04793
8. Salerno JL, Macko SA, Hallam SJ, Bright M, Won YJ, McKiness Z, and Van Dover CL (2005) Characterization of symbiont populations in life-history stages of mussels from chemosynthetic environments. *Biol Bull* 208:145-155
9. Hirose E, Adachi R, and Kuze K (2006) Sexual reproduction of the *Prochloron*-bearing ascidians, *Trididemnum cyclops* and *Lissoclinum bistratum*, in subtropical waters: seasonality and vertical transmission of photosymbionts. *J Mar Biol Assoc U K* 86:175-179
10. Sharp KH, Davidson SK, and Haygood MG (2007) Localization of '*Candidatus Endobugula sertula*' and the bryostatins throughout the life cycle of the bryozoan *Bugula neritina*. *ISME J* 1:693-702
11. Davidson SK, Stahl DA (2008) Selective recruitment of bacteria during embryogenesis of an earthworm. *ISME J* 2:510-518
12. Bright M, Bulgheresi S (2010) A complex journey: transmission of microbial symbionts. *Nat Rev Microbiol* 8:218-230. doi: 10.1038/nrmicro2262
13. Ruiz-Rodríguez M, Soler JJ, Martín-Vivaldi M, Martín-Platero AM, Méndez M, Peralta-Sánchez JM, Ananou S, Valdivia E, and Martínez-Bueno M (2014) Environmental factors shape the community of symbionts in the uropygial gland of hoopoes more than genetic factors. *Appl Environ Microbiol* 80:6714-6723
14. Douglas AE (1998) Host benefit and the evolution of specialization in symbiosis. *Heredity* 81:599-603
15. Herre EA (1993) Population structure and the evolution of virulence in nematode parasites of fig wasps. *Science* 259:1442-1445
16. Frank SA (1996) Models of Parasite Virulence. *Q Rev Biol* 71:37-78

- 460 17. Poulsen M, Bot ANM, Currie CR, Nielsen MG, and Boomsma JJ (2003)
461 Within-colony transmission and the cost of a mutualistic bacterium in the leaf-
462 cutting ant *Acromyrmex octospinosus*. *Funct Ecol* 17:260-269
- 463 18. Frank SA (1997) Models of symbiosis. *Am Nat* 150:s80-s99
- 464 19. Scheuring I, Yu DW (2012) How to assemble a beneficial microbiome in three
465 easy steps. *Ecol Lett* 15:1300-1307
- 466 20. Currie CR, Poulsen M, Mendenhall J, Boomsma JJ, and Billen J (2006)
467 Coevolved crypts and exocrine glands support mutualistic bacteria in fungus-
468 growing ants. *Science* 311:81-83
- 469 21. Quercia S, Candela M, Giuliani C, Turroni S, Luiselli D, Rampelli S, Brigidi P,
470 Franceschi C, Bacalini MG, Garagnani P, and Pirazzini C (2014) From lifetime
471 to evolution: escales of human gut microbiota adaptation. *Frontiers in*
472 *Microbiology* 5. doi: 10.3389/fmicb.2014.00587
- 473 22. Moeller AH, Li YY, Ngole EM, Ahuka-Mundeke S, Lonsdorf EV, Pusey AE,
474 Peeters M, Hahn BH, and Ochman H (2014) Rapid changes in the gut
475 microbiome during human evolution. *Proceedings of the National Academy of*
476 *Sciences of the United States of America* 111:16431-16435
- 477 23. Soler JJ, Martín-Vivaldi M, Ruiz-Rodríguez M, Valdivia E, Martín-Platero AM,
478 Martínez-Bueno M, Peralta-Sánchez JM, and Méndez M (2008) Symbiotic
479 association between hoopoes and antibiotic-producing bacteria that live in their
480 uropygial gland. *Funct Ecol* 22:864-871
- 481 24. Martín-Vivaldi M, Ruiz-Rodríguez M, Soler JJ, Peralta-Sanchez JM, Mendez
482 M, Valdivia E, Martin-Platero AM, and Martínez-Bueno M (2009) Seasonal,
483 sexual and developmental differences in hoopoe *Upupa epops* preen gland
484 morphology and secretions: evidence for a role of bacteria. *J Avian Biol* 40:191-
485 205
- 486 25. Rodríguez-Ruano SM (2015) Diversidad bacteriana en la glándula uropigial de
487 la abubilla: dinámica estacional y beneficios asociados. Dissertation, Granada
488 University
- 489 26. Rehsteiner U (1996) Abundance and habitat requirements of the Hoopoe *Upupa*
490 *epops* in Extremadura (Spain). *Ornithol Beobachter* 93:277-287
- 491 27. Barbaro L, Couzi L, Bretagnolle V, Nezan J, and Vetillard F (2008) Multi-scale
492 habitat selection and foraging ecology of the eurasian hoopoe (*Upupa epops*) in
493 pine plantations. *Biodivers Conserv* 17:1073-1087
- 494 28. Schaub M, Martinez N, Tagmann-Ioset A, Weisshaupt N, Maurer ML, Reichlin
495 TS, Abadi F, Zbinden N, Jenni L, and Arlettaz R (2010) Patches of bare ground
496 as a staple commodity for declining ground-foraging insectivorous farmland
497 birds. *PLoS ONE* 5:e13115. doi: 10.1371/journal.pone.0013115

- 498 29. Martín-Vivaldi M, Palomino JJ, Soler M, and Soler JJ (1999) Determinants of
499 reproductive success in the Hoopoe *Upupa epops*, a hole-nesting non-passerine
500 bird with asynchronous hatching. *Bird Study* 46:205-216
- 501 30. Cramp S. (1998) Cramp's the complete birds of the Western Palearctic.
502 Optimedia, Oxford University Press, Oxford
- 503 31. Cardinale M, Brusetti L, Quatrini P, Borin S, Puglia A, Rizzi A, Zanardini E,
504 Sorlini C, Corselli C, and Daffonchio D (2004) Comparison of different primer
505 sets for use in automated ribosomal intergenic spacer analysis of complex
506 bacterial communities. *Appl Environ Microbiol* 70:6147-6156
- 507 32. Ramette A (2009) Quantitative community fingerprinting methods for
508 estimating the abundance of operational taxonomic units in natural microbial
509 communities. *Appl Environ Microbiol* 75:2495-2505
- 510 33. Legendre P, Legendre L. (1998) Numerical Ecology. Elsevier Science,
511 Amsterdam
- 512 34. Merilä J (1996) Genetic variation in offspring condition: an experiment. *Funct*
513 *Ecol* 10:465-474
- 514 35. Lucas FS, Heeb P (2005) Environmental factors shape cloacal bacterial
515 assemblages in great tit *Parus major* and blue tit *P. caeruleus* nestlings. *J Avian*
516 *Biol* 36:510-516
- 517 36. Zuur AF, Ieno EN, , Smith GM. (2007) Analysing ecological data. Springer,
518 New York, NY
- 519 37. Statsoft Inc. (2011) STATISTICA (data analysis software system), version 10.
520 www.statsoft.com,
- 521 38. Soler JJ, Moreno J, and Potti J (2003) Environmental, genetic and maternal
522 components of immunocompetence of nestling pied flycatchers from a cross-
523 fostering study. *Evol Ecol Research* 5:259-272
- 524 39. Mousseau TA, Fox CW (1998) The adaptive significance of maternal effects.
525 *Trends Ecol Evol* 13:403-407
- 526 40. Wolf JB, Brodie III ED (1998) The coadaptation of parental and offspring
527 characters. *Evolution* 52:299-308
- 528 41. Wolf JB, Brodie III ED, Cheverud JM, Moore AJ, and Wade MJ (1998)
529 Evolutionary consequences of indirect genetic effects. *Trends Ecol Evol* 13:64-
530 69
- 531 42. Hansell M. (2000) Bird nests and construction behaviour. Cambridge University
532 Press, Cambridge
- 533 43. Walsh PT, Hansell M, Borello WD, and Healy SD (2009) Repeatability of nest
534 morphology in African weaver birds. *Biol Lett*

44. Prosser JI, Bohannan BJM, Curtis TP, Ellis RJ, Firestone MK, Freckleton RP, Green JL, Green LE, Killham K, Lennon JJ, Osborn AM, Solan M, van der Gast CJ, and Young JP (2007) The role of ecological theory in microbial ecology. *Nat Rev Microbiol* 5:384-392. doi: 10.1038/nrmicro1643
45. Ruiz-Rodríguez M, Valdivia E, Soler JJ, Martín-Vivaldi M, Martín-Platero AM, and Martínez-Bueno M (2009) Symbiotic bacteria living in the hoopoe's uropygial gland prevent feather degradation. *J Exp Biol* 212:3621-3626. doi: 10.1242/jeb.031336
46. Martín-Vivaldi M, Peña A, Peralta-Sánchez JM, Sánchez L, Ananou S, Ruiz-Rodríguez M, and Soler JJ (2010) Antimicrobial chemicals in hoopoe preen secretions are produced by symbiotic bacteria. *Proc R Soc Lond B* 277:123-130. doi: 10.1098/rspb.2009.1377
47. Martín-Vivaldi M, Soler JJ, Peralta-Sánchez JM, Arco L, Martín-Platero AM, Martínez-Bueno M, Ruiz-Rodríguez M, and Valdivia E (2014) Special structures of hoopoe eggshells enhance the adhesion of symbiont-carrying uropygial secretion that increase hatching success. *J Anim Ecol* 83:1289-1301. doi: 10.1111/1365-2656.12243

555 **Table 1.** Results of a PERMANOVA model explaining matrices of similarity among
 556 the bacterial communities found in the uropygial secretions of hoopoe nestlings. The
 557 model includes identity of nest of origin (genetic factor) and rearing (environmental
 558 factor) nested within nest of origin. Bold p-values are those lower than 0.05.

Factors	Pseudo-F	df	p	Permutations	% Explained variance
(a) Nest of origin	3.56	27	>0.001	9693	37.0
(b) Nest of rearing (nested in (a))	1.28	27	0.013	9737	14.9

559

560

Figure 1. Prevalence (%) of different bacterial OTUs (named by their length in terms of base pairs (bp)) found in more than 30% of samples from uropygial glands of hoopoe nestlings (N = 165) and females (N = 44).

Figure 2. Multidimensional space representation (PCoA) based on similarities of the OTUs communities harbored in uropygial secretions of hoopoe females and non-moved nestlings. The total variance explained is also shown (captured by the three axes = 33.1%).

Figure 3. Similarities in composition of bacterial communities (Jaccard's distance in percentage) among samples from cross-fostering experiments. (A) Similarity between uropygial secretions of nestlings that did not grow in their nests of origin (experimental nestlings) and genetic siblings reared in their native nests (between siblings), or foster siblings (between stepsiblings). (B) Similarities between microbiomes of the experimental moved nestlings and those of their genetic (mother - nestlings) or foster (stepmother - nestlings) mothers. Lines connect estimates of individuals from the same pair of cross-fostered nests.

579 Fig 1

580

581

582

583 Fig 2

584

585

586

587 Fig 3

589
590

589 **Appendix 1.** Prevalence (%) of different bacterial OTUs (named by their length in terms of base pairs (bp)) found in
 590 all sampled uropygial glands of nestlings (N=165) and females (N=44). Bold numbers show OTUs that were detected
 591 in more than 30% of samples from females or nestlings

OTU	Females	Nestlings	OTU	Females	Nestlings	OTU	Females	Nestlings
100	11.11	9.09	295	0.00	1.21	499	2.22	5.45
103	2.22	1.82	299	22.22	13.33	503	0.00	1.21
107	0.00	0.61	303	37.78	41.21	507	6.67	4.85
111	6.67	10.30	307	13.33	31.52	511	28.89	25.45
115	4.44	1.21	311	53.33	46.67	515	17.78	12.73
119	0.00	9.09	315	4.44	7.27	519	17.78	14.55
123	0.00	3.64	319	28.89	28.48	523	15.56	16.97
127	15.56	16.97	323	4.44	10.91	527	11.11	38.18
131	13.33	27.27	327	13.33	28.48	531	24.44	32.12
135	11.11	15.15	331	55.56	45.45	535	68.89	56.97
139	28.89	16.36	335	11.11	24.24	539	11.11	10.91
143	2.22	0.00	339	33.33	26.67	543	2.22	17.58
147	13.33	22.42	343	6.67	15.15	547	0.00	2.42
151	4.44	13.33	347	73.33	70.91	551	6.67	6.06
155	8.89	21.82	351	22.22	31.52	555	55.56	26.06
159	2.22	7.88	355	17.78	13.33	559	15.56	8.48
163	6.67	4.24	359	2.22	2.42	563	24.44	34.55
167	0.00	4.24	363	31.11	24.24	567	66.67	55.76
171	33.33	2.42	367	15.56	16.97	571	8.89	12.73
179	2.22	3.03	371	0.00	1.21	575	2.22	5.45
183	84.44	80.00	375	0.00	0.61	579	11.11	4.24
187	8.89	10.91	379	33.33	31.52	583	17.78	18.79
191	17.78	20.61	383	0.00	0.61	587	13.33	10.91
195	44.44	58.18	387	0.00	1.21	591	2.22	2.42
199	15.56	13.94	391	4.44	5.45	595	8.89	4.85
203	4.44	1.21	395	2.22	6.06	599	6.67	4.24
207	0.00	1.21	399	11.11	6.06	603	4.44	0.00
211	2.22	3.03	403	6.67	5.45	611	6.67	4.24
215	4.44	6.67	407	80.00	85.45	619	4.44	2.42
219	44.44	47.27	411	11.11	13.94	639	2.22	1.21
223	4.44	3.03	415	2.22	1.21	647	8.89	4.85
227	8.89	4.85	419	8.89	18.79	651	0.00	0.61
231	15.56	10.91	423	24.44	21.82	659	2.22	4.24
235	17.78	2.42	427	13.33	20.00	667	0.00	0.61
239	11.11	10.91	431	0.00	2.42	675	0.00	0.61
243	57.78	58.79	439	8.89	8.48	679	0.00	0.61
247	15.56	6.67	451	2.22	4.85	699	2.22	9.70
251	2.22	2.42	455	4.44	1.82	703	0.00	1.82
255	75.56	73.33	459	2.22	3.03	711	2.22	2.42
259	2.22	6.06	463	4.44	3.64	715	0.00	0.61
263	20.00	18.79	467	75.56	80.00	719	0.00	0.61
267	0.00	1.82	471	51.11	44.24	731	0.00	1.21
271	15.56	18.18	475	44.44	43.03	755	0.00	0.61
275	48.89	60.61	479	15.56	14.55	767	4.44	2.42
279	80.00	78.18	483	2.22	1.21	775	0.00	0.61
283	31.11	25.45	487	2.22	2.42	779	0.00	0.61
287	6.67	10.91	491	24.44	17.58	847	0.00	0.61
291	0.00	6.67	495	4.44	8.48			

592