Tectonic evolution of Variscan Iberia: Gondwana – Laurussia collision revisited

**Rubén Díez Fernández^{1,2*}, Ricardo Arenas¹, M. Francisco Pereira², Sonia Sánchez
Martínez¹, Richard Albert¹, Luis Miguel Martín Parra³, Francisco J. Rubio
Pascual³, Jerónimo Matas³**

¹*Departamento de Petrología y Geoquímica e Instituto de Geociencias (UCM, CSIC),
Universidad Complutense de Madrid. 28040 Madrid, Spain*

²*Instituto Dom Luiz, Departamento de Geociências, Escola de Ciências e Tecnologia,
Universidade de Évora, Apartado 94, 7001-554 Évora, Portugal*

³*Instituto Geológico y Minero de España, 28760 Tres Cantos, Madrid, Spain*

** Corresponding author at: Departamento de Petrología y Geoquímica e Instituto de
Geociencias (UCM, CSIC). Facultad de Geología, Universidad Complutense de
Madrid, C/ José Antonio Novais, no 2, 28040 Madrid, Spain. Tel.: +34 913944904; fax:
+34 915442535*

E-mail address: georuben@usal.es (R. Díez Fernández, corresponding author)

Keywords: European Geodynamics; Variscan Sutures; Allochthonous Complexes;
Iberian Massif; Paleozoic Tectonics; Pangea Amalgamation

Abstract

An integrated interpretation of the late Paleozoic structural and geochronological record of the Iberian Massif is presented and discussed under the perspective of a Gondwana-Laurussia collision giving way to the Variscan orogen. Compressional and extensional structures developed during the building of the Variscan orogenic crust of Iberia are linked together into major tectonic events operating at lithosphere scale. A review of the tectonometamorphic and magmatic evolution of the Iberian Massif reveals backs and forths in the overall convergence between Gondwana and Laurussia during the amalgamation of Pangea in late Paleozoic times. Stages dominated by lithosphere compression are characterized by subduction, both oceanic and continental, development of magmatic arcs, (over- and under-) thrusting of continental lithosphere, and folding. Variscan convergence resulted in the eventual transference of a large allochthonous set of peri-Gondwanan terranes, the Iberian Allochthon, onto the Gondwana mainland. The Iberian Allochthon bears the imprint of previous interaction between Gondwana and Laurussia, including their juxtaposition after the closure of the Rheic Ocean in Lower Devonian times. Stages governed by lithosphere extension are featured by the opening of two short-lived oceanic basins that dissected previous Variscan orogenic crust, first in the Lower-Middle Devonian, following the closure of the Rheic Ocean, and then in the early Carboniferous, following the emplacement of the peri-Gondwanan allochthon. An additional, major intra-orogenic extensional event in the early-middle Carboniferous dismembered the Iberian Allochthon into individual thrust stacks separated by extensional faults and domes. Lateral tectonics played an important role through the Variscan orogenesis, especially during the creation of new tectonic blocks separated by intracontinental strike-slip shear zones in the late stages of continental convergence.

49 **1. Introduction**

50 The Variscan orogen and its continuation into the Appalachian–Alleghanian
51 orogen have been the object of continuous rethinking and redefinition during the last
52 decades (Burg and Matte, 1978; Hatcher, 1978, 2002; Matte, 1986; Castro, 1987a;
53 Martínez Catalán et al., 1997; Franke, 2000; Ribeiro et al., 2007; Faure et al., 2008;
54 Ballèvre et al., 2009; Simancas et al., 2009; Kroner and Romer, 2013). These orogens
55 resulted from the late Paleozoic collision of Gondwana and Laurussia, and thus
56 represent a broad, axial suture zone right in the heart of Pangea (Bambach et al., 1980).

57

58 The Iberian Massif contains one of the most complete sections across the
59 Variscan orogen (Fig. 1), including several high-pressure metamorphic belts (Gil
60 Ibarguchi and Ortega Gironés, 1985; Mata and Munhá, 1986; Abalos et al., 1991b; De
61 Jong et al., 1991; Fonseca et al., 1993; Martínez Catalán et al., 1996; Rubio Pascual et
62 al., 2013b) and tectonic units with ophiolitic assemblages that separate tectonic slices of
63 continental crust (Arenas et al., 1986, 2007b; Crespo-Blanc and Orozco, 1988; Fonseca
64 and Ribeiro, 1993; Díaz García et al., 1999; Pedro et al., 2005; Pin et al., 2006; Sánchez
65 Martínez et al., 2009, 2012; Merinero et al., 2013, 2014). In the crystalline basement of
66 central and southern Europe, such combination has been classically considered as
67 indicative of multiple oceanic suture zones of Variscan age (Franke, 2000, 2006, 2014;
68 Tait et al., 2000; Matte, 2001; Von Raumer et al., 2003; Martínez Catalán et al., 2007;
69 Simancas et al., 2009).

70

71 Until recent times, cross-sections made to characterize the structure of the whole
72 orogen were strongly influenced by the recognition of strike-slip systems separating

73 tectonostratigraphic domains with opposite structural vergence (Burg et al., 1981;
74 Matte, 1986, 2001; Simancas et al., 2013). Taking that, together with the varied
75 geographic location of ophiolitic units, geodynamic models have put forward the
76 hypothesis of a collage of individualized continental micro-terrane variably dispersed
77 between Gondwana and Laurussia prior to the collision of the two latter (e.g.,
78 Winchester et al., 2002). However, new geochronological data show that the ophiolitic
79 ensembles consist of different age rocks which in some cases indicate opening of
80 ephemeral oceanic basins as Variscan orogenic crust was built (Azor et al., 2008;
81 Arenas et al., 2014b), thus casting doubt on the multi-terrane picture as a general pre-
82 orogenic feature. To this figure, most of the strike-slip shear zones that have been
83 interpreted as separating supposedly different terranes rather correspond to late
84 structures in the evolution of the Variscan orogen (Martínez Catalán, 2011; Gutiérrez-
85 Alonso et al., 2015). The main strike-slip systems postdate a well-established phase
86 dominated by tangential tectonics that includes development of continental subduction
87 systems, obduction of suture zones, and emplacement of large thrust nappes (Iglesias
88 Ponce de Leon and Choukroune, 1980; Gates et al., 1986; Martínez Catalán, 1990;
89 Hatcher, 2002).

90
91 In the Iberian Massif, the most influential strike-slip system in defining the
92 structure of the Variscan orogen has been the Coimbra-Córdoba shear zone (Fig. 1)
93 (Burg et al., 1981; Ribeiro et al., 2007). The stretching lineation associated with this
94 sinistral shear zone is subhorizontal (Pereira et al., 2008a, 2010a, 2010b), so movements
95 along this structure cannot explain alone the juxtaposition of eclogites, high-P
96 granulites, and blueschists to low-grade and low-pressure rocks unless tangential
97 tectonics or vertical extrusion existed prior to or alternating with lateral movements.

Yet, the vergence of major orogenic structures seems to change at both sides of the Coimbra-Córdoba shear zone, being to the E and NE in the block located north (Pérez-Estaún et al., 1991; Azor et al., 1994a) and to the SW in the block located south (Simancas et al., 2001). However, a careful revision of the whole tectonic evolution of each block prevents us from accepting such consensual assumption. For instance, top-to-the-E and -NE kinematics associated with tangential deformation has been also described in major structures of the block located south of the Coimbra-Córdoba shear zone (Araújo et al., 2005; Pereira et al., 2007; Rosas et al., 2008; Borrego, 2009; Ribeiro et al., 2010; Ponce et al., 2012), plus the age of structures showing opposite vergence across the Iberian Massif is different in most of the cases (see following sections).

For many years the tectonic evolution of the Iberian Massif has been presented and evaluated in a similar way to a double-blind procedure. Advances coming from its northwestern section were little considered in its southwestern part and the other way around. In the meantime, the amount of structural and geochronological data in both regions have increased noticeably, to a point that a structural correlation between these two sections of the Variscan orogen would allow a more precise integration of geological processes previously described elsewhere in the Iberian Massif.

Here we follow on to the discovery that the Iberian Massif contains allochthonous and autochthonous tectonic units that share fundamental features of their tectonostratigraphy and can be structurally correlated across the two blocks separated by the Coimbra-Córdoba shear zone, i.e. the pile of allochthonous units recognized in NW Iberia can be correlated with that of SW Iberia (Fig. 2; Díez Fernández and Arenas,

2015). Within this framework, this article presents an integrated model showing the Variscan evolution of the Iberian Massif by piecing together the structural history and chronology of major individual events registered across different sections of its orogenic crust.

2. Synthetic cross-section and tectonostratigraphy of the Iberian Massif

The Iberian Massif combines folds and thrust faults formed during Variscan compression with open structural domes and basins, granitization and normal faults that are a product of subsequent gravitational collapse and intra-orogenic extension (Simancas et al., 2001; Martínez Catalán et al., 2007; Pereira et al., 2009). Figure 3 shows a simplified, composite cross-section of the Iberian Massif highlighting eight major tectonostratigraphic elements of the Variscan orogen (Díez Fernández and Arenas, 2015), namely: (1) Cantabrian Zone foreland, (2) Iberian Autochthon, (3) Iberian Parautochthon, (4) Basal Allochthonous Units, (5) Allochthonous Ophiolitic Units, (6) Upper Allochthonous Units, (7) Beja-Acebuches Ophiolite, and (8) South Portuguese Zone. Gathering of 4, 5, and 6 will be also referred to as Iberian Allochthon (or simply Allochthon).

A simple restoration of the cross-section shown in Figure 3 (Fig. 4a) reveals two major suture zones featured with ophiolitic units with different protolith ages. The NNE-dipping suture represented by the Beja-Acebuches Ophiolite (Bard and Moine, 1979; Munhá et al., 1986; Quesada et al., 1994) cuts the suture marked by the Allochthonous Ophiolitic Units (Arenas et al., 1986) and divides the Iberian Massif in two major blocks. The northern block comprises the Cantabrian Zone, and the Iberian Autochthon, Parautochthon and Allochthon, all of which have Gondwanan affinity

(Fernández-Suárez et al., 2003; Robardet, 2003; Martínez Catalán et al., 2004; Díez Fernández et al., 2010, 2012b; Pastor-Galán et al., 2013; Albert et al., 2015a; Pereira, 2015). In the southern block, the South Portuguese Zone is considered a composite terrane located along the southern flank of Laurussia (Avalonia-Meguma) by the Middle Devonian (Lefort et al., 1988; Matte, 1991; Quesada et al., 1994; Simancas et al., 2005; Braid et al., 2011), but close to or juxtaposed with Gondwana since (at least) the Upper Devonian (Pereira et al., 2006b).

The Iberian Massif consists of sedimentary, plutonic and metamorphic rocks whose grade ranges from very low to catazonal. The Gondwanan series consist of Precambrian and Paleozoic marine sequences alternating with volcanic and plutonic rocks of variable age and abundance. These series are deformed into a thin-skinned fold and thrust belt with easterly propagation in the Cantabrian Zone (Pérez-Estaún et al., 1988), and into a crystalline thrust sheet characterized by large recumbent folds (e.g., Mondoñedo Nappe). This tectonic stacking was followed by the development of extensional faults and doming (Lugo dome) in the more external zones of the Gondwanan Variscan hinterland (West Asturian-Leonese Zone; Martínez Catalán et al., 2003). Towards more internal zones of the Variscan orogen, deformation affecting the Iberian Autochthon produces upright, overturned, and recumbent folds (Díez Balda, 1986; Macaya et al., 1991; Díaz Azpiroz et al., 2003; Dias et al., 2010; Díez Fernández et al., 2013b), which are underlain by a complex system of granite- and migmatite-cored dome structures (Escuder Viruete et al., 1994; Díez Balda et al., 1995; González del Tánago, 1995; Pereira et al., 2009; Díaz-Alvarado et al., 2012; Díez Fernández et al., 2012c; Arango et al., 2013; Rubio Pascual et al., 2013a).

A significant part of the Iberian Autochthon occurs under a huge thrust stack (Ribeiro et al., 1964; Ries and Shackleton, 1971). The lowermost set of thrust nappes is the Iberian Parautochthon (Farias et al., 1987; Ribeiro et al., 1990), which is restricted to the sections of the Variscan orogen that were closer to the Gondwana mainland and contains alternating volcanic and sedimentary rocks with remarkable lateral variability at regional scale (Dias da Silva et al., 2014). The lower contact of the overlying Basal Allochthonous Units traces the base of a large allochthonous ensemble, the Iberian Allochthon, transported onto the Iberian Autochthon and Parautochthon. Remnants of the Iberian Allochthon are found as klippen in the complexes of Cabo Ortegal, Órdenes, Malpica-Tui, Bragança and Morais in NW Iberia (Martínez Catalán et al., 2007), whereas in the Ossa-Morena Complex of SW Iberia, the base of the allochthonous ensemble is defined by high-P metamorphic “belts” (with equivalent protolith and metamorphic ages and lithological composition) that crop out at similar structural position across the Iberian Variscides (Díez Fernández and Arenas, 2015, 2016). The Basal and Upper Allochthonous Units are separated by a set of dismembered Allochthonous Ophiolitic Units.

The Basal and Upper Allochthonous Units comprise arc-related Precambrian and Cambrian sedimentary rock sequences intruded by or alternating with arc-related and later alkaline to peralkaline Cambrian and Ordovician igneous rocks. Most of their series, therefore, bear the imprint of Ediacaran and Early Paleozoic subduction and subsequent Cambro-Ordovician rifting (Arenas et al., 1986, 2009; Ribeiro and Floor, 1987; Liñán and Quesada, 1990; Quesada, 1990; Sánchez-García et al., 2003; Pereira et al., 2006a; Díez Fernández et al., 2010, 2015; Andonaegui et al., 2012). These sequences are succeeded by passive margin strata that covers up to the lowermost

Devonian in the Upper Allochthonous Units of the Ossa-Morena Complex (Robardet and Gutiérrez Marco, 2004), but it is lacking in the rest of the allochthonous complexes.

The Allochthonous Ophiolitic Units are made of mafic and ultramafic rocks and scarce metasedimentary rocks (see compilation by Arenas and Sánchez Martínez, 2015). These units can be divided in two groups according to protolith ages and chemical signature (Arenas et al., 2007a; Sánchez Martínez et al., 2009): a Cambrian-Ordovician group related to the closure of the Iapetus/Tornquist Ocean (Bazar ophiolite; Sánchez Martínez et al., 2012) and opening of the Rheic Ocean s.l. (Vila de Cruces, Izeda-Remondes, and Internal Ossa-Morena Zone ophiolites; Pin et al., 2006; Arenas et al., 2007b; Pedro et al., 2010), and an Lower-Middle Devonian group (Careón, Purrido, Moeche, and Morais-Talhinhas ophiolites; Díaz García et al., 1999; Pin et al., 2006; Sánchez Martínez et al., 2011) representing an ephemeral oceanic basin formed in early stages of the Variscan orogenesis (Arenas et al., 2014b). When these two groups contact with each other the Devonian counterparts occur on top.

The Beja-Acebuches Ophiolite is a rather continuous band of amphibolites, minor ultramafic rocks, mylonitic gabbros and a sheeted dike complex (Bard, 1977; Bard and Moine, 1979; Fonseca and Ribeiro, 1993; Quesada et al., 1994) that have been interpreted as a relict oceanic crust formed in a rifting context s.l. (Dupuy et al., 1979), either in a back-arc or intra-arc setting (Quesada et al., 1994), or in a mid-ocean ridge (Castro et al., 1996). Protoliths of this oceanic crust have been dated at Viséan, so it formed long after the onset of Variscan deformation (Azor et al., 2008).

The northern domains of the South Portuguese Zone contain an ensemble of metasedimentary rocks and lenses of metabasites (Pulo do Lobo Unit; Carvalho et al., 1976; Oliveira, 1990; Fonseca, 2005). The Pulo do Lobo Unit has been considered as an accretionary prism related to a north-dipping subduction zone (Eden and Andrews, 1990; Silva et al., 1990; Braid et al., 2010) or as a dismembered ophiolitic ensemble s.l. (Fonseca and Ribeiro, 1993) connected to the closure of the Beja-Acebuches oceanic basin (Quesada et al., 1994). Alternatively, other authors suggest it may represent an accretionary prism over a south-dipping subduction zone developed prior to the opening of the Beja-Acebuches oceanic basin (Azor et al., 2008). The general structure of this domain is an upright antiform (Silva et al., 1990; Martínez Poza et al., 2012; Pérez-Cáceres et al., 2015) and the age of its series could be as old as Silurian in the lower part of the stratigraphy (Braid et al., 2011), and have been dated at Middle-Upper Devonian in the intermediate part (Pereira et al., 2008b), and early Carboniferous in the overlying, discordant series (Santa Iria basin, Pereira et al., 2008b; Braid et al., 2011). The Iberian Pyrite Belt is located to the south of the Pulo do Lobo Unit and contains Upper Devonian sedimentary rocks and Upper Devonian to early Carboniferous volcanic strata covered by younger Carboniferous turbiditic series (Oliveira, 1990; Leistel et al., 1997; Pereira et al., 2012a). Its regional structure is defined by a southerly propagation of a fold and thrust belt (Silva et al., 1990).

3. Record of Variscan events

Table 1 is a synopsis that presents a summary of the main tectonic events that characterize the principal geotectonic zones of the Iberian Massif. It offers a general view of the time-based correlation of contrasted geological processes presented in this work (descriptions and citations in the text below). For comparison with isotopic ages,

ages obtained from fossil record have been converted into absolute ages following the IUGS International Chronostratigraphic Chart 2013 (Cohen et al., 2013).

3.1. Lower Devonian (Lochkovian-Emsian)

3.1.1. Upper Allochthonous Units of NW Iberia

The first phase of Variscan deformation in NW Iberia corresponds to a high-P/high-T tectonometamorphic event that is traceable across the lowermost structural levels of the Upper Allochthonous Units of Cabo Ortegal (Vogel, 1967; Gil Ibarguchi et al., 1990; Puelles et al., 2005; Albert et al., 2012), Órdenes (Arenas and Martínez Catalán, 2002; Gómez Barreiro, 2007), Bragança and Morais complexes (Marques et al., 1996). Altogether these structural levels define a high-P/high-T metamorphic belt characterized by variably-retrogressed eclogitic foliation formed in the course of dextral, west-directed continental subduction (Martínez Catalán et al., 1997; Ábalos et al., 2003). The high-P rocks occur below a thick series of siliciclastic rocks affected by Variscan intermediate-P metamorphism (Castiñeiras, 2005). Yet, the uppermost structural levels of this series still preserve the imprint of a previous Cambrian deformation (Díaz García et al., 2010). The high-P/high-T metamorphism has been dated at ca. 410-390 Ma (Santos Zalduegui et al., 1996; Ordóñez Casado et al., 2001; Fernández-Suárez et al., 2002, 2007). According to provenance analysis of their sedimentary sequences, both the upper and lower plate to this continental subduction zone represents a piece of Gondwanan continental crust located in the periphery of the West African Craton (Fernández-Suárez et al., 2003; Albert et al., 2015a, 2015b). This metamorphic belt occurs together with a set of ultramafic rocks considered as one of the world-class examples of heterogeneous upper mantle (Girardeau et al., 1989; Girardeau and Ibarguchi, 1991).

3.1.2. Upper Allochthonous Units of SW Iberia

The Precambrian to Lower Devonian series of the Upper Allochthonous Units of the Ossa-Morena Complex are deformed into a SW-verging train of recumbent folds (e.g., Monesterio Anticline) with associated axial planar foliation developed under Barrovian, low-T conditions (Vauchez, 1974, 1976; Chacón, 1979; Chacón et al., 1983; Apalategui et al., 1990; Quesada, 1990; Expósito Ramos, 2005). These folds are cut by south-directed thrusts (e.g., Monesterio Thrust; Eguíluz, 1987) and are covered by a syn-orogenic basin (Terena flysch), whose base rests discordant over the folds (Expósito et al., 2002). Some authors consider this flysch yet coeval with the SW-verging folds (Rocha et al., 2009; Araújo et al., 2013).

The base of the Terena flysch has been dated at Lower Devonian (Lochkovian; Piçarra, 1997; Pereira et al., 1998, 1999; Piçarra et al., 1998; Robardet and Gutiérrez Marco, 2004; Rocha et al., 2010). The youngest series that is bent into the recumbent folds contains intraformational conglomerates indicating basin instability and syn-orogenic deformation (Giese et al., 1994). Limestones in that series have been dated with fossils at Lochkovian-Emsian (Perdigão et al., 1982; Oliveira et al., 1991, 1992; Robardet et al., 1998), so the age of the SW-verging folds can be further constrained between ca. 419 Ma and ca. 393 Ma, and deformation preceding or coeval with fold nucleation probably started in the Lochkovian. Other age constraints exist for fold formation, such as K/Ar ages obtained from biotite associated with the axial planar foliation (385±11 Ma; Galindo et al., 1986, 1987) and the rejuvenation of Precambrian (protolith?) ages in amphibolites and metasedimentary rocks at ca. 400-390 Ma (Dallmeyer and Quesada, 1992). The later thrusts are covered by lowermost

Carboniferous sedimentary rocks, so these south-directed faults have been considered part of a continuous deformational process following fold amplification (Expósito et al., 2002).

3.2. Lower-Middle Devonian (Emsian-Eifelian-Givetian)

3.2.1. Upper Allochthonous Units of SW Iberia

Located to the north and near the Pulo do Lobo Unit, there are few outcrops of Emsian-Eifelian reef limestones with interbedded mafic calc-alkaline volcanic rocks that may represent a magmatic arc associated with the closure of the Rheic Ocean (Silva et al., 2011). Middle Devonian sedimentary rocks are very scarce or absent in the southern part of the Upper Allochthonous Units of SW Iberia, i.e. exposures south of the Coimbra-Córdoba Shear Zone (Robardet and Gutiérrez Marco, 2004).

Conversely, Givetian sedimentary rocks might exist in the block located north of the Coimbra-Córdoba shear zone, in the so-called Obejo-Valsequillo Domain (Sánchez Cela and Gabaldón, 1977a; Pérez Lorente, 1979; Rodríguez and Soto, 1979; Pardo and García Alcalde, 1996; García-López et al., 1999). A recent revision of the stratigraphy of the Obejo-Valsequillo Domain has not confirmed the occurrence of Middle Devonian sedimentary rocks (Matas et al., 2015a). Therefore, either the absence or scarcity of sediments of that age seems a common feature among the Upper Allochthonous Units of SW Iberia. The widespread Middle Devonian sedimentary gap has been explained by significant crustal uplift during Devonian tectonic activity (Giese et al., 1994).

3.2.2. Allochthonous Ophiolitic Units of NW Iberia

The protoliths of the Devonian Allochthonous Ophiolitic Units account for a short period of lithosphere extension and ocean-floor generation (Díaz García et al., 1999). Preliminary geochemical data led to place such extension in an intra-oceanic subduction setting (Sánchez Martínez et al., 2007), or as connected to incipient collisional processes (Pin et al., 2002). The isotopic signature of zircon in these ophiolites favors a setting that involves continental crust in their formation (Sánchez Martínez et al., 2011). A revision of all the geochemical, isotopic, and regional data available for these ophiolites in Iberia, considered a short-lived oceanic basin opened within a continental realm at ca. 400-395 Ma as their most likely setting (Arenas et al., 2014b). In any case, lithosphere extension occurred in a domain located between the paleogeographic realms of the Upper and Basal Allochthonous Units, both of which represent sections of Gondwana.

3.2.3. Iberian Autochthon and Cantabrian Zone

A series of Lower-Middle Devonian events distributed across the Iberian Massif have been integrated into an extensional setting affecting a large tract of the Gondwana margin at ca. 395 Ma (Gutiérrez-Alonso et al., 2008). These include alkaline volcanism in the Iberian Autochthon (Gutiérrez-Alonso et al., 2008) and Cantabrian Zone (Loeschke, 1983), increased subsidence in the Cantabrian Zone (Veselovski, 2004), and a sedimentary gap in the Iberian Autochthon (Puschmann, 1967).

3.3. Upper Devonian (Frasnian-Fammenian)

3.3.1. Upper Allochthonous Units of NW Iberia

The high-P/high-T record of the Upper Allochthonous Units is variably retrogressed as a consequence of a multi-stage Upper Devonian exhumation process

(Gómez Barreiro et al., 2007). Post-eclogitic contractional shearing was developed under amphibolite facies conditions and produced a widespread mylonitic foliation (Castiñeiras, 2005; Gómez Barreiro, 2007). The high-P/high-T units are separated from overlying intermediate-P units by a set of extensional detachments (Martínez Catalán et al., 2002; Castiñeiras, 2005), dated at ca. 375-371 Ma (Dallmeyer et al., 1997; Gómez Barreiro et al., 2006). Both, the regional mylonitic foliation and the extensional detachments, are affected by east-verging recumbent folds at a regional scale, both in the Órdenes Complex (Martínez Catalán et al., 2002; Gómez Barreiro, 2007; González Cuadra, 2007) and in the Cabo Ortegal Complex (Marcos et al., 1984; Ábalos et al., 2003; Albert et al., 2012). Therefore the nucleation of these recumbent folds occurred later than ca. 371 Ma. Besides the age of intervening extensional structures, the exhumation of the high-P and high-T units to amphibolite facies conditions has been estimated at ca. 380 Ma (Van Calsteren et al., 1979; Peucat et al., 1990; Dallmeyer et al., 1991; Santos Zalduegui et al., 1996; Valverde Vaquero and Fernández, 1996; Gómez Barreiro et al., 2006), whereas subsequent retrogression to greenschist facies conditions is dated at ca. 360-350 Ma (Peucat et al., 1990; Dallmeyer et al., 1997).

3.3.2. Upper Allochthonous Units of SW Iberia

The southern part of the metasedimentary series exposed in the Obejo-Valsequillo Domain are located in the hanging wall of a NE-directed thrust (Espiel thrust) and are deformed into a NE-verging train of recumbent folds (Martínez Poyatos et al., 2001; Martínez Poyatos, 2002). The youngest rocks affected by these folds are Devonian. Some authors propose the existence of Middle Devonian series (Givetian; Sánchez Cela and Gabaldón, 1977a; Pérez Lorente, 1979; Rodríguez and Soto, 1979; Pardo and García Alcalde, 1996), whereas other authors have also found Upper

Devonian strata (Frasnian-Famennian; Febrel and Saenz de Santa María, 1964; Herranz, 1985; Racheboeuf et al., 1986). On the contrary, Matas et al. (2015a) restrict the Upper Devonian series to the footwall of the Espiel thrust, and identify a stratigraphic hiatus spanning from the Middle to the Upper Devonian in the series affected by the NE-verging folds. These folds are covered by a discordant Carboniferous succession (Culm facies), whose base is dated at Tournaisian (ca. 359-347 Ma; Sánchez Cela and Gabaldón, 1977b; Garrote and Broutin, 1979; García Alcalde et al., 1984; Rodríguez et al., 1990). Amphibolites occurring in the Precambrian series and affected by these folds show rejuvenation of a Precambrian metamorphic imprint mainly during the Upper Devonian (Dallmeyer and Quesada, 1992). Consequently, the age of recumbent folding is mostly Upper Devonian-early Carboniferous (ca. 383-347 Ma).

3.3.3. Allochthonous Ophiolitic Units

The age of imbrication of the Cambrian-Ordovician and Devonian ophiolitic ensemble exposed in NW Iberia has been constrained, by means of $^{40}\text{Ar}/^{39}\text{Ar}$ dating of their low- to medium-T metamorphic fabrics (greenschist and amphibolite facies), to a range that extends from ca. 391 Ma to ca. 364 Ma (Peucat et al., 1990; Dallmeyer et al., 1997). The metamorphic grade decreases progressively down structure, as the ages of metamorphism become younger (Arenas et al., 2007a). Kinematic indicators for accretion consistently indicate top-to-the-Cantabrian foreland, i.e. subduction under the Upper Allochthonous Units (Arenas et al., 2007b; Gómez Barreiro et al., 2010b).

3.3.4. Basal Allochthonous Units of NW Iberia

The first deformation event recorded in the Basal Allochthonous Units relates to continental subduction (Gil Ibarguchi and Ortega Gironés, 1985; Martínez Catalán et

al., 1996). This tectonic event developed a penetrative fabric whose remnants occur in weakly or non-retrogressed lenses of high-P/low-intermediate-T rocks and as mineral trails within porphyroblasts grown during decompression (Díez Fernández and Martínez Catalán, 2012). Kinematic criteria associated with fabrics developed under high-P/low-intermediate-T metamorphic conditions show a consistent top-to-the-northeast shear-sense, which indicates dextral oblique subduction to the west in present-day coordinates (Díez Fernández et al., 2012a). Deformation took place under metamorphic conditions ranging from blueschist to eclogite facies (Munhá et al., 1984; Arenas et al., 1995, 1997; Gil Ibarguchi, 1995; Rubio Pascual et al., 2002; Rodríguez et al., 2003; López-Carmona et al., 2010, 2014), and has been consistently dated at ca. 380-370 Ma (Van Calsteren et al., 1979; Santos Zalduegui et al., 1995; Rodríguez et al., 2003; Abati et al., 2010).

Subsequent exhumation of these units was controlled by crustal-scale ductile thrusting (Fervenza thrust), which was accompanied by recumbent folding and attenuation of overlying lithosphere (tectonic and erosional). Early exhumation produced a mylonitic foliation throughout the high-P metamorphic belt. The kinematics of this event is consistent with east-directed tectonic transport (Díez Fernández et al., 2011). Deformation took place under high-P conditions, although in a clear decompressive path (Rodríguez et al., 2003; Díez Fernández et al., 2011). Dating of tectonic fabrics and partial melting formed in the course of subsequent exhumation under lower pressure conditions (amphibolite/greenschist facies) has yielded ages in the range ca. 360-346 Ma (Abati and Dunning, 2002; Rodríguez et al., 2003; López-Carmona et al., 2014). Therefore, the initial exhumation from peak-pressure conditions up to the lower crust occurred in the Famennian (ca. 370-360 Ma).

3.3.5. Basal Allochthonous Units of SW Iberia

In SW Iberia, the Basal Allochthonous Units crop out in two domains (Fig. 2). A northern domain made of continental crust, referred to as the Central Unit (Azor et al., 1994b), experienced high-P/low-to high-T metamorphism (Mata and Munhá, 1986; Eguiluz et al., 1990; Abalos et al., 1991b; Pereira and Apraiz, 2006; Pereira et al., 2010a). Most of this record is strongly retrogressed into amphibolite/greenschist facies rocks, within which only small retrogressed high-P granulite and eclogite pods and mineral relicts testifies for their subduction-related history. As a consequence, no kinematic criteria have been provided so far for the development of the high-P fabrics. This way, the current NE-dipping character of the post-eclogitic foliation has been taken as the sole criteria for a NE subduction polarity (Azor et al., 1994b).

The individual ages obtained for the high-P metamorphism in the Central Unit show some variation but complementary results. First attempts to date this deformation yielded very imprecise Silurian-Devonian ages that called for a Variscan event (427 ± 45 Ma; Schäfer et al., 1991). Subsequent surveys performed in eclogite boudins provided a minimum age of ca. 370-360 Ma for the high-P metamorphism in the Central Unit (Quesada and Dallmeyer, 1994). Finally, a more specific, yet imprecise age of ca. 380-350 Ma was obtained for this metamorphic event by means of U-Pb zircon dating (Ordóñez Casado, 1998). On the other hand, dating of mylonites formed during post-peak-P metamorphism yielded ages of ca. 355 Ma (García Casquero et al., 1988). If we consider all these data, the timing of continental subduction experienced by these rocks should be restricted to the Upper Devonian, probably ranging between ca. 380-360 Ma.

The Basal Allochthonous Units that are located in the southernmost domain of the Ossa-Morena Complex were recently referred to as Cubito-Moura Unit, which, similarly to the rest of Basal Allochthonous Units, consists of two juxtaposed sequences (Díez Fernández and Arenas, 2015). The upper sequence of this unit is known under the name of Cubito-Moura Schists (Fonseca et al., 1999). The lower sequence includes a series of metasedimentary rocks and orthogneisses referred to as Fuenteheridos group (Rubio Pascual et al., 2013b), or as the Serie Negra succession and Igneous-felsic dominated-sedimentary complex (only sections affected by high-P metamorphism; Chichorro, 2006; Chichorro et al., 2008; Rosas et al., 2008). Altogether these sequences represent a piece of continental crust subjected to high-P/low-intermediate-T metamorphism, as typified by evidence (blueschist and eclogite boudins and weakly-retrogressed mineral assemblages) of a first tectonic event (De Jong et al., 1991; Fonseca et al., 1993, 1999, 2004; Pedro, 1996; Leal et al., 1997; Moita, 1997; Booth-Rea et al., 2006; Ponce et al., 2012; Rubio Pascual et al., 2013b).

Strong retrogression affected the high-P rocks of the Cubito-Moura Unit during exhumation and no kinematic criteria extracted from direct observation of high-P fabrics have been reported so far for the subduction process. Yet, both NE and SW subduction polarities have been proposed (Fonseca et al., 1999; Díaz Azpiroz et al., 2004; Ribeiro et al., 2007; Rosas et al., 2008; Pin et al., 2008; Simancas et al., 2009; Rubio Pascual et al., 2013b). The age of high-P metamorphism obtained for the lower sequence of this unit is ca. 371 Ma (Moita et al., 2005), which is in agreement with an age of ca. 358 Ma obtained for post-peak-P metamorphism (Rosas et al., 2008).

Preservation of the structural record associated with the initial stages of exhumation is very rare in the Cubito-Moura Unit. One example corresponds to the development of boudinaged-folds affecting marbles and amphibolites (Rosas et al., 2002) under a top-to-the-SSW shear sense regime (Rosas et al., 2008). This particular shearing has been inferred older than a cooling age of ca. 358 Ma (Rosas et al., 2008), and consequently it should have been developed during the Famennian (ca. 371-358 Ma).

3.4. Early Carboniferous (Tournaisian-Viséan)

3.4.1. Upper Allochthonous and Ophiolitic Units of NW Iberia

A set of out-of-sequence-thrusts carried the Upper Allochthonous and Ophiolitic Units of NW Iberia approximately to their current position in the thrust pile (see extended description in Martínez Catalán et al., 2002). Out-of-sequence thrusting proceeded under low-T conditions (greenschist facies) and generated mylonites and ultramylonites with an associated top-to-the-southeast sense of motion. This thrust system moved the Upper Allochthonous Units over the Ophiolitic Allochthonous Units and affected thrust sheets of the previously (under)stacked ophiolites (Díaz García et al., 1999; Arenas et al., 2007b). The emplacement of the Upper Allochthonous Units generated duplexes with both hinterland-dipping and foreland-dipping horses, duplexes of antiformal stack type, as well as isolated horses made of both the Upper Allochthonous and Ophiolitic Allochthonous Units.

The out-of-sequence thrusts cut and partially rework extensional shear zones dated at ca. 375-371 Ma. They also cut the regional fabrics of the Upper Allochthonous Units associated with their retrogression to greenschist facies conditions dated at ca.

360-350 Ma. They might also affect some of the uppermost Basal Allochthonous Units, such as the Agualada Unit. The age of post-peak-P partial melting in that unit is ca. 346 Ma (Abati and Dunning, 2002), so the out-of-sequence thrusting is likely younger. This thrust system also cuts the Lalín-Forcarei Thrust (age estimated at ca. 340 Ma; Dallmeyer et al., 1997), which is structurally below and overprinted by an event of amphibolite facies metamorphism that is absent in the out-of-sequence thrusts. The out-of-sequence thrusts must be older than the extensional detachments of Pico Sacro and Bembibre-Ceán (Díez Fernández et al., 2012c), which cut all previous thrust faults and are bracketed between ca. 323-314 Ma (Martínez Catalán et al., 2002) and dated at ca. 337 Ma (López-Carmona et al., 2014), respectively. Fine-grained white micas extracted from a greenschist facies ultramylonitic gneiss located in an out-of-sequence thrust contact gave an $^{40}\text{Ar}/^{39}\text{Ar}$ cooling age of ca. 325 Ma. Considering all the previous data the most probable age for the major event of out-of-sequence thrusting is Viséan (ca. 346-337 Ma). However, a thrust-related phyllonite located at the base of the Cabo Ortegal Complex yielded an age of ca. 316 Ma (Dallmeyer et al., 1997), what certainly opens the possibility of later episodes of out-of-sequence thrusting in NW Iberia (Martínez Catalán et al., 2002).

3.4.2. Upper Allochthonous Units of SW Iberia

One of the most salient structures affecting the Iberian Allochthon exposed in the Obejo-Valsequillo Domain is a thrust fault that transports to the NE a NE-verging train of recumbent folds of Upper Devonian age (see section 3.3.2) onto Culm facies sedimentary rocks, the Espiel Thrust (Figs. 2 and 3; Martínez Poyatos et al., 2001; Martínez Poyatos, 2002). Some of the youngest series of its footwall may be upper Viséan or even early Serpukhovian (~340-330 Ma; Ortuño, 1971; Sánchez Cela and

Gabaldón, 1977b; Garrote and Broutin, 1979; Matas et al., 2015a), what places a reference age for the onset of thrusting. On the other hand, a regional analysis carried out in the syn-orogenic series, that occur both in the hanging wall and footwall of this fault, revealed that thrusting may have been active from the Viséan up to the upper Bashkirian (~340-315 Ma; Martínez Poyatos et al., 1998; Matas et al., 2014).

In the first description of the Espiel Thrust, Apalategui and Pérez-Lorente (1983) documented the existence of peridotites, amphibolites, mylonitic gneisses and phyllonites in close relation to some sections of its fault zone. These authors also suggested a correlation between those gneisses and the lithological ensemble that today constitutes the Basal Allochthonous Units bounding the Obejo-Valsequillo Domain to the south (the so-called Central Unit). The presence of peridotites (even if scarce) together with gneisses that have been considered as markers of a suture zone elsewhere, represent a singularity within the Upper Allochthonous Units of SW Iberia. Such singularity can be solved by considering the Espiel Thrust as a series of out-of-sequence structures that cut, bounded and transported upwards an underlying and previously structured suture zone consisting of high-P gneisses and ophiolitic rocks (peridotites and amphibolites).

NE of the Espiel Thrust, a group of reverse faults with top-to-the-east tectonic transport makes a tectonic imbricate and duplexes of continental crust within the Upper Allochthonous Units (Zalamea de la Serena imbricates; Castro, 1987b). Due to their similar geometry and kinematics, these faults have been considered as genetically related to the main thrust system that transported the NE-verging recumbent folds (Martínez Poyatos et al., 2001).

3.4.3. Basal Allochthonous Units of NW Iberia

The mylonitic shear zones developed during the Famennian (Fervenza Thrust; see section 3.3.4), together with the rest of the lithostratigraphy of the Basal Allochthonous Units, are affected by a regional-scale train of recumbent folds (Díez Fernández et al., 2011). The regional foliation in the Basal Allochthonous Units is axial planar to these folds (Díez Fernández, 2011) and developed under greenschist to amphibolite facies conditions (Rodríguez et al., 2003). The asymmetry depicted by this train and kinematic criteria observed in the axial planar foliation indicates top-to-the-Cantabrian Zone sense of shear (Díez Fernández and Martínez Catalán, 2012), which is in agreement with quartz crystal preferred orientation fabrics (Llana-Fúnez, 2002; Gómez Barreiro et al., 2010a; Fernández et al., 2011). Dating of this regional foliation has yielded Tournaisian ages in the range ca. 360-350 Ma (Rodríguez et al., 2003).

To the external parts of the hinterland, the regional foliation of the Basal Allochthonous Units is bent into a large recumbent anticline and minor associated folds (Carrio Anticline). The Lalín-Forcarei Thrust runs along the reverse limb of that major anticline, and produced ductile mylonitic deformation throughout the base of the Iberian Allochthon. Kinematic criteria show an eastward movement for this thrust, which accounts for the emplacement of the Iberian Allochthon onto the Iberian Parautochthon (Martínez Catalán et al., 1996). Isotopic ages for this nappe-fold structure are lacking, although fold nucleation must be younger than ca. 360-350 Ma (age of the folded foliation at a regional scale). A reference age for the Lalín-Forcarei Thrust is ca. 340 Ma, which is the timing of pervasive ductile deformation recorded by the Iberian

Parautochthon in response to the overriding of the Iberian Allochthon s.l. (Dallmeyer et al., 1997).

The uppermost part of the Basal Allochthonous Units of NW Iberia experienced high-T conditions and partial melting during post-eclogitic decompression (Arenas et al., 1997). Post-peak-P partial melting has been dated at ca. 346-341 Ma (Abati and Dunning, 2002).

3.4.4. Basal Allochthonous Units of SW Iberia

Ductile shearing associated with the exhumation of the Basal Allochthonous Units exposed in the Central Unit generated a penetrative foliation under amphibolite to greenschist facies conditions (Abalos et al., 1991a, 1991b), reaching the granulite facies in some sections (Pereira et al., 2010a). Although no major structures have been recognized so far within this unit, recumbent folding has been considered as a process related to the formation of the regional planar fabric (Azor, 1994). The kinematics shown by this fabric is top-to-the-NW, an orogen-parallel trend that may indicate a strong lateral shear component during and/or after exhumation (Azor et al., 1994b; Pereira et al., 2008a, 2010a).

The timing of exhumation show some variation in the Central Unit, but a Tournaisian-Viséan age can be taken as the most likely, if the whole set of geochronological data available is taken in consideration. Dating of biotite in gneisses yielded ages of ca. 335-330 Ma (Blatrix and Burg, 1981; Pereira et al., 2012b), in agreement with subsequent results obtained in muscovite (ca. 340-333 Ma; Dallmeyer and Quesada, 1992; Pereira et al., 2012b). The maximum age for the early (and warmer)

stages of exhumation can be vaguely constrained by the oldest ages obtained from amphibole in mafic rocks (ca. 370 Ma; Dallmeyer and Quesada, 1992), although dating of the regional fabric by Rb-Sr in mica provided an age of ca. 355 Ma (García Casquero et al., 1988). Additionally, some sections of the Central Unit experienced high-T conditions and eventual partial melting in the course of their exhumation to the mid-lower crust, a process that has been dated at ca. 340 Ma (Ordóñez Casado, 1998; Pereira et al., 2010a).

The Basal Allochthonous Units that occur to the south of the Ossa-Morena Complex (Cubito-Moura Unit) exhibit a widespread foliation developed under amphibolite to greenschist facies conditions (Pedro, 1996; Moita, 1997; Fonseca et al., 1999; Booth-Rea et al., 2006). No major structures associated with this fabric have been identified so far within this unit, although the generation of large recumbent folds has been tentatively proposed (Araújo and Ribeiro, 1995). In those sections where this foliation is best preserved, the analysis of kinematic criteria has provided a consistent top-to-the-northeast tectonic transport (Araújo et al., 2005; Rosas et al., 2008), whereas a top-to-the-east sense of shear has been deduced for those domains affected by subsequent tangential deformation (kinematics inferred after unfolding tectonic fabrics; Ponce et al., 2012). In both cases the tectonic transport includes a significant component directed to the foreland located in the Cantabrian Zone. The maximum age of this fabric is ca. 358 Ma (Rosas et al., 2008), whereas the intrusion of Variscan magmas in the region at ca. 350-340 Ma (Azor et al., 2008; Pin et al., 2008) could be taken as a reliable minimum age for this deformation.

3.4.5. Iberian Parautochthon and Autochthon of NW and Central Iberia

The onset of Variscan deformation in the Iberian Parautochthon is characterized by overturned to recumbent folds with axial planar foliation. These folds are overprinted by a pervasive flat-lying foliation. These two initial deformation events developed under prograde intermediate-P, Barrovian-type metamorphism and have been related to the juxtaposition of the Iberian Allochthon over the Iberian Parautochthon and then onto the Iberian Autochthon (Marquinez García, 1984; Farias et al., 1987; Ribeiro et al., 1990; Dias da Silva, 2014). Some sections of the Parautochthon were buried following a pressure gradient higher than classical Barrovian (Rubio Pascual et al., 2015). There are not isotopic age constraints for the earlier folds but the subsequent subhorizontal shearing has been dated at ca. 340 Ma (Dallmeyer et al., 1997).

The initial Variscan record of the Iberian Autochthon is rather similar. The first deformation produced a series of overturned to recumbent folds (Díez Balda, 1986; Macaya et al., 1991; Díez Fernández et al., 2013b), the vergence of which shows a radial pattern in relation to later oroclinal bends such as the Central Iberian arc (tectonic transport towards the Cantabrian Zone; Martínez Catalán et al., 2014). Early folding was accompanied by the development of a penetrative foliation under relatively low-grade metamorphic conditions. Subsequent subhorizontal shearing generated penetrative cleavages and phyllonites close to thrust faults. The basal contact of the Iberian Parautochthon has been interpreted as the most important of these thrusts (Ribeiro et al., 1990), and cuts previous folds (e.g., Marcos and Farias, 1999; Díez Montes, 2007). Similar thrusts exist within the Iberian Autochthon, but they seem to merge into a sole fault connected with the basal thrust of the Iberian Parautochthon (González Clavijo and Martínez Catalán, 2002; Dias da Silva, 2014). Dating of the axial planar foliation of the first folds yielded an age of ca. 354-347 Ma (Rubio Pascual et al., 2013a), which is in

agreement with the broad age range previously obtained for the onset of Variscan deformation in this domain (ca. 370-342 Ma; Bea et al., 2009). Subsequent thrusting is responsible for the establishment of Viséan syn-orogenic basins at the advancing front of the Iberian Allochthon (Martínez Catalán et al., 2016), and considered coeval with subhorizontal ductile shearing in the Iberian Parautochthon, dated at ca. 340 Ma (Dallmeyer et al., 1997).

The juxtaposition of the Iberian Allochthon and Parautochthon over the Autochthon produced minimum pressures of up to ~0.9 GPa in several domains of the latter (Escuder Viruete et al., 2000; Rubio Pascual et al., 2013a), reaching up to ~1.1-1.4 GPa in some cases (Barbero and Villaseca, 2000; Rubio Pascual et al., 2015). Crustal thickening in other parts seems much less pronounced (typically in the chlorite-biotite zone), thus suggesting an inhomogeneous thrust stack.

3.4.6. South Iberian Autochthon

The Iberian Autochthon of SW Iberia occurs in two domains (Fig. 2). Both, the northeastern and southwestern occurrences, experienced Barrovian-type metamorphism during the first phases of Variscan deformation, with peak-pressures of about 0.6-0.7 GPa (González del Tánago and Arenas, 1991; Díaz Azpiroz et al., 2006). Although the type and geometry of coeval structures is not well-established for the northeastern domain, large-scale recumbent folding has been proposed for the early stages of tectonic evolution in the southwestern domain (Díaz Azpiroz et al., 2003). As to the timing of deformation, geochronological data suggest a rejuvenation of Neoproterozoic ages that might have started at ca. 390 Ma in the northeastern domain, with a remarkable maximum at ca. 360-351 Ma (Dallmeyer and Quesada, 1992). The age range of ca. 341-

337 Ma obtained for tectonic fabrics from the southwestern domain of the Iberian Autochthon (Dallmeyer et al., 1993) provides additional support for a Tournaisian-Viséan age as the onset of main Variscan deformation in the Autochthon.

3.5. Early-late Carboniferous transition (Viséan-Bashkirian)

3.5.1. NW and Central Iberian Allochthon, Parautochthon, and Autochthon

The record of early Barrovian-type metamorphism in the Autochthon and Parautochthon of NW and Central Iberia is overprinted by a flat-flying foliation that mantles and dominates the internal structure of granite- and migmatite-cored domes (Martínez Catalán et al., 2014). Usually bounded by extensional detachments, these domes alternate with large, open upright synforms and structural basins that have preserved the allochthonous complexes at their cores (Fig. 3). This extensional overprinting did not affect the Iberian Allochthon homogeneously. The upper structural levels, occupied by the Upper and Ophiolitic Allochthonous Units, escaped widespread ductile deformation and were only involved into discrete detachment faults cutting across previous tectonic boundaries (Martínez Catalán et al., 2002). Near the domes, some sections of the Basal Allochthonous Units were strongly affected by extensional deformation, including pervasive ductile shearing, faulting, and heat transfer from underlying granitic massifs (Gómez Barreiro et al., 2010a; Díez Fernández et al., 2012c).

Deformation took place under mid- to low-P conditions, and produced a low-grade crenulation cleavage in the lower parts of the suprastructure that turns quickly into a schistosity and a high-grade gneissic and migmatitic banding in the infrastructure. The extensional detachments separating these two structural levels of the crust are

characterized by low-grade phyllonites and mylonites (Escuder Viruete et al., 1994, 1998; Díez Balda et al., 1995). Extensional ductile flow gave way to shearing and stretching of syn-kinematic granitoids (Díaz-Alvarado et al., 2012) as well as a profound distortion of previous structures via flattening (Díez Fernández et al., 2013b), sometimes resulting in large-scale isoclinal folding (Arango et al., 2013). Crustal attenuation varies depending on previous thickness, reaching decompression of the infrastructure up to the andalusite stability field in many cases. The kinematics of extensional flow reveals the non-coaxial character of deformation, which stretched the previous orogenic crust following divergent vectors, usually normal to those that governed previous crustal thickening in the hinterland (Díez Fernández et al., 2012c). Dating of migmatization and colder ductile deformation in different domes yields a Viséan-Bashkirian age of ca. 350-317 Ma (Escuder Viruete et al., 1998; Montero et al., 2004; Bea et al., 2006; Castiñeiras et al., 2008; Valle Aguado et al., 2008; Rubio Pascual et al., 2013a; López-Carmona et al., 2014), a range that confers a long-lasting nature to this lithosphere extension event. Despite that broad range, most of the domes seem to have experienced maximum thermal activity at ca. 340-320 Ma, whereas granite production abounds later (around 315-305 Ma; Valle-Aguado et al., 2005; Gutiérrez-Alonso et al., 2011; Martínez Catalán et al., 2014).

3.5.2. *SW Iberian Allochthon and Autochthon*

Southern Iberia is the site of Variscan metaluminous, alkaline to calc-alkaline magmatism, which is represented by gabbro-granodiorite suites including granites, granodiorites, diorites, tonalites, gabbros, gabbros with gabbroic and noritic cumulates, and mantle-derived ores (Capdevila et al., 1973; Aparicio et al., 1977). Some of the intrusives occurring just north of the Beja-Acebuches Ophiolite are characterized by a

boninitic signature (Castro et al., 1996), an imprint that may be related to subduction, similarly to what is proposed for the calc-alkaline volcanism occurring near that region (Toca da Moura-Cabrela basin, Santos et al., 1990; Oliveira et al., 1991; Quesada et al., 1994; Onézime et al., 2003). The geochemical and isotopic characteristics of some mafic to intermediate (calc-alkaline) intrusives located farther north have been explained as a combination of high heat flow and contamination by pelitic crustal material of a mafic magma chamber formed in the mid-crust (Casquet et al., 2001; Salman, 2002; Tornos et al., 2005), or as the variable combination of mantle-derived and crustal-derived magmas (Romeo et al., 2006; Moita et al., 2009; Pereira et al., 2015). The age of all this earlier magmatic event spans a range between ca. 350-335 Ma (Dallmeyer et al., 1993, 1995; Casquet et al., 2001; Romeo et al., 2006; Jesus et al., 2007; Pin et al., 2008; Pereira et al., 2009, 2015; Cambeses et al., 2015) and is coeval with volcanism, contributing to Viséan syn-orogenic sedimentary basins (Armendariz et al., 2008; Pereira et al., 2012b; Oliveira et al., 2013).

The middle and lower structural levels of the tectonic pile in SW Iberia (Autochthon and Basal Allochthonous Units) were overprinted by high-T/low-P metamorphism and pervasive ductile deformation (González del Tánago, 1995; Díaz Azpiroz et al., 2002; Pereira et al., 2007). Peak metamorphic conditions reached the granulite facies along the southern border of the Iberian Autochthon, which constitutes a salient thermal anomaly in this region (Díaz Azpiroz et al., 2004). The general structure of these high-grade domains corresponds to granite- and migmatite-cored domes flanked by low-angle normal faults (González del Tánago, 1995; Pereira et al., 2009), where the Iberian Allochthon is mostly restricted to their hanging wall. Ductile

deformation is widespread in the infrastructure, and was responsible for the generation of crenulation cleavages and mylonitic foliation.

The high-T/low-P deformation affected syn-orogenic granitoids (peraluminous granodiorite-granite suite) and previous folds, and was parted into subhorizontal and strike-slip shear zones parallel or subparallel to the structural trend of the orogen (Díaz Azpiroz et al., 2004; Pereira et al., 2009, 2012b, 2013). On the other hand, the Iberian Allochthon displays a number of low-angle normal faults that cut previous faults and folds (Azor et al., 1994b; Expósito et al., 2002). The age of these low-angle faults can be constrained to a range of ~345-315 Ma, as they are closely related to syn-orogenic deposits of uppermost Tournaisian to Viséan age but are affected by later upright folds during the Moscovian (Gabaldón and Quesada, 1986; Giese et al., 1994; Martínez Poyatos, 2002; Expósito Ramos, 2005). Dating of later syn-orogenic granitoids and migmatization in the lower structural levels of the Autochthon provides a similar time interval for the high-T event, which appears to show maximum thermal activity at ca. 335-325 Ma (Castro et al., 1999; Díaz Azpiroz et al., 2002; Pereira et al., 2009; Lima et al., 2011, 2013; Moita et al., 2015), somewhat younger than the earlier Variscan magmatism.

A special case is made for the Puente Génave-Castelo de Vide Detachment (see detailed description by Martín Parra et al., 2006), a low-angle normal fault that defines the northern contact of the Iberian Allochthon in SW Iberia (Díez Fernández and Arenas, 2015). Its fault line extends for a minimum of 400 km, whereas the fault zone is up to 150 m thick, dips to the SSW (30° mean dip), shows consistent top-to-the-SSW kinematics, and is made of graphite-bearing schist to the east. The movement of this

768 fault should be later than the thrust tectonics that emplaced the Iberian Allochthon in the
769 first place, estimated at ca. 340 Ma (see sections 3.4.2 and 3.4.4). A maximum age for
770 the Puente Génave-Castelo de Vide Detachment is provided by Variscan granitoids and
771 related rocks deformed into its fault zone (ca. 331 Ma; Larrea et al., 1999), while its
772 minimum age is given by Moscovian granites cutting across the fault zone and dated at
773 ca. 314 Ma and ca. 307 Ma (Carracedo et al., 2009; Solá et al., 2009).

774
775 Seismic and magnetotelluric data obtained in SW Iberia reveal the existence of a
776 ~140 km long reflective and low-resistivity body located at the mid-crust, the Iberian
777 Reflective Body (IRB; Simancas et al., 2003; Muñoz et al., 2008). IRB shows variable
778 thickness (up to 5 km), making its shape wavy in some places. This body displays its
779 maximum thickness to the SSW, wedges to the NNE, and dips about 5° to the SSW.
780 Current ideas on the origin of IRB defend a hybrid (tectonic-magmatic) model, i.e. a
781 layered, mantle-derived mafic intrusion in a detachment level (Simancas et al., 2003;
782 Carbonell et al., 2004). This hypothesis may explain the Variscan alkaline, calc-
783 alkaline, and metaluminous magmatism of the region (Carbonell et al., 2004; Cambeses
784 et al., 2015), but no surface expression of the detachment level has been proposed.

785
786 The prolongation of the Puente Génave-Castelo de Vide Detachment to the
787 south, following the seismic markers of the region (Simancas et al., 2003; Martínez
788 Poyatos et al., 2012), reaches the northern edge of IRB (Fig. 3), which would account
789 for the geophysical expression of this huge extensional shear zone, or at least that of a
790 layered magmatic body shaped into the shear planes of that shear zone, either after or
791 during intrusion. This correlation suggests that the extensional shear zone associated
792 with the detachment widens with depth, conferring a listric geometry to the detachment,

and reinforcing the influence of this fault on the current (down thrown) position of the Iberian Allochthon in SW Iberia (Díez Fernández and Arenas, 2015). Moreover, the prolongation of this fault further to the south, using as a guide the seismic markers (Simancas et al., 2003), unveils a potential structural correlation with the mid-crustal root of the basal decollement from which the south-directed thrusts of the South Portuguese Zone foreland may have derived. The age of these thrusts is uppermost Viséan to Moscovian (~330-307 Ma; Silva et al., 1990), a time interval that matches the chronology of the Puente Génave-Castelo de Vide Detachment (~330-314 Ma).

3.5.3. Beja-Acebuches Ophiolite and related rocks

There is a negligible age difference between the mafic protoliths of the Beja-Acebuches Ophiolite (ca. 340-332 Ma; Azor et al., 2008) and the subsequent ductile shearing that affected this ensemble at ca. 342-328 Ma (Dallmeyer et al., 1993; Castro et al., 1999). This ophiolitic unit is strongly overprinted by left-lateral, top-to-the-SW ductile shearing (South Iberian shear zone; Crespo-Blanc and Orozco, 1988, 1991) developed under low-P and mid- to low-T metamorphic conditions (Bard and Moine, 1979; Castro et al., 1996), although some of its sections seem to preserve early top-to-the-north shearing formed under a higher metamorphic grade (Fonseca and Ribeiro, 1993).

The regional fabrics developed throughout the Basal Allochthonous Units located just north of the Beja-Acebuches Ophiolite (age estimated at ca. 358-350 Ma, see section 3.4.4) and the high-grade rocks of the South Iberian Autochthon are affected by south- and southwest-verging folds and thrusts, which show a left-lateral component comparable to the structure of the Beja-Acebuches Ophiolite (Díaz Azpiroz et al., 2003;

Araújo et al., 2005; Borrego et al., 2005; Ponce et al., 2012). Folding was developed under greenschist facies conditions and produced additional flat-lying crenulation cleavages in the region.

3.6. Late Carboniferous (Bashkirian-Gzhelian)

3.6.1. Oroclines of the Iberian Massif

Some Variscan and pre-Variscan linear features that mark the structural trend of the Iberian Massif are curved into the shape of a plate-scale vertical fold to define a couple of oroclinal bends, namely the Ibero-Armorican arc and the Central Iberian arc (Fig. 1; Martínez Catalán, 2011). These arcs are delineated by some tectonostratigraphic domains of the Iberian Massif, by the first Variscan folds of the Iberian Autochthon, by low- and high-amplitude magnetic anomalies sourced from an unexposed crystalline basement (Aerden, 2004; Martínez Catalán, 2012), and by paleocurrents in Ordovician strata (Shaw et al., 2012). The structural grain and terranes of the Iberian Allochthon do not display such curved patterns for the case of the Central Iberian arc, either in NW or SW Iberia. However, the Iberian Allochthon occupies the core of the Central Iberian arc in the NW and flanks that orocline to the SW (Fig. 1). The southern boundary of this arc runs along the Puente Génave-Castelo de Vide Detachment, which appears to cut it at a high angle.

The nucleation of the Central Iberian arc is considered to have occurred later than ca. 335 Ma (age of the youngest folds affected by the arc), whereas its closure occurred at ca. 315-305 Ma (Martínez Catalán, 2011, 2012; Martínez Catalán et al., 2014). The age of the Puente Génave-Castelo de Vide Detachment allows further constraints on the age of this orocline, most of the vertical folding related to which

should be older than the detachment (ca. 330-314 Ma). The Ibero-Armorican arc is slightly younger (Martínez Catalán, 2011), its age being constrained by means of paleomagnetic data at ca. 304-295 Ma (Weil et al., 2010).

3.6.2. Strike-slip shear zones of the Iberian Massif

Except for the Ibero-Armorican arc, all the previous Variscan record is variably affected by a series of intracontinental, strike-slip shear zones and related structures (Martínez Catalán, 2011). NW, Central, and SW Iberia exhibit a combination of dextral and sinistral shear zones (Fig. 2). Yet, left-lateral movements dominate in SW Iberia (Burg et al., 1981; Crespo-Blanc and Orozco, 1988; Pereira and Silva, 2001; Pérez-Cáceres et al., 2015a), while the major strike-slip systems in NW and Central Iberia are dextral in most of the cases (Iglesias Ponce de Leon and Choukroune, 1980; Ribeiro et al., 1980).

The strike-slip systems include zones with variable intensity of shearing. Various types of subvertical mylonites, and pervasive ductile deformation in their cores, give way to more spaced subvertical crenulation cleavages, overprinting the previous record at both sides of the shear zones. At a larger scale, the lateral displacements of these strike-slip systems deflect previous geological features such as contacts or tectonic fabrics, whereas the subhorizontal shortening experienced by the two blocks of the shear zone is accommodated by upright regional folds, most of which amplifies former extensional domes and structural basins. This is the case of the Iberian Allochthon, which is located in the core of open structural basins and it is surrounded by migmatized basement cropping out in structural domes (e.g., Padrón dome; Díez Fernández et al.,

2012c). During the later stages of transcurrent shearing, many of the strike-slip systems evolved to subvertical faults with lateral and dip-slip motion.

The strike-slip shear zones show different relationships with the development of the Central Iberian arc. However, the trace of the upright folds associated with strike-slip deformation shows high correlation with the axial trace of this orocline, suggesting that eventual tightening of this vertical fold was accomplished by shortening related to strike-slip shearing (Martínez Catalán, 2011). Although not all the shear zones are coeval sensu stricto, their age and that of related folding as a whole is very consistent throughout the Iberian Massif, and ranges between ca. 315-305 Ma (Capdevila and Vialette, 1970; Martínez Poyatos et al., 1998; Rodríguez et al., 2003; Valle Aguado et al., 2005; Gutiérrez-Alonso et al., 2015).

3.7. South Portuguese Zone

The early Variscan deformation that is observed in the South Portuguese Zone is found in its northern section (Pulo do Lobo Unit). It consists of south verging folds developed under low-grade metamorphic conditions (Silva et al., 1990; Martínez Poza et al., 2012). Such folding affected both Silurian-Devonian and Frasnian series (Pereira et al., 2008b; Braid et al., 2011). Previous age estimations considered this deformation as Upper Devonian (Giese et al., 1999).

A subsequent phase of deformation in this region produced north- to south-southwest-verging folds and involved younger sedimentary series deposited (discordantly) on top of the previous folds (Silva et al., 1990, 2013; Fonseca, 2005; Martínez Poza et al., 2012). This younger series includes strata with ages ranging from

upper Famennian, Tournaisian, and up to Viséan, as evidenced by fossil (Pereira et al., 2008b; Matas et al., 2015b) and detrital zircon data (youngest age population at ca. 347 Ma; Braid et al., 2011). Thus, the age of the first folds in the Pulo do Lobo Unit can be better constrained between ca. 380-359 Ma.

The South Portuguese Zone was affected by extension and related bimodal magmatism during the Tournaisian (~356-346 Ma; Barrie et al., 2002; Dunning et al., 2002; Rosa et al., 2008; Valenzuela et al., 2011). Later deformation progressed in a thin-skinned fashion up to the Moscovian and propagated from the Beja-Acebuches Ophiolite to the south via thrusts and related folds (~330-305 Ma; Silva et al., 1990, 2013). According to geochronological data of lithologies affected and non-affected by this later phase of deformation, south-directed thrusting must have been older in the northern part of the South Portuguese Zone (Pulo do Lobo Unit), where its age is estimated at ca. 345-335 Ma (Gladney et al., 2014). These thrusts represent a major tectonic inversion in the region and cut across the north- to south-southwest-verging folds of the Pulo do Lobo Unit (Martínez Poza et al., 2012). If so, the age of the latter folds should be Viséan (~347-335 Ma).

Some sections of the pre-Upper Devonian series of the South Portuguese Zone experienced Variscan high-P and low-intermediate-T metamorphism (Rubio Pascual et al., 2013b). These series are affected by the first phase of deformation recognized in this zone (ca. 380-365 Ma) and its Carboniferous cover does not show such tectonometamorphic imprint. Therefore, the high-P metamorphism must be either coeval or previous to the early south-directed folding (Lower to Middle Devonian?).

4. Tectonic evolution of Variscan Iberia: model and discussion

Lateral tectonics played a role on Variscan deformation affecting the Iberian Massif, impossible to quantify in full, but qualifiable. Many of the major shear zones that occur in SW Iberia (either flat-lying or subvertical) include a left-lateral component. Consequently, the position of SW Iberia -or that of the several blocks associated with strike-slip structures- relative to Central and NW Iberia before orogenesis should be located more to the northwest in present-day coordinates, i.e. west of the NW Iberian section s.l. (either southwest, purely west, or northwest). Such general assumption has been made to construct the composite section shown in Figure 3, and is strongly supported by semi-quantitative estimations on the left-lateral displacement accumulated in SW Iberia through the Variscan orogenesis (e.g., Burg et al., 1981; Pereira et al., 1998; Pérez-Cáceres et al., 2015b). Despite such restoration along-strike may result imprecise (e.g., lateral intracontinental displacements accumulated in some particular faults might have exceeded several hundreds of kilometers), the impact on the qualitative reconstruction of Variscan tangential tectonics is probably minor, as suggested by the synchrony of tectonic events across the Iberian Massif (see below).

Figure 4a shows a simplified restoration of Variscan thrusts and strike-slip shear zones and provides a general picture of the pre-collisional paleogeography across the margin of Gondwana. This reconstruction acknowledges the following ideas on the Variscan and pre-Variscan evolution of the Iberian Massif: (i) the recognition of the Iberian Allochthon across the Coimbra-Córdoba shear zone (Díez Fernández and Arenas, 2015); (ii) the Upper Allochthonous Units represent a section of the margin of

Gondwana that was rifted from its mainland during the Cambrian-Ordovician (Gómez Barreiro et al., 2007), but remained attached or geographically close to it (at least) up to the Lower Devonian (Robardet, 2003; López-Guijarro et al., 2008; Arenas et al., 2014b); (iii) the Cambrian-Ordovician rifting shaped the margin of Gondwana into a series of continental microblocks connected by stretched lithosphere (Díez Fernández et al., 2015); (iv) the onset of Variscan deformation is Lower Devonian (ca. 410-395 Ma); (v) the suture zone represented by the Allochthonous Ophiolitic Units accounts for the closure of an ephemeral oceanic basin opened after the onset of Variscan deformation (i.e. a second-order suture zone; Arenas et al., 2014a); and (vi) the Beja-Acebuches Ophiolite is the suture of a transient oceanic basin that separated most of the Iberian Gondwana from Laurussia during the early Carboniferous (Azor et al., 2008). Finally, the initial Variscan evolution of the South Portuguese Zone may not be directly related to that of the rest of the Iberian Massif, i.e. this section of putative Laurussia did not face the sections of Gondwana preserved in Iberia until Upper Devonian-Carboniferous times (Braid et al., 2011). The evolution of pre-Upper Devonian sequences of the South Portuguese Zone might be associated with NeoAcadian events recorded in Meguma (Van Staal et al., 2009). The palynological assemblages (Pereira et al., 2006b) and detrital zircon populations (Pereira et al., 2012a) found in syn-orogenic deposits at both sides of the Beja-Acebuches Ophiolite indicate that the Iberian Allochthon and its relative autochthon were close to the South Portuguese Zone during the Upper Devonian and early Carboniferous, so large distances along-strike (if any) are not expected between these two domains after closure of major oceanic basins by the Lower Devonian, such as the Rheic Ocean.

The data compilation presented in section 3 shows the synchronous character of compressional and extensional deformation events across the Iberian Massif during the Variscan orogenesis (see also Table 1). Although timing, geometry, and tectonic polarity coincides in many cases after unfolding the late oroclinal bends, the following evolutionary model is also aimed to integrate both similar and contrasted structural and metamorphic record by using age reference lines. In order to keep our model as realistic as possible, sketches presented in Figures 4 and 5 show the geometry and location of actual structures of the Iberian Massif, a reference of which has been given in section 3. The series of sketches culminates with Figure 3, which represents a synthetic cross-section of the Iberian Massif today and therefore a good approximation to the eventual structure after Variscan deformation.

4.1. Initiation of Variscan Orogeny (Fig. 4b)

The onset of the Variscan orogeny took place in the Lower Devonian and produced a fragmentary record across the Iberian Massif. The first phases of deformation related to the interaction of Laurussia and Gondwana can be observed in the Upper Allochthonous Units of NW and SW Iberia, although the structures and associated metamorphism are strikingly different depending on the region.

A stratigraphic gap and basin instability are the first hints on deformational processes heralding the Variscan orogenesis in Iberia (ca. 420-410 Ma). These can be tracked in the Upper Allochthon of SW Iberia, and are followed by the formation of a SW-verging train of recumbent folds and thrusts in a sinistral transpressional regime (Expósito et al., 2002). This major crustal thickening event took place in the outermost section of the margin of Gondwana during the Pragian-Emsian (ca. 410-395 Ma),

991 following early events of syn-orogenic deposition associated with basement denudation
992 in Lochkovian times. Top-to-the-SW kinematics (in present-day coordinates) of Lower
993 Devonian structures implies underthrusting to the NE, i.e. subduction of Laurussia
994 under Gondwana, where a magmatic arc developed in Emsian-Eifelian times (Silva et
995 al., 2011). Shortening of the upper plate would be favored by the migration of the
996 subduction hinge toward the upper plate (Doglioni et al., 2007), thus allowing the
997 development of a shallower downgoing slab, as expected for upper plates of continental
998 origin (Lallemand et al., 2005).

999
1000 In Lower Devonian, a neck of stretched peri-Gondwanan lithosphere located
1001 inboard failed mechanically under the compressive regime derived from the interaction
1002 between Laurussia and Gondwana, thus creating an intra-Gondwana subduction zone
1003 for accommodating superimposed shortening throughout the continental margin. Intra-
1004 plate subduction was probably favored by a backstop effect exerted by thick (SW-
1005 verging folds) and more buoyant Variscan lithosphere located toward the Gondwana-
1006 Laurussia suture zone (Rheic suture). Lower Devonian continental subduction was
1007 oblique (dextral; Ábalos et al., 2003) and progressed up to high-P conditions under
1008 outboard sections of Gondwana (Albert et al., 2015b).

1009
1010 Some sections of the Upper Iberian Allochthon, such as the Obejo-Valsequillo
1011 Domain, seem to have escaped from penetrative Lower Devonian deformation. This is
1012 in agreement with its intermediate position across the Upper Allochthon inferred from
1013 restoration of Variscan thrusts (Fig. 4a), and confers a remarkable microplate-like entity
1014 to the whole Upper Iberian Allochthon during this stage. The apparent lack of
1015 widespread shortening affecting the upper plate of this subduction zone (uppermost

allochthonous units) suggest that the subduction hinge remained relatively stationary, as expected for the onset of subduction zones (Doglioni et al., 2007). In this regard, sinistral lateral components acting over the external parts of the Upper Allochthon (sinistral transpression during SW-vergent folding), combined with coeval dextral movements affecting its inboard sections (oblique continental subduction), depict an overall setting of northwards escape tectonics for the case of the Upper Allochthon “microplate”.

4.2. Opening of a Devonian intra-Gondwana basin (Fig. 4c)

The geochemical signature of the Lower-Middle Devonian rocks of the Ophiolitic Allochthonous Units indicate that there were physical conditions for the opening of a marine basin following Late Devonian continental subduction (Arenas et al., 2014a). This interpretation ties into the coeval extensional record of the Iberian Autochthon (alkaline magmatism and basin subsidence; Gutiérrez-Alonso et al., 2008). Simple orthogonal restoration of the allochthonous pile reveals the location of the spreading center of this basin between the Upper Allochthonous Units and the pair constituted by the Cambrian Allochthonous Ophiolites and the Basal Allochthonous Units. According to the stratigraphic record, neither sediments were laid down at that time in the continental margins of that basin, nor do thick sedimentary series exist within the Lower-Middle Devonian ophiolites. Whether or not the lack of Middle Devonian sedimentary and volcanic rocks in the continental counterparts of the Iberian Allochthon is associated with deformation and denudation (emerged areas?), the absence of such stratigraphic record in all this domain may indicate a broad thermal uplift in relation to ridge inception.

The Lower Devonian high-P metamorphic belt that is preserved in the lowermost structural position of the Upper Allochthonous Units of NW Iberia was developed at ca. 410-390 Ma. This is virtually the same age (somewhat older) as that of the mafic protoliths of the Lower-Middle Devonian Allochthonous Ophiolitic Units (ca. 400-395 Ma), which are tectonically juxtaposed right underneath that high-P metamorphic belt. Exhumation of high- to ultra-high-P metamorphic rocks today is observed in regions subjected to high-rates of lithosphere extension and coeval ocean basin formation, such as the Woodlark rift (Davies and Warren, 1988; Wallace et al., 2004). Previously deep-seated high-P metamorphic rocks in these cases can reach lower crustal levels, and then the upper crust, in less than 3 and 5 Ma, respectively (Gordon et al., 2012). On the grounds of modern analogues, we propose that initial exhumation of the Lower Devonian high-P metamorphic rocks was strongly controlled (probably fuelled) by the opening of the intra-Gondwana oceanic basin shortly after their burial. Extension of the upper plate, triggered by a subduction hinge migrating away from the upper plate (Doglioni et al., 2007), could have facilitated both a fast exhumation and the opening of a basin, which could have then evolved as one of pull-apart type under dominant transcurrent movements (Arenas et al., 2014a).

4.3. Closure of the intra-Gondwana Devonian basin (Fig. 4d)

In Upper Devonian times, renewed convergence between Gondwana and Laurussia led to the closure of Middle Devonian oceanic domains. The tectonic polarity for this event was ruled again by thicker Variscan crust located outboard mainland Gondwana, i.e. subduction to the W and SW in present-day coordinates. Understacking of young (Devonian) oceanic crust under the Upper Allochthonous Units was followed by accretion of older (Cambrian-Ordovician) tracts of transitional crust (Arenas et al.,

2007a), and then by subduction of continental crust at ca. 380-370 Ma (Basal Allochthonous Units; Martínez Catalán et al., 1996). Regarding the latter process, insertion of more buoyant lithosphere under the (previous) Lower Devonian high-P metamorphic belt caused further exhumation of the bottom members of the Upper Allochthonous Units via tectonic denudation, which coupled to east-verging folding in response to simple shearing at the base of the upper plate (Gómez Barreiro et al., 2007). These processes continued the initial decompression experienced by the Lower Devonian high-P metamorphic rocks under the Lower-Middle Devonian rifting setting.

The Upper Devonian continental (intra-Gondwana) subduction system was formed with an angle of inclination between 15° and 30° (Alcock et al., 2005) and absorbed ongoing dextral convergence (Díez Fernández et al., 2012a). Initial exhumation within this system (Basal Allochthonous Units) was driven by crustal-scale ductile thrusting directed to the Gondwana mainland at ca. 370-360 Ma (e.g., Fervenza Thrust). Tangential deformation at this stage was concentrated on the upper part of the subducted plate, and it was likely coeval with further sinking of continental lithosphere. Ductile thrusting was assisted by erosion in the upper plate and it also forced the generation of a master, and/or a series of normal faults on top of the overthrusting high-P nappes (Díez Fernández et al., 2011). In this regard, top-to-Laurussia shear sense components of Famennian age affecting the Basal Allochthonous Units (e.g., older than ca. 358 Ma; Rosas et al., 2008) may account for normal, flat-lying shearing at the onset of decompression in response to upthrusting and extrusion of deep-seated continental nappes.

4.4. Development of the Iberian Allochthon (Figs. 5a and 5b)

Continental convergence remained during the latest Devonian and early Carboniferous (ca. 360-350 Ma) and was absorbed by W to SW (present-day coordinates) underthrusting of Gondwanan crust. Superimposed shortening probably created new contractional shear zones below the Upper Devonian subduction-exhumation channel. The onset of deformation in the Parautochthon and Autochthon of NW Iberia represents the transition from a purely continental subduction setting (recorded in the Basal Allochthonous Units) to a continent-continent collisional scenario.

At this stage, the progressive diminishing of initial high-P gradients down structure through the Variscan nappes favors a model of underthrusting of progressively thicker continental lithosphere. Protracted accretion of more buoyant continental crust to the base of the Upper Devonian subduction-exhumation system led to its progressive rotation about an horizontal axis and, consequently, to its deactivation. The early response to that exhumation process was the nucleation and propagation to the Gondwanan foreland of a train of recumbent folds within the high-P metamorphic belt and in its relative autochthon (Iberian Parautochthon and Autochthon). Convergence at this stage was also accompanied by dextral lateral movements, as indicated by tectonic fabrics associated with fold development in NW Iberia (Díez Fernández and Martínez Catalán, 2012) and probably the top-to-the-NW kinematics (Azor et al., 1994b) that dominated the exhumation process in the currently NE-dipping (originally SW-dipping) Central Unit.

Continuous constriction of the mantle wedge over the former subduction channel produced an eventual mechanical coupling between the Basal Allochthonous Units and

the rest of the Iberian Allochthon, which, from this point on, would absorb general shear deformation associated with ongoing underthrusting more efficiently (ca. 350-340 Ma). In the Upper Allochthonous Units, the existence of a train of recumbent folds with comparable age, trend and vergence than those observed in the Basal Allochthonous Units supports this idea. Fold propagation across the Upper Allochthonous Units progressed toward Laurussia, reaching the lower parts of the Obejo-Valsequillo Domain shortly afterwards. However, some of those folds in the Upper Allochthonous Units were probably nucleated during the Upper Devonian, prior to the aforementioned mechanical coupling. This may be the case of the recumbent folds affecting the Lower Devonian high-P metamorphic belt exposed in NW Iberia, for which subsequent general shear after coupling would have produced additional amplification of their initial (overturned?) recumbent geometry. All these processes attest for an orogenic shortening that is propagating more pervasively into the lower plate, but that was already affecting the upper plate since the onset of continental subduction. This transition is observed in advanced stages of continental collision following a stage more dominated by subduction (Doglioni et al., 2006, 2007).

Large-scale ductile thrusts, such as the Lalín-Forcarei Thrust and the basal thrust of the Iberian Parautochthon, represent advanced stages of the accretion of mainland Gondwana under the Iberian Allochthon and Parautochthon, respectively (ca. 340 Ma). These structures moved the Iberian Allochthon onto inner domains of Gondwana in the first place (Martínez Catalán et al., 1996), and are responsible for the juxtaposition of the Iberian Parautochthon onto the Iberian Autochthon (Ribeiro et al., 1990). The prolongation of these broad ductile shear zones toward Laurussia is possible through the (top-to-the-Cantabrian Zone) strongly sheared sequences and tectonic fabrics that

dominate the internal structure of the Basal Allochthonous Units of Iberia. The general mylonitic character of these fabrics accounts for pervasive ductile deformation along the lower structural levels of the Iberian Allochthon. A progressive ductile drag and stretching of the Basal Allochthonous Units during underthrusting would have conferred its apparent far-traveled nature to the Iberian Allochthon. During this process, the Allochthonous Ophiolites may have acquired some of its tectonically dismembered appearance. Such a broad ductile drag explains the great lateral continuity of the Upper Devonian high-P metamorphic belt across the Iberian Massif (Díez Fernández and Arenas, 2015), as well as the generation of unusually large allochthonous terranes like the Iberian Allochthon in a collisional orogeny.

Underthrusting continued during the Tournaisian-Viséan (ca. 340-330 Ma). However, the absence of regional, east to northeast verging folds of that age in the Upper Allochthonous Units located south of the Coimbra-Córdoba shear zone, suggests that simple shearing at the base of the Iberian Allochthon (if any) did not trigger folding in its outboard-most sections. In turn, continental convergence at this stage was accommodated by the nucleation of discrete reverse faults cutting across the upper plate. Among them we find the set out-of-sequence thrusts that bring pieces of an underlying suture zone to internal sections of the Upper Allochthonous Units, in the Obejo-Valsequillo Domain (Espiel Thrust; Apalategui and Pérez-Lorente, 1983; Martínez Poyatos et al., 2001). These type of faults have been also described in NW Iberia (Martínez Catalán et al., 2002), and altogether they depict an overthrusting event that transported most of the Iberian Allochthon further inboard Gondwana, thus enhancing its far-traveled nature. According to geological data, this out-of-sequence thrusting event was accomplished by taking the Upper Devonian, intra-Gondwana

suture zone and its major tectonic boundaries as primary detachment levels (e.g., Díez Fernández et al., 2013a).

Due to limited structural and tectonostratigraphic record, the role of Laurussia in the course of all this continental accretion remains uncertain. However, the development of coeval folds (south-) vergent to its mainland in the South Portuguese Zone (Martínez Poza et al., 2012) suggests that the backstop effect exerted by the Rheic suture between Gondwana and Laurussia –dipping to Gondwana since Lower Devonian times– remained active up to the lowermost Carboniferous, at both sides of the suture zone. In this scenario, the late amplification and development of south-directed thrusts affecting the south-verging folds of the Upper Allochthonous Units of SW Iberia might have occurred during the Upper Devonian through the early Carboniferous.

In the South Portuguese Zone, folding of sedimentary series postdating the onset of the Rheic suture probably represents backs and forths in the far-field interaction between Gondwana and Laurussia, as demonstrated by alternating compressional and extensional events affecting the inner sections of Gondwana (see evolutionary model). Subsequent folds verging toward Gondwana attest to a switch in the inclination of the reference shear planes. The age and structural polarity of these folds fit the timing and kinematics of ongoing crustal underthrusting under the Iberian Allochthon. Thus, we speculate that understacking of Gondwanan lithosphere might have surpassed and interplayed with the Rheic suture by the Tournaisian-Viséan.

Some of the lithosphere extension and related magmatism observed in the South Portuguese Zone (not represented in Fig. 5) occurred in the early Carboniferous (ca.

360-330 Ma). At that age, a progressive Laurussia-directed underthrusting of Gondwanan lithosphere must have produced an incremental constriction in the mantle wedge resting over the Upper Devonian continental subduction system. Such constriction implies a lateral extrusion of that portion of mantle toward Laurussia, i.e. toward the South Portuguese Zone. A readjustment like this in the mantle lithosphere under Gondwana could have led to diffuse asthenosphere upwelling, extension, and magmatism in the Laurussian side of the orogen during the course of ongoing convergence. Some of the lowermost Carboniferous magmatism in SW Iberia might be explained by this large-scale mechanism, which might also have contributed to subsequent thermal anomalies in that region.

4.5. Opening of the Beja-Acebuches basin and the onset of orogenic collapse (Fig. 5c)

The Viséan is a stage of major changes in the dynamics of the Variscan orogen. A former period ruled by convergence between Gondwana and Laurussia gives way to a new phase characterized by intra-orogenic extensional activity (Simancas et al., 2006; Pereira et al., 2012b). Two main processes stand out: the opening of the Beja-Acebuches basin (named after the Beja-Acebuches Ophiolite) and the start of orogenic gravitational collapse.

Though speculative, a look into the mantle topography before the switch to an extensional regime may offer a tectonic perspective about the origin of the latter. In our model, Viséan extension followed the underthrusting of Gondwanan crust toward Laurussia. Regardless of the amount of crustal material seated under the Iberian Allochthon, such tectonic polarity favors a thicker crustal root toward Gondwana, i.e. a higher mantle topography toward Laurussia (Fig. 5b). The constriction and lateral

extrusion of the mantle resting below peri-Gondwana, in response to a protracted underthrusting of mainland Gondwana, may have also favored such a higher mantle topography.

The numerous occurrences of mafic to intermediate magmatism of Tournaisian-Viséan age in SW Iberia have been related to a large-scale extensional event (Simancas et al., 2003), in which the opening of the Beja-Acebuches basin, floored with mafic and some ultramafic rocks, represents an eloquent proof of lithosphere necking (Azor et al., 2008). In this process, the mantle certainly played a role, as shown by varied petrological evidence from coeval mafic to intermediate magmatism (e.g., Moita et al., 2009; Pereira et al., 2009, 2015; Cambeses et al., 2015). But there is no consensus on whether or not extension was triggered by thermal anomalies in the mantle (e.g., plumes; Simancas et al., 2006), by subduction (Bard, 1977; Quesada et al., 1994), by a process of transcurrent slab break-off after collision (Pin et al., 2008), or due to intracontinental rifting in a transtensional (pull-apart?) setting (Bard, 1977; Azor et al., 2008; Cambeses et al., 2015). A higher mantle topography toward Laurussia, as suggested before, not only would imply a major thermal anomaly in the region, but also explains the location of the Beja-Acebuches basin, which could have been opened using this broad “lithosphere orogenic neck” as a trigger. The upwelling of the asthenosphere was probably responsible for the decompressional melting of the lithospheric mantle, which had already been metasomatized by a subducted slab (Rheic Ocean) leading to the generation of mafic parental magmas (Pereira et al., 2015). Simultaneously, the underplating of mafic magmas caused partial melting of continental crust. Time-equivalent and mantle-influenced magmatic activity in other parts of the Iberian Massif was apparently not related to the opening of additional oceanic basins (e.g., Pyrite belt;

Mitjavila et al., 1997; Martin-Izard et al., 2016), so lithosphere extension at this stage was probably heterogeneous (additional minor lithosphere necks might have existed) and/or the far-field influence of the mantle high here suggested was not restricted to the Gondwana-Laurussia suture.

A simple restoration of Variscan thrusts in SW Iberia indicates that the opening of the Beja-Acebuches basin cut off the tectonic pile culminated by the Iberian Allochthon. This implies that the former Rheic suture between Gondwana and Laurussia in Iberia became part of a different plate than the rest of the orogen. The actual location of that suture zone is a matter of debate, because traces of the Rheic Ocean crust are yet to be found. The terrane capable of sourcing sediments dispersed on both sides of the Rheic suture is interpreted to have been completely removed by erosion in SW Iberia (Pereira et al., 2012a). In this regard, erosion of rift shoulders during the opening of the Beja-Acebuches basin and/or subsequent crustal understacking during its closure are two likely mechanisms capable of hiding most of the previous orogenic record associated with the Rheic suture. This is particularly true for the lower plate to the north-dipping suture of the Beja-Acebuches basin. Interestingly, that plate is the one where the initial suture between Gondwana and Laurussia was located after the intracontinental rifting that gave way to the Beja-Acebuches basin. Therefore, even if separating Gondwanan and Laurussian domains, the suture of the Beja-Acebuches basin should be considered as a reworked one.

Lessening of the gravitational disequilibrium created after the transference of the Iberian Allochthon onto the Gondwana mainland was conducted by extensional detachments in the upper part of the tectonic pile. The former understacking, thickening

and pressurization of its relative autochthon favored the thermal disequilibrium of the orogenic crust. This crust started to flow laterally and melt fertile crustal layers as response to thermal reequilibration, giving rise to migmatitic domes across the Iberian Autochthon and to felsic (crust-derived) magmatism after a period of thermal maturation (Alcock et al., 2009; Pereira et al., 2009; Martínez Catalán et al., 2014). The consequent crustal extension forced the mantle to rise to compensate for lithosphere attenuation. The opening of the Beja-Acebuches basin was roughly coincidental with the initiation of the gravitational collapse of the Variscan thrust pile. With this perspective, we think that positive feedback probably existed between the early stages of thermal and gravitational reequilibration of the orogen and the lithosphere extension that led to the opening of that basin.

Thermal models suggest that the overriding of the Iberian Allochthon can explain alone the extensional collapse of the Variscan crust and the generation of abundant (crust-derived) magmatism of Serpukhovian-Bashkirian age (Alcock et al., 2015). Yet, the mafic to intermediate (alkaline, calc-alkaline, and metaluminous) magmatism of SW Iberia is slightly older (Tournaisian-Viséan) and shows fair mantle input. Consequently, the mechanism(s) in place for the development of such mafic to intermediate magmatism (e.g., plumes, transient arc, slab break-off, and/or orogenic transtension) could have also contributed to the collapse of the orogenic hinterland in the first place. In this regard, the existence of a higher mantle topography toward Laurussia would explain not only older (mantle-induced?) extensional activity in SW Iberia but also larger (petrological) mantle contributions in the lack of a thick crustal root underneath this region. Remarkably, mantle contributions diminish toward the north and northeast of the Iberian Massif, where extensional syn-orogenic magmatism is

slightly younger and is clearly dominated by crustal sources (e.g., Villaseca et al., 1998). That petrological and geochronological trend across the orogen accords well with the existence of a thicker crustal root toward Gondwana, supports the existence of irregular mantle topography before extension, and is consistent with protracted underthrusting of Gondwanan crust toward Laurussia.

4.6. The collapse of Variscan orogenic crust (Fig. 5d)

Once the cohesion of the orogen was lost to its thermal re-equilibrium, the gravitational collapse gained importance through Serpukhovian-Bashkirian times. This stage is characterized by extensive felsic magmatism, which occurred preferentially in areas subjected to severe denudation, i.e. under the Iberian Allochthon. Extensional faults in the upper crust drove further tectonic denudation (e.g., Pico Sacro and Puente Génave – Castelo de Vide detachments), whereas lower crustal flow distributed vertical flattening and lithosphere attenuation across the orogen. Regions dominated by felsic magmatism of Serpukhovian-Bashkirian age occur in the core of dome structures (e.g., Padrón dome), revealing the contribution of diapiric flow to the reequilibration process.

The extensional collapse of the Variscan orogen has been classically viewed as a syn-convergent process (Franke, 2000). One of the main reasons sustaining this idea in the case of Iberia is that extension was preceded and followed by indisputable phases of continental convergence (Martínez Catalán et al., 2002). Indeed, thermal and gravitational re-equilibration were not acting alone on the overthickened crust. The onset of the Central Iberian arc has been framed in this stage too (Martínez Catalán, 2012). Orogen-parallel extensional flow dominated the gravitational re-equilibration of the hinterland, and has been interpreted as the result of ongoing oblique plate

movements in the course of orogenic collapse (Díez Fernández et al., 2012c). Additionally, the development of strike-slip shear zones coeval with extension, furthers the role exerted by lateral tectonics at this stage (Pereira et al., 2009). Any of the aforementioned structural records could have been developed either under plate-scale transtension and/or transpression. There is, however, a major geodynamic event on which convergence setting at plate scale relies during Serpukhovian-Bashkirian times, the closure of the Beja-Acebuches basin.

Dipping under the Iberian Allochthon and Autochthon, the Beja-Acebuches Ophiolite has an age of accretion (ca. 342-328 Ma) that matches the age of the onset of Variscan orogenic collapse. Hence, the subduction of oceanic lithosphere formed in the Beja-Acebuches basin provides a convergence geodynamic setting under which ongoing gravitational collapse must have evolved. The closure of the oceanic domain represented in the Beja-Acebuches Ophiolite has been widely considered as related to a subduction process (e.g., Munhá et al., 1986; Eden and Andrews, 1990; Silva et al., 1990; Fonseca and Ribeiro, 1993; Quesada et al., 1994; Simancas et al., 2003; Díaz Azpiroz et al., 2006; Braid et al., 2010), although part of the tectonic evolution of this domain could be also related to an obduction event (Pérez-Cáceres et al., 2015a). Recent findings of lawsonite-bearing rocks in the Pulo do Lobo Unit (Rubio Pascual et al., 2013b) suggest the formation of a pressure-dominated metamorphic belt during the accretion of the Beja-Acebuches Ophiolite, thus providing additional support to models that acknowledge subduction as a driving mechanism during the closure of the Beja-Acebuches basin.

The development of SW-verging folds and thrusts affecting the previous record, both in the Iberian Allochthon and Autochthon of SW Iberia during the Serpukhovian-Bashkirian, can be explained by a NE-directed tectonic polarity for the closure of the Beja-Acebuches basin. Convergence at this stage probably occurred in a transpressional setting, as suggested by sinistral lateral movements along major tectonic boundaries of SW Iberia (Crespo-Blanc, 1992). In this scenario, convergence may have also facilitated reactivation of previous thrusts, particularly in the upper crust (e.g., out-of-sequence thrusts with Serpukhovian-Bashkirian age). Later pronounced extension within the orogenic hinterland facilitated the widening of former sedimentary basins over the Variscan allochthonous nappes (e.g., Los Pedroches basin).

The opening of the Beja-Acebuches basin also covers the start of the orogenic collapse (see section 4.5). Either a mantle upwelling in response to the inception of the Beja-Acebuches basin, and/or the subsequent consumption of that same basin by NE-directed subduction, are two expected contributors of deep-sourced material to bear on the orogenic collapse. The ca. 328-317 Ma calc-alkaline to adakitic-like magmatism (Pavia pluton; Lima et al., 2013) lying to the north of the Beja-Acebuches Ophiolite may represent the product of such later subduction. In this way, we find older (and much more abundant) evidence of such crustal growth toward SW Iberia (closer to the Beja-Acebuches Ophiolite; e.g., Pereira et al., 2009; Cambeses et al., 2015) than to Central and NW Iberia (toward the advancing front of Variscan allochthonous nappes; e.g., Dias et al., 2002; Rodríguez et al., 2007). This makes a petro-geochronological trend that may express either the lag in the rise of the mantle after maximum crustal thickening (earlier in SW Iberia by favorable mantle topography after rifting), and/or the arrival of mantle-derived melts related to a downgoing oceanic tract that sinks

progressively to the northeast (consumption of the Beja-Acebuches basin), i.e. toward the advancing front of Variscan allochthonous nappes. That sector of the orogen would be equivalent to a broad back-arc region relative to the subduction zone closing the Beja-Acebuches basin. Eventual migrations of its subduction hinge (e.g., Doglioni et al., 2007) might explain transient extensional or compressional regimes affecting that section of the orogen.

The east- and northeast-directed collapse of the eastern orogenic hinterland is roughly contemporaneous with the early stages of east-directed thrusting in the western part of the foreland of the Cantabrian Zone (Dallmeyer et al., 1997; Martínez Catalán et al., 2003). Such coupling between orogen-perpendicular hinterland extension and foreland compression in the Gondwanan flank of the orogen has its equivalent in the Laurussian side (South Portuguese Zone). South-directed extension along the Puente Génave-Castelo de Vide Detachment is coeval with the southerly propagation of thrusts and folds in the foreland of the South Portuguese Zone. Therefore the lateral spreading of the orogenic crust has been a fundamental cause for triggering Variscan shortening across foreland basins at both sides of the orogen (Cantabrian and South Portuguese zones).

4.7. Late strike-slip tectonics (Fig. 3)

Convergence persisted during the orogenic collapse, which, in turn, waned as extensional flow reduced gradients of potential energy. Thermal equilibrium was not fully achieved in the process, since syn-orogenic magmatism remained throughout the Bashkirian and Moscovian (Pereira et al., 2009, 2015; Martínez Catalán et al., 2014; Cambeses et al., 2015). Subhorizontal extension was eventually outpaced by

superimposed subhorizontal compression in Moscovian times. Some of the magmatism at this stage occurred in close relation to strike-slip shear zones (e.g., Aranguren et al., 1997; Valle Aguado et al., 2005; Carracedo et al., 2009), which accommodated most of the lateral components of convergence along their central parts and distributed shortening in their tectonic blocks, thus producing open upright folds all over the Variscan hinterland. Transcurrent deformation reworked previous thrusts and normal faults and partly redrew the map of tectonic blocks. Dextral and sinistral strike-slip shear zones acted together and created escape tectonics settings at local scale (Iglesias Ponce de Leon and Choukroune, 1980). Such settings in the hinterland were coeval with further propagation of thrusts and folds to both the Gondwanan and Laurussian forelands. Oroclinal bending of the orogen occurred in the course of all of this deformation, starting from the Serpukhovian-Bashkirian extensional collapse and culminating with the folding of the latest syn-orogenic deposits of the Gondwanan foreland.

5. Conclusions

Strike-slip deformation during the Moscovian segmented the hinterland of the Variscan orogen into new tectonic blocks, partly different from those operating during previous convergence processes between Gondwana and Laurussia. Becoming fully aware of this particular switch in the architecture of the orogen (even if it was transitional) is essential for understanding the common structural history linked to previous tangential tectonics at both sides of major transcurrent shear zones, such as the Coimbra-Córdoba shear zone. Shear zones accommodating large amounts of tangential deformation, transported pieces of continental and oceanic crust located at the periphery of Gondwana that were affected by previous Variscan deformation. These shear zones

are envisaged as the rulers during the early stages of Pangea amalgamation, which, however, did not seal Gondwana and Laurussia once for all. Inception of short-lived oceanic basins following periods of convergence provides solid evidence on a complex amalgamation process in southern Europe, hardly explainable by a single collisional process.

Based on our integration of structural and geochronological data, the Variscan tectonic evolution of the Iberian Massif can be summarized as follows (Paleozoic geographic coordinates):

1. Following the closure of the Rheic Ocean, Gondwana and Laurussia collided in the Lower Devonian. Kinematics of major structures that developed toward the most external margin of Gondwana support that Laurussia was the lower plate to the Rheic suture.
2. Contraction over the margin of Gondwana initiated an intra-Gondwana, continental subduction zone dipping to the north, which progressively spread under the Rheic suture.
3. A transient period of extension after continental subduction led to the opening of an intra-Gondwana oceanic basin in the Lower-Middle Devonian. Such intra-orogenic rifting coupled with the initial exhumation of high-P rocks within the former continental subduction system.
4. Closure of the intra-Gondwana basin in the Upper Devonian caused the accretion to the north of Devonian oceanic crust, then Cambrian-Ordovician transitional crust, and finally the subduction of inner sections of Gondwana to the north.

- 1439 5. Continuous convergence between Gondwana and Laurussia during the early
1440 Carboniferous was accommodated by underthrusting of Gondwanan lithosphere
1441 to the north, below the peri-Gondwanan domain that had been previously
1442 involved in the collisional orogenesis. Protracted underthrusting locked the
1443 Upper Devonian intra-Gondwana subduction first, and then forced the
1444 mechanical coupling between the lower and upper tectonic plate. Coeval
1445 shearing throughout the orogenic crust generated a series of south-directed folds
1446 and a series of extensional faults in the upper plate. Ductile drag exerted by the
1447 lower plate extended the Upper Devonian subduction system and the intra-
1448 Gondwana suture zone under the upper plate, thus shaping this whole ensemble
1449 of peri-Gondwanan terranes into a set of allochthonous units.
- 1450 6. Further convergence nucleated a system of out-of-sequence thrusts, which
1451 reworked the intra-Gondwana suture in the course of its obduction onto the
1452 Gondwana mainland.
- 1453 7. Rifting of the resulting overthickened crust led to the opening of a short-lived
1454 oceanic basin (Beja-Acebuches Ophiolite) near the Gondwana-Laurussia suture
1455 zone formed in the Lower Devonian.
- 1456 8. Intra-continental extension was followed by or coeval with the gravitational
1457 collapse and thermal re-equilibration of the orogen, which remained active up to
1458 the late Carboniferous. Continental convergence resumed shortly afterwards, and
1459 forced the closure of newly-formed oceanic basins. Deformation propagated via
1460 thrusts and folds toward the mainland of both Gondwana and Laurussia and
1461 favored the reactivation of former thrusts.
- 1462 9. Lithosphere extension in the hinterland was progressively replaced by strike-slip
1463 deformation. Oroclinal bending of the orogen started in this transition. Lateral

1464 tectonics at this stage was manifested in discrete, subvertical shear bands and in
1465 the upright folding of previous flat-lying structures. Foreland propagation of
1466 deformation continued during this period. Variscan deformation concluded with
1467 the development of late oroclinal bends affecting the whole structural grain of
1468 the orogen.

1470 **6. Acknowledgments**

1471 We are indebted to insightful comments and input provided by Dr. W. Franke
1472 and Dr. C. Doglioni. Financial support has been provided by the Spanish project
1473 CGL2012-34618. Rubén Díez Fernández appreciates financial support from Ministerio
1474 de Economía y Competitividad (Spain) through its Juan de la Cierva postdoctoral
1475 program (JCI-2012-11967). This work is a contribution to IGCP project 648
1476 (Supercontinent Cycle and Global Geodynamics) and IDL Research Group 5.

1478 **7. References**

1480 Abalos, B., Eguiluz, L., Gil Ibarguchi, J.I., 1991a. Evolución tectono-metamórfica del
1481 Corredor Blastomilonítico de Badajoz-Córdoba. II: Las unidades alóctonas y
1482 trayectorias PTt. Boletín Geológico y Minero 102-5, 617-671.

1483 Abalos, B., Gil Ibarguchi, J.I., Eguiluz, L., 1991b. Cadomian subduction, collision and
1484 Variscan transpression in the Badajoz-Cordoba Shear Belt, Southwest Spain.
1485 Tectonophysics 199, 51-72.

1486 Ábalos, B., Puellas, P., Gil Ibarguchi, J.I., 2003. Structural assemblage of high-pressure
1487 mantle and crustal rocks in a subduction channel (Cabo Ortegal, NW Spain). Tectonics
1488 22, 1006.

- 1489 Abati, J., Dunning, G.R., 2002. Edad U-Pb en monacitas y rutilos de los paragneisses de
1490 la Unidad de Aqualada (Complejo de Ordenes, NW del Macizo Ibérico). *Geogaceta* 32,
1491 95-98.
- 1492 Abati, J., Gerdes, A., Fernández-Suárez, J., Arenas, R., Whitehouse, M.J., Díez
1493 Fernández, R., 2010. Magmatism and early-Variscan continental subduction in the
1494 northern Gondwana margin recorded in zircons from the basal units of Galicia, NW
1495 Spain. *Geological Society of America Bulletin* 122, 219-235.
- 1496 Aerden, D.G.A.M., 2004. Correlating deformation in Variscan NW-Iberia using
1497 porphyroblasts; implications for the Ibero-Armorican Arc. *Journal of Structural*
1498 *Geology* 26, 177-196.
- 1499 Albert, R., Arenas, R., Sánchez-Martínez, S., Gerdes, A., 2012. The eclogite facies
1500 gneisses of the Cabo Ortegal Complex (NW Iberian Massif): Tectonothermal evolution
1501 and exhumation model. *Journal of Iberian Geology* 38, 389-406.
- 1502 Albert, R., Arenas, R., Gerdes, A., Sánchez Martínez, S., Fernández-Suárez, J.,
1503 Fuenlabrada, J.M., 2015a. Provenance of the Variscan Upper Allochthon (Cabo Ortegal
1504 Complex, NW Iberian Massif). *Gondwana Research* 28, 1434-1448.
- 1505 Albert, R., Arenas, R., Gerdes, A., Sánchez Martínez, S., Marko, L., 2015b. Provenance
1506 of the high-P and high-T unit of the Cabo Ortegal Complex (NW Iberian Massif).
1507 *Journal of Metamorphic Geology* 33, 959-979.
- 1508 Alcock, J.E., Arenas, R., Martínez Catalán, J.R., 2005. Shear stress in subducting
1509 continental margin from high-pressure, moderate-temperature metamorphism in the
1510 Ordenes Complex, Galicia, NW Spain. *Tectonophysics* 397, 181-194.
- 1511 Alcock, J.E., Martínez Catalán, J.R., Arenas, R., Díez Montes, A., 2009. Use of thermal
1512 modeling to assess the tectono-metamorphic history of the Lugo and Sanabria gneiss
1513 domes, Northwest Iberia. *Bulletin de la Societe Geologique de France* 180, 179-197.
- 1514 Alcock, J.E., Martínez Catalán, J.R., Rubio Pascual, F.J., Montes, A.D., Díez
1515 Fernández, R., Gómez Barreiro, J., Arenas, R., Dias da Silva, Í., González Clavijo, E.,

- 1516 2015. 2-D thermal modeling of HT-LP metamorphism in NW and Central Iberia:
1517 Implications for Variscan magmatism, rheology of the lithosphere and orogenic
1518 evolution. *Tectonophysics* 657, 21-37.
- 1519 Andonaegui, P., Castiñeiras, P., González Cuadra, P., Arenas, R., Sánchez Martínez, S.,
1520 Abati, J., Díaz García, F., Martínez Catalán, J.R., 2012. The Corredoiras orthogneiss
1521 (NW Iberian Massif): Geochemistry and geochronology of the Paleozoic magmatic
1522 suite developed in a peri-Gondwanan arc. *Lithos* 128–131, 84-99.
- 1523 Apalategui, O., Pérez-Lorente, F., 1983. Nuevos datos en el borde meridional de la zona
1524 centro ibérica : el dominio Obejo-Valsequillo-Puebla de la Reina. *Studia Geológica*
1525 *Salmanticensia* 18, 193-200.
- 1526 Apalategui, O., Eguiluz, L., Quesada, C., 1990. Ossa-Morena zone: structure, in:
1527 Dallmeyer, R.D., Martínez García, E. (Eds.), *Pre-Mesozoic Geology of Iberia*. Springer-
1528 Verlag, Berlin, Germany, pp. 280-292.
- 1529 Aparicio, A., Barrera, J.L., Casquet, C., Peinado, M., Tinao, J.M., 1977. El Plutonismo
1530 hercínico post-metamórfico en el SO del macizo hespérico (España). *Boletín Geológico*
1531 *y Minero* 88, 497-500.
- 1532 Arango, C., Díez Fernández, R., Arenas, R., 2013. Large-scale flat-lying isoclinal
1533 folding in extending lithosphere: Santa María de la Alameda dome (Central Iberian
1534 Massif, Spain). *Lithosphere* 5, 483-500.
- 1535 Aranguren, A., Larrea, F., Carracedo, M., Cuevas, J., Tubía, J., 1997. The Los
1536 Pedroches Batholith (Southern Spain): Polyphase Interplay between Shear Zones in
1537 Transtension and Setting of Granites, in: Bouchez, J.L., Hutton, D.H.W., Stephens,
1538 W.E. (Eds.), *Granite: From Segregation of Melt to Emplacement Fabrics*. Springer
1539 Netherlands, pp. 215-229.
- 1540 Araújo, A., Ribeiro, A., 1995. Tangential transpressive strain regime in the Évora-
1541 Aracena Domain (Ossa-Morena Zone). *Boletín Geológico y Minero* 106, 111-117.

- 1542 Araújo, A., Fonseca, P.E., Munhá, J.M., Moita, P., Pedro, J., Ribeiro, A., 2005. The
1543 Moura Phyllonitic Complex: an accretionary complex related with obduction in the
1544 Southern Iberia Variscan Suture. *Geodinamica Acta* 18, 375-388.
- 1545 Araújo, A., Piçarra, J., Borrego, J., Pedro, J., Oliveira, T., 2013. As regiões central e sul
1546 da Zona de Ossa Morena, in: Dias, R., Araújo, A., Terrinha, P., Kullberg, J.C. (Eds.),
1547 *Geologia de Portugal*. Escolar Editora, pp. 151-172.
- 1548 Arenas, R., Martínez Catalán, J.R., 2002. Prograde development of corona textures in
1549 metagabbros of the Sobrado unit (Ordenes Complex, northwestern Iberian Massif), in:
1550 Martínez Catalán, J.R., Hatcher, R.D., Arenas, R., Díaz García, F. (Eds.), *Variscan-*
1551 *Appalachian Dynamics: The building of the late Paleozoic basement*. Geological
1552 Society of America Special Paper, pp. 73-88, doi: 10.1130/1130-8137-2364-1137.1173.
- 1553 Arenas, R., Gil Ibarguchi, J.I., González Lodeiro, F., Klein, E., Martínez Catalán, J.R.,
1554 Ortega Gironés, E., Pablo Maciá, J.G.d., Peinado, M., 1986. Tectonostratigraphic units
1555 in the complexes with mafic and related rocks of the NW of the Iberian Massif.
1556 *Hercynica* 2, 87-110.
- 1557 Arenas, R., Rubio Pascual, F.J., Díaz García, F., Martínez Catalán, J.R., 1995. High-
1558 pressure micro-inclusions and development of an inverted metamorphic gradient in the
1559 Santiago-schists (Órdenes-Complex, NW Iberian Massif, Spain) - Evidence of
1560 subduction and syncollisional decompression. *Journal of Metamorphic Geology* 13,
1561 141-164.
- 1562 Arenas, R., Abati, J., Martínez Catalán, J.R., Díaz García, F., Rubio Pascual, F.J., 1997.
1563 P-T evolution of eclogites from the Agualada unit (Ordenes complex, northwest Iberian
1564 Massif, Spain): Implications for crustal subduction. *Lithos* 40, 221-242.
- 1565 Arenas, R., Martínez Catalán, J.R., Sánchez Martínez, S., Díaz García, F., Abati, J.,
1566 Fernández-Suárez, J., Andonaegui, P., Gómez Barreiro, J., 2007a. Paleozoic ophiolites
1567 in the Variscan suture of Galicia (northwest Spain): distribution, characteristics and
1568 meaning, in: Hatcher, R.D., Carlson, M.P., McBride, J.H., Martínez Catalán, J.R. (Eds.),

1569 4-D Framework of Continental Crust. Geological Society of America Memoir, Boulder,
1570 Colorado, pp. 425-444.

1571 Arenas, R., Martínez Catalán, J.R., Sánchez Martínez, S., Fernández-Suárez, J.,
1572 Andonaegui, P., Pearce, J.A., Corfú, F., 2007b. The Vila de Cruces ophiolite: A
1573 remnant of the early Rheic Ocean in the Variscan suture of Galicia (northwest Iberian
1574 Massif). *The Journal of Geology* 115, 129-148.

1575 Arenas, R., Sánchez Martínez, S., Castineiras, P., Jeffries, T.E., Díez Fernández, R.,
1576 Andonaegui, P., 2009. The basal tectonic melange of the Cabo Ortegal Complex (NW
1577 Iberian Massif): a key unit in the suture of Pangea. *Journal of Iberian Geology* 35, 85-
1578 125.

1579 Arenas, R., Díez Fernández, R., Sánchez Martínez, S., Gerdes, A., Fernández-Suárez, J.,
1580 Albert, R., 2014a. Two-stage collision: Exploring the birth of Pangea in the Variscan
1581 terranes. *Gondwana Research* 25, 756-763.

1582 Arenas, R., Sánchez Martínez, S., Gerdes, A., Albert, R., Díez Fernández, R.,
1583 Andonaegui, P., 2014b. Re-interpreting the Devonian ophiolites involved in the
1584 Variscan suture: U–Pb and Lu–Hf zircon data of the Moeche Ophiolite (Cabo Ortegal
1585 Complex, NW Iberia). *International Journal of Earth Sciences* 103, 1385-1402.

1586 Arenas, R., Sánchez Martínez, S., 2015. Variscan ophiolites in NW Iberia: Tracking lost
1587 Paleozoic oceans and the assembly of Pangea. *Episodes* 38, 315-333.

1588 Armendariz, M., López-Guijarro, R., Quesada, C., Pin, C., Bellido, F., 2008. Genesis
1589 and evolution of a syn-orogenic basin in transpression: Insights from petrography,
1590 geochemistry and Sm-Nd systematics in the Variscan Pedroches basin (Mississippian,
1591 SW Iberia). *Tectonophysics* 461, 395-413.

1592 Azor, A., 1994. Evolución tectonometamórfica del límite entre las zonas Centroibérica
1593 y de Ossa-Morena (Cordillera Varisca, SO de España). Universidad de Granada,
1594 Granada, p. 312.

1595 Azor, A., González Lodeiro, F., Martínez Poyatos, D., Simancas, J., 1994a. Regional
1596 significance of kilometric-scale north-east vergent recumbent folds associated with east
1597 to south-east directed shear on the southern border of the Central Iberian Zone
1598 (Hornachos-Oliva region, Variscan belt, Iberian Peninsula). *Geologische Rundschau* 83,
1599 377-387.

1600 Azor, A., Lodeiro, F.G., Simancas, J.F., 1994b. Tectonic evolution of the boundary
1601 between the Central Iberian and Ossa-Morena zones (Variscan belt, southwest Spain).
1602 *Tectonics* 13, 45-61.

1603 Azor, A., Rubatto, D., Simancas, J.F., González Lodeiro, F., Martínez Poyatos, D.,
1604 Martín Parra, L.M., Matas, J., 2008. Rheic Ocean ophiolitic remnants in southern Iberia
1605 questioned by SHRIMP U-Pb zircon ages on the Beja-Acebuches amphibolites.
1606 *Tectonics* 27, TC5006.

1607 Ballèvre, M., Bosse, V., Ducassou, C., Pitra, P., 2009. Palaeozoic history of the
1608 Armorican Massif: Models for the tectonic evolution of the suture zones. *Comptes*
1609 *Rendus Geoscience* 341, 174-201.

1610 Bambach, R.K., Scotese, C.R., Ziegler, A.M., 1980. Before Pangea - Geographies of the
1611 Paleozoic World. *American Scientist* 68, 26-38.

1612 Barbero, L., Villaseca, C., 2000. Eclogite facies relics in metabasites from the Sierra de
1613 Guadarrama (Spanish Central System) P-T estimations and implications for the
1614 Hercynian evolution. *Mineralogical Magazine* 64, 815-836.

1615 Bard, J.P., 1977. Signification tectonique des metatholeites d'affinite abyssale de la
1616 ceinture metamorphique de basse pression d'Aracena (Huelva, Espagne). *Bulletin de la*
1617 *Societe Geologique de France Series 7 Vol. XIX*, 385-393.

1618 Bard, J.P., Moine, B., 1979. Acebuches amphibolites in the Aracena hercynian
1619 metamorphic belt (southwest Spain): Geochemical variations and basaltic affinities.
1620 *Lithos* 12, 271-282.

1621 Barrie, T.C., Amelin, Y., Pascual, E., 2002. U–Pb Geochronology of VMS
1622 mineralization in the Iberian Pyrite Belt. *Mineral. Deposita* 37, 684-703.

1623 Bea, F., Montero, P., Gonzalez Lodeiro, F., Talavera, C., Molina, J.F., Scarrow, J.H.,
1624 Whitehouse, M.J., Zinger, T., 2006. Zircon thermometry and U-Pb ion-microprobe
1625 dating of the gabbros and associated migmatites of the Variscan Toledo Anatectic
1626 Complex, Central Iberia. *Journal of the Geological Society* 163, 847-855.

1627 Bea, F., Pesquera, A., Montero, P., Torres-Ruiz, J., Gil-Crespo, P.P., 2009. Tourmaline
1628 $^{40}\text{Ar}/^{39}\text{Ar}$ chronology of tourmaline-rich rocks from Central Iberia dates the main
1629 Variscan deformation phases. *Geologica Acta* 7, 399-412.

1630 Blatrix, P., Burg, J.P., 1981. $^{40}\text{Ar}/^{39}\text{Ar}$ dates from Sierra Morena (southern Spain):
1631 Variscan metamorphism and Cadomian Orogeny. *Neues Jahrbuch Mineral Monatshefte*
1632 10, 470-478.

1633 Booth-Rea, G., Simancas, J.F., Azor, A., Azañón, J.M., González-Lodeiro, F., Fonseca,
1634 P., 2006. HP-LT Variscan metamorphism in the Cubito-Moura schists (Ossa-Morena
1635 Zone, southern Iberia). *Comptes Rendus Geoscience* 338, 1260-1267.

1636 Borrego, J., 2009. Cartografia Geológico-Estrutural de um sector da Zona de Ossa-
1637 Morena (Subsector de Estremoz - Barrancos-Ficalho) e sua interpretação Tectónica.
1638 Universidade de Évora, Évora, p. 479.

1639 Borrego, J., Araújo, A., Fonseca, P., 2005. A geotraverse through the south and central
1640 sectors of the Ossa-Morena Zone in Portugal (Iberian Massif). *Journal of the Virtual*
1641 *Explorer* 19, 1-16.

1642 Braid, J.A., Murphy, J.B., Quesada, C., 2010. Structural analysis of an accretionary
1643 prism in a continental collisional setting, the Late Paleozoic Pulo do Lobo Zone,
1644 Southern Iberia. *Gondwana Research* 17, 422-439.

1645 Braid, J.A., Murphy, J.B., Quesada, C., Mortensen, J., 2011. Tectonic escape of a
1646 crustal fragment during the closure of the Rheic Ocean: U-Pb detrital zircon data from

- 1647 the Late Palaeozoic Pulo do Lobo and South Portuguese zones, southern Iberia. *Journal*
1648 *of the Geological Society* 168, 383-392.
- 1649 Burg, J.P., Matte, P., 1978. A cross section through the French Massif Central and the
1650 scope of its Variscan geodynamic evolution. *Zeitschrift der Deutschen Geologischen*
1651 *Gesellschaft* 129, 429-440.
- 1652 Burg, J.P., Iglesias, M., Laurent, P., Matte, P., Ribeiro, A., 1981. Variscan
1653 intracontinental deformation: The Coimbra-Cordoba shear zone (SW Iberian Peninsula).
1654 *Tectonophysics* 78, 161-177.
- 1655 Cambeses, A., Scarrow, J.H., Montero, P., Molina, J.F., Moreno, J.A., 2015. SHRIMP
1656 U–Pb zircon dating of the Valencia del Ventoso plutonic complex, Ossa-Morena Zone,
1657 SW Iberia: Early Carboniferous intra-orogenic extension-related ‘calc-alkaline’
1658 magmatism. *Gondwana Research* 28, 735-756.
- 1659 Capdevila, R., Vialette, Y., 1970. Estimation radiométrique de l'âge de la deuxième
1660 phase tectonique hercynienne en Galice moyenne (Nordouest de l'Espagne). *Comptes*
1661 *Rendus Académie des Sciences Paris* 270, 2527-2530.
- 1662 Capdevila, R., Corretgé, L.G., Floor, P., 1973. Les granitoides varisques de la meseta
1663 ibérique. *Bulletin de la Société Géologique de France* 7, 209-228.
- 1664 Carbonell, R., Simancas, F., Juhlin, C., Pous, J., Pérez-Estaún, A., Gonzalez-Lodeiro,
1665 F., Muñoz, G., Heise, W., Ayarza, P., 2004. Geophysical evidence of a mantle derived
1666 intrusion in SW Iberia. *Geophysical Research Letters* 31, L11601.
- 1667 Carracedo, M., Paquette, J.L., Olazabal, A.A., Santos Zalduegui, J.F., de Madinabeitia,
1668 S.G., Tiepolo, M., Gil Ibarguchi, J.I., 2009. U-Pb dating of granodiorite and granite
1669 units of the Los Pedroches batholith. Implications for geodynamic models of the
1670 southern Central Iberian Zone (Iberian Massif). *International Journal of Earth Sciences*
1671 98, 1609-1624.
- 1672 Carvalho, D., Correia, H., Inverno, C., 1976. Contribuição para o conhecimento
1673 geológico do Grupo de Ferreira-Ficalho. Suas relações com a Faixa Piritosa e o Grupo

- 1674 de Pulo do Lobo. IV Reunião de Geologia do Oeste Peninsular. Salamanca - Coimbra,
1675 1976. Memórias y Notícias, Coimbra 82, 145-169.
- 1676 Casquet, C., Galindo, C., Tornos, F., Velasco, F., Canales, A., 2001. The Aguablanca
1677 Cu–Ni ore deposit (Extremadura, Spain), a case of synorogenic orthomagmatic
1678 mineralization: age and isotope composition of magmas (Sr, Nd) and ore (S). Ore
1679 Geology Reviews 18, 237-250.
- 1680 Castiñeiras, P., 2005. Origen y evolución tectonotermal de las unidades de O Pino y
1681 Cariño (Complejos Alóctonos de Galicia). Nova Terra 28, 1-279.
- 1682 Castiñeiras, P., Villaseca González, C., Barbero González, L., Martín Romera, C., 2008.
1683 SHRIMP U-Pb zircon dating of anatexis in high-grade migmatite complexes of Central
1684 Spain: implications in the Hercynian evolution of Central Iberia. International Journal
1685 of Earth Sciences 97, 35-50.
- 1686 Castro, A., 1987a. Implicaciones de la zona Ossa-Morena y dominios equivalentes en el
1687 modelo geodinámico de la Cadena Hercinica europea. Estudios Geológicos 43, 249-
1688 260.
- 1689 Castro, A., 1987b. Los granitoides deformados de la banda del Guadamez (La Serena,
1690 Badajoz), in: Bea, F., Carnicero, A., Gonzalo, J.C., López-Plaza, M., Rodríguez Alonso,
1691 M.D. (Eds.), Geología de los granitoides y rocas asociadas del Macizo Hespérico. Libro
1692 Homenaje a L.C. García de Figuerola. Ed. Rueda, pp. 413-426.
- 1693 Castro, A., Fernández, C., de la Rosa, J., Moreno-Ventas, I., Rogers, G., 1996.
1694 Significance of MORB-derived Amphibolites from the Aracena Metamorphic Belt,
1695 Southwest Spain. Journal of Petrology 37, 235-260.
- 1696 Castro, A., Fernández, C., El-Hmidi, H., El-Biad, M., Díaz, M., de la Rosa, J., Stuart,
1697 F., 1999. Age constraints to the relationships between magmatism, metamorphism and
1698 tectonism in the Aracena metamorphic belt, southern Spain. International Journal of
1699 Earth Sciences 88, 26-37.

- 1700 Chacón, J., 1979. Estudio geológico del sector central del anticlinorio Portalegre-
1701 Badajoz-Córdoba (Macizo Ibérico Meridional). Universidad de Granada, p. 728.
- 1702 Chacón, J., Oliveira, V., Ribeiro, A., Oliveira, J.T., 1983. La estructura de la Zona de
1703 Ossa-Morena. Geologia de España. Libro Jubilar J.M. Rios 1, 490-504.
- 1704 Chichorro, M., 2006. Tectonic evolution of Montemor-o-Novo Shear zone (SW Ossa
1705 Morena Zone - Santiago do Escoural - Cabrela Área). Universidade de Évora, Évora,
1706 Portugal, p. 569.
- 1707 Chichorro, M., Pereira, M.F., Díaz-Azpiroz, M., Williams, I.S., Fernández, C., Pin, C.,
1708 Silva, J.B., 2008. Cambrian ensialic rift-related magmatism in the Ossa-Morena Zone
1709 (Évora-Aracena metamorphic belt, SW Iberian Massif): Sm-Nd isotopes and SHRIMP
1710 zircon U-Th-Pb geochronology. Tectonophysics 461, 91-113.
- 1711 Cohen, K.M., Finney, S.C., Gibbard, P.L., Fan, J.-X., 2013. The ICS International
1712 Chronostratigraphic Chart. Episodes 36, 199-204.
- 1713 Crespo-Blanc, A., 1992. Structure and kinematics of a sinistral transpressive suture
1714 between the Ossa-Morena and the South Portuguese Zones, South Iberian Massif.
1715 Journal of the Geological Society 149, 401-411.
- 1716 Crespo-Blanc, A., Orozco, M., 1988. The southern Iberian shear zone: a major
1717 boundary in the Hercynian folded belt. Tectonophysics 148, 221-227.
- 1718 Crespo-Blanc, A., Orozco, M., 1991. The boundary between the Ossa- Morena and
1719 Southportuguese zones (Southern Iberian Massif): A major suture in the European
1720 Hercynian Chain. Geologische Rundschau 80, 691-702.
- 1721 Dallmeyer, R.D., Quesada, C., 1992. Cadomian vs. Variscan evolution of the Ossa-
1722 Morena zone (SW Iberia): field and $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age constraints. Tectonophysics
1723 216, 339-364.

- 1724 Dallmeyer, R.D., Ribeiro, A., Marques, F., 1991. Polyphase Variscan emplacement of
1725 exotic terranes (Morais and Bragança Massifs) onto Iberian successions: Evidence from
1726 $^{40}\text{Ar}/^{39}\text{Ar}$ mineral ages. *Lithos* 27, 133-144.
- 1727 Dallmeyer, R.D., Fonseca, P.E., Quesada, C., Ribeiro, A., 1993. $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age
1728 constraints for the tectonothermal evolution of a Variscan suture in southwest Iberia.
1729 *Tectonophysics* 222, 177-194.
- 1730 Dallmeyer, R.D., García Casquero, J.L., Quesada, C., 1995. $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age
1731 constraints on the emplacement of the Burguillos del Cerro Igneous complex (Ossa-
1732 Morena zone, SW Iberia). *Boletín Geológico y Minero* 106, 203-214.
- 1733 Dallmeyer, R.D., Martínez Catalán, J.R., Arenas, R., Gil Ibarguchi, J.I., Gutiérrez-
1734 Alonso, G., Farias, P., Bastida, F., Aller, J., 1997. Diachronous Variscan tectonothermal
1735 activity in the NW Iberian Massif: Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ dating of regional fabrics.
1736 *Tectonophysics* 277, 307-337.
- 1737 Davies, H.L., Warren, R.G., 1988. Origin of eclogite-bearing, domed, layered
1738 metamorphic complexes ("core complexes") in the D'entrecasteaux Islands, Papua New
1739 Guinea. *Tectonics* 7, 1-21.
- 1740 De Jong, G., Dalstra, H., Boorder, H., Savage, J.F., 1991. Blue amphiboles, Variscan
1741 dformation and plate tectonics in the Beja Massif, South Portugal. *Comunicações dos*
1742 *Serviços Geologicos de Portugal* 77, 59-64.
- 1743 Dias, G., Simões, P.P., Ferreira, N., Leterrier, J., 2002. Mantle and crustal sources in the
1744 genesis of late-Hercynian granitoids (NW Portugal): geochemical and Sr-Nd isotopic
1745 constraints. *Gondwana Research* 5, 287-305.
- 1746 Dias, R., Coke, C., Moreira, N., 2010. Deformação Varisca heterogénea no eixo Marão
1747 – Foz Côa (autóctone da Zona Centro Ibérica); implicações para a estrutura regional. e-
1748 *Terra* 11, 1-4.
- 1749 Dias da Silva, I., 2014. Geología de las Zonas Centro Ibérica y Galicia-Trás-os-Montes
1750 en la parte oriental del Complejo de Morais, Portugal/España. *Nova Terra* 45, 1-424.

1751 Dias da Silva, Í., Linnemann, U., Hofmann, M., González-Clavijo, E., Díez-Montes, A.,
1752 Martínez Catalán, J.R., 2014. Detrital zircon and tectonostratigraphy of the
1753 Parautochthon under the Morais Complex (NE Portugal): implications for the Variscan
1754 accretionary history of the Iberian Massif. *Journal of the Geological Society*, doi:
1755 10.1144/jgs2014-005.

1756 Díaz-Alvarado, J., Fernández, C., Díaz-Azpiroz, M., Castro, A., Moreno-Ventas, I.,
1757 2012. Fabric evidence for granodiorite emplacement with extensional shear zones in the
1758 Variscan Gredos massif (Spanish Central System). *Journal of Structural Geology* 42,
1759 74-90.

1760 Díaz Azpiroz, M., Fernández, C., Castro, A., 2002. El evento de fusión parcial en el
1761 dominio continental de la banda metamórfica de Aracena (Macizo Ibérico meridional):
1762 condicionantes estructurales, geoquímicos e isotópicos. *Revista de la Sociedad*
1763 *Geológica de España* 15, 27-39.

1764 Díaz Azpiroz, M., Fernández, C., Castro, A., 2003. Estructura y evolución tectónica del
1765 dominio continental de la banda metamórfica de Aracena (Macizo Ibérico Meridional).
1766 *Revista de la Sociedad Geológica de España* 16, 167-184.

1767 Díaz Azpiroz, M., Castro, A., Fernández, C., López, S., Fernández Caliani, J.C.,
1768 Moreno-Ventas, I., 2004. The contact between the Ossa Morena and the South
1769 Portuguese zones. Characteristics and significance of the Aracena metamorphic belt, in
1770 its central sector between Aroche and Aracena (Huelva). *Journal of Iberian Geology* 30,
1771 23-51.

1772 Díaz Azpiroz, M., Fernández, C., Castro, A., El-Biad, M., 2006. Tectonometamorphic
1773 evolution of the Aracena metamorphic belt (SW Spain) resulting from ridge-trench
1774 interaction during Variscan plate convergence. *Tectonics* 25, TC1001.

1775 Díaz García, F., Arenas, R., Martínez Catalán, J.R., del Tanago, J.G., Dunning, G.R.,
1776 1999. Tectonic evolution of the Careon ophiolite (northwest Spain). A remnant of
1777 oceanic lithosphere in the Variscan belt. *The Journal of Geology* 107, 587-605.

1778 Díaz García, F., Sánchez Martínez, S., Castiñeiras, P., Fuenlabrada, J.M., Arenas, R.,
1779 2010. A peri-Gondwanan arc in NW Iberia. II: Assessment of the intra-arc
1780 tectonothermal evolution through U-Pb SHRIMP dating of mafic dykes. *Gondwana*
1781 *Research* 17, 352-362.

1782 Díez Balda, M.A., 1986. El Complejo Esquisto-Grauváquico, las series paleozoicas y la
1783 estructura hercínica al Sur de Salamanca. Universidad de Salamanca, Salamanca, p.
1784 162.

1785 Díez Balda, M.A., Martínez Catalán, J.R., Ayarza, P., 1995. Syn-collisional extensional
1786 collapse parallel to the orogenic trend in a domain of steep tectonics - The Salamanca
1787 detachment zone (Central Iberian Zone, Spain). *Journal of Structural Geology* 17, 163-
1788 182.

1789 Díez Fernández, R., 2011. Evolución estructural y cinemática de una corteza continental
1790 subducida: la Unidad de Malpica-Tui (NO del Macizo Ibérico). *Nova Terra* 40, 1-228.

1791 Díez Fernández, R., Martínez Catalán, J.R., 2012. Stretching lineations in high-pressure
1792 belts: the fingerprint of subduction and subsequent events (Malpica–Tui complex, NW
1793 Iberia). *Journal of the Geological Society* 169, 531-543.

1794 Díez Fernández, R., Arenas, R., 2015. The Late Devonian Variscan suture of the Iberian
1795 Massif: A correlation of high-pressure belts in NW and SW Iberia. *Tectonophysics* 654,
1796 96-100.

1797 Díez Fernández, R., Arenas, R., 2016. Reply to Comment on “The Late Devonian
1798 Variscan suture of the Iberian Massif: A correlation of high-pressure belts in NW and
1799 SW Iberia”. *Tectonophysics* 670, 155-160.

1800 Díez Fernández, R., Martínez Catalán, J.R., Gerdes, A., Abati, J., Arenas, R.,
1801 Fernández-Suárez, J., 2010. U-Pb ages of detrital zircons from the Basal allochthonous
1802 units of NW Iberia: Provenance and paleoposition on the northern margin of Gondwana
1803 during the Neoproterozoic and Paleozoic. *Gondwana Research* 18, 385-399.

1804 Díez Fernández, R., Martínez Catalán, J.R., Arenas, R., Abati, J., 2011. Tectonic
 1805 evolution of a continental subduction-exhumation channel: Variscan structure of the
 1806 basal allochthonous units in NW Spain. *Tectonics* 30, TC3009.

1807 Díez Fernández, R., Martínez Catalán, J.R., Arenas, R., Abati, J., 2012a. The onset of
 1808 the assembly of Pangaea in NW Iberia: Constraints on the kinematics of continental
 1809 subduction. *Gondwana Research* 22, 20-25.

1810 Díez Fernández, R., Martínez Catalán, J.R., Arenas, R., Abati, J., Gerdes, A.,
 1811 Fernández-Suárez, J., 2012b. U–Pb detrital zircon analysis of the lower allochthon of
 1812 NW Iberia: age constraints, provenance and links with the Variscan mobile belt and
 1813 Gondwanan cratons. *Journal of the Geological Society* 169, 655-665.

1814 Díez Fernández, R., Martínez Catalán, J.R., Gómez Barreiro, J., Arenas, R., 2012c.
 1815 Extensional flow during gravitational collapse: a tool for setting plate convergence
 1816 (Padrón migmatitic dome, Variscan belt, NW Iberia). *The Journal of Geology* 120, 83-
 1817 103.

1818 Díez Fernández, R., Foster, D.A., Gómez Barreiro, J., Alonso-García, M., 2013a.
 1819 Rheological control on the tectonic evolution of a continental suture zone: the Variscan
 1820 example from NW Iberia (Spain). *International Journal of Earth Sciences* 102, 1305-
 1821 1319.

1822 Díez Fernández, R., Gómez Barreiro, J., Martínez Catalán, J.R., Ayarza, P., 2013b.
 1823 Crustal thickening and attenuation as revealed by regional fold interference patterns:
 1824 Ciudad Rodrigo basement area (Salamanca, Spain). *Journal of Structural Geology* 46,
 1825 115-128.

1826 Díez Fernández, R., Pereira, M.F., Foster, D.A., 2015. Peralkaline and alkaline
 1827 magmatism of the Ossa-Morena zone (SW Iberia): Age, source, and implications for the
 1828 Paleozoic evolution of Gondwanan lithosphere. *Lithosphere* 7, 73-90.

1829 Díez Montes, A., 2007. La geología del Dominio “Ollo de Sapo” en las comarcas de
 1830 Sanabria y Terra do Bolo. *Nova Terra* 34, 1-494.

- 1831 Doglioni, C., Carminati, E., Cuffaro, M., 2006. Simple kinematics of subduction zones.
1832 International Geology Review 48, 479-493.
- 1833 Doglioni, C., Carminati, E., Cuffaro, M., Scrocca, D., 2007. Subduction kinematics and
1834 dynamic constraints. Earth-Science Reviews 83, 125-175.
- 1835 Dunning, G.R., Díez Montes, A., Matas, J., Martín Parra, L.M., Almarza, J., Donaire,
1836 M., 2002. Geocronología U/Pb del volcanismo ácido y granitoides de la Faja Pirítica
1837 Ibérica (Zona Surportuguesa). Geogaceta 32, 127-130.
- 1838 Dupuy, C., Dostal, J., Bard, J.P., 1979. Trace element geochemistry of paleozoic
1839 amphibolites from S. W. Spain. Tschermaks Petr. Mitt. 26, 87-93.
- 1840 Eden, C.P., Andrews, J.R., 1990. Middle to upper Devonian melanges in SW Spain and
1841 their relationship to the Meneage formation in south Cornwall. Proceedings of the
1842 Ussher Society 7, 217-222.
- 1843 Eguíluz, L., 1987. Petrogénesis de rocas ígneas y metamórficas en el Antiforme
1844 Burguillos-Monesterio, Macizo Ibérico Meridional. Universidad del País Vasco, p. 694.
- 1845 Eguiluz, L., Abalos, B., Gil Ibarguchi, J.I., 1990. Eclogitas de la banda de Cizalla
1846 Badajoz-Cordoba (Suroeste de España). Datos petrográficos y significado geodinámico.
1847 Geogaceta 7, 28-31.
- 1848 Escuder Viruete, J.E., Arenas, R., Martínez Catalán, J.R., 1994. Tectonothermal
1849 evolution associated with Variscan crustal extension in the Tormes gneiss dome (NW
1850 Salamanca, Iberian massif, Spain). Tectonophysics 238, 117-138.
- 1851 Escuder Viruete, J., Hernáiz Huerta, P.P., Valverde-Vaquero, P., Rodríguez Fernández,
1852 R., Dunning, G., 1998. Variscan syncollisional extension in the Iberian Massif:
1853 structural, metamorphic and geochronological evidence from the Somosierra sector of
1854 the Sierra de Guadarrama (Central Iberian Zone, Spain). Tectonophysics 290, 87-109.

- 1855 Escuder Viruete, J.E., Indares, A., Arenas, R., 2000. P-T paths derived from garnet
1856 growth zoning in an extensional setting: an example from the Tormes Gneiss Dome
1857 (Iberian Massif, Spain). *Journal of Petrology* 41, 1489-1515.
- 1858 Expósito Ramos, I., 2005. Evolución estructural de la mitad septentrional de la Zona de
1859 Ossa-Morena y su relación con el límite Zona Ossa-Morena/Zona Centroibérica. *Nova*
1860 *Terra* 27, 1-286.
- 1861 Expósito, I., Simancas, J.F., González Lodeiro, F., Azor, A., Martínez Poyatos, D.J.,
1862 2002. La estructura de la mitad septentrional de la Zona de Ossa-Morena: deformación
1863 en el bloque inferior de un cabalgamiento cortical de evolución compleja. *Revista de la*
1864 *Sociedad Geológica de España* 15, 3-14.
- 1865 Farias, P., Gallastegui, G., González Lodeiro, F., Marquínez García, J., Martín-Parra,
1866 L.M., Martínez Catalán, J.R., Pablo Maciá, J.G.d., Rodríguez-Fernández, L.R., 1987.
1867 Aportaciones al conocimiento de la litoestratigrafía y estructura de Galicia Central.
1868 *Mem. Museo e Lab. Miner. Geol., Fac. Ciencias, Univ. Porto* 1, 411-431.
- 1869 Faure, M., Bé Mézème, E., Cocherie, A., Rossi, P., Chemenda, A., Boutelier, D., 2008.
1870 Devonian geodynamic evolution of the Variscan Belt, insights from the French Massif
1871 Central and Massif Armoricain. *Tectonics* 27, TC2005.
- 1872 Febrel, T., Saenz de Santa María, J., 1964. El Devoniano al S del Batolito de Los
1873 Pedroches en las provincias de Córdoba y Badajoz [Nota preliminar de las Hojas N° 856
1874 (Maguilla) y N° 857 (Valsequillo)]. *Notas y Comunicaciones del IGME* 73, 51-60.
- 1875 Fernández, F.J., Díaz García, F., Marquínez, J., 2011. Kinematics of the Forcarei
1876 Synform (NW Iberian Variscan belt), in: Poblet, J., Lisle, R.J. (Eds.), *Kinematic*
1877 *Evolution and Structural Styles of Fold-and-Thrust Belts*. Geological Society, London,
1878 *Special Publications*, pp. 185-201, doi: 110.1144/sp1349.1110.
- 1879 Fernández-Suárez, J., Corfu, F., Arenas, R., Marcos, A., Martínez Catalán, J.R., Díaz
1880 García, F., Abati, J., Fernández, F.J., 2002. U-Pb evidence for a polyorogenic evolution
1881 of the HP-HT units of the NW Iberian Massif. *Contributions to Mineralogy and*
1882 *Petrology* 143, 236-253.

- 1883 Fernández-Suárez, J., Díaz García, F., Jeffries, T.E., Arenas, R., Abati, J., 2003.
1884 Constraints on the provenance of the uppermost allochthonous terrane of the NW
1885 Iberian Massif: inferences from detrital zircon U-Pb ages. *Terra Nova* 15, 138-144.
- 1886 Fernández-Suárez, J., Arenas, R., Abati, J., Martínez Catalán, J.R., Whitehouse, M.J.,
1887 Jeffries, T.E., 2007. U-Pb chronometry of polymetamorphic high-pressure granulites:
1888 An example from the allochthonous terranes of the NW Iberian Variscan belt.
1889 *Geological Society of America Memoirs* 200, 469-488.
- 1890 Fonseca, P.E., 2005. O terreno acrecionário do Pulo do Lobo: implicações
1891 geodinâmicas da sutura com a Zona de Ossa-Morena (SW da Cadeia Varisca Ibérica).
1892 *Cadernos del Laboratorio Xeolóxico de Laxe* 30, 213-222.
- 1893 Fonseca, P., Ribeiro, A., 1993. Tectonics of the Beja-Acebuches ophiolite: A major
1894 suture in the Iberian Variscan Foldbelt. *Geologische Rundschau* 82, 440-447.
- 1895 Fonseca, P., Araújo, A., Leal, N., Munhá, J.M., 1993. Variscan Glaucophane Eclogites
1896 in the Ossa Morena Zone. XII Reunião de Geologia do Oeste Peninsular. Évora, 20-24
1897 de Setembro de 1993, *Terra Abstracts*, supplement n. 6 to *Terra Nova* 5, 11-12.
- 1898 Fonseca, P., Munhá, J., Pedro, J., Rosas, F., Moita, P., Araújo, A., Leal, N., 1999.
1899 Variscan ophiolites and high-pressure metamorphism in southern Iberia. *Ofioliti* 24,
1900 259-268.
- 1901 Fonseca, P.E., Fonseca, M.M., Munhá, J.M., 2004. Ocorrência de aragonite em
1902 mármore da região de Alvito-Viana do Alentejo (Zona de Ossa-Morena): significado
1903 geodinâmico. *Cuadernos do Laboratorio Xeolóxico de Laxe* 29, 79-96.
- 1904 Franke, W., 2000. The mid-European segment of the Variscides: Tectonostratigraphic
1905 units, terrane boundaries and plate tectonic evolution, in: Franke, W., Haak, V., Oncken,
1906 O., Tanner, D. (Eds.), *Orogenic Processes: Quantification and Modelling in the*
1907 *Variscan Belt*. Geological Society, London, Special Publications, pp. 35-61, doi:
1908 10.1144/GSL.SP.2000.1179.1101.1105.

- 1909 Franke, W., 2006. The Variscan orogen in Central Europe: construction and collapse,
1910 in: Gee, D.G., Stephenson, R.A. (Eds.), European lithosphere dynamics. Geological
1911 Society of London, Memoirs, pp. 333-343, doi: 10.1144/GSL.MEM.2006.032.01.20.
- 1912 Franke, W., 2014. Topography of the Variscan orogen in Europe: failed–not collapsed.
1913 International Journal of Earth Sciences 103, 1471-1499.
- 1914 Gabaldón, V., Quesada, C., 1986. Exemples de bassins houillers limniques du sud-ouest
1915 de la Péninsule Ibérique: évolution sédimentaire et contrôle structural. Mémoires de la
1916 Société Géologique de France 149, 27-36.
- 1917 Galindo, C., Casquet, C., Portugal Ferreira, M.R., Regencio Macedo, C.A., 1986. O
1918 Complexo Plutónico de Táliga-Barcarrota - Um Complexo intrusivo com Idades
1919 Caledónica e Hercínica. Maleo. Boletim Informativo da Sociedade Geológica de
1920 Portugal. II Congresso Nacional de Geologia de Portugal, Lisboa Volumen 2, numero
1921 13, 22.
- 1922 Galindo, C., Casquet, C., Portugal Ferreira, M.R., Regencio Macedo, C.A., 1987.
1923 Geocronología del Complejo plutónico Táliga-Barcarrota (CPTB). Badajoz, España, in:
1924 Bea, F., Carnicero, A., Gonzalo, J.C., López-Plaza, M., Rodríguez Alonso, M.D. (Eds.),
1925 Geología de los granitoides y rocas asociadas del Macizo Hespérico. Libro Homenaje a
1926 L.C. García de Figuerola. Ed. Rueda, pp. 385-392.
- 1927 García-López, S., Sanz López, J., Pardo Alonso, M.V., 1999. Conodontos
1928 (bioestratigrafía, biofacies y paleotemperaturas) de los sinclinales de Almadén y
1929 Guadalmez (Devónico-Carbonífero Inferior), Zona Centroibérica meridional, España.
1930 Revista española de paleontología (Homenaje Prof. J. Truyols), 161-172.
- 1931 García Alcalde, J.L., Arbizu, M.A., Pardo Alonso, M.V., García López, S., 1984. El
1932 límite Devónico-Carbonífero en el área de Guadalmez-Santa Eufemia (Provincias de
1933 Ciudad Real y Córdoba, Sierra Morena, España). I Congreso Geológico de España,
1934 Segovia, Resúmenes 1, 421-430.

- 1935 García Casquero, J.L., Priem, H.N.A., Boelrijk, N.A.I.M., Chacon, J., 1988. Isotopic
1936 dating of the mylonitization of the Azuaga Group in the Badajóz-Córdoba belt, SW
1937 Spain. *Geologische Rundschau* 77, 483-489.
- 1938 Garrote, A., Broutin, J., 1979. Le bassin toumaisien de Benajarafe (Province de
1939 Cordoue, Espagne). *Géologie et premières données paléobotaniques et palynologiques*.
1940 *Comptes Rendus du 104 Congrès national des Sociétés savantes (Bordeaux)* 1, 175-184.
- 1941 Gates, A.E., Simpson, C., Glover, L., 1986. Appalachian Carboniferous dextral strike-
1942 slip faults - An example from Brookneal, Virginia. *Tectonics* 5, 119-133.
- 1943 Giese, W., Hoegen, R.V., Hollmann, G., Walter, R., 1994. Geology of the southwestern
1944 Iberian Meseta I. The Palaeozoic of the Ossa Morena Zone north and south of the
1945 Olivenza-Monesterio Anticline (Huelva province, SW Spain). *Neues Jahrbuch für*
1946 *Geologie und Paläontologie* 192, 293-331.
- 1947 Giese, U., Hoymann, K.-H., Glodny, J., Kramm, U., Dallmeyer, R.D., 1999. Age
1948 constraints for the tectonometamorphic evolution of the Pulo do Lobo zone in SW
1949 Spain. *Zeitschrift der Deutschen Geologischen Gesellschaft* 150, 565-582.
- 1950 Gil Ibarguchi, J.I., 1995. Petrology of jadeite metagranite and associated orthogneiss
1951 from the Malpica-Tuy allochthon (Northwest Spain). *European Journal of Mineralogy*
1952 7, 403-415.
- 1953 Gil Ibarguchi, J.I., Ortega Gironés, E., 1985. Petrology, structure and geotectonic
1954 implications of glaucophane-bearing eclogites and related rocks from the Malpica Tuy
1955 (MT) Unit, Galicia, Northwest Spain. *Chemical Geology* 50, 145-162.
- 1956 Gil Ibarguchi, J.I., Mendiá, M., Girardeau, J., Peucat, J.J., 1990. Petrology of eclogites
1957 and clinopyroxene garnet metabasites from the Cabo-Ortega complex (Northwestern
1958 Spain). *Lithos* 25, 133-162.
- 1959 Girardeau, J., Ibarguchi, J.I.G., 1991. Pyroxenite-Rich Peridotites of the Cabo Ortega
1960 Complex (Northwestern Spain): Evidence for Large-Scale Upper-Mantle Heterogeneity.
1961 *Journal of Petrology Special Volume*, 135-154.

- 1962 Girardeau, J., Gil Ibarguchi, J.I., Ben Jamaa, N., 1989. Evidence for a heterogeneous
1963 upper mantle in the Cabo Ortegal Complex, Spain. *Science* 245, 1231-1233.
- 1964 Gladney, E.R., Braid, J.A., Murphy, J.B., Quesada, C., McFarlane, C.R.M., 2014. U–Pb
1965 geochronology and petrology of the late Paleozoic Gil Marquez pluton: magmatism in
1966 the Variscan suture zone, southern Iberia, during continental collision and the
1967 amalgamation of Pangea. *International Journal of Earth Sciences* 103, 1433-1451.
- 1968 Gómez Barreiro, J., 2007. La Unidad de Fornás: Evolución tectonometamórfica del SO
1969 del Complejo de Órdenes. *Nova Terra* 32, 1-291.
- 1970 Gómez Barreiro, J., Wijbrans, J.R., Castineiras, P., Martínez Catalán, J.R., Arenas, R.,
1971 Díaz García, F., Abati, J., 2006. $^{40}\text{Ar}/^{39}\text{Ar}$ laserprobe dating of mylonitic fabrics in a
1972 polyorogenic terrane of NW Iberia. *Journal of the Geological Society* 163, 61-73.
- 1973 Gómez Barreiro, J., Martínez Catalán, J.R., Arenas, R., Castiñeiras, P., Abati, J., Díaz
1974 García, F., Wijbrans, J.R., 2007. Tectonic evolution of the upper allochthon of the
1975 Órdenes Complex (northwestern Iberian Massif): structural constraints to a
1976 polyorogenic peri- Gondwanan terrane, in: Linnemann, U., Nance, R.D., Kraft, P.,
1977 Zulauf, G. (Eds.), *The evolution of the Rheic Ocean: from Avalonian-Cadomian active*
1978 *margin to Alleghenian-Variscan collision*. Geological Society of America Special
1979 Paper, pp. 315-332, doi: 310.1130/2007.2423(1115).
- 1980 Gómez Barreiro, J., Martínez Catalán, J.R., Díez Fernández, R., Arenas, R., Díaz
1981 García, F., 2010a. Upper crust reworking during gravitational collapse: The Bemibre-
1982 Pico Sacro detachment system (NW Iberia). *Journal of the Geological Society* 167, 769-
1983 784.
- 1984 Gómez Barreiro, J., Martínez Catalán, J.R., Prior, D., Wenk, H.-R., Vogel, S., Díaz
1985 García, F., Arenas, R., Sánchez Martínez, S., Lonardelli, I., 2010b. Fabric Development
1986 in a Middle Devonian Intraoceanic Subduction Regime: The Careón Ophiolite
1987 (Northwest Spain). *The Journal of Geology* 118, 163-186.
- 1988 González Clavijo, E., Martínez Catalán, J.R., 2002. Stratigraphic record of preorogenic
1989 to synorogenic sedimentation, and tectonic evolution of imbricate units in the Alcañices

- 1990 synform (northwestern Iberian Massif), in: Martínez Catalán, J.R., Hatcher, R.D.,
1991 Arenas, R., Díaz García, F. (Eds.), Variscan-Appalachian Dynamics: The building of
1992 the late Paleozoic basement. Geological Society of America Special Paper, pp. 17-35,
1993 doi: 10.1130/1130-8137-2364-1137.1117.
- 1994 González Cuadra, P., 2007. La Unidad de Corredoiras (Complejo de Órdenes, Galicia):
1995 Evolución estructural y Metamórfica. Nova Terra 33, 1-254.
- 1996 González del Tánago, J., 1995. El núcleo metamórfico de Sierra Albarrana y su campo
1997 de pegmatitas graníticas asociado, Macizo Ibérico, Córdoba. Nova Terra 12, 1-713.
- 1998 González del Tánago, J., Arenas, R., 1991. Anfibolitas granatíferas de Sierra Albarrana,
1999 Córdoba : termobarometría e implicaciones para el desarrollo del metamorfismo
2000 regional. Revista de la Sociedad Geológica de España 4, 251-269.
- 2001 Gordon, S.M., Little, T.A., Hacker, B.R., Bowring, S.A., Korchinski, M., Baldwin, S.L.,
2002 Kylander-Clark, A.R.C., 2012. Multi-stage exhumation of young UHP–HP rocks:
2003 Timescales of melt crystallization in the D’Entrecasteaux Islands, southeastern Papua
2004 New Guinea. Earth and Planetary Science Letters 351–352, 237-246.
- 2005 Gutiérrez-Alonso, G., Murphy, J.B., Fernández-Suárez, J., Hamilton, M.A., 2008.
2006 Rifting along the northern Gondwana margin and the evolution of the Rheic Ocean: A
2007 Devonian age for the El Castillo volcanic rocks (Salamanca, Central Iberian Zone).
2008 Tectonophysics 461, 157-165.
- 2009 Gutiérrez-Alonso, G., Fernández-Suárez, J., Jeffries, T.E., Johnston, S.T., Pastor-Galán,
2010 D., Murphy, J.B., Franco, M.P., Gonzalo, J.C., 2011. Diachronous post-orogenic
2011 magmatism within a developing orocline in Iberia, European Variscides. Tectonics 30,
2012 TC5008.
- 2013 Gutiérrez-Alonso, G., Collins, A.S., Fernández-Suárez, J., Pastor-Galán, D., González-
2014 Clavijo, E., Jourdan, F., Weil, A.B., Johnston, S.T., 2015. Dating of lithospheric
2015 buckling: $^{40}\text{Ar}/^{39}\text{Ar}$ ages of syn-orocline strike–slip shear zones in northwestern Iberia.
2016 Tectonophysics 643, 44-54.

2017 Hatcher, R.D., 1978. Tectonics of the western Piedmont and Blue Ridge, southern
2018 Appalachians: Review and Speculation. *American Journal of Science* 278, 276-304.

2019 Hatcher, R.D., 2002. Alleghanian (Appalachian) orogeny, a product of zipper tectonics:
2020 Rotational transpressive continent-continent collision and closing of ancient oceans
2021 along irregular margins, in: Martínez Catalán, J.R., Hatcher, R.D., Arenas, R., Díaz
2022 García, F. (Eds.), *Variscan-Appalachian Dynamics: the Building of the Late Paleozoic*
2023 *Basement. Geological Society of America Special Paper*, pp. 199-208, doi:
2024 110.1130/1130-8137-2364-1137.1199.

2025 Herranz, P., 1985. El Precámbrico y su cobertera paleozoica en la región centro-oriental
2026 de la provincia de Badajoz. *Universidad Complutense de Madrid, Madrid*, p. 1220.

2027 Iglesias Ponce de Leon, M., Choukroune, P., 1980. Shear zones in the iberian arc.
2028 *Journal of Structural Geology* 2, 63-68.

2029 Jesus, A.P., Munhá, J., Mateus, A., Tassinari, C., Nutman, A.P., 2007. The Beja
2030 Layered Gabbroic Sequence (Ossa-Morena Zone, Southern Portugal): geochronology
2031 and geodynamic implications. *Geodinamica Acta* 20, 139-157.

2032 Kroner, U., Romer, R.L., 2013. Two plates — Many subduction zones: The Variscan
2033 orogeny reconsidered. *Gondwana Research* 24, 298-329.

2034 Lallemand, S., Heuret, A., Boutelier, D., 2005. On the relationships between slab dip,
2035 back-arc stress, upper plate absolute motion and crustal nature in subduction zones.
2036 *Geochemistry Geophysics Geosystems* 6, Q09006.

2037 Larrea, F.J., Carracedo, M., Alonso, A., Ortega, L.A., Menéndez, M., 1999. Granitoides
2038 postcolisionales emplazados en situaciones extensionales: el stock de Santa Elena (zona
2039 Centroibérica, España). *XV reun. geol. oeste peninsular, Extended Abstracts*, 147-157.

2040 Leal, N., Pedro, J., Moita, P., Fonseca, P., Araújo, A., Munhá, J., Araújo, A., Pereira,
2041 M.F., 1997. *Metamorfismo nos sectores meridionais da Zona de Ossa-Morena:*
2042 *Atualização dos Conhecimentos, Estudo sobre a geologia de Ossa-Morena (Maciço*

2043 Ibérico). Homenagem ao Prof. Francisco Gonçalves. Universidade de Évora, Évora, pp.
 2044 119-131.

2045 Lefort, J.P., Max, M.D., Roussel, J., 1988. Geophysical evidence for the location of the
 2046 NW boundary of Gondwanaland and its relationship with two older satellite sutures.
 2047 Geological Society, London, Special Publications 38, 49-60.

2048 Leistel, J.M., Marcoux, E., Thiéblemont, D., Quesada, C., Sánchez, A., Almodóvar,
 2049 G.R., Pascual, E., Sáez, R., 1997. The volcanic-hosted massive sulphide deposits of the
 2050 Iberian Pyrite Belt Review and preface to the Thematic Issue. Mineral. Deposita 33, 2-
 2051 30.

2052 Lima, S.M., Corfu, F., Neiva, A.M.R., Ramos, J.M.F., 2011. Dissecting complex
 2053 magmatic processes: an in-depth U-Pb study of the Pavia pluton, Ossa-Morena Zone,
 2054 Portugal. Journal of Petrology 53, 1887-1911.

2055 Lima, S.M., Neiva, A.M.R., Ramos, J.M.F., 2013. Adakitic-like magmatism in western
 2056 Ossa-Morena Zone (Portugal): Geochemical and isotopic constraints of the Pavia
 2057 pluton. Lithos 160-161, 98-116.

2058 Liñán, E., Quesada, C., 1990. Ossa-Morena Zone: 2. Stratigraphy. Rift phase, in:
 2059 Dallmeyer, R.D., Martínez García, E. (Eds.), Pre-Mesozoic Geology of Iberia. Springer-
 2060 Verlag, Berlin, Germany, pp. 259–266.

2061 Loeschke, J., 1983. Igneous and pyroclastic rocks in Devonian and Lower
 2062 Carboniferous strata of the Cantabrian Mountains (NW Spain). Neues Jahrbuch für
 2063 Geologie und Paläeontologie. Monatshefte 7, 419-439.

2064 López-Carmona, A., Abati, J., Reche, J., 2010. Petrologic modeling of chloritoid-
 2065 glaucophane schists from the NW Iberian Massif. Gondwana Research 17, 377-391.

2066 López-Carmona, A., Abati, J., Pitra, P., Lee, J.W., 2014. Retrogressed lawsonite
 2067 blueschists from the NW Iberian Massif: P–T–t constraints from thermodynamic
 2068 modelling and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology. Contributions to Mineralogy and Petrology
 2069 167, 1-20.

2070 López-Guijarro, R., Armendariz, M., Quesada, C., Fernández-Suárez, J., Murphy, J.B.,
 2071 Pin, C., Bellido, F., 2008. Ediacaran-Palaeozoic tectonic evolution of the Ossa Morena
 2072 and Central Iberian zones (SW Iberia) as revealed by Sm-Nd isotope systematics.
 2073 *Tectonophysics* 461, 202-214.

2074 Llana-Fúnez, S., 2002. Quartz c-axis texture mapping of a Variscan regional foliation
 2075 (Malpica-Tui Unit, NW Spain). *Journal of Structural Geology* 24, 1299-1312.

2076 Macaya, J., González-Lodeiro, F., Martínez-Catalán, J.R., Alvarez, F., 1991.
 2077 Continuous deformation, ductile thrusting and backfolding of cover and basement in the
 2078 Sierra de Guadarrama, Hercynian orogen of central Spain. *Tectonophysics* 191, 291-
 2079 309.

2080 Marcos, A., Farias, P., 1999. La estructura de las láminas inferiores del Complejo de
 2081 Cabo Ortegal y su autóctono relativo (Galicia, NW de España). *Trabajos de Geología*
 2082 21, 201-220.

2083 Marcos, A., Marquínez, J., Pérez-Estaún, A., Pulgar, J.A., Bastida, F., 1984. Nuevas
 2084 aportaciones al conocimiento de la evolución tectonometamórfica del Complejo de
 2085 Cabo Ortegal (NW de España). *Cuadernos do Laboratorio Xeolóxico de Laxe* 7, 125-
 2086 137.

2087 Marques, F.O., Ribeiro, A., Munhá, J.M., 1996. Geodynamic evolution of the
 2088 Continental Allochthonous Terrane (CAT) of the Bragança Nappe Complex, NE
 2089 Portugal. *Tectonics* 15, 747-762.

2090 Marquínez García, J.L., 1984. La geología del área esquistosa de Galicia Central
 2091 (Cordillera Herciniana, NW de España). *Memorias del Instituto Geológico y Minero de*
 2092 *España* 100, 1-231.

2093 Martin-Izard, A., Arias, D., Arias, M., Gumiel, P., Sanderson, D.J., Castañón, C.,
 2094 Sanchez, J., 2016. Ore deposit types and tectonic evolution of the Iberian Pyrite Belt:
 2095 From transtensional basins and magmatism to transpression and inversion tectonics. *Ore*
 2096 *Geology Reviews* 79, 254-267.

2097 Martín Parra, L.M., González Lodeiro, F., Martínez Poyatos, D., Matas, J., 2006. The
 2098 Puente Génave–Castelo de Vide Shear Zone (southern Central Iberian Zone, Iberian
 2099 Massif): geometry, kinematics and regional implications. *Bulletin de la Société*
 2100 *Géologique de France* 177, 191-202.

2101 Martínez Catalán, J.R., 1990. A non-cylindrical model for the northwest Iberian
 2102 allochthonous terranes and their equivalents in the Hercynian belt of Western Europe.
 2103 *Tectonophysics* 179, 253-272.

2104 Martínez Catalán, J.R., 2011. Are the oroclinal of the Variscan belt related to late
 2105 Variscan strike-slip tectonics? *Terra Nova* 23, 241-247.

2106 Martínez Catalán, J.R., 2012. The Central Iberian arc, an orocline centered in the
 2107 Iberian Massif and some implications for the Variscan belt. *International Journal of*
 2108 *Earth Sciences* 101, 1299-1314.

2109 Martínez Catalán, J.R., Arenas, R., Díaz García, F., Rubio Pascual, F.J., Abati, J.,
 2110 Marquínez García, J., 1996. Variscan exhumation of a subducted paleozoic continental
 2111 margin: The basal units of the Ordenes Complex, Galicia, NW Spain. *Tectonics* 15,
 2112 106-121.

2113 Martínez Catalán, J.R., Arenas, R., Díaz García, F., Abati, J., 1997. Variscan
 2114 accretionary complex of northwest Iberia: Terrane correlation and succession of
 2115 tectonothermal events. *Geology* 25, 1103-1106.

2116 Martínez Catalán, J.R., Díaz García, F., Arenas, R., Abati, J., Castineiras, P., González
 2117 Cuadra, P., Gómez Barreiro, J., Rubio Pascual, F.J., 2002. Thrust and detachment
 2118 systems in the Ordenes Complex (northwestern Spain): Implications for the Variscan-
 2119 Appalachian geodynamics, in: Martínez Catalán, J.R., Hatcher, R.D., Arenas, R., Díaz
 2120 García, F. (Eds.), *Variscan-Appalachian Dynamics: the Building of the Late Paleozoic*
 2121 *Basement. Geological Society of America Special Paper*, pp. 163-182, doi:
 2122 10.1130/1130-8137-2364-1137.1163.

2123 Martínez Catalán, J.R., Arenas, R., Díez Balda, M.A., 2003. Large extensional
 2124 structures developed during emplacement of a crystalline thrust sheet: the Mondoñedo
 2125 nappe (NW Spain). *Journal of Structural Geology* 25, 1815-1839.

2126 Martínez Catalán, J.R., Fernández-Suárez, J., Jenner, G.A., Belousova, E., Díez Montes,
 2127 A., 2004. Provenance constraints from detrital zircon U-Pb ages in the NW Iberian
 2128 Massif: implications for Palaeozoic plate configuration and Variscan evolution. *Journal*
 2129 *of the Geological Society* 161, 463-476.

2130 Martínez Catalán, J.R., Arenas, R., Díaz García, F., Gómez Barreiro, J., González
 2131 Cuadra, P., Abati, J., Castiñeiras, P., Fernández-Suárez, J., Sánchez Martínez, S.,
 2132 Andonaegui, P., González Clavijo, E., Díez Montes, A., Rubio Pascual, F.J., Valle
 2133 Aguado, B., 2007. Space and time in the tectonic evolution of the northwestern Iberian
 2134 Massif. Implications for the Variscan belt, in: Hatcher, R.D., Carlson, M.P., McBride,
 2135 J.H., Martínez Catalán, J.R. (Eds.), *4-D Framework of Continental Crust*. Geological
 2136 Society of America Memoir, Boulder, Colorado, pp. 403-423.

2137 Martínez Catalán, J.R., Rubio Pascual, F.J., Díez Montes, A., Díez Fernández, R.,
 2138 Gómez Barreiro, J., Dias da Silva, Í., González Clavijo, E., Ayarza, P., Alcock, J.E.,
 2139 2014. The late Variscan HT/LP metamorphic event in NW and Central Iberia:
 2140 relationships to crustal thickening, extension, orocline development and crustal
 2141 evolution. *Geological Society, London, Special Publications* 405, 225-247.

2142 Martínez Catalán, J.R., González Clavijo, E., Meireles, C., Díez Fernández, R., Bevis,
 2143 J., 2016. Relationships between syn-orogenic sedimentation and nappe emplacement in
 2144 the hinterland of the Variscan belt in NW Iberia deduced from detrital zircons.
 2145 *Geological Magazine* 153, 38-60.

2146 Martínez Poyatos, D.J., 2002. Estructura del borde meridional de la Zona Centroibérica
 2147 y su relación con el contacto entre las Zonas Centroibérica y de Ossa-Morena. *Nova*
 2148 *Terra* 18, 1-295.

2149 Martínez Poyatos, D., Simancas, J.F., Azor, A., González Lodeiro, F., 1998. Evolution
2150 of a Carboniferous piggyback basin in the southern Central Iberian Zone (Variscan Belt,
2151 SE Spain). *Bulletin de la Société Géologique de France* 169, 573-578.

2152 Martínez Poyatos, D., González Lodeiro, F., Azor, A., Simancas, J.F., 2001. La
2153 estructura de la Zona Centroibérica en la región de Los Pedroches (Macizo Ibérico
2154 meridional). *Revista de la Sociedad Geológica de España* 14, 147-160.

2155 Martínez Poyatos, D., Carbonell, R., Palomeras, I., Simancas, J.F., Ayarza, P., Martí,
2156 D., Azor, A., Jabaloy, A., González Cuadra, P., Tejero, R., Martín Parra, L.M., Matas,
2157 J., González Lodeiro, F., Pérez-Estaún, A., García Lobón, J.L., Mansilla, L., 2012.
2158 Imaging the crustal structure of the Central Iberian Zone (Variscan Belt): The
2159 ALCUDIA deep seismic reflection transect. *Tectonics* 31, TC3017.

2160 Martínez Poza, A.I., Martínez Poyatos, D., Simancas, F., Azor, A., 2012. La estructura
2161 varisca de la Unidad del Pulo do Lobo (SO del Macizo Ibérico) en las transversales de
2162 Aroche y Rosal de la Frontera (Huelva). *Geogaceta* 52, 21-24.

2163 Mata, J., Munhá, J., 1986. Geodynamic significance of high-grade metamorphic rocks
2164 from Degolados-Campo Maior (Tomar-Badajoz-Córdoba shear zone). *Maleo* 2, 28.

2165 Matas, J., Martín Parra, L., Montes Santiago, M., 2014. Un olistostroma con cantos y
2166 bloques del Paleozoico Inferior en la cuenca carbonífera del Guadalquivir (Córdoba).
2167 Parte I: Estratigrafía y marco geodinámico varisco. *Revista de la Sociedad Geológica de*
2168 *España* 27, 11-26.

2169 Matas, J., Martín Parra, L.M., Martínez Poyatos, D., 2015a. Mapa y Memoria del Mapa
2170 Geológico Nacional a escala 1:200.000 (MAGE200) nº 69 (Pozoblanco). Instituto
2171 Geológico y Minero de España.

2172 Matas, J., Martín Parra, L.M., Rubio Pascual, F.J., Roldán, F.J., Martín-Serrano, A.,
2173 Alonso-Chaves, F., Salazar Rincón, A., 2015b. Mapa y Memoria del Mapa Geológico
2174 Nacional a escala 1:200.000 (MAGE200) nº 74-75 (Sevilla-Puebla de Guzmán).
2175 Instituto Geológico y Minero de España.

2176 Matte, P., 1986. Tectonics and plate tectonics model for the Variscan belt of Europe.
2177 Tectonophysics 126, 329-374.

2178 Matte, P., 1991. Accretionary history and crustal evolution of the Variscan belt in
2179 Western Europe. Tectonophysics 196, 309-337.

2180 Matte, P., 2001. The Variscan collage and orogeny (480-290 Ma) and the tectonic
2181 definition of the Armorica microplate: a review. Terra Nova 13, 122-128.

2182 Merinero, R., Lunar, R., Menor, L.O., García, R.P., Monterrubio, S., Gervilla, F., 2013.
2183 Hydrothermal palladium enrichment in podiform chromitites of Calzadilla de los Barros
2184 (SW Iberian Peninsula). The Canadian Mineralogist 51, 387-404.

2185 Merinero, R., Lunar, R., Ortega, L., Piña, R., Monterrubio, S., Gervilla, F., 2014. Zoned
2186 chromite records multiple metamorphic episodes in the Calzadilla de los Barros
2187 ultramafic bodies (SW Iberian Peninsula). European Journal of Mineralogy 26, 757-
2188 770.

2189 Mitjavila, J., Martí, J., Soriano, C., 1997. Magmatic Evolution and Tectonic Setting of
2190 the Iberian Pyrite Belt Volcanism. Journal of Petrology 38, 727-755.

2191 Moita, P., 1997. Caracterização petrográfica e geoquímica do metamorfismo de alta
2192 pressão no sector de Viana do Alentejo-Alvito (Zona de Ossa-Morena). GeoFCUL,
2193 Universidade de Lisboa, Lisboa, p. 158.

2194 Moita, P., Munhá, J., Fonseca, P.E., Pedro, J., Tassinari, C.C.G., Araújo, A., Palacios,
2195 T., 2005. Phase equilibria and geochronology of Ossa-Morena eclogites, XIV Semana
2196 de Geoquímica, VIII Congresso de Geoquímica dos Países de Língua Portuguesa, pp.
2197 463-466.

2198 Moita, P., Santos, J.F., Pereira, M.F., 2009. Layered granitoids: interaction between
2199 continental crust recycling processes and mantle-derived magmatism. Examples from
2200 the Évora Massif (Ossa–Morena Zone, southwest Iberia, Portugal). Lithos 111, 125-
2201 141.

- 2202 Moita, P., Santos, J.F., Pereira, M.F., Costa, M.M., Corfu, F., 2015. The quartz-dioritic
2203 Hospitais intrusion (SW Iberian Massif) and its mafic microgranular enclaves —
2204 Evidence for mineral clustering. *Lithos* 224-225, 78-100.
- 2205 Montero, P., Bea, F., Zinger, T.F., Scarrow, J.H., Molina, J.F., Whitehouse, M.J., 2004.
2206 55 million years of continuous anatexis in Central Iberia: single-zircon dating of the
2207 Peña Negra Complex. *Journal of the Geological Society* 161, 255-263.
- 2208 Munhá, J., Ribeiro, A., Ribeiro, M.L., 1984. Blueschists in the Iberian Variscan Chain
2209 (Trás-os-Montes: NE Portugal). *Comunicações Serviço Geológico de Portugal* 70, 31-
2210 53.
- 2211 Munhá, J., Oliveira, J.T., Ribeiro, A., Oliveira, V., Quesada, C., Kerrich, R., 1986.
2212 Beja-Acebuches Ophiolite: characterization and geodynamic significance. Maleo,
2213 *Boletim Informativo da Sociedade Geológica de Portugal* 2, 1-31.
- 2214 Muñoz, G., Mateus, A., Pous, J., Heise, W., Monteiro Santos, F., Almeida, E., 2008.
2215 Unraveling middle-crust conductive layers in Paleozoic Orogens through 3D modeling
2216 of magnetotelluric data: The Ossa-Morena Zone case study (SW Iberian Variscides).
2217 *Journal of Geophysical Research: Solid Earth* 113, B06106.
- 2218 Oliveira, J.T., 1990. South Portuguese Zone: (1) Introduction, (2) Stratigraphy and
2219 synsedimentary tectonism, in: Dallmeyer, R.D., Martínez García, E. (Eds.), *Pre-*
2220 *Mesozoic Geology of Iberia*. Springer- Verlag, Berlin, Germany, pp. 333-347.
- 2221 Oliveira, J.T., Oliveira, V., Piçarra, J.M., 1991. Traços gerais da evolução tectono-
2222 estratigráfica da Zona de Ossa Morena, em Portugal: síntese crítica do estado actual dos
2223 conhecimentos. *Comunicações Serviço Geológico de Portugal* 77, 3-26.
- 2224 Oliveira, J.T., Pereira, E., Piçarra, J.M., Young, T., Romano, M., 1992. O Paleozoico
2225 Inferior de Portugal: síntese da estratigrafia e da evolução paleogeografica, in: Gutiérrez
2226 Marco, J.C., Saavedra, J., Rábano, I. (Eds.), *Paleozóico inferior de Ibero-América*.
2227 *Universidad de Extremadura*, pp. 359-375.

- 2228 Oliveira, J.T., Relvas, J., Pereira, Z., Munhá, J., Matos, J., Barriga, F., Rosa, C., 2013. O
2229 Complexo Vulcano-Sedimentar de Toca da Moura-Cabrela (Zona de Ossa Morena):
2230 evolução tectono-estratigráfica e mineralizações associadas, in: Dias, R., Araújo, A.,
2231 Terrinha, P., Kullberg, J.C. (Eds.), *Geologia de Portugal*. Escolar Editora, pp. 621-645.
- 2232 Onézime, J., Charvet, J., Faure, M., Bourdier, J.-L., Chauvet, A., 2003. A new
2233 geodynamic interpretation for the South Portuguese Zone (SW Iberia) and the Iberian
2234 Pyrite Belt genesis. *Tectonics* 22, 1027.
- 2235 Ordóñez Casado, B., 1998. Geochronological studies of the Pre-Mesozoic basement of
2236 the Iberian Massif: the Ossa Morena zone and the Allochthonous Complexes within the
2237 Central Iberian zone. Swiss Federal Institute of Technology, Zürich, Switzerland, p.
2238 235.
- 2239 Ordóñez Casado, B., Gebauer, D., Schafer, H.J., Gil Ibarguchi, J.I., Peucat, J.J., 2001. A
2240 single Devonian subduction event for the HP/HT metamorphism of the Cabo Ortegal
2241 complex within the Iberian Massif. *Tectonophysics* 332, 359-385.
- 2242 Ortuño, M.G., 1971. Middle Westphalian strata in South-West Spain. *Proceedings of*
2243 *the VII Congr. Intern. Strat. Géol. Carbonif. (Sheffield, 1967)* 3, 1275-1292.
- 2244 Pardo, M.V., García Alcalde, J.L., 1996. El Devónico de la Zona Centroibérica. *Revista*
2245 *española de Paleontología Número Extraordinario*, 72-81.
- 2246 Pastor-Galán, D., Gutiérrez-Alonso, G., Murphy, J.B., Fernández-Suárez, J., Hofmann,
2247 M., Linnemann, U., 2013. Provenance analysis of the Paleozoic sequences of the
2248 northern Gondwana margin in NW Iberia: Passive margin to Variscan collision and
2249 orocline development. *Gondwana Research* 23, 1089-1103.
- 2250 Pedro, J., 1996. Estudo do metamorfismo de alta pressão na área de Safira (Montemor-
2251 o-Novo) Zona de Ossa Morena. *GeoFCUL*, Universidade de Lisboa, Lisboa, p. 69.
- 2252 Pedro, J.C., Araújo, A., Fonseca, P.E., Munhá, J.M., 2005. Sequências ofiolíticas
2253 internas da zona de Ossa-Morena: implicações geodinâmicas na evolução da Cadeia
2254 Varisca Ibérica. *Cuadernos do Laboratorio Xeolóxico de Laxe* 30, 235-258.

2255 Pedro, J., Araújo, A., Fonseca, P., Tassinari, C., Ribeiro, A., 2010. Geochemistry and
 2256 U-Pb zircon age of the internal Ossa-Morena zone ophiolite sequences: a remnant of
 2257 Rheic ocean in SW Iberia. *Ophioliti* 35, 117-130.

2258 Perdigo, J.C., Oliveira, J.T., Ribeiro, A., 1982. Carta Geológica de Portugal na escala
 2259 1/50000. Notícia explicativa da folha 44-B, Barrancos. *Servicios Geológicos de*
 2260 *Portugal*, 49 pp.

2261 Pereira, M.F., 2015. Potential sources of Ediacaran strata of Iberia: a review.
 2262 *Geodinamica Acta* 27, 1-14.

2263 Pereira, M.F., Silva, J.B., 2001. A new model for the Hercynian Orogen of Gondwanan
 2264 France and Iberia: discussion. *Journal of Structural Geology* 23, 835-838.

2265 Pereira, M.F., Apraiz, A., 2006. High-pressure mafic granulites in the Coimbra-Córdoba
 2266 shear zone (Campo Maior Unit, northeast Alentejo): metamorphic and
 2267 geothermobarometric characterization. VII Congreso Nacional de Geologia, Portugal,
 2268 89-92.

2269 Pereira, M.F., Chichorro, M., Linnemann, U., Eguiluz, L., Silva, J.B., 2006a. Inherited
 2270 arc signature in Ediacaran and Early Cambrian basins of the Ossa-Morena Zone (Iberian
 2271 Massif, Portugal): Paleogeographic link with European and North African Cadomian
 2272 correlatives. *Precambrian Research* 144, 297-315.

2273 Pereira, M.F., Silva, J.B., Chichorro, M., Moita, P., Santos, J.F., Apraiz, A., Ribeiro, C.,
 2274 2007. Crustal growth and deformational processes in the northern Gondwana margin:
 2275 Constraints from the Évora Massif (Ossa-Morena Zone, southwest Iberia, Portugal), in:
 2276 Linnemann, U., Nance, R.D., Kraft, P., Zulauf, G. (Eds.), *The evolution of the Rheic*
 2277 *Ocean: From Avalonian-Cadomian active margin to Alleghenian-Variscan collision*.
 2278 *Geological Society of America Special Paper*, pp. 333-358, doi:
 2279 10.1130/2007.2423(1116).

2280 Pereira, M.F., Apraiz, A., Silva, J.B., Chichorro, M., 2008a. Tectonothermal analysis of
 2281 high-temperature mylonitization in the Coimbra-Cordoba shear zone (SW Iberian

2282 Massif, Ouguela tectonic unit, Portugal): Evidence of intra-continental transcurrent
 2283 transport during the amalgamation of Pangea. *Tectonophysics* 461, 378-394.

2284 Pereira, M.F., Chichorro, M., Williams, I.S., Silva, J.B., Fernández, C., Díaz-Azpíroz,
 2285 M., Apraiz, A., Castro, A., 2009. Variscan intra-orogenic extensional tectonics in the
 2286 Ossa-Morena Zone (Évora-Aracena-Lora del Río metamorphic belt, SW Iberian
 2287 Massif): SHRIMP zircon U-Th-Pb geochronology. Geological Society, London, Special
 2288 Publications 327, 215-237.

2289 Pereira, M.F., Apraiz, A., Chichorro, M., Silva, J.B., Armstrong, R.A., 2010a.
 2290 Exhumation of high-pressure rocks in northern Gondwana during the Early
 2291 Carboniferous (Coimbra-Cordoba shear zone, SW Iberian Massif): Tectonothermal
 2292 analysis and U-Th-Pb SHRIMP in-situ zircon geochronology. *Gondwana Research* 17,
 2293 440-460.

2294 Pereira, M.F., Silva, J.B., Drost, K., Chichorro, M., Apraiz, A., 2010b. Relative timing
 2295 of transcurrent displacements in northern Gondwana: U-Pb laser ablation ICP-MS
 2296 zircon and monazite geochronology of gneisses and sheared granites from the western
 2297 Iberian Massif (Portugal). *Gondwana Research* 17, 461-481.

2298 Pereira, M.F., Chichorro, M., Johnston, S.T., Gutiérrez-Alonso, G., Silva, J.B.,
 2299 Linnemann, U., Hofmann, M., Drost, K., 2012a. The missing Rheic Ocean magmatic
 2300 arcs: Provenance analysis of Late Paleozoic sedimentary clastic rocks of SW Iberia.
 2301 *Gondwana Research* 22, 882-891.

2302 Pereira, M.F., Chichorro, M., Silva, J.B., Ordóñez-Casado, B., Lee, J.K.W., Williams,
 2303 I.S., 2012b. Early carboniferous wrenching, exhumation of high-grade metamorphic
 2304 rocks and basin instability in SW Iberia: Constraints derived from structural geology
 2305 and U-Pb and ^{40}Ar - ^{39}Ar geochronology. *Tectonophysics* 558-559, 28-44.

2306 Pereira, M.F., Chichorro, M., Fernández, C., Silva, J.B., Matias, F.V., 2013. The role of
 2307 strain localization in magma injection into a transtensional shear zone (Variscan belt,
 2308 SW Iberia). *Journal of the Geological Society* 170, 93-105.

- 2309 Pereira, M.F., Chichorro, M., Moita, P., Santos, J.F., Solá, A.M.R., Williams, I.S.,
2310 Silva, J.B., Armstrong, R.A., 2015. The multistage crystallization of zircon in calc-
2311 alkaline granitoids: U-Pb age constraints on the timing of Variscan tectonic activity in
2312 SW Iberia. *International Journal of Earth Sciences* 104, 1167-1183.
- 2313 Pereira, Z., Piçarra, J.M., Oliveira, J.T., 1998. Palinomorfos do Devónico Inferior da
2314 região de Barrancos (Zona de Ossa-Morena). *Comunicações do V Congresso Nacional*
2315 *de Geologia* 84, A-18/A-20.
- 2316 Pereira, Z., Piçarra, J.M., Oliveira, J.T., 1999. Lower Devonian palynomorphs from the
2317 Barrancos region, Ossa-Morena Zone, Portugal. *Bollettino Della Societa Geologica*
2318 *Italiana* 38, 239-245.
- 2319 Pereira, Z., Oliveira, V., Oliveira, J.T., 2006b. Palynostratigraphy of the Toca da Moura
2320 and Cabrela Complexes, Ossa Morena Zone, Portugal. *Geodynamic implications.*
2321 *Review of Palaeobotany and Palynology* 139, 227-240.
- 2322 Pereira, Z., Matos, J., Fernandes, P., Oliveira, J.T., 2008b. Palynostratigraphy and
2323 systematic palynology of the Devonian and Carboniferous successions of the South
2324 Portuguese Zone, Portugal. *Memórias Geológicas do Instituto Nacional de Engenharia,*
2325 *Tecnologia e Inovação* 34, 1-146.
- 2326 Pérez-Cáceres, I., Martínez Poyatos, D., Simancas, J.F., Azor, A., 2015a. The elusive
2327 nature of the Rheic Ocean suture in SW Iberia. *Tectonics* 34, 2429-2450.
- 2328 Pérez-Cáceres, I., Simancas, J.F., Martínez Poyatos, D., Azor, A., González Lodeiro, F.,
2329 2015b. Oblique collision and deformation partitioning in the SW Iberian Variscides.
2330 *Solid Earth Discussions* 7, 3773-3815.
- 2331 Pérez-Estaún, A., Bastida, F., Alonso, J.L., Marquínez, J., Aller, J., Alvarez-Marrón, J.,
2332 Marcos, A., Pulgar, J.A., 1988. A thin-skinned tectonics model for an arcuate fold and
2333 thrust belt: The Cantabrian Zone (Variscan Ibero-Armorican Arc). *Tectonics* 7, 517-
2334 537.

- 2335 Pérez-Estaún, A., Martínez-Catalán, J.R., Bastida, F., 1991. Crustal thickening and
2336 deformation sequence in the footwall to the suture of the Variscan belt of northwest
2337 Spain. *Tectonophysics* 191, 243-253.
- 2338 Pérez Lorente, F., 1979. Geología de la Zona de Ossa-Morena al norte de Córdoba
2339 (Pozoblanco-Belmez-Villaviciosa). Universidad de Granada, Granada, p. 340.
- 2340 Peucat, J.J., Bernardgriffiths, J., Gil Ibarguchi, J.I., Dallmeyer, R.D., Menot, R.P.,
2341 Cornichet, J., Deleon, M.I.P., 1990. Geochemical and geochronological cross-section of
2342 the deep Variscan crust - The Cabo-Ortega high-pressure nappe (Northwestern Spain).
2343 *Tectonophysics* 177, 263-292.
- 2344 Piçarra, J.M., 1997. Nota sobre a descoberta de graptólitos do Devónico Inferior na
2345 Formação de Terena, em Barrancos (Zona de Ossa-Morena), in: Araújo, A., Pereira,
2346 M.F. (Eds.), *Estudos sobre a Geologia da Zona de Ossa-Morena (Maciço Ibérico)*. Livro
2347 de Homenagem ao Prof. Francisco Gonçalves, Évora, pp. 27-36.
- 2348 Piçarra, J.M., Pereira, Z., Oliveira, J.T., 1998. Novos dados sobre a idade da sucessão
2349 Slúrico-Devónica do Sinclinal de Terena, na região de Barrancos. Implicações
2350 geodinâmicas, in: Portugal, I.G.e.M.y.S.G.d. (Ed.), *Comunicações Actas do V*
2351 *Congresso Nacional de Geologia*, pp. A15-A16.
- 2352 Pin, C., Paquette, J.L., Santos Zalduegui, J.F., Gil Ibarguchi, J.I., 2002. Early devonian
2353 suprasubduction-zone ophiolite related to incipient collisional processes in the Western
2354 Variscan Belt: The Sierra de Careón unit, Ordenes Complex, Galicia, in: Martínez
2355 Catalán, J.R., Hatcher, R.D., Arenas, R., Díaz García, F. (Eds.), *Variscan-Appalachian*
2356 *Dynamics: the Building of the Late Paleozoic Basement*. Geological Society of America
2357 *Special Paper*, pp. 57-71, doi: 10.1130/1130-8137-2364-1137.1157.
- 2358 Pin, C., Paquette, J.L., Abalos, B., Santos, F.J., Gil Ibarguchi, J.I., 2006. Composite
2359 origin of an early Variscan transported suture: Ophiolitic units of the Morais Nappe
2360 Complex (north Portugal). *Tectonics* 25, TC5001.

- 2361 Pin, C., Fonseca, P.E., Paquette, J.-L., Castro, P., Matte, P., 2008. The ca. 350 Ma Beja
2362 Igneous Complex: A record of transcurrent slab break-off in the Southern Iberia
2363 Variscan Belt? *Tectonophysics* 461, 356-377.
- 2364 Ponce, C., Simancas, J.F., Azor, A., Martínez Poyatos, D.J., Booth-Rea, G., Expósito,
2365 I., 2012. Metamorphism and kinematics of the early deformation in the Variscan suture
2366 of SW Iberia. *Journal of Metamorphic Geology* 30, 625-638.
- 2367 Puelles, P., Abalos, B., Gil Ibarguchi, J.I., 2005. Metamorphic evolution and
2368 thermobaric structure of the subduction-related Bacariza high-pressure granulite
2369 formation (Cabo Ortegal Complex, NW Spain). *Lithos* 84, 125-149.
- 2370 Puschmann, H., 1967. Zum Problem der Schichtlücken im Devon der Sierra Morena
2371 (Spanien). *Geologische Rundschau* 56, 528-542.
- 2372 Quesada, C., 1990. Precambrian successions in SW Iberia: their relationship to
2373 'Cadomian' orogenic events, in: Lemos, D.R., Strachan, R.A., Topley, C.G. (Eds.), *The*
2374 *Cadomian Orogeny*, pp. 353-362.
- 2375 Quesada, C., Dallmeyer, R.D., 1994. Tectonothermal evolution of the Badajoz-Cordoba
2376 shear zone (SW Iberia) - characteristics and $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age constraints.
2377 *Tectonophysics* 231, 195-213.
- 2378 Quesada, C., Fonseca, P.E., Munhá, J.M., Oliveira, J.T., Ribeiro, A., 1994. The Beja-
2379 Acebuches Ophiolite (southern Iberia Variscan fold belt): geological characterization
2380 and geodynamic significance. *Boletín Geológico y Minero* 105, 3-49.
- 2381 Racheboeuf, P.R., Lethiers, F., Babin, C., Rolfe, W.D.I., Marez, E., 1986. Les faunes du
2382 Dévonien supérieur d'Alange (province de Badajoz. Sud-Ouest de l'Espagne). *Géologie*
2383 *Méditerranéenne* 12-13, 37-47.
- 2384 Ribeiro, A., Cramez, C., Almeida Rebelo, J., 1964. Sur la structure de Trás-os-Montes
2385 (Nord-Est du Portugal). *Comptes Rendues Académie des Sciences Paris* 258, 263-265.

- 2386 Ribeiro, A., Pereira, E., Severo, L., 1980. Análise da deformação da zona de
2387 cisalhamento Porto-Tomar na transversal de Oliveira de Aeméis. Comunicações dos
2388 Serviços Geológicos de Portugal 66, 3-9.
- 2389 Ribeiro, A., Pereira, E., Dias, R., 1990. Structure in the northwest of the Iberian
2390 Peninsula, in: Dallmeyer, R.D., Martínez García, E. (Eds.), Pre-Mesozoic geology of
2391 Iberia. Springer-Verlag, Berlin, Germany, pp. 220-236.
- 2392 Ribeiro, A., Munhá, J., Dias, R., Mateus, A., Pereira, E., Ribeiro, L., Fonseca, P.,
2393 Araújo, A., Oliveira, T., Romão, J., Chaminé, H., Coke, C., Pedro, J., 2007.
2394 Geodynamic evolution of the SW Europe Variscides. *Tectonics* 26, TC6009.
- 2395 Ribeiro, A., Munhá, J., Fonseca, P.E., Araújo, A., Pedro, J.C., Mateus, A., Tassinari, C.,
2396 Machado, G., Jesus, A., 2010. Variscan ophiolite belts in the Ossa-Morena Zone
2397 (Southwest Iberia): Geological characterization and geodynamic significance.
2398 *Gondwana Research* 17, 408-421.
- 2399 Ribeiro, M.L., Floor, P., 1987. Magmatismo peralcalino no Macico Hesperico: Sua
2400 distribuico e significado geodinamico, geologia de los granitoides y rocas asociadas del
2401 Macizo Hesperico, in: Bea, F., Carnicero, A., Gonzalo, J.C., López-Plaza, M.,
2402 Rodríguez Alonso, M.D. (Eds.), *Geología de los granitoides y rocas asociadas del*
2403 *Macizo Hespérico. Libro Homenaje a L.C. García de Figuerola*. Ediciones Rueda,
2404 Madrid, pp. 211-222.
- 2405 Ries, A.C., Shackleton, R.M., 1971. Catazonal Complexes of North-West Spain and
2406 North Portugal, Remnants of a Hercynian Thrust Plate. *Nature Physical Science* 234,
2407 65-79.
- 2408 Robardet, M., 2003. The Armorica 'microplate': fact or fiction? Critical review of the
2409 concept and contradictory palaeobiogeographical data. *Palaeogeography,*
2410 *Palaeoclimatology, Palaeoecology* 195, 125-148.
- 2411 Robardet, M., Gutiérrez Marco, J.C., 2004. The Ordovician, Silurian and Devonian
2412 sedimentary rocks of the Ossa-Morena Zone (SW Iberian Peninsula, Spain). *Journal of*
2413 *Iberian Geology* 30, 73-92.

- 2414 Robardet, M., Piçarra, J.M., Storch, P., Gutiérrez Marco, J.C., Sarmiento, G.N., 1998.
2415 Ordovician and Silurian stratigraphy and faunas (graptolites and conodonts) in the Ossa
2416 Morena Zone of SW Iberian Peninsula (Portugal and Spain). *Temas Geologicos-*
2417 *Mineros ITGE* 23, 289-318.
- 2418 Rocha, R.C., Araújo, A., Borrego, J., Fonseca, P.E., 2009. Transected folds with
2419 opposite patterns in Terena Formation (Ossa Morena Zone, Portugal): anomalous
2420 structures resulting from sedimentary basin anisotropies. *Geodinamica Acta* 22, 157-
2421 163.
- 2422 Rocha, R., Pereira, Z., Araújo, A., 2010. Novos dados bioestratigráficos (miosporos) na
2423 Formação de Terena - Implicações para a interpretação estrutural (Rio Ardila,
2424 Barrancos). *e-Terra* 17 (VIII Congresso Nacional de Geologia), 1-4.
- 2425 Rodríguez, J., Cosca, M.A., Gil Ibarguchi, J.I., Dallmeyer, R.D., 2003. Strain
2426 partitioning and preservation of $^{40}\text{Ar}/^{39}\text{Ar}$ ages during Variscan exhumation of a
2427 subducted crust (Malpica-Tui complex, NW Spain). *Lithos* 70, 111-139.
- 2428 Rodríguez, J., Gil Ibarguchi, J.I., Paquette, J.L., 2007. Sincronía del magmatismo
2429 varisco en el Macizo Ibérico: nuevas edades U-Pb en granitoides de la región de
2430 Finisterre (La Coruña, España). *XV Semana - VI Congreso Ibérico de Geoquímica*, Vila
2431 Real, Portugal, DVD-ROM, 146-149.
- 2432 Rodríguez, L., Mira, M., Ortega, E., 1990. Mapa Geológico, Hoja 833 (Hinojosa del
2433 Duque), Serie MAGNA, 1/50.000. Instituto Geológico y Minero de España, Madrid.
- 2434 Rodríguez, S., Soto, F., 1979. Nuevos datos sobre los corales rugosos del Devónico de
2435 la Sierra del Pedroso. *Estudios Geológicos* 35, 345-354.
- 2436 Romeo, I., Lunar, R., Capote, R., Quesada, C., Piña, R., Dunning, G.R., Ortega, L.,
2437 2006. U-Pb age constraints on Variscan magmatism and Ni-Cu-PGE metallogeny in the
2438 Ossa-Morena zone (SW Iberia). *Journal of the Geological Society* 163, 837-846.
- 2439 Rosa, D.R.N., Finch, A.A., Andersen, T., Inverno, C.M.C., 2008. U-Pb geochronology
2440 of felsic volcanic rocks hosted in the Gafo Formation, South Portuguese Zone: the

2441 relationship with Iberian Pyrite Belt magmatism. *Mineralogical Magazine* 72, 1103-
 2442 1118.

2443 Rosas, F., Marques, F.O., Luz, A., Coelho, S., 2002. Sheath folds formed by drag
 2444 induced by rotation of rigid inclusions in viscous simple shear flow: nature and
 2445 experiment. *Journal of Structural Geology* 24, 45-55.

2446 Rosas, F.M., Marques, F.O., Ballèvre, M., Tassinari, C., 2008. Geodynamic evolution
 2447 of the SW Variscides: Orogenic collapse shown by new tectonometamorphic and
 2448 isotopic data from western Ossa-Morena Zone, SW Iberia. *Tectonics* 27, doi:
 2449 10.1029/2008TC002333.

2450 Rubio Pascual, F., Arenas, R., Díaz García, F., Martínez Catalán, J.R., Abati, J., 2002.
 2451 Contrasting high-pressure metabasites from the Santiago unit (Ordenes complex,
 2452 northwestern Iberian massif, Spain), in: Martínez Catalán, J.R., Hatcher, R.D., Arenas,
 2453 R., Díaz García, F. (Eds.), *Variscan-Appalachian dynamics: The building of the Late*
 2454 *Paleozoic basement*. Geological Society of America Special Paper, pp. 105-124, doi:
 2455 110.1130/1130-8137-2364-1137.1105.

2456 Rubio Pascual, F.J., Arenas, R., Catalán, J.R.M., Fernández, L.R.R., Wijbrans, J.R.,
 2457 2013a. Thickening and exhumation of the Variscan roots in the Iberian Central System:
 2458 Tectonothermal processes and $^{40}\text{Ar}/^{39}\text{Ar}$ ages. *Tectonophysics* 587, 207-221.

2459 Rubio Pascual, F.J., Matas, J., Martín Parra, L.M., 2013b. High-pressure metamorphism
 2460 in the Early Variscan subduction complex of the SW Iberian Massif. *Tectonophysics*
 2461 592, 187-199.

2462 Rubio Pascual, F.J., Martín Parra, L.M., Díez Montes, A., Díez Fernández, R.,
 2463 Gallastegui, G., Valverde Vaquero, P., Rodríguez Fernández, L.R., Heredia, N., 2015.
 2464 Metamorphic records of partial subduction and continental collision in and around the
 2465 parautochthon of the NW Iberian Massif. The Variscan belt: correlations and plate
 2466 dynamics. Special meeting of the French & Spanish Geological Societies (Rennes).
 2467 *Géologie de la France* 1, 123-124.

- 2468 Salman, K., 2002. Estudio petrológico, geoquímico y geocronológico de los granitoides
2469 del área Monesterio-Cala, Zona de Ossa-Morena (Macizo Ibérico). Universidad de
2470 Granada, Granada, Spain, p. 232.
- 2471 Sánchez-García, T., Bellido, F., Quesada, C., 2003. Geodynamic setting and
2472 geochemical signatures of Cambrian–Ordovician rift-related igneous rocks (Ossa-
2473 Morena Zone, SW Iberia). *Tectonophysics* 365, 233-255.
- 2474 Sánchez Cela, V., Gabaldón, V., 1977a. Mapa Geológico, Hoja 831 (Zalamea de la
2475 Serena), Serie MAGNA, 1/50.000. Instituto Geológico y Minero de España, Madrid.
- 2476 Sánchez Cela, V., Gabaldón, V., 1977b. Mapa Geológico, Hoja 856 (Maguilla), Serie
2477 MAGNA, 1/50.000. Instituto Geológico y Minero de España, Madrid.
- 2478 Sánchez Martínez, S., Arenas, R., Díaz García, F., Martínez Catalán, J.R., Gómez
2479 Barreiro, J., Pearce, J.A., 2007. Careon ophiolite, NW Spain: Suprasubduction zone
2480 setting for the youngest Rheic Ocean floor. *Geology* 35, 53-56.
- 2481 Sánchez Martínez, S., Arenas, R., Fernández-Suárez, J., Jeffries, T.E., 2009. From
2482 Rodinia to Pangaea: ophiolites from NW Iberia as witness for a long-lived continental
2483 margin, in: Murphy, J.B., Keppie, J.D., Hynes, A.J. (Eds.), *Ancient Orogens and*
2484 *Modern Analogues*. Geological Society, London, Special Publications, pp. 317-341,
2485 doi: 310.1144/SP1327.1114.
- 2486 Sánchez Martínez, S., Arenas, R., Gerdes, A., Castiñeiras, P., Potrel, A., Fernández-
2487 Suárez, J., 2011. Isotope geochemistry and revised geochronology of the Purrido
2488 Ophiolite (Cabo Ortegal Complex, NW Iberian Massif): Devonian magmatism with
2489 mixed sources and involved Mesoproterozoic basement. *Journal of the Geological*
2490 *Society* 168, 733-750.
- 2491 Sánchez Martínez, S., Gerdes, A., Arenas, R., Abati, J., 2012. The Bazar Ophiolite of
2492 NW Iberia: a relic of the Iapetus–Tornquist Ocean in the Variscan suture. *Terra Nova*
2493 24, 283-294.

- 2494 Santos, J.F., Andrade, A.S., Munhá, J., 1990. Magmatismo orogénico varisco no limite
2495 meridional da Zona de Ossa-Morena. Comunicações dos Serviços Geológicos de
2496 Portugal 76, 91-124.
- 2497 Santos Zalduegui, J.F., Scharer, U., Gil Ibarguchi, J.I., 1995. Isotope constraints on the
2498 age and origin of magmatism and metamorphism in the Malpica-Tuy allochthon,
2499 Galicia, NW Spain. Chemical Geology 121, 91-103.
- 2500 Santos Zalduegui, J.F., Scharer, U., Gil Ibarguchi, J.I., Girardeau, J., 1996. Origin and
2501 evolution of the Paleozoic Cabo Ortegal ultramafic-mafic complex (NW Spain): U-Pb,
2502 Rb-Sr and Pb-Pb isotope data. Chemical Geology 129, 281-304.
- 2503 Schäfer, H.J., Gebauer, D., Nägler, T.F., 1991. Evidence for Silurian eclogite and
2504 granulite facies metamorphism in the Badajoz-Córdoba Shear belt, SW Spain. Terra
2505 Nova 3, suppl. 6, 11.
- 2506 Shaw, J., Johnston, S.T., Gutiérrez-Alonso, G., Weil, A.B., 2012. Oroclines of the
2507 Variscan orogen of Iberia: Paleocurrent analysis and paleogeographic implications.
2508 Earth and Planetary Science Letters 329–330, 60-70.
- 2509 Silva, J.C., Mata, J., Moreira, N., Fonseca, P.E., Jorge, R.C.G.S., Machado, G., 2011.
2510 Evidence for a Lower Devonian subduction zone in the southeastern boundary of the
2511 Ossa-Morena-Zone. Livro de actas VIII Congresso Ibérico de Geoquímica, Instituto
2512 Politécnico de Castelo Branco, 295-299.
- 2513 Silva, J.B., Oliveira, J.T., Ribeiro, A., 1990. South Portuguese zone. Structural outline,
2514 in: Dallmeyer, R.D., Martínez García, E. (Eds.), Pre-Mesozoic Geology of Iberia.
2515 Springer- Verlag, Berlin, Germany, pp. 348-362.
- 2516 Silva, J.B., Pereira, M.F., Chichorro, M., 2013. Estrutura das áreas internas da Zona Sul
2517 Portuguesa, no contexto do Orógeno Varisco, in: Dias, R., Araújo, A., Terrinha, P.,
2518 Kullberg, J.C. (Eds.), Geologia de Portugal. Escolar Editora, pp. 767-786.

2519 Simancas, J.F., Martínez Poyatos, D., Expósito, I., Azor, A., González Lodeiro, F.,
 2520 2001. The structure of a major suture zone in the SW Iberian Massif: the Ossa-
 2521 Morena/Central Iberian contact. *Tectonophysics* 332, 295-308.

2522 Simancas, J.F., Carbonell, R., González Lodeiro, F., Pérez Estaún, A., Juhlin, C.,
 2523 Ayarza, P., Kashubin, A., Azor, A., Martínez Poyatos, D., Almodóvar, G.R., Pascual,
 2524 E., Sáez, R., Expósito, I., 2003. Crustal structure of the transpressional Variscan orogen
 2525 of SW Iberia: SW Iberia deep seismic reflection profile (IBERSEIS). *Tectonics* 22,
 2526 1062.

2527 Simancas, J.F., Tahiri, A., Azor, A., Lodeiro, F.G., Poyatos, D.J.M., Hadi, H.E., 2005.
 2528 The tectonic frame of the Variscan–Alleghanian orogen in Southern Europe and
 2529 Northern Africa. *Tectonophysics* 398, 181-198.

2530 Simancas, J.F., Carbonell, R., González Lodeiro, F., Pérez Estaún, A., Juhlin, C.,
 2531 Ayarza, P., Kashubin, A., Azor, A., Martínez Poyatos, D., Sáez, R., Almodóvar, G.R.,
 2532 Pascual, E., Flecha, I., Martí, D., 2006. Transpressional collision tectonics and mantle
 2533 plume dynamics: the Variscides of southwestern Iberia. Geological Society, London,
 2534 *Memoirs* 32, 345-354.

2535 Simancas, J.F., Azor, A., Martínez-Poyatos, D., Tahiri, A., El Hadi, H., González-
 2536 Lodeiro, F., Pérez-Estaún, A., Carbonell, R., 2009. Tectonic relationships of Southwest
 2537 Iberia with the allochthons of Northwest Iberia and the Moroccan Variscides. *Comptes*
 2538 *Rendus Geoscience* 341, 103-113.

2539 Simancas, J.F., Ayarza, P., Azor, A., Carbonell, R., Martínez Poyatos, D., Pérez-Estaún,
 2540 A., González Lodeiro, F., 2013. A seismic geotraverse across the Iberian Variscides:
 2541 Orogenic shortening, collisional magmatism, and orocline development. *Tectonics* 32,
 2542 417-432.

2543 Solá, A.R., Williams, I.S., Neiva, A.M.R., Ribeiro, M.L., 2009. U–Th–Pb SHRIMP
 2544 ages and oxygen isotope composition of zircon from two contrasting late Variscan
 2545 granitoids, Nisa-Albuquerque batholith, SW Iberian Massif: Petrologic and regional
 2546 implications. *Lithos* 111, 156-167.

- 2547 Tait, J., Schätz, M., Bachtadse, V., Soffel, H., 2000. Palaeomagnetism and Palaeozoic
2548 palaeogeography of Gondwana and European terranes. Geological Society, London,
2549 Special Publications 179, 21-34.
- 2550 Tornos, F., Casquet, C., Relvas, J.M.R.S., 2005. Transpressional tectonics, lower crust
2551 decoupling and intrusion of deep mafic sills: A model for the unusual metallogensis of
2552 SW Iberia. Ore Geology Reviews 27, 133-163.
- 2553 Valenzuela, A., Donaire, T., Pin, C., Toscano, M., Hamilton, M.A., Pascual, E., 2011.
2554 Geochemistry and U–Pb dating of felsic volcanic rocks in the Riotinto–Nerva unit,
2555 Iberian Pyrite Belt, Spain: crustal thinning, progressive crustal melting and massive
2556 sulphide genesis. Journal of the Geological Society 168, 717-732.
- 2557 Valverde Vaquero, P., Fernández, F.J., 1996. Edad de enfriamiento U/Pb en rutilos del
2558 Gneis de Chímparra (Cabo Ortegal, NO de España). Geogaceta 20, 475-478.
- 2559 Valle Aguado, B., Azevedo, M.R., Schaltegger, U., Martínez Catalán, J.R., Nolan, J.,
2560 2005. U-Pb zircon and monazite geochronology of Variscan magmatism related to syn-
2561 convergence extension in Central Northern Portugal. Lithos 82, 169-184.
- 2562 Valle Aguado, B., Azevedo, M., Nolan, J., Burgess, R., 2008. $^{40}\text{Ar}/^{39}\text{Ar}$ Age Constraints
2563 for the D₂ Variscan Extension in the Porto-Viseu Metamorphic Belt (Portugal).
2564 Geochimica et Cosmochimica Acta 72, Supplement 970.
- 2565 Van Calsteren, P.W.C., Boelrijk, N., Hebeda, E.H., Priem, H.N.A., Dentex, E.,
2566 Verdurmen, E.A.T., Verschure, R.H., 1979. Isotopic dating of older elements (including
2567 the Cabo Ortegal mafic-ultramafic complex) in the Hercynian orogen of NW Spain -
2568 Manifestations of a presumed Early Paleozoic mantle-plume. Chemical Geology 24, 35-
2569 56.
- 2570 Van Staal, C.R., Whalen, J.B., Valverde-Vaquero, P., Zagorevski, A., Rogers, N., 2009.
2571 Pre-Carboniferous, episodic accretion-related, orogenesis along the Laurentian margin
2572 of the northern Appalachians, in: Murphy, J.B., Keppie, J.D., Hynes, A.J. (Eds.),
2573 Ancient orogens and modern analogues. Geological Society of London Special
2574 Publication, pp. 271-316.

2575 Vauchez, A., 1974. Etude tectonique et microtectonique d'un secteur de la Chaîne
 2576 Hercynienne Sud-Iberique. Les nappes et plis couches de la Region de Fregenal.
 2577 Université des Sciences et Techniques du Languedoc, p. 184.

2578 Vauchez, A., 1976. Les structures hercyniennes dans la Region de Fregenal-Oliva de la
 2579 Frontera. (Badajoz, España). Comunicações dos Serviços Geológicos de Portugal 40,
 2580 261-267.

2581 Veselovski, Z., 2004. Integrated numerical modelling of a polyhistory basin, Southern
 2582 Cantabrian Basin (Palaeozoic, NW-Spain). Heidelberg University, Heidelberg,
 2583 Germany, p. 225.

2584 Villaseca, C., Barbero, L., Rogers, G., 1998. Crustal origin of Hercynian peraluminous
 2585 granitic batholiths of Central Spain: petrological, geochemical and isotopic (Sr, Nd)
 2586 constraints. Lithos 43, 55-79.

2587 Vogel, D.E., 1967. Petrology of an eclogite- and pyrigarnite-bearing polymetamorphic
 2588 rock complex at Cabo Ortegal, NW Spain. Leidse Geologische Mededelingen 40, 121-
 2589 213.

2590 Von Raumer, J.F., Stampfli, G.A., Bussy, F., 2003. Gondwana-derived microcontinents
 2591 - the constituents of the Variscan and Alpine collisional orogens. Tectonophysics 365,
 2592 7-22.

2593 Wallace, L.M., Stevens, C., Silver, E., McCaffrey, R., Lortung, W., Hasiata, S.,
 2594 Stanaway, R., Curley, R., Rosa, R., Taugaloidi, J., 2004. GPS and seismological
 2595 constraints on active tectonics and arc-continent collision in Papua New Guinea:
 2596 Implications for mechanics of microplate rotations in a plate boundary zone. Journal of
 2597 Geophysical Research: Solid Earth 109, B05404.

2598 Weil, A., Gutiérrez-Alonso, G., Conan, J., 2010. New time constraints on lithospheric-
 2599 scale oroclinal bending of the Ibero-Armorican Arc: a palaeomagnetic study of earliest
 2600 Permian rocks from Iberia. Journal of the Geological Society 167, 127-143.

Winchester, J.A., Pharaoh, T.C., Verniers, J., 2002. Palaeozoic amalgamation of Central Europe: an introduction and synthesis of new results from recent geological and geophysical investigations. Geological Society, London, Special Publications 201, 1-18.

Figure Captions

Fig. 1. Zonation of the Variscan orogen after Martínez Catalán et al. (2007) and Díez Fernández and Arenas (2015). The locations of the Coimbra-Córdoba Shear Zone and the oroclinal bends of the orogen are shown.

Fig. 2. Geological map showing the main zones of the Iberian Massif (after Díez Fernández and Arenas, 2015). Abbreviations: AF — Azuaga Fault; BToIP — Basal Thrust of the Iberian Parautochthon; BAO — Beja-Acebuches Ophiolite; CA — Carvalhal Amphibolites; CF — Canaleja Fault; CMU — Cubito-Moura Unit; CO — Calzadilla Ophiolite; CU — Central Unit; EsT — Espiel Thrust; ET—Espina Thrust; HF— Hornachos Fault; IOMZO —Internal Ossa-Morena Zone Ophiolites; J-PCSZ — Juzbado-Penalva do Castelo Shear Zone; LFT — Lalín-Forcarei Thrust; LPSZ — Los Pedroches Shear Zone; LLSZ — Llanos Shear Zone; MLSZ — Malpica-Lamego Shear Zone; MF — Matachel Fault; OF — Onza Fault; OVD — Obejo-Valsequillo Domain; PG-CVD — Puente Génave-Castelo de Vide Detachment; PRSZ— Palas de Rei Shear Zone; PTSZ — Porto-Tomar Shear Zone; RF — Riás Fault; SISZ —South Iberian Shear Zone; VF — Viveiro Fault; ZSI — Zalamea de la Serena Imbricates.

Fig. 3. Composite cross-section of major tectonic elements of the Iberian Massif (after Díez Fernández and Arenas, 2015). Abbreviations follow Figure 2. The location of the Iberian Reflective Body is shown.

2627

2628 **Fig. 4.** Idealized Variscan evolution of the Iberian Massif during the Devonian (see text
2629 for explanation). Circled numbers refer to specific structures and basins of the Iberian
2630 Massif. (a) Simplified pre-collisional paleogeography across the margin of Gondwana
2631 after restoration of Variscan deformation. Note the model is at 50% scale relative to the
2632 rest of the drawings (b) Tectonic setting showing the onset of Variscan deformation in
2633 the peri-Gondwana realm after the closure of the Rheic Ocean. Note the north-directed
2634 tectonic escape proposed for the peri-Gondwanan domain. (c) Opening of an intra-
2635 Gondwanan oceanic basin in the Lower-Middle Devonian. (d) Closure of the intra-
2636 Gondwana basin by accretion of different tectonic slices of oceanic crust (ophiolitic
2637 units) and followed by subduction of Gondwanan continental crust under the previously
2638 stacked ophiolites.

2639

2640 **Fig. 5.** Idealized Variscan evolution of the Iberian Massif during the Carboniferous (see
2641 text for explanation). Circled numbers refer to specific structures and basins of the
2642 Iberian Massif (numbering continues list of Figure 4). (a) Thrust and fold nappe
2643 tectonics during the early stages of the emplacement of the Iberian Allochthon.
2644 Propagation of Gondwana-verging folds in the Allochthon accompanied the
2645 underthrusting of the Iberian Autochthon and Parautochthon. (b) Climax of Gondwanan
2646 lithosphere underthrusting and onset of out-of-sequence thrusts. Note the proposed
2647 mantle topography near the Rheic suture that separates Gondwana from Laurussia. Time
2648 lines in sections b and c overlap each other because they both represent processes that
2649 may have occurred simultaneously. (c) Inception of the Beja-Acebuches basin,
2650 beginning of the orogenic extensional collapse and advance of out-of-sequence thrusts.
2651 (d) Closure of the Beja-Acebuches basin, widespread collapse of the orogen and

2652 reactivation of out-of-sequence thrusts. Along-strike movements are not represented for
2653 extensional faults. Location and kinematics of later strike-slip shear zones is shown.
2654 Abbreviations: IRB — Iberian Reflective Body; SZ — Shear Zone.

2655

2656 **Table 1.** Summary of the main tectonic events recognized on each geotectonic zone of
2657 the Iberian Massif during the Variscan orogeny. Dashed lines show a time-based
2658 correlation. Names in capital letters refer to the nomenclature utilized in this work for
2659 tectonic integration, while the commonly used regional names are shown in grey boxes
2660 below (the terms autochthon and allochthon inside parentheses refer to the
2661 allochthonous or autochthonous nature after Díez Fernández and Arenas, 2015).

2662

2663

EUROPEAN VARISCIDES

(CROPPING OUT / COVERED)

LAURUSSIAN TERRANES

- South-Portuguese Zone
- Unexposed basement

PERI-GONDWANA

- Uppermost allochthonous units
- High-P/high-T units

OPHIOLITES

- Intra-Gondwana, allochthonous ophiolites
- Beja-Acebuches ophiolite

GONDWANA

Continental edge:

- basal allochthonous units
- Upper sequence
- Lower sequence
- Lithospheric mantle

Mainland

- Parautochthonous sequences
- Autochthonous sequences
- Variscan granitoids and molten crust
- Variscan mafic rocks
- Unexposed basement of Gondwana