Marginal-Fermi-liquid behavior from two-dimensional Coulomb interaction

J. González

Instituto de Estructura de la Materia, Consejo Superior de Investigaciones Científicas, Serrano 123, 28006 Madrid, Spain

F. Guinea

Instituto de Ciencia de Materiales, Consejo Superior de Investigaciones Científicas, Cantoblanco, 28049 Madrid, Spain

M. A. H. Vozmediano

Escuela Politécnica Superior, Universidad Carlos III, Butarque 15 Leganés, 28913 Madrid, Spain

(Received 29 June 1998)

A full, nonperturbative renormalization group analysis of interacting electrons in a graphite layer is performed, in order to investigate the deviations from Fermi-liquid theory that have been observed in the experimental measures of a linear quasiparticle decay rate in graphite. The electrons are coupled through Coulomb interactions, which remain unscreened due to the semimetallic character of the layer. We show that the model flows towards the noninteracting fixed point for the whole range of couplings, with logarithmic corrections which signal the marginal character of the interaction separating Fermi-liquid and non-Fermi-liquid regimes. [S0163-1829(99)50204-X]

During recent years there has been important progress in understanding the properties of quantum electron liquids in dimension $D < 3$. One of the most fruitful approaches in this respect springs from the use of renormalization group (RG) methods, in which the different liquids are characterized by several fixed points controlling the low-energy properties. The Landau theory of the Fermi liquid in dimension $D > 1$ can be taken as a paradigm of the success of this program. It has been shown that, at least in the continuum limit, a system with isotropic Fermi surface and regular interactions is capable of developing a fixed point in which the interaction remains stable in the infrared.¹

The question of whether different critical points may arise at dimension $D = 2$ is now a subject of debate.²⁻⁵ From the perspective of the RG approach, one of the premises leading to the Fermi-liquid fixed point should be relaxed in order to flow to a different universality class. In the case of models proposed to understand the electronic properties of copper-oxide superconductors, the high anisotropy of the Fermi surface⁶ may play an important role in the anomalous behavior of the normal as well as of the superconducting state.⁷ On the other hand, a possible source of non-Fermi-liquid behavior may arise in systems with singular interactions.⁸ In the case of the Coulomb interaction screened by the Fermi sea, a solution by means of bosonization methods has shown that no departures from Fermi-liquid behavior arise at $D = 2$ and 3.⁹ It has been also shown for the conventional screened interaction that only potentials as singular as $V(q) \sim 1/|q|^{2D-2}$ can lead to a different electron liquid.^{8,10} The system of electrons with gauge interactions is quite different in that respect. It is known that in this case the system shows non-Fermi-liquid behavior for $D \leq 3$, manifested in properties like the specific heat or anomalous electron field dimensions.¹¹⁻¹³

In the present work, we address the applicability of the notion of Fermi-liquid fixed point to two-dimensional systems with unscreened Coulomb interaction. The absence of

screening requires the vanishing of the density of states at the Fermi level. Semimetals show this behavior, while retaining a gapless electronic spectrum. The existence of a well-defined continuum at low energies permits the existence of nontrivial scaling properties.¹⁴ The most remarkable example of this kind is given by the two-dimensional sheet of graphite, which has a vanishing density of states at the Fermi level.¹⁵ Recent photoemission experiments in graphite, at intermediate energies, show a decay rate of quasiparticles proportional to their energy.¹⁶ This represents a clear deviation with respect to the behavior in metals, which follow the conventional Fermi-liquid picture with quasiparticle lifetimes proportional to the inverse of the energy square, with, at most, logarithmic corrections. A description in terms of an effective field theory model has shown that the electronic interactions within the graphite layers are mainly responsible for the anomalous properties measured in the experiment.¹⁷

We apply RG techniques to investigate whether the mentioned anomalous behavior can be understood as a marginal deviation from Fermi-liquid theory, or rather it points towards a different universality class realized in the graphite sheet. We recall that the low-energy electronic excitations of the latter at half-filling are concentrated around two Fermi points at the corners of the hexagonal Brillouin zone, where the dispersion relation is well approximated by two cones in contact at the apex. The effective field theory is given therefore by a pair of Dirac fermions, with a Coulomb potential that remains unscreened due to the vanishing density of states at the Fermi points. The effective Hamiltonian can be written in the form¹⁸

$$\begin{aligned}
 H = & -iv_F \int d^2r \Psi^\dagger(\mathbf{r}) \boldsymbol{\sigma} \cdot \nabla \Psi(\mathbf{r}) \\
 & + \frac{e^2}{8\pi} \int d^2r_1 \int d^2r_2 \Psi^\dagger(\mathbf{r}_1) \\
 & \times \Psi(\mathbf{r}_1) \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \Psi^\dagger(\mathbf{r}_2) \Psi(\mathbf{r}_2), \quad (1)
 \end{aligned}$$

where $\Psi(\mathbf{r})$ is a two-dimensional Dirac spinor and $\sigma \equiv (\sigma_x, \sigma_y)$. Such effective field theory provides a good starting point for a RG analysis since, given that the scaling dimension of the $\Psi(\mathbf{r})$ field is -1 (in length units), the four-fermion Coulomb interaction turns out to be scale invariant, at this level, with a dimensionless coupling constant e^2 .

In order to address the existence of a different universality class, besides the trivial noninteracting phase, a nonperturbative approach has to be adopted, since the former can only be revealed by a nontrivial fixed point in coupling constant space. In the following, we will implement a GW approximation in the computation of the self-energy properties. This is more easily achieved in the present model by replacing the four-fermion term in Eq. (1) by the interaction with an auxiliary scalar field used to propagate the Coulomb interaction. The effective Hamiltonian can be rewritten in the form

$$H = -iv_F \int d^2r \Psi^\dagger(\mathbf{r}) \sigma \cdot \nabla \Psi(\mathbf{r}) + e \int d^2r \Psi^\dagger(\mathbf{r}) \Psi(\mathbf{r}) \phi(\mathbf{r}), \quad (2)$$

where the scalar field $\phi(\mathbf{r})$ has the propagator

$$\langle i T \phi(\mathbf{r}, t) \phi(\mathbf{r}', t') \rangle = \frac{1}{4\pi} \delta(t - t') \frac{1}{|\mathbf{r} - \mathbf{r}'|}. \quad (3)$$

In this framework we will introduce the GW approximation by taking into account the quantum corrections to the ϕ propagator due to particle-hole excitations of the Fermi sea. This kind of approximation has proven to be adequate to the description of the crossover from Fermi-liquid to Luttinger-liquid behavior upon lowering the dimension from $D=2$ to 1, capturing the relevant physical processes in the electron system.¹⁹ Therefore, it seems also appropriate to uncover any possible fixed point, different from that of Fermi-liquid theory, in the case of the system with unscreened Coulomb interaction.

The perturbative analysis of our model shows, in fact, the existence of a free fixed point that is stable in the infrared limit.¹⁴ This can be understood from the nontrivial scaling of the model with respect to variations of the bandwidth cutoff E_c , that is needed to regulate the divergent contribution of virtual processes to observable quantities. In the perturbative regime the Fermi velocity v_F grows steadily as the cutoff E_c is reduced and, as long as the electron charge e is not renormalized at the one-loop level, the effective coupling constant e^2/v_F flows to zero in the low-energy effective theory.

The most interesting point, however, concerns the analysis of the model away from the perturbative regime. With regard to the direct application to the graphite system the weak coupling results are of little use, since the bare coupling in the graphite sheet has an estimated value $e^2/v_F \sim 10$. Some possibly relevant effects, like, for instance, the renormalization of the quasiparticle weight, have to be consistently understood in a nonperturbative framework.

The polarization tensor built out of particle-hole processes is given by

$$i\Pi(\mathbf{k}, \omega_k) = i \frac{e^2}{8} \frac{\mathbf{k}^2}{\sqrt{v_F^2 \mathbf{k}^2 - \omega_k^2}}. \quad (4)$$

One can check that the higher-order corrections to $\Pi(\mathbf{k}, \omega_k)$ are independent of E_c , after renormalization of the three-point vertex and the electron self-energy. This comes from the fact that, after subtraction of all the divergent dependences on E_c coming from these objects, the structure of the Feynman diagrams for Π remains the same as for the one-loop order.²⁰ We may consider formally the result (4) as the leading order in a $1/N$ approximation. The fact that Π does not scale itself with E_c ensures that subleading corrections do not become relevant as the cutoff approaches the Fermi level, and that one can rely on the random-phase approximation (RPA) for the low-energy description of the theory.

By using the dressed propagator of the interaction in the RPA,

$$\langle \phi(\mathbf{k}, \omega_k) \phi(-\mathbf{k}, -\omega_k) \rangle = \frac{-i}{2|\mathbf{k}| + \frac{e^2}{8} \frac{\mathbf{k}^2}{\sqrt{v_F^2 \mathbf{k}^2 - \omega_k^2}}}, \quad (5)$$

one is able to perform a partial sum of perturbation theory in the computation of the self-energy $\Sigma(\mathbf{k}, \omega_k)$ and the three-point vertex $\Gamma(\mathbf{k}, \omega_k; \mathbf{p}, \omega_p)$. These two objects are related, since as long as the scalar potential ϕ is introduced by the minimal prescription $i\partial_0 \rightarrow i\partial_0 + e\phi$, a Ward identity similar to that of quantum electrodynamics can be derived:

$$\frac{\partial}{\partial \omega_k} \Sigma(\mathbf{k}, \omega_k) = \Gamma(\mathbf{k}, \omega_k; \mathbf{k}, \omega_k). \quad (6)$$

Thus the renormalization of Γ exactly matches the electron wave function renormalization as computed from the self-energy Σ . This leads to the relation between charge and ϕ -field renormalization factors $Z_e Z_\phi = 1$. The no-scaling of the polarization tensor means that $Z_\phi = 1$, which in turn implies the absence of charge renormalization in the model.

We focus then on the self-energy in GW approximation

$$i\Sigma(\mathbf{k}, \omega_k) = i2e^2 \int \frac{d^2p}{(2\pi)^2} \frac{d\omega}{2\pi} \frac{\omega_k - \omega + v_F \sigma \cdot (\mathbf{k} - \mathbf{p})}{v_F^2 (\mathbf{k} - \mathbf{p})^2 - (\omega_k - \omega)^2} \times \frac{-i}{2|\mathbf{p}| + \frac{e^2}{8} \frac{\mathbf{p}^2}{\sqrt{v_F^2 \mathbf{p}^2 - \omega^2}}}. \quad (7)$$

The imaginary part of the self-energy coming from Eq. (7) has been computed elsewhere.¹⁷ In this paper we are interested in the computation of the real part of $\Sigma(\mathbf{k}, \omega_k)$, which provides information about the nontrivial scaling of the quasiparticle weight and the Fermi velocity in the low-energy limit.

The terms linear in ω_k and $\mathbf{k}$ in the self-energy (7) display a logarithmic dependence on the high-energy cutoff E_c . Upon integration of the frequency from $-\infty$ to $+\infty$ and placing the bandwidth cutoff E_c in momentum space, $v_F |\mathbf{p}| < E_c$, the coefficients of the logarithmically divergent con-

tributions can be computed in terms of elementary functions of $g \equiv e^2/(16v_F)$. The renormalization of the electron propagator turns out to be given by

$$\begin{aligned} \frac{1}{G} &= \frac{1}{G_0} - \Sigma \approx Z^{-1}(\omega_k - v_F \boldsymbol{\sigma} \cdot \mathbf{k}) \\ &- Z^{-1} \omega_k \frac{8}{\pi^2} \left(g^2 + (2 - g^2) \left(1 - \frac{\arcsin g}{g \sqrt{1 - g^2}} \right) \right) \log E_c \\ &- Z^{-1} \omega_k \frac{8}{\pi} \frac{1}{g} \left(\frac{1 - g^2/2}{\sqrt{1 - g^2}} - 1 \right) \log E_c \\ &+ Z^{-1} v_F \boldsymbol{\sigma} \cdot \mathbf{k} \frac{8}{\pi^2} \left(1 - \frac{\sqrt{1 - g^2}}{g} \arccos g \right) \log E_c \\ &- Z^{-1} v_F \boldsymbol{\sigma} \cdot \mathbf{k} \frac{4}{\pi} \frac{1}{g} (1 - \sqrt{1 - g^2}) \log E_c, \end{aligned} \quad (8)$$

where $Z^{1/2}$ represents the scale of the bare electron field compared to that of the cutoff-independent electron field

$$\Psi_{\text{bare}}(E_c) = Z^{1/2} \Psi. \quad (9)$$

In the RG approach, we require the cutoff independence of the renormalized Green function, since this object leads to observable quantities in the quantum theory. For this purpose, Z and v_F have to be understood as cutoff-dependent effective parameters, that reflect the behavior of the quantum theory as $E_c \rightarrow 0$ and more states are integrated out from high-energy shells of the band. In this respect, the parameter v_F that appears in the ϕ -field propagator is also supposed to scale with the cutoff E_c . This agrees with the self-consistent character of the GW approximation, in which the parameters that enter in the polarization tensor have to match those obtained from the renormalized quasiparticle propagator. Phrased in the RG framework, the parameter v_F in the boson propagator has to approach, as $E_c \rightarrow 0$, the value of the Fermi velocity at the fixed point of the model.

We get the RG flow equations

$$E_c \frac{d}{dE_c} \log Z(E_c) = - \frac{8}{\pi^2} \left(2 + \frac{2 - g^2}{g} \frac{\arccos g}{\sqrt{1 - g^2}} \right) + \frac{8}{\pi} \frac{1}{g}, \quad (10)$$

$$E_c \frac{d}{dE_c} v_F(E_c) = - \frac{8}{\pi^2} v_F \left(1 + \frac{\arccos g}{g \sqrt{1 - g^2}} \right) + \frac{4}{\pi} v_F \frac{1}{g}. \quad (11)$$

Given that the electron charge e is not renormalized, we may write down the flow equation for the effective coupling constant $g = e^2/(16v_F)$:

$$E_c \frac{d}{dE_c} g(E_c) = \frac{8}{\pi^2} \left(g + \frac{\arccos g}{\sqrt{1 - g^2}} \right) - \frac{4}{\pi}. \quad (12)$$

In the weak-coupling regime, one may check that the renormalization of both the Fermi velocity and the electron wave function takes place in the expected direction. The quasiparticle weight Z at small g is smaller than the bare value mea-

FIG. 1. Flow of the coupling constant for different bare values.

sured before integration of high-energy modes. The Fermi velocity v_F flows to higher values in the infrared, and the density of states around the Fermi energy decreases, as a consequence of screening effects. This ensures the consistency of the weak-coupling phase, where the results of perturbation theory become increasingly reliable in the low-energy limit. However, the most important point concerns the possible existence of a different phase at large values of g . In this respect, the flow equations (10) and (12) can be analytically continued to values $g > 1$, by simple use of the formula $\arccos z = i \log(z - i\sqrt{1 - z^2})$. The flows of the coupling constant and the electron wave function are then differentiable across $g = 1$, which shows that the apparent singularity at this point has no physical meaning.

The right-hand side of Eq. (12) is a monotonous function of g , taking into account the mentioned analytic continuation. This means that there is no phase different from that of the perturbative regime, and that the strong-coupling regime is connected to it through RG transformations. The RG flow represented in Fig. 1 shows that the perturbative regime is attained at low energies, starting from fairly large bare values of the coupling constant.

The present analysis is relevant to the phenomenology of the graphite layers. It shows that, even in such a system with unconventional quasiparticle lifetimes, the low-energy behavior is governed by a fixed point which can be described as a Fermi liquid, as Z tends to a constant in the infrared (see Fig. 2), unlike for the line of nontrivial fixed points which characterize Luttinger liquids. On the other hand, the scaling behavior near the fixed point should give a more precise estimate of the quasiparticle lifetimes measured experimentally. Taking into account the results of Ref. 17 and the flow of the coupling constant in the infrared, the imaginary part of the self-energy has to behave in the form $\text{Im} \Sigma \sim g^2 \omega \sim \omega / \log^2(\omega)$. This expression of $\text{Im} \Sigma$ is consistent with the low-energy limit of the real part $\text{Re} \Sigma \sim g^2 \omega \log(\omega) \sim \omega / \log(\omega)$, as this actually matches the asymptotic behavior dictated by the Kramers-Kronig relation. We note, however, that the logarithmic correction to the linear quasiparticle decay rate cannot be discerned at present in the experiments, as these only cover one order of magnitude in energy about the Fermi level.¹⁶

Our study stresses the anomalous screening properties of

FIG. 2. Wave function renormalization for bare coupling constant $g = 5$ (thick line) and $g = 1$ (thin line).

the Coulomb interaction in low-dimensional systems. This fact has also been put forward recently in a different framework, pointing out that in one- and two-dimensional systems the screening of the long-range interactions goes in the direction of reducing the electron correlations.²¹ In the RG approach we see how this effect arises under the form of a renormalization of v_F . Such nontrivial scaling of the Fermi velocity in the infrared seems to be also present in systems with gauge interactions.^{12,13}

To summarize, semimetals described by the two-dimensional Dirac equation, such as a graphite layer, show significant differences with respect to the properties of standard Fermi liquids with (screened) Coulomb interactions. In the present problem, the quasiparticle lifetime goes like $\sim \log^2(\omega)/\omega$, while the enhancement of the Fermi velocity implies the vanishing of the effective coupling in the infrared. We expect the same behavior in three-dimensional zero-gap semiconductors, which are also described by an effective Dirac equation.

In the context of more general long-range interactions, the system with a Dirac sea also behaves differently with respect to the conventional Fermi sea, as in the former an interaction $V(q) \sim 1/|q|^{1+\epsilon}$ already departs from the Fermi-liquid universality class for $\epsilon > 0$. In our RG framework, the interaction becomes relevant no matter how small ϵ may be, and a nontrivial fixed point can be found away from the origin within the ϵ expansion.

The logarithmic behavior that we have found also affects some thermodynamic quantities like the specific heat or the susceptibility, which pick up logarithmic corrections as $T \rightarrow 0$. These are the signature of the marginal character of the interaction, which has in our model the precise degree of singularity to separate regimes with Fermi-liquid ($\epsilon < 0$) and non-Fermi-liquid behavior ($\epsilon > 0$).

We would like to thank C. Castellani for useful remarks on the paper. This work has been partially supported by the Ministerio de Educación y Cultura Grant No. PB96-0875.

¹R. Shankar, Rev. Mod. Phys. **66**, 129 (1994).

²P. W. Anderson, Phys. Rev. Lett. **64**, 1839 (1990); **65**, 2306 (1990).

³J. Feldman, M. Salmhofer, and E. Trubowitz, J. Stat. Phys. **84**, 1209 (1996).

⁴K. Yokoyama and H. Fukuyama, J. Phys. Soc. Jpn. **66**, 529 (1997).

⁵C. J. Halboth and W. Metzner, Phys. Rev. B **57**, 8873 (1998).

⁶Z.-X. Shen *et al.*, Science **267**, 343 (1995); K. Gofron *et al.*, Phys. Rev. Lett. **73**, 3302 (1994).

⁷J. González, F. Guinea, and M. A. H. Vozmediano, Phys. Rev. Lett. **79**, 3514 (1997).

⁸P.-A. Bares and X.-G. Wen, Phys. Rev. B **48**, 8636 (1993).

⁹A. Houghton, H.-J. Kwon, J. B. Marston, and R. Shankar, J. Phys.: Condens. Matter **6**, 4909 (1994).

¹⁰C. Castellani and C. Di Castro, Physica C **235-240**, 99 (1994); C. Castellani, C. Di Castro, and A. Maccarone, Phys. Rev. B **55**, 2676 (1997).

¹¹J. Gan and E. Wong, Phys. Rev. Lett. **71**, 4226 (1993).

¹²C. Nayak and F. Wilczek, Nucl. Phys. B **417**, 359 (1994).

¹³S. Chakravarty, R. E. Norton, and O. F. Syljuasen, Phys. Rev. Lett. **74**, 1423 (1995).

¹⁴J. González, F. Guinea, and M. A. H. Vozmediano, Mod. Phys. Lett. B **7**, 1593 (1994); Nucl. Phys. B **424**, 595 (1994); J. Low Temp. Phys. **99**, 287 (1994).

¹⁵J. González, F. Guinea, and M. A. H. Vozmediano, Nucl. Phys. B **406**, 771 (1993).

¹⁶S. Xu *et al.*, Phys. Rev. Lett. **76**, 483 (1996).

¹⁷J. González, F. Guinea, and M. A. H. Vozmediano, Phys. Rev. Lett. **77**, 3589 (1996).

¹⁸We stick from now on to one of the Fermi points, which provides all the relevant physics since the Coulomb interaction becomes singular at small momentum transfer.

¹⁹C. Castellani, C. Di Castro, and W. Metzner, Phys. Rev. Lett. **72**, 316 (1994).

²⁰Technically, the Feynman diagrams for Π may be computed in dimensional regularization, giving rise to a well-behaved Γ function of half-integer negative argument.

²¹J. van den Brink and G. A. Sawatzky, cond-mat/9802132 (unpublished).