**Use of Wildlife Rehabilitation Centres in Pathogen Surveillance: A Case Study in
White Storks (*Ciconia Ciconia*)**

Camacho, MariaCruz ¹; Hernández, Jose Manuel ²; Lima-Barbero, Jose Francisco¹,
Höfle, Ursula ^{1*}

¹SaBio working group, Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-
UCLM-JCCM), Ronda de Toledo 12, 13005 Ciudad Real, Spain.

Camacho, M.C: Mariacruz.Camacho@uclm.es; Höfle, U: Ursula.Hofle@uclm.es;
Lima-Barbero, J.F.: josefranvet@gmail.com

² Plaza de España, Ciudad Real, Spain. Hernández, J.M.: jmhdez@telefonica.net

*Corresponding author: Ursula Höfle, SaBio Instituto de Investigación en Recursos
Cinegéticos. Ronda de Toledo 12, 13005 Ciudad Real, Spain. E-mail:
Ursula.Hofle@uclm.es; Tlf.: 0034-926 295450, Fax: 0034-926 295451

Running title: Pathogen surveillance in nestling and injured white storks.

Abstract

More than 70% of new human pathogens are zoonotic and many originate from the wildlife reservoir. Wildlife rehabilitation centres (WRC) are an easily accessible source for sample and data collection for preventive surveillance, but data collected this way may be biased. We use white storks (*Ciconia ciconia*) as a model to compare pathogen prevalence obtained in the field and WRC. We address factors that may affect disease prevalence data like origin, the age group and the “diseased” state of WRC admissions. In this study we compared prevalence of *E. coli* and *Salmonella* spp. in the digestive tract; antibodies against West Nile virus, avian influenza and Newcastle disease virus, and antimicrobial resistance patterns of *E. coli* between nestling and adult wild storks established in different habitats (n=90) and storks admitted to two different WRC (n=30) in the same region.

When age groups and colonies of origin were disregarded, the mean enterobacteria (*E.coli*, *Salmonella*) and viral antibody prevalence of the wild population (n= 90) were similar to prevalence observed in the individuals admitted to WRC (n= 30). However, in fledgling juvenile storks admitted to WRC, the prevalence of *Salmonella* spp. (13.3%), *E. coli* showing resistance to cefotaxime (37.9%) and against two antimicrobials at once (41.4%) were more similar to the prevalence in stork nestlings from landfill-associated colonies (7.9%, 37.1% and 48.6%, respectively for prevalence of *Salmonella* spp. and *E.coli* displaying, cefotaxime resistance and resistance against two antimicrobials), and significantly higher than in colonies located in natural habitats (0%; 10.5% and 15.8%, respectively).

Thus, pathogen surveillance in individuals from an abundant species admitted to WRC is useful to monitor overall mean prevalence, but for certain pathogens may not be sufficient to detect differences between local populations. In addition, the ecology of the

tested species and the specific temporal, spatial and age group distribution of WRC admissions have to be taken into account.

Keywords: Antimicrobial resistance; *Enterobacteriaceae*; landfills; white stork; surveillance; wildlife rehabilitation centre.

Introduction

Most human pathogens are zoonotic and many originate from a wildlife reservoir (Jones et al., 2008). Especially birds are frequently considered reservoirs of pathogens for domestic animals and humans. Avian Influenza viruses and *Flaviviruses* such as West Nile and Usutu virus are maintained in avian reservoirs (Jones et al., 2008). Urban birds may carry zoonotic bacteria (Palomo et al., 2013; Vazquez et al., 2010). Due to their ability to travel long distances and forage in humanized environments birds may recirculate pathogens into the human or domestic animal populations. Consequently, surveillance and monitoring of pathogens in avian wildlife has become an important preventive tool (Stallknecht et al., 2007). Wildlife rehabilitation centres (WRC) can be a valuable resource for surveillance and monitoring of the prevalence of certain pathogens in wildlife. Stitt et al. (2007) documented that in comparison with other surveillance methods WRC received a wider variety of taxa from a greater geographical area. Sick and injured wildlife received at the WRC might be more likely to suffer from diseases, facilitating pathogen emergence detection (Wendell et al., 2002; Kelly and Sleeman, 2003; Wobeser, 2006; Nemeth et al., 2008). A recent nation-wide study in Australia using data on wildlife diseases from WRC associated with zoos, concluded that this information could improve the capacity for the detection of emerging pathogens in national surveillance programs (Cox-Witton et al., 2014).

Spain has many private and public funded WRC (56 in mainland Spain) that could become a powerful tool in monitoring certain risk pathogens. The study of potential zoonotic pathogens presumably prevalent across the Spanish geography, in a widely distributed conspicuous species that is admitted frequently to WRC, such as the white stork (*Ciconia ciconia*), allows us to test this hypothesis. Almost all WRC of mainland Spain receive white storks regularly, but concentrated in the summer months, as most stork casualties are juvenile birds.

The white stork is a widely distributed species in Europe. In Spain the population has experienced a significant increase in recent decades because of the use of landfills as predictable food source (Blanco, 1996; Schulz, 1999). In addition, the Iberian Peninsula has become the wintering area for many storks from northern Europe (Tortosa et al., 2002). Juvenile white storks frequently suffer accidents and are admitted to WRC, especially during the first few months after fledging. At the same time, only a few weeks prior to fledging banding takes place in many colonies, thus creating an interesting scenario for cost effective testing of the same species in the field and after admission to WRC within a short time period. Using the white stork, and some of the pathogens to which it may be exposed as a model, our goal was to provide a better understanding of the usefulness of WRC as sampling points for disease surveillance in wild birds. More specifically, we wanted to explore if and how factors such as the colony of origin, age group and the “diseased” status among others of WRC admitted white storks affected the detection of the surveyed pathogens.

Materials and Methods

Ethical statement

In this study, all applicable international, national, and/or institutional guidelines for the care and ethical use of animals were followed. Capture, ringing, radio-tagging and sampling of adult and nestling white storks in the field and at WRC was approved and authorised under permit from the regional government, the Junta de Comunidades de Castilla – La Mancha. JCCM (avp_13_037_aut).

Study area

The study area is located in the province of Ciudad Real (39° 0' 0" N, 4° 0' 0" W) in the south-centre of the Iberian peninsula, with a mean altitude of 629m. The area has a Mediterranean climate with very hot summers and cold winters, with average annual temperatures between 13 and 14° C and an annual rainfall of 438 mm, concentrated in autumn and spring.

Study populations

We sampled free-living white storks (nestlings and adults) in four different colonies with a different degree of exposure to human residues, and juvenile (fledgling) white storks upon admission to WRC (Table 1). One of the colonies (n° 1) was located in a National Park, the second (n° 2) in an open oak forest with extensive sheep farming, while the other two colonies (n° 3, n° 4) were associated with urban waste landfills. One of these (n° 4) is directly associated with an open active landfill site, while colony n° 3 is located at a landfill site that was sealed in 2007. Satellite tracking data from two adults captured and tagged at each colony in May 2013 (data not shown) confirmed foraging in natural habitats during nestling raising for colonies one and two. Foraging in colony three included both natural habitat and an open active landfill site located at approximately 45km flight distance from the colony. Foraging in colony four was limited to the landfill.

Nine adult white storks were trapped at the nest and sampled, three at colony n° 4 and two each at colonies one to three. For the wider purpose of this study we distinguish two subgroups within the “juvenile stork” group: “nestlings” are juvenile storks yet unable to fly and thus bound to the nest dependent on provisioning by their parents. This is the group of juvenile wild storks sampled in the field. The juvenile wild storks sampled upon admission at the WRC are “fledglings”, this is they have left the nest and are thus mobile and mostly forage on their own.

Between the four colonies we sampled 81 white stork nestlings of approximately 40 to 50 days of age (Table 1). Adult white storks were sampled in May, while stork nestlings were sampled in June. Fledgling juvenile storks (n=30) were sampled upon admission to two different WRC (15 each) during June-September of the same year. This is the time of the year with the highest number of admissions of storks to WRC, as the storks leave the nest (fledge) and are most vulnerable. Individuals in WRC were sampled on the day of arrival at the centre. The WRC sample also included one nestling that was sampled 48 hours after admission due to its young age and deteriorated condition.

Given the population of storks and the number of WRC in Spain our sample size is relatively small. For comparison of pathogen surveillance results within a logistically feasible way we designed the study to include two replicas for each category of storks (WRC, landfill exposed, natural habitat) in a cluster sampling approach. We wanted to approach field and WRC samples from the same geographical study area. While the number of storks admitted to the two WRC under study is generally high, in the study year the WRC admitted storks included in the study (n=30) were the total of admissions. Moreover, some of our sample colonies in the study area are very much smaller than others a fact that we accepted, as we wanted to study colonies that clearly evidence the actually existing strong differences with view to nesting habitat and foraging behaviour.

136

137 *Sampling*

138 Blood and cloacal swab specimens were obtained from all sampled birds (n=120). We
139 used sterile 25G needles and 5ml syringes to obtain blood samples (5 ml) from the
140 brachial vein, that were immediately transferred to sterile tubes with lithium heparin as
141 anticoagulant. Cloacal swabs were taken using sterile cotton swabs in AMIES transport
142 medium (Deltalab, Barcelona, Spain).

143 All samples were kept at 4° C until arrival at the laboratory and processed less than 12
144 hours after collection. Body weight, tarsus and head-bill length were recorded in free-
145 living nestlings (n=81).

146 *Laboratory analysis*

147 In the blood samples we first determined the haematocrit and measured total solids in
148 the plasma fraction using a hand held refractometer. The rest of the sample was
149 centrifuged for the separation of cells and plasma that were frozen in several aliquots for
150 further analysis.

151 We used one of the plasma samples for detection of antibodies against avian influenza
152 virus (AIV), Newcastle disease (ND, avian Paramyxovirus -1) and West Nile virus
153 (WNV). For this the samples were thawed, inactivated for 30 minutes at 56° C, and
154 tested using commercial blocking ELISAs (Ingezim Influenza A 1.0. FLU.K.3,
155 Ingenasa, Madrid, España; ID Screen, Newcastle competition, and ID Screen, West Nile
156 Competition, IDVet, Montpellier, France) according to the manufacturer's instructions.

157 Cloacal swabs in AMIES transport medium (Deltalab, Barcelona, Spain) were cultured
158 for the detection of *E. coli*. and *Salmonella* spp. For isolation of *E. coli* cloacal swabs

were plated on MacConkey agar (Scharlab S.L., Barcelona, Spain) and were incubated at 37° C for 24 hours. Phenotypic pattern of antimicrobial resistance of *E. coli* strains was assessed by culture in MacConkey media supplemented with 16µg/ml gentamicin (Sigma-Aldrich Chemical, Madrid, Spain), 4µg/ml cefotaxime (Sigma-Aldrich Chemical, Madrid, Spain) or 4µg/ml enrofloxacin (Sigma-Aldrich Chemical, Madrid, Spain) respectively. After incubation at 37° C for 24 hours, colonies morphologically compatible with *E. coli* were recorded as evidence of phenotypic resistance to the tested antimicrobial.

The choice of different antimicrobials was based on the frequency of use in Spain. Gentamicin and enrofloxacin are used in livestock and pets. Cefotaxime and similar cephalosporins are administered to humans. The concentrations used are those recommended by the National Antimicrobial Resistance Monitoring System (NARMS, Food and drug administration, Centres for disease control and prevention, United States Department of Agriculture, 2010).

Isolation of *Salmonella* spp. was performed according to the standard ISO 6579 method (2002). Colonies considered *Salmonella* spp. by their morphology were collected in ultrapure water for DNA extraction and confirmation of the identification as *Salmonella* spp. by amplification of the *invA* gene using the protocol described by Rahn et al. (1992). Briefly, amplification was carried out in a reaction mixture of a total volume of 30 µl containing: 2 µl of DNA; 0.2 µl Enzyme (5U/ µl); 3 µl of *invA*-L (0.01 mM) primer; 3 µl *invA*-R primer (0.01 mM); 3 µl Buffer (10x); 0.6 µl dNTPs (10 mM); 0.9 µl MgCl₂ (50 mM); 17.3 µl ddH₂O. The reaction was performed in a Techne thermal cycler TC-512 (Techne Inc. Cambridge, UK) using the following protocol: 94 ° C/3min; 40 cycles of 95 ° C/30s, 55 ° C/30s., 72 ° C/30s; and a final cycle of 72° C/10 min. The samples were subjected to electrophoresis in a 2% agarose gel. Bands were stained

using GelRed™ (Biotium, Hayward, USA) and visualized in a UV transilluminator (UVitec Ltd. Cambridge, UK).

Similar to *E. coli* we studied the phenotypic pattern of antimicrobial resistance in strains of *Salmonella* spp., by plating colonies on XLD agar supplemented with the same antibiotics and concentrations used in the evaluation of *E. coli* antimicrobial resistance. The plates were incubated at 37° C and examined for growth of colonies compatible with *Salmonella* spp. 24 hours later.

Statistical analysis

We calculated the body condition of white stork nestlings and adults sampled in the field according to the scaled mass index proposed by Peig and Green (2009). This index can be computed as: $M = M_i (L_0/L_i)^{bsma}$, where M_i and L_i are the body mass and the structural size measurement (tarsus length) of each individual respectively; $bsma$ is the scaling exponent estimated by the standardised major axis (SMA) regression of M on L , L_0 is the arithmetic mean value for the study population; M is the predicted body mass for individual i when the structural size body measure is standardized to L_0 . The scaling exponent “ $bsma$ ” has been calculated indirectly by dividing the slope from an ordinary least squares (OLS) regression (bols) by the Pearson’s correlation coefficient r . Normal distribution of the continuous variables (body condition index M , haematocrit, plasma total solids) was confirmed using the Shapiro Wilks test prior to further analysis. We compared body condition, haematocrit and plasma total solids between natural and landfill associated field populations using a Generalized linear model (GzLM) with the colony and the nest of origin as random factors in order to account for the expected dependency of individuals within each colony and the of sibling nestlings within each nest. We used Fisher’s exact test ($p \leq 0.05$) to explore differences in the prevalence of pathogens and antibodies between adult storks and nestlings. Finally we used

Generalized mixed linear models (GLMM) with a binary response and a logit regression and the colony and the nest of origin as random factors, to compare differences in prevalence of pathogens and antibodies among natural and landfill colonies and between white stork nestlings studied in the field and fledglings analysed upon admission to WRC. In addition to *E. coli* prevalence in the cloaca of individuals, we compared the prevalence of antimicrobial resistance of *E. coli* between the groups, namely the prevalence of *E. coli* resistant to gentamicin, enrofloxacin and cefotaxime, as well as the prevalence of resistance phenotypes against 2 and 3 antimicrobials at the same time. We also compared the prevalence of *Salmonella* spp. and WNV antibodies. All analysis were carried out using SPSS statistical software, version 19.0 (IBM®, SPSS Inc., Chicago, USA).

Results

Overall prevalence of *E. coli* in free-living (adult and nestling) white storks was 85.6% (77 out of 90) while prevalence of *Salmonella* spp. was 4.4% (4 out of 90). Prevalence of *E.coli* in WRC admitted storks was 96.7% (27 out of 30) and *Salmonella* spp. prevalence 13.3% (4 out of 30). Antibodies against AIV were not detected in the field. An adult and a nestling from two different nests situated on one of the landfill site colonies had antibodies against ND and only one adult white stork each from both natural and landfill site habitats had antibodies against WNV. While *E. coli* was isolated significantly more frequently in nestlings (90.1%, 73 out of 81) than in adult individuals (44.4%, 4 out of 9) (Fisher's exact test, $p=0.003$), *Salmonella* spp. was detected more frequently in adult white storks (Table 1).

White stork nestlings from landfill colonies ($M= 3327.18$, 95% CI: 3140.39-3513.97) were in significantly better body condition than individuals established in natural colonies ($M=2962.02$, 95% CI: 2830.54-3093.53; GzLM, $p=0.005$, $\beta= -0.226$,

SE=40.01). *E. coli* was present in both natural and landfill associated colonies, while *Salmonella* spp. was not isolated in natural colonies but was detected in one adult (20%, 1 out of 5) and three nestlings (7.9%, 3 out of 38) from landfill site associated colonies (GLMM, $p=0.003$) (Table 1). All *Salmonella* spp. strains isolated were susceptible to all antimicrobials tested. Due to the small number of isolates, we could not statistically analyse antimicrobial resistance patterns in salmonella. Phenotypic antimicrobial resistance of *E. coli* was significantly more prevalent in landfill site colonies than in natural colonies for two antimicrobials at once (48.6%, 17 out of 35, GLMM, $p=0.005$) and marginally significantly more prevalent for cefotaxime (37.1%, 13 out of 35, GLMM, $p=0.053$, Table 1, Figure 1).

For comparison of prevalences between the field and WRC we did not consider adults as only juvenile (fledgling and one nestling) storks were admitted to the WRC in our study period. Comparing white stork nestlings sampled in the field and fledgling white storks admitted to WRC, the prevalence of pathogens and antibodies studied did not differ significantly. However, if we considered storks from natural colonies and landfill site associated colonies separately, and compared pathogen prevalence to that in WRC admitted storks, significant differences became evident (Table 1, Figure 1). Thus prevalence of *Salmonella* spp. was similar in fledgling white storks sampled in WRC (13.3%, 4 out of 30) and nestlings from landfill site colonies (7.9%, 3 out of 38) while it was absent in colonies in natural habitats. *E. coli* resistant to cefotaxime was detected significantly more frequently in fledgling white storks at WRC (37.9%, 11 out of 29) and nestlings from colonies in landfill sites (37.1%, 13 de 35) than in nestlings from natural colonies (10.5%, 4 out of 38) (GLMM, $p=0.003$, Fig. 1). We also found resistance phenotypes against two different antibiotics significantly more frequently in nestlings from landfill colonies (48.6%, 17 out of 35), and fledglings from WRC

(41.4%, 12 out of 29) than in individuals from natural colonies (15.8%, 6 out of 38) (GLMM, $p=0.048$, Fig. 1). Antibodies against AIV and WNV were detected in one individual each admitted to WRC, while antibodies against ND, the only antibodies that had been detected in free-living nestlings were absent in WRC admitted storks (Table 1).

Discussion

In this study, we use the white stork to explore the pathogen-carrier-status information obtained in juvenile birds in the field and in WRC.

Our study evidences that at least during the breeding season significant differences in physiological condition and pathogen carrier status exist between colonies within a population. Nestlings in landfill colonies have a higher body condition index than those in natural colonies, probably due to the availability of abundant food in the vicinity, but are apparently more exposed to potential bacterial pathogens.

Carriage, of *E. coli* is similar between colonies, potentially because this *Enterobacteriaceae* is widely distributed in all types of environments and is part of the intestinal flora of white storks (Han et al., 2011). Natural colonies selected for our study are located in areas with extensive livestock farming, thus contact with livestock residues during foraging is possible. More frequent detection of *E. coli* in nestlings, could be related to a more diverse flora in the adults, longer cooling prior to culture in samples from adult birds and the comparatively smaller sample size in adult white storks.

In contrast, isolation of *Salmonella* spp. only from storks from landfill colonies may be due to the contact with human residues. In fact, the absence of *Salmonella* spp. in natural populations was previously observed in a study performed by Vlahovic et al.

(2004) in Croatia. We used the cited ISO 6579 method despite the risk of missing some of the true prevalence of *Salmonella* spp., because we wanted to apply a commonly used and recognised method for *Salmonella* spp. surveillance that is applied in numerous reference laboratories for *Salmonella* spp. serotypes that are of interest as zoonosis. Similarly, a higher prevalence of antimicrobial resistance patterns of the *E. coli* isolates to cefotaxime and resistance to two different antimicrobials was observed in storks exposed to human residues in the colonies associated with landfills. This has been previously observed in other species such as gulls exposed to human residues (Camarda et al., 2007).

The mean overall prevalence of *E.coli* and *Salmonella* spp. of white storks sampled in the field was similar to that detected in white storks admitted to the WRC, showing that data collected at the WRC reflects the situation in the field. However when considering nestlings from the different habitat types separately, the prevalence of *E. coli* and *Salmonella* spp. and phenotypes of antimicrobial resistance observed in storks admitted to WRC were similar to those observed in nestlings sampled in colonies associated with landfills and significantly higher than those observed in nestlings from colonies in natural habitats. That is, our results suggest that while during the nestling period the breeding habitat determines the diet and pathogen exposure of nestlings, this may change rapidly after fledging. Thus, fledged young storks from either nesting habitat may forage in landfills. This is further supported by satellite transmitter data that shows that adult storks from natural habitats use landfills after the end of the breeding season and by observation of long-distance identification rings from nestlings from natural habitats at nearby landfill sites (data not shown). Unfortunately, none of the patients in the WRC carried a ring or transmitter so that we were unable to determine its colony of origin. We also have to take into account that an age difference of one to two months

existed between nestling white storks sampled at the colony and juveniles sampled at the WRC. During this period fledging and a change from provisioning by the parent storks to independent foraging occur, two stressful events that could have a negative effect on the condition and immune status of the fledgling white storks and increase susceptibility to certain agents. In addition, the fact that storks admitted to WRC are generally fledgling or first year birds has to be taken into account during pathogen surveillance, as for example pathogens that adult storks may become exposed to during migration and wintering may not be detected in nestling or juvenile birds.

Viral seroprevalence is more difficult to assess as it can show exposure in the past, as well as miss recently infected seroconverting individuals, but it has been shown to be a useful tool in surveillance of viral activity within a population (e.g. Alba et al., 2014).

We used three avian viruses, frequently detected in wild birds across Europe and with a seasonal variation in their prevalence, all of them of surveillance interest either due to their zoonotic nature (AIV, WNV) or due to their importance for poultry (ND). AIV and ND are directly transmitted, while WNV is vector born and dependent on mosquito vector abundance and activity. WNV epidemics in the study area take place in late summer and early autumn, when bird migration is at its peak and mosquito populations are greatest, while peaks of AIV and ND prevalence are presumably associated with wintering (Perez-Ramirez et al., 2010).

Overall WNV seroprevalence, was low, but similar between storks sampled in the field and WRC. This is in contrast to the observation by Randall et al. (2012) who concluded that WNV exposure in individuals admitted to rehabilitation centres was not representative of the seroprevalence of WNV in natural populations of the same species. However, as the author stated, the low prevalence of this virus in the sampling area, working with too small an area or too many different orders of birds, may have led to

his results. In a study on WNV in Southern Spain, López et al. (2011) found a similar seroprevalence of WNV in the same species in the field and upon admission to WRC. Taking a closer look at our data we can see that in the field only adults are seropositive, consistent with exposure to the virus throughout their lives, perhaps through their migratory routes, as shown in other studies (Figuerola et al., 2007). In contrast, antibodies against WNV found in a juvenile stork admitted to a WRC could have been acquired in the field after fledging. Maternal antibodies transferred via egg yolk have been shown to be detectable for over one month (Komar, 2001; Gibbs et al., 2005). Thus, as stork nestlings fledge at the age of approximately 60 (58-64) days, it is possible but not very likely that these antibodies were of maternal origin.

These results show that the age group of a particular species that is admitted to WRC has to be taken into account when assessing the information WRC samples give of the situation in the field. As an example, while exposure of white storks to WNV may be more likely in adults, storks admitted to WRC are more frequently juveniles than adults. We only detected AIV antibodies in a juvenile white stork at a WRC. The time when we conducted our study may have influenced our findings. Previous studies show that AIV prevalence peaks in October and November (Pérez-Ramirez et al., 2010), when thousands of migratory birds come to Spain for wintering. However the peak of admissions of white storks to WRC is in summer and the bulk of these are recently fledged juvenile white storks. Nevertheless exposure of white storks to AIV appears to be low. In a previous study of our group only one of 129 faecal samples of white storks collected in the field tested positive for LPAIV and H11N9 was identified (Pérez-Ramirez et al., 2010). In Germany, out of over 600 nestlings, 103 fecal samples and 88 dead storks only three LPAIV positive faecal samples and two HPAIV H5N1 positive

dead storks were found (Müller et al., 2009). Also, antibodies against H5/H7 subtypes of AIV could not be detected in white storks from WRC in Germany (Kaleta and Kummerfeld, 2012). Antibodies against ND virus were detected in a nestling and an adult in a landfill colony. None of the storks admitted to WRC was positive. In contrast, Stenzel et al. (2008) found a seroprevalence of 20% in white storks from WRC in Poland and a long term study on white storks in WRC in Germany revealed haemagglutinating antibodies against NDV in 16 of 191 samples and carriage of virulent aPMV-1 virus in four individuals (Kaleta and Kummerfeld, 2012). In Spain, in a study on avian Paramyxovirus surveillance in semi-free-ranging birds 17 sera from white storks tested negative for antibodies against aPMV-1, but antibodies against aPMV- 8 and aPMV-9 were found in haemagglutination inhibition tests (Esperón et al., 2014). The significant differences in pathogen prevalence between the (in this study very few) adult and nestling wild storks, and the availability in WRC of only juveniles also illustrates how WRC origin sample and data sets are often biased towards the juvenile population of a species, which on the other hand may be of advantage for the detection of pathogen emergence.

Conclusions

Sample collection from wildlife casualties admitted to WRC can be a useful method for specific pathogen surveillance if the associated bias mediated by factors such as the diseased status, age group, and spatial or temporal distribution of admissions are taken into account. These associated confounding factors are most likely strongly related to the ecology of the species sampled that should receive an important consideration in surveillance programs. In our example, juvenile white storks admitted to WRC can be used as indicators of the emergence of *Enterobacteriaceae* carrying antimicrobial resistance mechanisms in wild populations but may be less good indicators for

circulation of WNV, NDV and AIV in the field, due to the timing of the peak of admissions.

Acknowledgements

This work was funded by the projects: RTA2011-00111-C03-02 and ERTA2013-0013 funded by the Instituto Nacional de Investigaciones Agrarias, INIA. María Cruz Camacho is a fellow of the Junta de Comunidades de Castilla – La Mancha (JCCM).

We thank the staff of SEO-Birdlife (Javier de la Puente) for help in capturing adult individuals and radio-tagging of these, and the staff of WRC the Chaparillo and CERI for providing samples of storks admitted as patients, especially Elena Crespo Junquera and Amalia García Crespo.

Conflict of interest statement

The authors declare that they have no conflict of interest.

References

- Alba A, Allepuz A, Napp S, Soler M, Selga I, Aranda C, Casal J, Pages N, Hayes EB, Busquets N., 2014. Ecological surveillance for West Nile in Catalonia (Spain), learning from a five-year period of follow-up. *Zoonoses Public Health* 61,181-91.
- Blanco, G., 1996. Population dynamics and communal roosting of white storks foraging at a Spanish refuse dump. *Colonial Waterbirds* 19, 273-276.
- Camarda, A., Circella, E., Giovanardi, D., Pennelli, D., Battista, P., Campagnari, E., Bruni, G. & Tagliabue, G., 2007. Avian Pathogenic *Escherichia coli* in Audouin gulls (*Larus audouinii*). Could they affect the surviving of the bird colonies? *Italian Journal of Animal Science* 6, 317-320.

404 Cox-Witton, K., Reiss, A., Woods, R., Grillo, V., Baker, R.T., Blyde, D.J., Boardman,
 405 W., Cutter, S., Lacasse, C., McCracken, H., Pyne, M., Smith, I., Vitali, S., Vogelnest,
 406 L., Wedd, D., Phillips, M., Bunn, C., Post, L., 2014. Emerging Infectious Diseases in
 407 Free-Ranging Wildlife—Australian Zoo Based Wildlife. Hospitals Contribute to National
 408 Surveillance. PLoS ONE 9(5): e95127. doi:10.1371/journal.pone.0095127

409 Esperón, F., Vázquez, B., Sánchez, A., Fernández-Piñero, J., Yuste, M., Neves, E.,
 410 Muñoz, M. J., 2014. Seroprevalence of paramyxoviruses in synanthropic and semi-free-
 411 range birds. Avian Diseases 58, 306-308.

412 Figuerola, J., Jiménez-Clavero, M.A., Rojo, G., Gómez-Tejedor C.& Soriguer, R.,
 413 2007. Prevalence of West Nile virus neutralizing antibodies in colonial aquatic birds in
 414 southern Spain. Avian Pathology 36, 209–212.

415 Gibbs, S.E.J., Hoffman, D.M., Stark, L.M., Marlenee, N.L., Blitvich, B.J., Beaty, B.J.
 416 and Stallknecht, D.E., 2005. Persistence of antibodies to West Nile Virus in naturally
 417 infected rock pigeons (*Columba livia*). Clinical and Diagnostic Laboratory
 418 Immunology 12, 665–667. doi:10.1128/CDLI.12.5.665-667.2005.

419 Peig, J., Green, A.J., 2009. New perspectives for estimating body condition from
 420 mass/length data: the scaled mass index as an alternative method. Oikos 118, 1883–
 421 1891.

422 Jones, K.E., Patel, N.G., Levy, M.A., Storeygard, A., Balk, D., Gittleman, J.L., Daszak,
 423 P., 2008. Global trends in emerging infectious diseases. Nature 451(7181):990-3. doi:
 424 10.1038/nature06536.

425 Kaleta, E.F., & Kummerfeld, N., 2012. Isolation of herpesvirus and Newcastle disease
 426 virus from white storks (*Ciconia ciconia*) maintained at four rehabilitation centres in

427 northern Germany during 1983 to 2001 and failure to detect antibodies against avian
 428 influenza A viruses of subtypes H5 and H7 in these birds. *Avian Pathology* 41, 383-389.
 429 Komar, N. 2001. West Nile virus surveillance using sentinel birds. *Annals of New York*
 430 *Academy of Sciences* 951, 58–73.
 431 López, G., Jiménez-Clavero, M.A., Vázquez, A., Soriguer, R., Gómez-Tejedor, C.,
 432 Tenorio, A., & Figuerola, J., 2011. Incidence of West Nile virus in birds arriving in
 433 wildlife rehabilitation centres in southern Spain. *Vector-Borne and Zoonotic Diseases*
 434 11, 285-290.
 435 Müller, T., Hlinak, A., Freuling, C., Muhle, Ru., Engelhardt, A., Globig, A., Schulze, C.,
 436 Starick, E., Eggers, U., Sass, B., Wallschlager, D., Teifke, J., Harder, T., Conraths, F.J.,
 437 2009. Virological Monitoring of White Storks (*Ciconia ciconia*) for Avian Influenza.
 438 *Avian Diseases* 53, 578-584.
 439 Palomo, G., Campos, M.J., Ugarte, M., Porrero, M.C., Alonso, J.M., Borge, C., Píriz,
 440 S., 2013. Dissemination of antimicrobial-resistant clones of *Salmonella enterica* among
 441 domestic animals, wild animals, and humans. *Foodborne Pathogens and Disease* 10(2),
 442 171-176.
 443 Pérez-Ramirez, E., Gerrikagoitia, X., Barral, M., & Höfle, U., 2010. Detection of low
 444 pathogenic avian influenza viruses in wild birds in Castilla-La Mancha (south central
 445 Spain). *Veterinary Microbiology* 146, 200-208.
 446 Rahn, S.A., De Grandis, R.C., Mcewen. S.A., Galan, J.E., Ginocchio, C., Curtiss, R., &
 447 Gyles, C.L., 1992. Amplification of an inv A gene sequence of *Salmonella typhimurium*
 448 by polymerase chain reaction as a specific method of detection of salmonella. *Molecular*
 449 *and Cellular Probes* 6, 271-279.

450 Randall, N.J., Blitvich, B.J., Blanchong, J.A., 2012. Efficacy of wildlife rehabilitation
 451 centres in surveillance and monitoring of pathogen activity: A case study with west Nile
 452 virus. *Journal of Wildlife Diseases* 48, 646-653.

453 Schulz, H., 1999. The world population of the White Stork (*Ciconia ciconia*) Results of
 454 the 5th International White Stork Census 1994/95. In, H. Schulz (Ed.): – White Storks
 455 on the up? Pp. 351-365 Proceedings of the International Symposium on the White Stork
 456 (Hamburg, 1996). NABU. Bonn.

457

458 Stallknecht, D.E., 2007. Impediments to wildlife disease surveillance, research, and
 459 diagnostics. *Current Topics in Microbiology and Immunology* 315, 445–461.

460 Stenzel, T., Tykałowski, B., Mazur-Lech, B., & Koncicki, A., 2008. Infections in
 461 wildlife birds - results of serological screening. *Bulletin of the Veterinary Institute in*
 462 *Pulawy* 52, 63-66.

463 Stitt, T., J. Mountfield, and C. Stephen., 2007. Opportunities and obstacles to collecting
 464 wildlife disease data for public health purposes: Results of a pilot study on Vancouver
 465 Island, British Columbia. *Canadian Veterinary Journal* 48, 83–90.

466 Tortosa, F.S., Caballero, J.M. & Reyes López, J., 2002. Effect of rubbish dumps on
 467 breeding success in the white stork in southern Spain. *Waterbirds* 25, 39-43.

468 Vlahovic, K., Matica, B., Bata, I., Pavlak, M., Pavićić, Z., Popović, M., Nejedli, S.,
 469 Dović, A., 2004. *Campylobacter*, *Salmonella* and *Chlamydia* in free living birds of
 470 Croatia. *Eur. J. Wildl. Res.* 50, 127–132. DOI 10.1007/s10344-004-0052-1

471 Wendell, M.D., Sleeman, J.M., & Kratz, G., 2002. Retrospective study of morbidity and
472 mortality of raptors admitted to the Colorado State University Veterinary Teaching
473 Hospital during 1995–1998. *Journal of Wildlife Disease* 38, 101–106.

474 Wobeser, G.A., 2006. *Essentials of disease in wild animals*. Blackwell Publishing
475 Professional, Ames, Iowa, USA, Ames, Iowa, 243 pp. ISBN 0-8138-0589-9.

476

Table 1. Prevalence of *E. coli*, and *Salmonella* spp., *E.coli* with antimicrobial resistance to gentamicin (G), cefotaxime (C) and enrofloxacin (E), multiresistance against 2 (MR2) and 3 (MR3) antimicrobials, and seroprevalence of West Nile (WNV), Newcastle (ND) and Avian Influenza (AI) virus in white storks by age, habitat and colonies (National Park (1), open oak forest with extensive sheep farming (2), closed landfill (3), active landfill (4)) and WRC

Population	Age	PREVALENCE n/N (%)		PREVALENCE OF ANTIMICROBIAL RESISTANCE PATTERNS n/N (%)					SEROPREVALENCE n/N (%)		
		<i>E. coli</i>	<i>Salmon.</i>	G	C	E	MR3	MR2	WNV	ND	IA
NATURAL	1	Nestling	13/15 86.8%	0/15 0%	7/13 53.8%	2/13 15.4%	2/13 15.4%	1/13 7.7%	3/13 23.1%	0/15 0%	0/15 0%
		Adult	0/2 0%	0/2 0%	0/0 0%	0/0 0%	0/0 0%	0/0 0%	0/2 0%	0/2 0%	0/2 0%
	2	Nestling	25/28 89.3%	0/28 0%	5/25 20%	2/25 8%	11/25 44%	2/25 8%	3/25 12%	0/28 0%	0/28 0%
		Adult	1/2 50%	0/2 0%	1/1 100%	0/1 0%	0/1 0%	0/1 0%	2/2 100%	0/2 0%	0/2 0%
LANDFILL	3	Nestling	27/30 90%	2/30 6.7%	17/27 63%	8/27 29.6%	15/27 55.5%	5/27 18.5%	13/27 48.1%	0/30 0%	1/30 3.3%
		Adult	1/2 50%	0/2 0%	0/1 0%	0/1 0%	0/1 0%	0/1 0%	2/2 100%	1/2 50%	0/2 0%
	4	Nestling	8/8 100%	1/8 12.5%	6/8 75%	5/8 62.5%	3/8 37.5%	2/8 25%	4/8 50%	0/8 0%	0/8 0%
		Adult	2/3 66.7%	1/3 33.3%	0/2 0%	0/2 0%	0/2 0%	0/2 0%	0/3 0%	0/3 0%	0/3 0%
Total Free-living	Nestling & Adult	77/90 85.6%	4/90 4.4%	36/77 46.7%	17/77 22%	31/77 40.2%	10/77 13%	23/77 29.9%	4/90 4.4%	2/90 2.2%	0/90 0%
WRC	Fledg-ling	29/30 96.7%	4/30 13.3%	13/29 44.8%	11/29 37.9%	12/29 41.4%	6/29 20.7%	12/29 41.4%	1/27 3.7%	0/27 0%	1/27 3.7%

Fig. 1 Prevalence of antimicrobial resistance patterns to *E. coli* in white storks analyzed in natural and landfill colonies and WRC (G = Gentamicin, E = Enrofloxacin, C = Cefotaxime; MR = multiresistance against two (MR2) or three antibiotics (MR3). Significant differences ($p < 0.05$) are marked with “*”