1 EFFECTS OF FLOW REGULATION AND NON-NATIVE SPECIES ON FEEDING
2 HABITS OF EURASIAN OTTER *LUTRA LUTRA* IN MEDITERRANEAN
3 TEMPORARY RIVERS
4

5 J. BUENO-ENCISO,^{a*} F. DÍAZ-RUIZ,^b D. ALMEIDA,^c and P. FERRERAS^b
6

7 ^a *Facultad de Ciencias Ambientales y Bioquímica, Universidad de Castilla-La Mancha,*
8 *Av. Carlos III s/n, 45071 Toledo, Spain*

9 ^b *Instituto de Investigación en Recursos Cinegéticos (IREC, UCLM-JCCM-CSIC),*
10 *Ronda de Toledo s/n, 13071 Ciudad Real, Spain*

11 ^c *Centre for Conservation Ecology and Environmental Change, Bournemouth*
12 *University, Poole, Dorset BH12 5BB, U.K.*
13
14

15 *Correspondence to: J. Bueno-Enciso, Facultad de Medio Ambiente, Universidad de
16 Castilla-La Mancha, Av. Carlos III s/n, 45071 Toledo, Spain.

17 E-mail: jbuenoenciso@gmail.com
18
19
20
21
22
23
24
25

ABSTRACT

In Mediterranean temporary rivers, ecological resources greatly fluctuate due to the high hydrological variability throughout the year. However, flow regulation prevents this natural regime and commonly entails associated non-native species, which change the structure of aquatic communities. Nonetheless few studies have tested the interaction of these two disruptive factors (flow regulation and non-native species) and their synergistic effects on the Eurasian otter (*Lutra lutra*) diet at the river-scale. The aim of this study was to compare the seasonal feeding habits of the otter between a temporary non-regulated stretch and two regulated stretches invaded by non-native species in a Mediterranean water course. The Bullaque River (Guadiana River basin, central Spain) was seasonally sampled for otter spraints and prey abundance assessed from December 2009 to November 2010. Three stretches were considered: High (source, non-regulated), Medium (transition, regulated) and Low (confluence, regulated). Diet varied from native prey in the High stretch (amphibians, insects and endemic cyprinids) to non-native species in the Low stretch (red-swamp crayfish *Procambarus clarkii* and pumpkinseed sunfish *Lepomis gibbosus*). Seasonally, ingested biomass of native prey increased in spring. Diet was more diverse in the High stretch. Otter neutrally selected native cyprinids in the high stretch throughout the year; whereas crayfish was selected in the other two stretches. Overall results showed flow regulation and non-native species have increased prey availability for the otter; however this paper highlights the importance of maintaining natural regimes in Mediterranean temporary rivers to conserve native communities and thus least-impacted food webs in Iberian freshwaters.

KEY WORDS: bioinvasion; diet; Iberian Peninsula; *Lutra lutra*; prey selection;
temporary rivers

SHORT TITLE: Diet of Eurasian otters in Mediterranean temporary rivers

INTRODUCTION

Temporary rivers are highly represented in Mediterranean regions because of the influence of climate, characterized by cool, wet winters and summer droughts (LeHouórou, 1990). The discharge regime generally follows this rainfall pattern, with torrential floods usually occurring in autumn and winter and minimum flow and severe droughts in summer (Gasith and Resh, 1999). As a consequence, ecological resources greatly fluctuate within the year in Mediterranean temporary rivers (Larned *et al.*, 2010) and species living there are highly specialized to cope with this high hydrological and resource variability (Williams, 1996).

Flow regulation, resulting from the construction of dams, reduces the effect of droughts and prevents natural flooding (Nilsson *et al.*, 2005; Wang *et al.*, 2011). In Mediterranean temporary rivers, flow regulation changes composition and structure of aquatic communities such as fish (Godinho and Ferreira, 2000; Grown and Grown, 2001), becoming dominated by non-native species (Pedroso *et al.*, 2007; Basto *et al.*, 2011). In the Iberian Peninsula, non-native fishes are increasing their ranges and this promotes the decline of the endemic fish fauna, both in their ranges and abundances through a variety of biotic interactions (see Leunda, 2010 for a comprehensive review). Among non-native fishes, two centrarchids, pumpkinseed sunfish *Lepomis gibbosus* (L., 1758) and largemouth bass *Micropterus salmoides* (Lacépède, 1802), are among the most widespread species in the Iberian Peninsula (e.g. Blanco-Garrido *et al.*, 2008). Also an invasive crustacean, the red-swamp crayfish *Procambarus clarkii* (Girard, 1852), is very widespread in the Iberian Peninsula and has deeply altered the Iberian freshwater ecosystems where it has been introduced (Geiger *et al.*, 2005), particularly in relation to food web structure (Tablado *et al.*, 2010). Nonetheless few studies have tested the interaction of these two disruptive factors (i.e. flow regulation and non-native

species) and their synergistic effects on the diet of a top-predator, including seasonal and spatial variation.

The Eurasian otter *Lutra lutra* (L., 1758) is a top-predator and key-species in the aquatic community of European inland waters (Ruiz-Olmo and Jiménez, 2009; Clavero *et al.*, 2010; Almeida *et al.*, 2012a), that contributes to maintain the ecological balance of freshwater ecosystem (Chanin *et al.*, 2003; Miranda *et al.*, 2008). In Mediterranean temporary rivers, summer droughts pose a handicap for otters (Ruiz-Olmo *et al.*, 2007), since it is the most important limiting factor in their distribution and abundance, because of the great fluctuation of prey availability (Prenda *et al.*, 2001; Ruiz-Olmo *et al.*, 2001). As a consequence, this seasonal factor affects otter breeding, carrying capacity and mortality (Kruuk and Carss, 1996; Ruiz-Olmo and Delibes, 1998; Ruiz-Olmo and Jiménez, 2009).

The aim of the present study was to assess the effects of flow regulation and the associated non-native species on the feeding habits of a top-predator, the Eurasian otter, in Mediterranean temporary rivers. For this purpose, we studied the feeding habits of the otter throughout a year along a partially regulated Mediterranean river of the Iberian Peninsula. Specifically, we seasonally compared diet, trophic diversity, prey availability and prey selection between three stretches in the Bullaque River (central Spain). We hypothesized that flow regulation and non-native species will affect the feeding habits of the otter and predicted that: (i) otters will feed more on red-swamp crayfish in regulated stretches, because though less energetic is easier to capture; (ii) otter trophic diversity will be higher in the non-regulated stretch, due to its higher seasonal variability and (iii) otters will still select native prey over non-native.

METHODS

Study area

The field sampling was carried out in the 1019 km² Bullaque River catchment (altitude: 550–620 m.a.s.l.), located in the Guadiana River basin (central Spain, 39°11'N - 4°15'O, Figure 1). The area is characterised by a continental Mediterranean climate, with rainfall from late autumn to spring (500–800 mm), whereas summer is hot and dry. Annual mean temperature ranges between 9 and 14°C. The lowest temperatures are recorded in December (–5°C) and the highest in August (43°C), (Almeida, 2008). Land use is mainly characterised by agricultural activity (e.g. corn and wheat crops, pastures for raising cattle and sheep). The Bullaque River (94 km length) includes a dam and a reservoir called Torre de Abraham (Figure 1). Upstream the reservoir, the river has an intermittent flow regime with seasonal flooding (autumn, winter) and severe droughts (summer); downstream the dam, the river has a regulated flow regime with weak seasonal fluctuations. Regarding biota, the particular invertebrate communities, fish assemblages and riparian vegetation of Bullaque River are well described in Almeida *et al.* (2012b, 2013).

Field sampling

We considered three stretches in the river according to their different hydrological and ecological characteristics: 1. *High stretch*: from the source of the river and its headwaters to the reservoir, included a tributary, the Milagro River (Figure 1). Both watercourses are lotic and oligotrophic temporary rivers, with narrow and medium-high speed flow; they are strongly affected by the climate conditions, becoming intermittent with a succession of pools of variable dimensions with little or no water flow in summer. Flow discharge is similar for both water courses, ranging between 0.3 m³/s in summer and 0.9 in winter. Their bank vegetation is well preserved.

150 2. *Medium stretch*: it begins at the outlet of the reservoir and it includes the transition of
151 the river. It is characterized by a wider and deeper channel, and it is less influenced by
152 the meteorological conditions, keeping a minimum ecological flow all the year. It flows
153 along an area submitted to an intensive agrarian and cattle exploitation, which have
154 converted the gallery forest in isolated stains of ash trees (*Fraxinus angustifolia*; Vahl,
155 1804) and Mediterranean scrubland (mainly *Crataegus monogyna*, Jacq, 1775; *Rubus*
156 sp.; *Rosa* sp. and *Cistus* sp.). Its flow discharge range between 0.9 m³/s in autumn and
157 1.2 in winter. 3. *Low stretch*: which is the confluence of the river, and is influenced by
158 the Guadiana River where it flows. The volume of flow here is maximum as it receives
159 water from more tributaries than the medium stretch. It is highly eutrophicated, because
160 of the agriculture runoff from the medium stretch and also because the river is naturally
161 dammed in many parts of this stretch, creating semi-permanent floodplains where water
162 is practically stagnant. This part of the river presents a discharge regime that oscillates
163 between 1.1 m³/s in autumn and 1.5 in winter. In this stretch of the river, the agrarian
164 activity is lower but it is more urbanized and has more human presence. See Almeida *et*
165 *al.*, 2013 for a more detailed description of the discharge regime profiles.

166 With the aim of assessing the variation of the otter diet throughout a year and
167 between stretches, we searched monthly for otter faeces (referred to as ‘spraints’
168 hereafter), from December 2009 to November 2010. According to the methodology
169 proposed by Ruiz-Olmo and Delibes (1998), we selected four sampling sites (600 m
170 river length) per stretch (Figure 1) and monthly collected 5–6 spraints per site whenever
171 it was possible; we grouped those spraints per stretch and season (3 months), resulting
172 in 60 spraints per group for diet analysis, a sample size higher than in other studies in
173 Mediterranean Rivers (Miranda *et al.*, 2006; Marques *et al.*, 2007; Novais *et al.*, 2010).
174 We did not collect more spraints per site to avoid disturbing natural sprainting

behaviour of otters. Also, only fresh spraints were collected to reduce loss of prey remains after defecation and to ensure regular presence of otters in the site (Almeida *et al.*, 2012b). In total, we collected 731 spraints for diet analysis.

In order to assess prey availability, fish and crayfish biomasses (measured as kg ha⁻¹) were estimated once each season at each sampling site by using block nets and electrofishing (2000 W DC generator at 200–250 V, 2–3 A) in an upstream direction, following the removal sampling without replacement or Zippin's method (1956), with three passes made (sampling time for each pass 20–30 minutes). Fish and crayfish were immediately immersed in an innocuous solution of anaesthetic (MS-222 at 0.1g L⁻¹), identified to the species level, counted and weighed (± 0.1 g). Fish were kept in a tank and supplied with oxygen (two aerators Aera, portable battery pump) until fully recovery before releasing them. All field procedures were complied with animal use and care regulations of Europe and Spain (specific Licence Code: DGPF/MRP-2010 for Scientific Field Research in Castilla-La Mancha, Spain). Electrofishing was performed by trained personnel (i.e. the holder of the Licence, D.A.), who had already sampled for fish by electrofishing in the same study area for previous projects (e.g. Almeida *et al.* 2009; Almeida *et al.*, 2012b).

Dietary analysis

The diet of the otter was determined by analysing spraints and identifying indigestible parts of the food intake (e.g. bones, scales, hair and feathers). These analyses were done in the laboratory, following a standard methodology described by Beja (1997). Food items were identified to the lowest possible taxonomic level using a dedicated reference collection of scales and hair, as well as published literature (Day, 1966; Gállego and Alemany, 1985; Teerink, 1991; Prenda and Granado-Lorencio,

1992a; Conroy *et al.*, 1993; Chinery, 1997; Prenda *et al.*, 1997; Miranda and Escala, 2002). The minimum number of individuals of each prey item present in a spraint was estimated by integrating the number, position (left–right) and relative size of diagnostic hard structures (mainly vertebrae, pharyngeal arches and scales for fish, and endopodites/exopodites and telson for red-swamp crayfish).

Each identified prey item was considered as an ‘occurrence’, and we calculated four dietary indices commonly used in carnivore diet studies (Klare *et al.*, 2011). The *Frequency of Occurrence* (FO, percentage of spraints in which a prey item was present), *Relative Frequency of Occurrence* (RFO, percentage of the total number of occurrences corresponding to a certain prey item), the *Percentage of Numbers* (%N, : total number of individuals corresponding to a certain prey item / total number of individuals) and the *Percentage of Ingested Biomass* (%Biomass, multiplying the total number of individuals corresponding to a certain prey item by their average weight in the environment). Average weight for fish and crayfish were calculated from averaging the weights of conspecifics from the electrofishing sampling. Thus, for the calculations of %Biomass in each spraint, we used the average weight of the fishes and crayfishes electro-fished in the same sampling site and season; as we saw in a previous study done in the same river (D. Almeida, pers. Observ.), that the average weights of prey in the environment were similar to those captured by the otter (Almeida *et al.*, 2012b). For the rest of prey items, we assigned the following weights: insects, 1g; amphibians, 10 g; reptilians (only one species, *Natrix maura*), 50 g; rodent species, 20 g; rest of mammals and birds, 100g (Beja, 1996). We calculated the Shannon index (H') for trophic diversity. All scientific and common names of freshwater fishes have been checked according to Leunda *et al.* (2009).

Prey selection

As recommended by Lechowicz (1982), prey selection (preferences) for crayfish and the main fish species consumed was evaluated using the Vanderploeg and Scavia (1979) normalised electivity index (ε_i):

$$\varepsilon_i = [\alpha_i - (1/n)] / [\alpha_i + (1/n)], \text{ where } \alpha_i = (r_i/p_i) / \sum_{i=1} (r_i/p_i)$$

where r_i is the proportional abundance of prey i in the diet (Ingested Biomass _{i} /Ingested Biomass_{Total}), p_i is the proportional abundance of prey i in the environment (from electrofishing data), n is the number of prey types included in the analysis and α is the Manly-Chesson's alpha (Chesson, 1978). The electivity values range from -1 (negative selection) to 1 (positive selection), and zero implies neutral selection.

Statistical analyses

To simplify the analytical models, data were pooled per season: winter (December–February), spring (March–May), summer (June–August) and autumn (September–November). Thus, the statistical power of the remaining sources of variation is increased, which would otherwise be seriously compromised. We tested if our spatial and temporal (stretch and season) sub-sample size was representative of the spectrum of the otter diet in the river by plotting the cumulative curve of new resource items by number of sampled spraints after randomization (Marques *et al.*, 2007).

To assess the spatial and temporal interactions, as well as the variations in feeding habits and prey availability, we performed General Linear Models (GLMs: factorial and univariate ANOVAs) with the percentage of ingested biomass, H' , prey biomasses and ε_i as dependent variables, and stretch and season as factors, followed by

post-hoc tests (Tukey-Kramer honestly significant difference, HSD test). To test whether electivity significantly deviated from 0, one sample Student's *t*-test was used with Bonferroni corrections. The percentage of ingested biomass was used for statistical analysis because it is the index best reflecting the relative importance of food items in carnivore diet (Klare *et al.*, 2011), and the only one that could be compared with prey availability. The other diet indexes were also provided in order to enable comparisons with studies using them but not the ingested biomass.

For statistical analyses, proportions and electivity indices were arcsine transformed, whereas the remaining variables were log₁₀ transformed. Assumptions of normality of distributions and homogeneity of variance were verified using Shapiro-Wilk and Levene tests, respectively. All statistical analyses were performed with STATISTICA 7.0 (Statsoft INC., Tulsa, OK, USA). The significance level was set at $\alpha = 0.05$.

RESULTS

The cumulative curves of the number of prey items in otter diet per number of analysed spraints show that the size of the sub-samples collected in each stretch (figure 2) and season, as well as in the whole river, were representative of the otter diet.

Red-swamp crayfish was the main prey consumed by the otter across the river and throughout the year, representing about 40% of the total ingested biomass, followed by fishes with 30%. Among fishes, non-native northern pike *Esox lucius* (L., 1758) contributed with the greatest biomass to the diet, followed by other non-native fishes, the pumpkinseed sunfish and the common carp *Cyprinus carpio* (L., 1758). The third prey category in importance was amphibians with 14% of the total ingested biomass. Other noticeable prey was the common moorhen *Gallinula chloropus* (L., 1758). The

remaining prey items can be considered as of minor importance (any of them with <2% of ingested biomass, Table I). The diet showed significant spatial variation ($F_{30, 20} = 5.77$, $p < 0.001$). The High stretch was characterized by a higher consumption of endemic cyprinids such as southern straight-mouth nase *Pseudochondrostoma willkommii* (Steindachner, 1866) and calandino *Squalius alburnoides* (Steindachner, 1866), as well as amphibians and insects (mostly diving beetles) (Table II). In the Low stretch, the otter fed on a high proportion of pumpkinseed sunfish, reptilians and red-swamp crayfish. In the Medium stretch the otter had intermediate trophic characteristics between the other two stretches. There, the consumption of calandino was higher while the intake of reptilians, amphibians and insects were not. Even the consumption of pumpkinseed sunfish was lower than in the Low stretch (Table II). Seasonally, the overall composition of the otter diet also varied significantly ($F_{45, 30.49} = 1.87$, $p = 0.04$), in particular the Iberian arched-mouth nase *Iberochondrostoma lemmingii* (Steindachner, 1866) was more consumed in spring than in autumn (Table II). Focusing on the seasonal diet variation in each stretch separately, there were differences between them. While in the Medium and Low stretches prey items did not vary seasonally, they did in the High stretch ($F_{24, 3.5} = 47.38$, $p < 0.01$). In particular pumpkinseed sunfish and insects were more consumed in autumn than in the rest of the year.

Trophic diversity varied significantly along the river ($F_{2, 24} = 20.03$, $p < 0.001$), being maximum in the High stretch in contraposition with the Medium and Low stretches, where no differences were found (Figure 3). Regarding spatial variation, trophic diversity did not significantly vary between seasons ($F_{3, 24} = 2.09$, $p = 0.13$).

Results of electrofishing are shown in Table III. Total available biomass of fish and crayfish increased from the High stretch (mean = 12.79 kg ha⁻¹, SE = 5.58) to the Low stretch (mean = 28.32 kg ha⁻¹, SE = 9.85), although differences among stretches

were not statistically significant ($F_{2, 45} = 1.12, p = 0.33$). Conversely, composition of prey availability significantly differed between stretches ($F_{24, 50} = 2.23, p < 0.01$), with significant differences of biomass of southern Iberian spined-loach *Cobitis paludica* (De Buen, 1930) along the stream, being higher in the High stretch than in the rest of the river (Table IV). Total available biomass also varied seasonally ($F_{3, 44} = 9.49, p < 0.001$), being highest in summer (mean= 50.12 kg ha⁻¹, SE = 12.86) followed by spring (mean= 19.36 kg ha⁻¹, SE = 4.32), winter (mean= 4.91 kg ha⁻¹, SE = 3.44) and finally autumn (mean= 3.42 kg ha⁻¹, SE = 1.17) (Tukey test, $p < 0.05$). For particular prey categories, the introduced red-swamp crayfish was much more abundant in spring and in summer than in the rest of the year, opposite to calandino, which resulted much more abundant in autumn and winter (Table IV). Also, Iberian arched-mouth nase showed seasonal variations being more abundant in winter than in the rest of the year (Table IV). Available biomass of non-native Eastern mosquitofish *Gambusia holbrooki* (Girard, 1859) also varied seasonally, reaching a peak in summer (Table IV). The interaction between these two factors, Stretch and Season, was not statistically significant for any species ($F_{72, 141.82} = 0.93, p = 0.63$).

Globally, the otter showed significant negative electivity for most prey items in the whole river and over the year (t -tests, $p < 0.001$) with the only exception of red-swamp crayfish, which was neutrally selected (t -test, $p > 0.05$). However, the electivity index showed differences both spatially and seasonally ($F_{18, 32} = 3.59, p < 0.001$ and $F_{27, 47.37} = 4.65, p < 0.001$; respectively), the interaction between these two factors also was statistically significant ($F_{54, 86} = 2.17, p < 0.001$). Within stretches, southern straight-mouth nase was neutrally selected in the High stretch while it was avoided in the rest of the river (Table V), contrary to red-swamp crayfish which was avoided in the High stretch and neutrally selected in the rest of the river (Table V). Also, calandino showed

spatial variation, being more selected in the medium stretch than in the rest of the river (Table V). Seasonally, pumpkinseed sunfish was neutrally selected in summer and in autumn, in contrast to winter, when it was completely avoided (Table V). Red-swamp crayfish was highly positively selected in winter and avoided in spring and summer, being neutrally selected in autumn (Table V).

DISCUSSION

Considering the whole river and throughout the year, otter diet resembles that typical of regulated water courses and reservoirs, i.e. based on red-swamp crayfish and non-native fish species (López-Nieves and Hernando, 1984; Adrián and Moreno, 1986; Delibes and Adrián, 1987; Pedroso and Santos-Reis, 2006; Sales-Luis *et al.*, 2007), instead of that typical of Mediterranean temporary rivers (Ruiz-Olmo *et al.*, 1998; Bartolomé, 2001; Clavero *et al.*, 2003). Possibly the opportunistic predator behaviour of the otter (Balestrieri *et al.*, 2013), allows them to feed on the most abundant prey (Erlinge, 1969, Taastrom and Jacobsen, 1999). In our study, red-swamp crayfish, the most abundant prey in the river, was also the main otter prey. As in other places where the red-swamp crayfish has been introduced, it has become a major prey for a variety of predators (Delibes and Adrián, 1987; Peris *et al.*, 1994; Tablado *et al.*, 2010), and it has been hypothesized to be a key factor in the recovery of otter populations in the Iberian Peninsula (Ruiz-Olmo and Delibes, 1998). For this role as a factor of otter recovery, red-swamp crayfish should be also abundant during the critical periods of the year (i.e. winter floods and summer droughts) (Beja, 1996). In our study, red-swamp crayfish was highly consumed throughout the year, even in winter when it usually stays in burrows (Niquette and D'Abramo, 1991; Correia and Ferreira, 1995). Among fishes, the second prey category in importance, non-native species contributed with the highest proportion

of biomass to the otter diet in the whole river. Non-native species are related to reservoirs and regulated flows (Godinho *et al.*, 1998; Basto *et al.*, 2011), where they interact with, and force the decline of native species becoming the dominant species (Leunda, 2010) and also the major prey for otters (López-Nieves and Hernando, 1984; Adrián and Moreno, 1986; Pedroso and Santos-Reis, 2006; Sales-Luis *et al.*, 2007). Our results are in concordance with the preferences for prey type by otters (Ruiz-Olmo, 1995). Thus, among non-native species, the northern pike was the most important prey for the otter in terms of biomass. Also the higher abundance of pumpkinseed sunfish and red-swamp crayfish in the river may make them easier prey to capture than endemic cyprinids, considered preferred prey for the otter in Mediterranean rivers of the Iberian Peninsula (López-Nieves and Hernando, 1984; Callejo and Delibes, 1987; Ruiz-Olmo *et al.*, 1989; Ruiz-Olmo, 1995; Bartolomé, 2001; Morales *et al.*, 2004), despite the anti-predator body structures of the pumpkinseed sunfish (Blanco-Garrido *et al.*, 2008) or the less energetic contribution of the red-swamp crayfish to otter diet (Beja, 1996).

Is well known that otters are selective in prey size (Lanszki *et al.*, 2001) and consequently %Biomass is biased according to this behavior. To mitigate this source of error, studies assessing %Biomass incorporate regression equations to accurately calculate the actual weight of the specimens consumed found from the indigestible parts in the spraints (Prenda and Granado-Lorencio, 1992b; Copp and Kováč, 2003). However we did not use such regression equations since in a previous study (Almeida *et al.*, 2012b) it was seen that the average weight of the fishes and crayfishes electro-fished were similar to those selected by the otter in this river (D. Almeida, pers. Observ.).

The spatial variation in otter diet reflects the effect of the reservoir and flow regulation. In the High stretch, hardly influenced by the reservoir, the diet of the otter is similar to that described for Mediterranean temporary rivers; based on endemic

cyprinids and high amounts of invertebrates and amphibians (Adrián and Delibes 1987; Ruiz-Olmo, 1995; Bartolomé, 2001; Ruiz-Olmo *et al.*, 2002). Conversely, the diet of the otter in the regulated Medium and Low stretches resembles that described in reservoirs regarding non-native species (López-Nieves and Hernando, 1984; Adrián and Moreno, 1986; Pedroso and Santos-Reis, 2006; Sales-Luis *et al.*, 2007; Basto *et al.*, 2011). Seasonally and considering the river as a whole, otter diet did not show any significant variation; but if we focus on each stretch separately, the differences come to light. In particular, within the High stretch, diet varied seasonally whereas it did not in the rest of the river. This result highlights the temporary flow regime of the High stretch, which presents different prey types in each season due to its fluctuating ecological conditions (Gasith and Resh, 1999). This contrasts with the stability of the ecosystem downstream the reservoir, where most of the prey types are available for otters throughout the year. According with the optimal foraging theory, generalist predators change prey foraging patterns according to their profitability (Ferrerías *et al.*, 2011). The high consumption of pumpkinseed sunfish and insects in autumn in the High stretch may be due to the scarcity of endemic cyprinids after the summer drought, and possibly to the use of the close reservoir as an alternative source of prey in the dry season (Basto *et al.*, 2011). Except for calandino, whose abundance significantly increased (Table IV), the availability of the rest of cyprinid species decreased in autumn (Table III), as it occurs in other Mediterranean streams (Mas-Marti *et al.*, 2010). This increase in the availability of calandino was followed by a non-significant increment in its consumption after the summer drought. However, an increase in the percentage of biomass for a particular prey type in the river is not necessarily related to an increase in its availability for the otter. Small pools where otters fish during summer in the High stretch increase their volume with the autumn rains and torrential flows drag fishes to

400 them, which result in an increase of available biomass (according to electrofishing) in
401 these habitats. However, an increase in depth and water volume in the pools can hinder
402 the capture of fishes by the otter (Barrientos *et al.*, 2003; Kruuk, 2006; Almeida *et al.*,
403 2012b). Trophic diversity varied spatially in accordance with the negative relationship
404 with water flow stability (Clavero *et al.*, 2003). The higher values of H' in the High
405 stretch than in the rest of the river are possibly due to the harsher environmental
406 conditions, which force the otter to prey on less profitable prey such as amphibians or
407 insects (Clavero *et al.*, 2008; Román, 2011).

408 Our results of prey selection show that, except for the red-swamp crayfish,
409 almost all prey items were used below their availability and no species was positively
410 selected, similar to Almeida *et al.* (2012a). These results differ from the trophic
411 behaviour stated for the otter; which establish that otters consume each different prey
412 item according to their particular availability (Clavero *et al.*, 2003; Remonti *et al.*,
413 2010). Our results are unusual and may be explained by the wide availability of a
414 variety of prey items in the river (i.e. pumpkinseed, red-swamp crayfish, calandino or
415 southern straight-mouth nase) and also by the generalist and opportunistic trophic
416 behaviour of the otter; which allows it to prey on most of them and as a result, the
417 encounter rate with one prey item in particular is divided among the wide variety of
418 available prey items; resulting in a prey selection under the particular availability of
419 each prey item in particular.

420 Although with our data we could not fit a clear functional response related to
421 red-swamp crayfish consumption, otters could be displaying a type II or III functional
422 response (Holling, 1959) in their last steps. This would imply that crayfish availability
423 had passed a threshold beyond which the otter searches actively for it. Nevertheless, this
424 does not happen in the High stretch, possibly because the availability of the invasive

crustacean in this part of the river is lower and otters search for other more profitable prey, mostly native cyprinids (such as the case of southern straight-mouth nase). This is due to its higher abundance or because they are easily captured by the otter in that part of the river (Barrientos *et al.*, 2003). Seasonally, otters increased the selection of pumpkinseed sunfish during the warmer seasons, possibly because this species availability increased during these months (Almeida *et al.*, 2009). The highly positive selection of red-swamp crayfish in winter may be overestimated because of its burrowing behaviour in the cold months of autumn and winter (Niquette and D'Abramo, 1991; Correia and Ferreira, 1995), which makes them less vulnerable to sampling methods. However, otters seem not to have problems in their capture, according to the lack of seasonal variation in crayfish consumption. The apparent avoidance of this type of prey in spring and summer should be due to the red-swamp crayfish consumption rate by the otter has reached an asymptote which is independent of the higher density of red-swamp crayfish populations during these seasons in comparison with the rest of the year.

Even though prey biomass did not significantly differ between stretches, it increased from the source to the mouth of the river, with prey biomass doubling between the High stretch and the Low stretch considering all year (142 kg ha^{-1} and 350 kg ha^{-1} respectively, data obtained from Table III), as it happens in other rivers (Townsend *et al.*, 2003; Davey and Kelly, 2007; Magalhães *et al.*, 2007). This is because Medium and Low stretches maintain a minimum flow throughout the year due to the flow regulation by the dam downstream, which mitigates or even neutralize the effect of seasonal floods or droughts. This is opposite to the High stretch, where the narrow and shallow channel limits its prey carrying capacity. But the High stretch also undergoes seasonal floods which drag the biota, and seasonal droughts, which reduce

habitat availability and suitability in several ways, resulting in a lower fish biomass (Gasith and Resh, 1999; Mas-Martí *et al.*, 2010). The species composition slightly varied between stretches which is in concordance with other studies (Matthews and Marsh-Matthews, 2003; Aparicio and Vargas, 2004; Magalhães *et al.*, 2007). Even so, a trend in its distribution can be appreciated, with the dominance of non-native species near the confluence of the river possibly because of the presence of the reservoir upstream, which acts as a source of non-native species (Godinho *et al.*, 1998) and the influence of the outflow in the Guadiana River. Seasonally, the differences found in available prey biomass can be explained above all by the availability increase of red-swamp crayfish in summer and, in a lesser extent, by the peak of availability of the pumpkinseed sunfish in the warm months of the year (Table III).

Our study reveals the dual effect of both flow regulation and non-native species in natural Mediterranean temporary rivers. On the one hand flow regulation increases the water availability throughout the year, which allows the maintenance and growth of non-native population species normally at expense of the native fish species (Leunda, 2010). This eventually results beneficial for otters and other generalist predators (Tablado *et al.*, 2010), since the carrying capacity of the river for them increases, due to the increase of prey availability. Otters also likely improved their breeding success and diminished their mortality rate (Kruuk and Carss, 1996; Ruiz-Olmo and Delibes, 1998; Ruiz-Olmo and Jiménez, 2009). On the other hand, these changes deeply modified the natural ecological processes of Mediterranean temporary rivers, i.e. changes in the composition and structure of aquatic communities and also in their natural trophic web; so it is detrimental in a broader conservation context. Thus, conservation efforts should give priority to preserve the Mediterranean temporary rivers, because they harbour a wider variety of species than Mediterranean regulated rivers, making them valuable

habitats, and avoid disturbing natural feeding behaviours of predatory species in freshwater environments (Basto *et al.*, 2011). Our non-regulated stretch maintains otter presence throughout the year, and more important, maintains the natural diet of the mustelid in this region. In view of the radical changes caused by flow regulation in temporary rivers, from species to ecosystem level, such type of actuations should be carried out only in unavoidable situations and the management should be aimed to preserve the original biota, avoiding the introduction of non-native species. Mitigating measures such as potamodromus fish-ways in the dams should be also implemented.

ACKNOWLEDGEMENTS

Special thanks to Máximo Florín and his team for showing us the study area facilitating the fieldwork. This study has been possible thanks to the funding to the project: “Convenio entre Confederación Hidrográfica del Guadiana y la UCLM para el desarrollo de estudios de investigación y seguimiento de las medidas ambientales de restauración fluvial del río Bullaque”. Also thanks to Javier Fernández de Simón for his invaluable assistance during laboratory work. J.B.E. held a pre-doctoral fellowship from Junta de Comunidades de Castilla-La Mancha.

REFERENCES

- Adrián M I, Moreno S. 1986. Notas sobre la alimentación de la nutria (*Lutra lutra*) en el embalse de Matavacas (Huelva). *Doñana Acta Vertebrata* **13**: 189–191.
- Adrián M I, Delibes M. 1987. Food habits of the otter (*Lutra lutra*) in two habitats of the Doñana National Park , SW Spain . *Journal of Zoology* **212**: 399–406.

498 Almeida D. 2008. *Ecología y conservación de la fauna fluvial en el Parque Nacional de*
 499 *Cabañeros: efectos de la degradación del hábitat y de la introducción de especies*
 500 *exóticas*. PhD Thesis, Complutense University of Madrid: Madrid.

501 Almeida D, Almodóvar A, Nicola GG, Elvira B. 2009. Feeding tactics and body
 502 condition of two introduced populations of pumpkinseed *Lepomis gibbosus*: taking
 503 advantages of human disturbances? *Ecology of Freshwater Fish* **18**: 15–23. DOI:
 504 10.1111/j.1600-0633.2008.00317.x.

505 Almeida D, Copp GH, Masson L, Miranda R, Murai M, Sayer CD. 2012a. Changes in
 506 the diet of a recovering Eurasian otter population between the 1970s and 2010.
 507 *Aquatic Conservation: Marine and Freshwater Ecosystems* **22**: 26–35. DOI:
 508 10.1002/aqc.1241.

509 Almeida D, Barrientos R, Merino-Aguirre R, Angeler DG. 2012b. The role of prey
 510 abundance and flow regulation in the marking behaviour of Eurasian otters in a
 511 Mediterranean catchment. *Animal Behaviour* **84**: 1475–1482. DOI:
 512 10.1016/j.anbehav.2012.09.020.

513 Almeida D, Merino-Aguirre R, Angeler DG. 2013. Benthic invertebrate communities in
 514 regulated Mediterranean streams and least-impacted tributaries. *Limnologia* **43**: 34–
 515 42. DOI: 10.1016/j.limno. 2012.02.003.

516 Aparicio E, Vargas MJ. 2004. Influence of hydrological variability on fish populations
 517 in Fuirosos stream. *IV Trobada d'estudiosos del Montnegre i el Corredor*. Barcelona:
 518 Diputació de Barcelona. Monografies, núm 4: 119–122 (in Catalan, abstract in
 519 English).

520 Balestrieri A, Remonti L, Vezza P, Prigioni G, Copp GH. 2013. Do non-native fish as
 521 prey favour the conservation of the threatened indigenous Eurasian otter?.
 522 *Freshwater Biology* **58**: 995–1007. DOI: 10.1111/fw.b.12102.

523 Barrientos R, Gil T, Hernando J, Iglesias A, Jiménez MD, Navarro E. 2003. Influencia de
524 un embalse sobre la distribución y alimentación de la nutria (*Lutra lutra* L.) en el río
525 Riaza (Segovia). *Galemys* **15**: 81–90.

526 Bartolomé MA. 2001. Alimentación de la Nutria (*Lutra lutra*) en el río Bergantes
527 (Castellón). *Dugastella* **1**: 35–42.

528 Basto MP, Pedroso NM, Mira A, Santos-Reis M. 2011. Use of small and medium-sized
529 water reservoirs by otters in a Mediterranean ecosystem. *Animal Biology* **61**: 75–94.
530 DOI: 10.1163/157075511x554455.

531 Beja PR. 1996. An analysis of otter *Lutra lutra* predation on introduced American
532 crayfish *Procambarus clarkii* in Iberian streams. *Journal of Applied Ecology* **33**:
533 1156–1170.

534 Beja PR. 1997. Predation by marine-feeding otters (*Lutra lutra*) in south-west Portugal
535 in relation to fluctuating food resources. *Journal of Zoology* **242**: 503–518.

536 Blanco-Garrido F, Prenda J, Narváez M. 2008. Eurasian otter (*Lutra lutra*) diet and prey
537 selection in Mediterranean streams invaded by centrarchid fishes. *Biological*
538 *Invasions* **10**: 641–648. DOI: 10.1007/s10530-007-9158-1.

539 Callejo A, Delibes M. 1987. Dieta de la nutria *Lutra lutra* (Linnaeus, 1758) en la cuenca
540 del alto Ebro, norte de España. *Miscellanea Zoologica* **11**: 353–362.

541 Chanin P. 2003. *Ecology of the European Otter*. Conserving Natura 2000. Rivers
542 Ecology Series No. 10. English Nature, Peterborough.

543 Chesson J. 1978. Measuring preference in selective predation. *Ecology* **59**: 211–215.

544 Chinery M. 1997. *Guía de campo de los insectos de España y de Europa*. Ediciones
545 Omega, S.A.: Barcelona.

546 Clavero M, Prenda J, Delibes M. 2003. Trophic diversity of the otter (*Lutra lutra* L.) in
547 temperate and Mediterranean freshwater habitats. *Journal of Biogeography* **30**: 761–
548 769. DOI: 10.1046/j.1365-2699.2003.00865.x.

549 Clavero M, Prenda J, Blanco-Garrido F, Delibes M. 2008. Hydrological stability and
550 otter trophic diversity: a scale-insensitive pattern?. *Canadian Journal of Zoology-*
551 *Revue Canadienne De Zoologie* **86**: 1152–1158. DOI: 10.1139/Z08-094.

552 Clavero M, Hermoso V, Brotons L, Delibes M. 2010. Natural, human and spatial
553 constraints to expanding populations of otters in the Iberian Peninsula. *Journal of*
554 *Biogeography* **37**: 2345–2357. DOI: 10.1111/j.1365-2699.2010.02377.x.

555 Conroy JWH, Watt J, Weeb JB, Jones A. 1993. A guide to the identification of pray
556 remains in Otter Spraint. *Occasional Publications of the Mammal Society*. No. 16.
557 The Mammal Society, London.

558 Copp GH, Kováč V. 2003. Biometric relationships between body size and bone lengths
559 in fish prey of the Eurasian otter *Lutra lutra*: chub *Leuciscus cephalus* and perch
560 *Perca fluviatilis*. *Folia Zoologica* **52**: 109–112.

561 Correia AM, Ferreira O. 1995. Burrowing behaviour of the introduced red swamp
562 crayfish *Procambarus clarkii* (Decapoda: Cambaridae) in Portugal. *Journal of*
563 *Crustacean Biology* **15**: 248–257.

564 Davey AJH, Kelly DJ. 2007. Fish community responses to drying disturbances in an
565 intermittent stream: a landscape perspective. *Freshwater Biology* **52**: 1719–1733.
566 DOI: 10.1111/j.1365-2427.2007.01800.x.

567 Day MG. 1966. Identification of hair and feather remains in the gut a faeces of stoats
568 and weasels. *Journal of Zoology* **148**: 201–217. DOI: 10.1111/j.1469-
569 7998.1966.tb02948.x.

570 Delibes M, Adrián MI. 1987. Effects of crayfish introduction on otter *Lutra lutra* food
 571 in the Doñana National Park, SW Spain. *Biological Conservation* **42**: 153–159. DOI:
 572 10.1016/0006-3207(87)90021-8.

573 Erlinge S. 1969. Food habits of the otter *Lutra lutra* L. and the mink *Mustela vison*
 574 Schreber in a trout water in southern Sweden. *Oikos* **20**: 1–7.

575 Ferreras P, Travaini A, Zapata SC, Delibes M. 2011. Short-term responses of
 576 mammalian carnivores to a sudden collapse of rabbits in Mediterranean Spain. *Basic*
 577 *and Applied Ecology* **12**: 116–124. DOI: 10.1016/j.baae.2011.01.005.

578 Gállego L, Alemany A. 1985. Vertebrados ibéricos, roedores y lagomorfos. Imprés a
 579 Balears: España.

580 Gasith A, Resh VH. 1999. Streams in Mediterranean climate regions: Abiotic influences
 581 and biotic responses to predictable seasonal events. *Annual Review of Ecology and*
 582 *Systematics* **30**: 51–81.

583 Geiger W, Alcorlo P, Baltanás A, Montes C. 2005. Impact of an introduced Crustacean
 584 on the trophic webs of Mediterranean wetlands. *Biological Invasions* **7**: 49–73. DOI:
 585 10.1007/s10530-004-9635-8.

586 Godinho FN, Ferreira MT. 2000. Composition of endemic fish assemblages in relation
 587 to exotic species and river regulation in a temperate stream. *Biological Invasions* **2**:
 588 231–244. DOI: 10.1023/A:1010022123669.

589 Godinho FN, Ferreira MT, Castro MI. 1998. Fish assemblage composition in relation to
 590 environmental gradients in Portuguese reservoirs. *Aquatic Living Resources* **11**: 325–
 591 334. DOI: 10.1016/S0990-7440(98)80003-X.

592 Gowns IO, Gowns JE. 2001. Ecological effects of flow regulation on
 593 macroinvertebrates and periphytic diatom assemblages in the Hawkesbury- Nepean

594 River, Australia. *Regulated Rivers: Research & Management* **17**: 275–293.
 595 DOI: 10.1002/rrr.622.

596 Holling CS. 1959. The functional components of predation as revealed by a study of
 597 small mammal predation of the European pine sawfly. *The Canadian Entomologist*
 598 **91**: 293–320.

599 Klare U, Kamler JF, Macdonald DW. 2011. A comparison and critique of different scat-
 600 analysis methods for determining carnivore diet. *Mammal Review* **41**: 294–312. DOI:
 601 10.1111/j.1365-2907.2011.00183.x.

602 Kruuk H. 2006. *Otters: Ecology, Behaviour and Conservation*. Oxford University
 603 Press: Oxford.

604 Kruuk H, Carss DN. 1996. *Costs and benefits of fishing by a semi-aquatic carnivore*.
 605 The otter *Lutra lutra*. In Aquatic predators and their prey: 10–16. Greenstreet, S. P.
 606 R. & Tasker, M. L. (Eds). Fishing News Books.

607 Lanszki J, Körmendi S, Hancz C, Martin TG. 2002. Examination of some factors
 608 affecting selection of fish prey by otters (*Lutra lutra*) living by eutrophic fish ponds.
 609 *Journal of Zoology* **255**: 97–103. DOI: 10.1017/S0952836901001145.

610 Larned ST, Datry T, Arscott DB, Tockner K. 2010. Emerging concepts in temporary-
 611 river ecology. *Freshwater Biology* **55**: 717–738. DOI: 10.1111/j.1365-
 612 2427.2009.02322.x.

613 Lechowicz MJ. 1982. The sampling characteristics of electivity indices. *Oecologia* **52**:
 614 22–30. DOI: 10.1007/BF00349007.

615 LeHouórou HN. 1990. Global change: vegetation, ecosystems and land use in the
 616 southern Mediterranean basin by the mid twenty-first century. *Israel Journal of*
 617 *Botany* **39**: 481–508. DOI: 10.1080/0021213X.1990.10677170.

- Leunda PM. 2010. Impacts of non-native fishes on Iberian freshwater ichthyofauna: current knowledge and gaps. *Aquatic Invasions* **5**: 239–262. DOI: 10.3391/ai.2010.5.3.03.
- Leunda PM, Elvira B, Ribeiro F, Miranda R, Oscoz J, Alves MJ, Collares-Pereira MJ. 2009. International Standardization of Common Names for Iberian Endemic Freshwater Fishes. *Limnetica* **28**: 189–202.
- López-Nieves P, Hernando JA. 1984. Food habits of the otter in the Central Sierra Morena (Cordoba , Spain). *Acta Theriologica* **29**: 383–401.
- Magalhães MF, Beja PR, Schlosser IJ, Collares-Pereira MJ. 2007. Effects of multi-year droughts on fish assemblages of seasonally drying Mediterranean streams. *Freshwater Biology* **52**: 1494–1510. DOI: 10.1111/j.1365-2427.2007.01781.x.
- Marques C, Rosalino LM, Santos-Reis M. 2007. Otter predation in a trout fish farm of Central–east Portugal: Preference for ‘fast-food’?. *River Research and Applications* **23**: 1147–1153. DOI: 10.1002/rra.1037.
- Mas-Marti E, Garcia-Berthou E, Sabater S, Tomanova S, Munoz I. 2010. Comparing fish assemblages and trophic ecology of permanent and intermittent reaches in a Mediterranean stream. *Hydrobiologia* **657**: 167–180. DOI: 10.1007/s10750-010-0292-x.
- Matthews WJ, Marsh-Matthews E. 2003. Effects of drought on fish across axes of space, time and ecological complexity. *Freshwater Biology* **48**: 1232–1253.
- Miranda R, Escala MC. 2002. Guía de identificación de restos óseos de los Ciprínidos presentes en España. Escamas, opérculos, cleitros y arcos faríngeos. Publicaciones de Biología de la Universidad de Navarra. *Serie Zoológica* **28**: 1–239.

- Miranda R, García-Fresca C, Barrachina P. 2006. Summer prey size selection by European otter *Lutra lutra* in Mediterranean habitats. *Mammalia* **70**: 315–318. DOI: 10.1515/MAMM.2006.059.
- Miranda R, Coop GH, Williams J, Beyer K, Gozlan RE. 2008. Do Eurasian otters *Lutra lutra* (L.) in the Somerset Levels prey preferentially on non-native fish species?. *Fundamental and Applied Limnology* **172**: 339–347. DOI: 10.1127/1863-9135/2008/0172-0339.
- Morales JJ, Lizana M, Acera F. 2004. Ecología trófica de la nutria paleártica *Lutra lutra* en el Río Francia (Cuenca del Tajo, Salamanca). *Galemys* **16** (2): 57–77.
- Nilsson C, Reidy CA, Dynesius M, Revenga C. 2005. Fragmentation and flow regulation of the world's large river systems. *Science* **308**: 405–408. DOI: 10.1126/science.1107887.
- Niquette DJ, D'Abramo LR. 1991. Population dynamics of red swamp crayfish, *Procambarus clarkii* (Girard, 1991) and white river crayfish, *P. acutus acutus* (Girard, 1852), cultured in earthen ponds. *Journal of Shellfish Research* **10**: 179–186.
- Novais A, Sedlmayr A, Moreira-Santos M, Goncalves F, Ribeiro R. 2010. Diet of the otter *Lutra lutra* in an almost pristine Portuguese river: seasonality and analysis of fish prey through scale and vertebrae keys and length relationships. *Mammalia* **74**: 71–81. DOI: 10.1515/MAMM.2010.010.
- Pedroso NM, Santos-Reis M. 2006. Summer diet of Eurasian otters in large dams of south Portugal. *Hystrix* **17**: 117–128. DOI: 10.4404/hystrix-17.2–4369.
- Pedroso NM, Sales-Luis T, Santos-Reis M. 2007. Use of Aguieira Dam by Eurasian otters in Central Portugal. *Folia Zoologica* **56**: 365–377.

664 Peris SJ, Briz FJ, Campos F. 1994. Recent changes in the food of the Grey Heron *Ardea*
665 *cinerea* in central-west Spain. *Ibis* **136**: 488–489. DOI: 10.1111/j.1474-
666 919X.1994.tb01126.x.

667 Prenda J, Granado-Lorencio C. 1992a. Claves de identificación de *Barbus bocagei*,
668 *Chondrostoma polylepis*, *Leuciscus pyrenaicus* y *Ciprinus carpio* mediante algunas
669 de sus estructuras óseas. *Doñana, Acta Vertebrata* **19**: 25–36.

670 Prenda J, Granado-Lorencio C. 1992b. Biometric-analysis of some cyprinid bones of
671 prey fishes to estimate the original lengths and weights. *Folia Zoologica* **41**: 175–
672 185.

673 Prenda J, Freitas D, Santos-Reis M, Collares-Pereira MJ. 1997. Guía para la
674 identificación de restos óseos pertenecientes a algunos peces comunes en las aguas
675 continentales de la Península Ibérica para el estudio de la dieta de depredadores
676 ictiófagos. *Doñana, Acta Vertebrata* **24**: 155–180.

677 Prenda J, Lopez-Nieves P, Bravo R. 2001. Conservation of otter (*Lutra lutra*) in a
678 Mediterranean area: the importance of habitat quality and temporal variation in water
679 availability. *Aquatic Conservation: Marine and Freshwater Ecosystems* **11**: 343–
680 355. DOI: 10.1002/aqc.454.

681 Remonti L, Prigioni C, Balestrieri A, Sgroso S, Priore G. 2010. Eurasian otter (*Lutra*
682 *lutra*) prey selection in response to a variation of fish abundance. *Italian Journal of*
683 *Zoology* **77**: 331–338. DOI: 10.1080/11250000903229809.

684 Román J. 2011. What do otters eat where there is no fish?. *Mammalian Biology* **76**:
685 237–239. DOI: 10.1016/j.mambio.2010.08.005.

686 Ruiz-Olmo J. 1995. Observations on the predation behaviour of the otter *Lutra lutra* in
687 NE Spain. *Acta Theriologica* **40**: 175–180.

- Ruiz-Olmo J, Delibes M. 1998. La nutria (*Lutra lutra* L.) en España ante el horizonte del año 2000. SECEM, Málaga.
- Ruiz-Olmo J, Jimenez J. 2009. Diet diversity and breeding of top predators are determined by habitat stability and structure: a case study with the Eurasian otter (*Lutra lutra* L.). *European Journal of Wildlife Research* **55**: 133–144. DOI: 10.1007/s10344-008-0226-3.
- Ruiz-Olmo J, Jordán G, Gosálbez J. 1989. Alimentación de la nutria (*Lutra lutra* L., 1758) en el nordeste de la Península Ibérica. *Doñana Acta Vertebrata* **16** (2): 227–237.
- Ruiz-Olmo J, Jiménez J, Margalida A. 1998. Capture and consumption of prey of the Otter (*Lutra lutra*) in Mediterranean freshwater habitats of the Iberian Peninsula. *Galemys* **10** (número especial): 209–226.
- Ruiz-Olmo J, López-Martín JM, Palazón S. 2001. The influence of fish abundance on the otter (*Lutra lutra*) populations in Iberian Mediterranean habitats. *Journal of Zoology* **254**: 325–336. DOI: 10.1017/S0952836901000838.
- Ruiz-Olmo J, Olmo-Vidal JM, Manas S, Batet A. 2002. The influence of resource seasonality on the breeding patterns of the Eurasian otter (*Lutra lutra*) in Mediterranean habitats. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **80**: 2178–2189. DOI: 10.1139/z02-186.
- Ruiz-Olmo J, Jiménez J, Chacon W. 2007. The importance of ponds for the otter (*Lutra lutra*) during drought periods in Mediterranean ecosystems: A case study in Bergantes River. *Mammalia* **71**: 16–24. DOI: 10.1515/MAMM.2007.003.
- Sales-Luis T, Pedroso NM, Santos-Reis M. 2007. Prey availability and diet of the Eurasian otter (*Lutra lutra*) on a large reservoir and associated tributaries. *Canadian Journal of Zoology* **85**: 1125–1135. DOI: 10.1139/z07-087.

- Taaström HM, Jacobsen L. 1999. The diet of otters (*Lutra lutra* L.) in Danish freshwater habitats: comparisons of prey fish populations. *Journal of Zoology* **248**: 1–13. DOI: 10.1111/j.1469-7998.1999.tb01017.x.
- Tablado Z, Tella JL, Sanchez-Zapata JA, Hiraldo F. 2010. The Paradox of the Long-Term Positive Effects of a North American Crayfish on a European Community of Predators. *Conservation Biology* **24**: 1230–1238. DOI: 10.1111/j.1523-1739.2010.01483.x.
- Teerink BJ. 1991. *Hair of west-european mammals. Atlas and identification key*. Cambridge University Press: Cambridge.
- Townsend CR, Begon M, Harper JL. 2003. *Essentials of ecology*, 2nd edition. Blackwell Publishing co.: Oxford.
- Vanderploeg HA, Scavia D. 1979. Calculation and use of selectivity coefficients of feeding: zooplankton grazing. *Ecological Modelling* **7**: 135–149.
- Wang L, Infante D, Lyons J, Stewart J, Cooper A. 2011. Effects of dams in river networks on fish assemblages in non-impoundment sections of rivers in Michigan and Wisconsin, USA. *River Research and Applications* **27**: 473–487. DOI: 10.1002/rra.1356.
- Williams DD. 1996. Environmental constraints in temporary fresh waters and their consequences for the insect fauna. *Journal of the North American Benthological Society* **15**: 634–650.
- Zippin C. 1956. An evaluation of the removal method of estimating animal populations. *Biometrics* **2**: 163–189.

Table I. Overall diet composition of the otter in Bullaque River. The total number of individuals (*n*) of each prey item is shown

Prey category	n	FO	RFO	%N	%Biomass
<i>Pseudochondrostoma willkommii</i>	201	1.78	11.36	2.62	0.48
<i>Iberochondrostoma lemmingii</i>	51	18.74	1.09	10.24	0.13
<i>Squalius pyrenaicus</i>	34	3.28	2.01	1.73	0.25
<i>Squalius alburnoides</i>	134	10.4	6.36	6.84	0.58
<i>Luciobarbus spp.</i>	6	0.68	0.84	0.31	0.16
<i>Cyprinus carpio</i>	7	0.96	0.57	0.36	2.69
<u><i>Cyprinidae</i></u>	<u>433</u>	<u>35.84</u>	<u>22.23</u>	<u>22.09</u>	<u>4.29</u>
<i>Lepomis gibbosus</i>	236	22.57	13.61	12.03	4.73
<i>Micropterus salmoides</i>	1	0.14	0.08	0.05	0.21
<u><i>Centrarchidae</i></u>	<u>237</u>	<u>22.71</u>	<u>13.69</u>	<u>12.08</u>	<u>4.94</u>
<i>Gambusia holbrooki</i>	1	0.14	0.08	0.05	0
<i>Esox lucius</i>	10	1.37	0.87	0.51	19.45
<i>Cobitis paludica</i>	46	5.88	3.6	2.34	3.23
<u>Other fishes</u>	<u>57</u>	<u>7.39</u>	<u>4.55</u>	<u>2.9</u>	<u>22.68</u>
FISHES	727	65.94	40.47	37.07	31.91
<i>Arvicola sapidus</i>	4	0.55	0.33	0.2	0.78
<i>Oryctolagus cuniculus</i>	1	0.14	0.08	0.05	0.97
<i>Apodemus sylvaticus</i>	1	0.14	0.08	0.05	0.19
MAMMALS	6	0.83	0.49	0.31	1.94
<i>Alectoris rufa</i>	1	0.14	0.08	0.05	0.97
<i>Gallinula chloropus</i>	5	0.68	0.42	0.25	4.86
Other birds	2	0.27	0.17	0.1	1.94
BIRDS	8	1.09	0.67	0.41	7.77
REPTILIANS (<i>Natrix maura</i>)	4	0.55	0.33	0.2	1.94
AMPHIBIANS (<i>Pelophylax perezi</i>)	29	3.42	2.17	1.48	14.1
<i>Procambarus clarkii</i>	1017	85.36	52.02	51.81	40.68
Insects	169	6.02	3.68	8.61	1.64
Spiders	2	0.27	0.17	0.11	0.24
OTHER INVERTEBRATES	171	6.29	3.85	8.72	1.88
Total of preys	1962				
Number of samples	731				

743 Table II. Seasonal and spatial variations of otter diet (percentage of ingested biomass) in Bullaque River. Significant factors (St, stretch; S,
744 season) for the Factorial ANOVA and subsequent univariate ANOVAs on prey categories are shown. Season: SWi, winter; SSp, spring; SSu,
745 summer; and SAu, autumn. Stretch: StH, High; StM; Medium; and StL; Low. Average ranks of levels with different superscripts are significantly
746 different (Tukey test, $p < 0.05$). Results are means $\pm$ SE in the same order as the average ranks of levels. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Factor	Wilk's λ	d.f.	Prey items	F	d.f	Tukey test	%Biomass
Season (S)	0.02*	48	<i>I. lemmingii</i>	2.87	3	SSp ¹ >SAu ² / SWi ^{1,2} , SSu ^{1,2}	1.61 $\pm$ 0.1 > 0 / 0.24 $\pm$ 0.026, 0.48 $\pm$ 0.04
Stretch (St)	0.012***	50	<i>P. willcommii</i>	9.69***	2	StH ¹ >StM ² , StL ²	2.79 $\pm$ 1.27 > 0.27 $\pm$ 0.16, 0.12 $\pm$ 0.06
			<i>S. alburnoides</i>	5.25*	2	StH ¹ , StM ¹ >StL ²	0.96 $\pm$ 0.27, 1.02 $\pm$ 0.54 > 0.06 $\pm$ 0.03
			<i>L. gibbosus</i>	12.62***	2	StL ¹ >StH ² , StM ²	15.81 $\pm$ 4.2 > 1.65 $\pm$ 0.99, 2.64 $\pm$ 0.85
			Reptilians	5.33**	2	StL ¹ >StH ² , StM ²	3.91 $\pm$ 1.75 > 0, 0
			Amphibians	8.24**	2	StH ¹ >StM ² , StL ²	25.37 $\pm$ 6.4 > 4.38 $\pm$ 4.38, 2.72 $\pm$ 1.84
			<i>P. clarkii</i>	10.10***	2	StH ¹ <StM ² , StL ²	21.65 $\pm$ 5.12 < 66.11 $\pm$ 8.33, 54.88 $\pm$ 7.72
			Insects	8.86***	2	StH ¹ >StM ² , StL ²	10.28 $\pm$ 4.64 > 0, 0

752 Table III: Prey availability in each stretch and season. Results are given in Biomass (kg ha⁻¹).

753

	Winter			Spring			Summer			Autumn		
	High	Medium	Low	High	Medium	Low	High	Medium	Low	High	Medium	Low
<i>Pseudochondrostoma willkommii</i>	0	0.01	0.07	0	0	0	0	0	0	0	0.02	0
<i>Iberochondrostoma lemmingii</i>	0.09	0.41	0.12	0.07	0.2	0.29	0.02	0.01	0.05	0.02	0.02	0
<i>Squalius pyrenaicus</i>	0.13	0.02	3.2	3.7	0.41	0.86	0.86	0.01	0.17	0.52	0	0.02
<i>Squalius alburnoides</i>	1.2	2.98	3.04	2.25	2.14	2.54	1.18	1.99	3.11	3.09	1.15	2.72
<i>Luciobarbus spp.</i>	0.27	0	0.03	1.36	0	0.1	6.27	0	1	0	2.64	0.22
<i>Cyprinus carpio</i>	0	0.76	0.39	0	0	1982.5	0.63	12.39	1.22	0	0	2.09
<i>Ciprinidae</i>	<u>1.69</u>	<u>3.42</u>	<u>6.4</u>	<u>7.38</u>	<u>2.75</u>	<u>3.78</u>	<u>8.32</u>	<u>2.01</u>	<u>4.33</u>	<u>3.63</u>	<u>3.82</u>	<u>2.95</u>
<i>Lepomis gibbosus</i>	0.01	1.27	0.84	0.12	4.67	0	0	0.22	5.83	0	0.95	1.35
<i>Micropterus salmoides</i>	0	0	8.89	0	0	0.45	0	4.75	0.25	0	0	0.17
<i>Centrarchidae</i>	<u>0.01</u>	<u>1.27</u>	<u>9.73</u>	<u>0.12</u>	<u>4.67</u>	<u>0.45</u>	<u>0</u>	<u>4.97</u>	<u>6.08</u>	<u>0</u>	<u>0.95</u>	<u>1.53</u>
<i>Gambusia holbrooki</i>	0	0	0	0.05	0.46	0.2	0.1	0.1	3.96	0	0	0
<i>Cobitis paludica</i>	0.51	0.2	1.07	6.26	1.96	0.21	0.57	0.09	2.73	0.06	0.21	0
<i>Ameiurus melas</i>	0	0	0	0	0	0.14	0.01	0	0	0	0	0.03
<i>Other fishes</i>	<u>0.51</u>	<u>0.2</u>	<u>1.07</u>	<u>6.3</u>	<u>2.42</u>	<u>0.55</u>	<u>0.68</u>	<u>0.19</u>	<u>6.69</u>	<u>0.06</u>	<u>0.21</u>	<u>0.03</u>
FISHES	2.21	4.88	17.19	13.8	9.84	4.78	9	7.16	17.11	3.69	4.98	4.5
<i>Procambarus clarkii</i>	0	0	0	28.32	104.92	70.28	84.33	75.26	235.96	0.5	0.4	0.15

754 Table IV. Seasonal and spatial variations of prey availability (biomass, kg ha⁻¹) in Bullaque River. Significant factors (St, stretch; S, season) for
755 the Factorial ANOVA and subsequent univariate ANOVAs on prey categories are shown. Season: SWi, winter; SSp, spring; SSu, summer; and
756 SAu, autumn. Stretch: StH, High; StM; Medium; and StL; Low. Average ranks of levels with different superscripts are significantly different
757 (Tukey test, $p < 0.05$). Results are means $\pm$ SE in the same order as the average ranks of levels. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Factor	Wilk's λ	d.f.	Prey items	F	d.f.	Tukey test	%Biomass
Season (S)	0.059***	36	<i>G. holbrooki</i>	4.82**	3	SSu ¹ >SWi ² , SAu ² /SSp ^{1,2}	5.56 $\pm$ 0.001 > 0, 0 / 1.34 $\pm$ 0.001
			<i>I. lemmingii</i>	8.07***	3	SWi ¹ >SSp ² , SSu ² , SAu ²	1.22 $\pm$ 0.003 > 0.82 $\pm$ 0.001, 0.29 $\pm$ 0.001, 0.17 $\pm$ 0.001
			<i>S. alburnoides</i>	6.23**	3	SAu ¹ >SSp ^{2,3} , SSu ³ /SWi ^{1,2}	6.69 $\pm$ 0.011 > 6.2 $\pm$ 0.006, 3.39 $\pm$ 0.002 / 5.72 $\pm$ 0.008
			<i>P. clarkii</i>	31.68***	3	SSp ¹ , SSu ¹ >SWi ² , SAu ²	186.55 $\pm$ 0.008, 497.61 $\pm$ 0.01 > 0, 3.51 $\pm$ 0.005
Stretch (St)	0.23**	24	<i>C. paludica</i>	5.36**	2	StH ¹ >StM ² , StL ²	8.12 $\pm$ 0.0071 > 3.86 $\pm$ 0.0014, 5.12 $\pm$ 0.007

764 Table V. Seasonal and spatial variation in otter electivity index (ϵ_i) in Bullaque River. Significant factors (St, stretch; S, season) and interactions
765 for the Factorial ANOVA and subsequent univariate ANOVAs on prey items are shown. Season: SWi, winter; SSp, spring; SSu, summer; and
766 SAu, autumn. Stretch: StH, High; StM; Medium; and StL; Low. Average ranks of levels with different superscripts are significantly different
767 (Tukey test, $p < 0.05$). Results are means $\pm$ SE in the same order as the average ranks of levels. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Factor	Wilk's λ	d.f.	Prey items	F	d.f	Tukey test	Mean values $\pm$ SE
Season (S)	0.023***	27	<i>L. gibbosus</i>	4.41*	3	SWi ¹ <SSu ² , SAu ² /SSp ^{1,2}	-1 $\pm$ 0 < 0 $\pm$ 0.22, -0.13 $\pm$ 0.08 / -0.34 $\pm$ 0.18
			<i>P. clarkii</i>	8.26***	3	SWi ¹ >SSp ² , SSu ² /SAu ^{1,2}	0.79 $\pm$ 0.01 > -0.61 $\pm$ 0.1, -0.53 $\pm$ 0.09 / 0.03 $\pm$ 0.1
Stretch (St)	0.11***	18	<i>P. willcommii</i>	7.18**	2	StH ¹ >StM ² , StL ²	0.24 $\pm$ 0.12 > -0.54 $\pm$ 0.17, -0.64 $\pm$ 0.16
			<i>S. alburnoides</i>	3.84*	2	StM ¹ >StH ² , StL ²	-0.88 $\pm$ 0.05 > -1 $\pm$ 0, -1 $\pm$ 0
			<i>P. clarkii</i>	4.5*	2	StH ¹ <StM ² /StL ^{1,2}	-0.56 $\pm$ 0.01 < 0.3 $\pm$ 0.13 / 0.02 $\pm$ 0.09
S $\times$ St	0.013***	54	<i>L. gibbosus</i>	3.42*	6		
			<i>P. clarkii</i>	4.96**	6		

768

769

Figure 1. Map of the study area with sampling sites represented as follows: H, High stretch; M, Medium stretch and L, Low stretch

Figure 1. Cumulative frequency of resource items against increase in sample size of otter spraints in the three stretches considered and throughout the year.

Figure 2. Spatial and seasonal variation of otter trophic diversity (Shannon index, H') in Bullaque River. Vertical bars indicate SE. Means marked with different letters are significantly different from one to another (Tukey test, $p > 0.05$)