Broad wintering range and intercontinental migratory divide within a core population of the near threatened pallid harrier.

J. Terraube^{1,2*}, F. Mougeot³, T. Cornulier⁴, A. Verma⁵, A. Gavrilov⁶, B. Arroyo².

1-Natural Research Ltd, Banchory Business Centre, Burn O'Bennie Road, Banchory, AB31 5ZU, UK.

2-Instituto de Investigación en Recursos Cinegéticos, CSIC-UCLM-JCCM, Ronda de Toledo s/n, 13005 Ciudad Real, Spain.

3-Estación Experimental de Zonas Áridas (EEZA-CSIC), Ctra. de Sacramento s/n La Cañada de San Urbano, 04120 Almería, Spain.

4-Institute for Biological and Environmental Sciences, University of Aberdeen, Tillydrone Avenue, Aberdeen AB24 2TZ, UK.

5-Guru Ghasidas University, Department of Forestry, Wildlife and Environmental Sciences, Bilaspur, Chhatis Garh, 495 009 India.

6-Institute of Zoology, Akademgorodok, 480060 Almaty, Kazakhstan.

* Corresponding author: Julien.Terraube@uclm.es.

Running head: Migratory connectivity in central asian pallid harriers.

ABSTRACT

Aim To identify the migration routes and wintering grounds of the core populations of the near threatened pallid harrier *Circus macrourus* and highlight conservation needs associated with these phases of the annual cycle.

Location Breeding area: north-central Kazakhstan; Wintering areas: Sahel belt (Burkina Faso to Ethiopia) and north-west India.

Methods We used ring recovery data from Kazakhstan and satellite tracking data from 2007-2008 on 6 adults breeding in north-central Kazakhstan to determine migration routes and locate wintering areas. In addition, one first-year male was tagged in winter 2007-2008 in India.

Results Data evidenced an inter-continental migratory divide within the core pallid harrier population, with birds wintering in either Africa or India. The six individuals tagged in north-central Kazakhstan followed a similar route (west of the Caspian Sea and Middle East) towards east Africa, before spreading along the Sahel belt to winter either in Sudan, Ethiopia, Niger or Burkina Faso. Spring migration followed a shorter, more direct route, with marked inter-individual variation. The bird tagged in India spent the summer in central Kazakhstan. Half of the signal losses (either because of failure or bird mortality) occurred on the wintering areas and during migration.

Main conclusions Our study shows that birds from one breeding area may winter over a strikingly broad range within and across continents. The intercontinental migratory divide of pallid harriers suggests the coexistence of distinct migratory strategies within

the core breeding population, a characteristic most likely shared by a number of threatened species in central Asia. Conservation strategies for species like the pallid harrier therefore require considering very large spatial scales with possibly area-specific conservation issues. We highlight urgent research priorities in order to effectively inform the conservation of these species.

Keywords: Conservation, satellite telemetry, migration strategy, migratory divide, wintering areas, Kazakhstan, *Circus macrourus*.

INTRODUCTION

The study of migration is crucial for fully understanding the population ecology and conservation needs of migrant species (Webster & Marra, 2005). Bird populations of long distance migrants breeding in Eurasia and wintering in sub-Saharan Africa have experienced stronger population declines than short-distance migrants or resident species (Sanderson et al., 2006). Suggested causes include a combination of harsher climatic conditions and agricultural intensification in the wintering areas, leading to widespread land degradation and increased mortality during long migrations (e.g. Newton, 2008). Additionally, breeding success may be influenced both by the quality of the breeding habitat and by the carry-over influence of winter habitat quality on body condition and arrival dates (Norris et al., 2003; Gunnarsson et al., 2005). As a consequence, the effective conservation of migratory populations depends on the understanding of the factors that influence an individual's fitness through the entire annual cycle. Satellite tracking is a powerful tool for investigating large scale animal movements and migratory pathways (Meyburg et al., 1998; Kjellén et al., 2001; Robinson et al., 2010; see Guan & Higuélin, 2000 for a review), and increasingly available for smaller species (Strandberg et al., 2009).

Many Eurasian species have their wintering quarters split between India and Africa (del Hoyo et al., 1994), where they may experience different human and environmental pressures (Naidoo & Iwamura, 2007). However, the migratory pathways and connectivity between breeding and wintering areas is largely unknown for most populations. This is the case of the pallid harrier (*Circus macrourus*), a long-distance migrant whose core breeding populations (where the species is most abundant) are

located in north central Kazakhstan and south-western Siberia (del Hoyo et al., 1994). Substantial population declines of this poorly known species have been reported in many breeding areas, particularly in peripheral populations west of the Ural River (Davygora & Belik, 1994; Kostin, 1983; Nankinov et al., 1991; Birdlife International, 2003). As a result, this species is now classified as ‘Near Threatened’ (Birdlife International, 2000). The main limiting factors include breeding habitat degradation and reduced prey availability through land-use changes, as well as a decrease in the availability and quality of suitable wintering areas and illegal hunting on migration routes and bottlenecks (van Maanen et al., 2001; Birdlife International, 2003). However, most information available to date essentially relates to breeding ecology (Bragin, 2003; Terraube et al., 2009). Pallid harriers are highly dependent on small rodents during the breeding period and core breeding populations may be vulnerable to recent land-use changes in central Asia, causing the loss of high-quality vole-rich habitats such as the natural steppe and set-asides (Terraube et al., 2011). However, part of the negative trend observed in pallid harrier’s populations during the last decades may be linked to ‘wintering area effects’. It is therefore necessary to better understand pallid harrier ecology outside breeding areas, starting with the identification of migration routes and wintering areas.

To date, information on pallid harrier migration is mostly anecdotal (observations at bottleneck migration points; Corso & Cardelli, 2004; Panuccio & Agostini, 2006). Its main wintering grounds are open grasslands and agricultural areas throughout the Indian subcontinent, the savannah belt in Africa, south of the Sahara through to the eastern African savannahs. The migration routes and wintering areas of the core breeding

populations remain enigmatic and of particular interest, with any or both of the African and Indian continents being potential wintering areas.

In this paper, we first review Kazakh ringing recovery data to study pallid harrier migration patterns. Second, we report on satellite telemetry data and document for the first time the migration routes and wintering areas of pallid harriers from a known origin (north-central Kazakhstan, within the core breeding area). Finally, we discuss the implications of our findings for the conservation of this and other migratory species of conservation concern in central Asia.

METHODS

Ringing and recovery data of pallid harriers in and from Kazakhstan (hereafter KZ) were obtained from the Institute of Zoology, Almaty (KZ). For the satellite tracking study, we trapped breeding pallid harriers in the Naurzum and Turgai regions (51°N, 64°E), located in north-central KZ (Terraube et al., 2009). One adult pallid harrier female was captured in July 2007 in the Turgai region and eight adult pallid harriers (four males and four females) were captured in June and July 2008 in Naurzum (c.100 km north of Turgai). All birds were trapped using a mist net and a stuffed corsac fox (*Vulpes corsac*) as a decoy, placed at the proximity of active nests, or with a nest-trap placed above the nestlings when older than 10 days. In addition to the previous birds, one first-year male was captured in the Velavadar National Park near Bhavnagar, Gujarat State, north India, in March 2008 (Verma, in press), as part of a research project set up in India, in order to study the winter ecology of this species.

Birds were fitted with a 11.5 g solar Northstar PTT transmitter attached using a Teflon harness. The transmitter and harness weighed approximately 4% and 3% of the male and female weights. This extra-weight could negatively impact on survival, but the same tags fitted onto the closely related and smaller Montagu's harrier (23% lighter than pallid harrier) did not have any noticeable effects on survival (Trierweiler et al., 2007; Treirweiler & Koks, 2009; Limiñana et al., 2007). After release at the capture site, birds quickly resumed normal activities (foraging, provisioning of young, etc...).

Harrier movements were tracked using the Argos satellite system which provides location estimates coupled with nominal location accuracy, the location class ('LC') (Meyburg & Fuller, 2007). Location classes 0,1, 2 and 3 indicate that the reported location lies (with 68.23% confidence) within $>1\text{km}$, $\leq 1\text{km}$, $\leq 350\text{ m}$ and $\leq 150\text{ m}$ of the true location, respectively (Hays et al., 2001). Location classes A, B, C and Z are not associated to reported accuracy and may thus be less reliable, but may be also useful for long-distance tracks (Trierweiler, 2010). We used one location per day, selecting the highest accuracy class available (including lower class locations if they were the only available) after having excluded aberrant data via filtering (following Trierweiler & Klaassen, pages-112-123 in Trierweiler, 2010). When more than one location of similar accuracy was available for a given duty cycle, we selected the first one.

Out of the nine birds tagged in north-central Kazakhstan, only six provided useful data to study migration (three stopped transmitting shortly after PTT setup, possibly because of loss or transmitter failure). For all individuals, we reported here the first year of tracking data.

Birds were considered to initiate migration when leaving their breeding areas or wintering areas. The onset of migration trips was very obvious since, in all cases they were initiated with movements of $>100 \text{ km.day}^{-1}$. Similarly, the end of migration trip (and change to wintering behaviour) was very obvious as daily movements changed to $<20 \text{ km.day}^{-1}$. All results are presented as means $\pm$ SD.

Habitats (natural vs agricultural or anthropised) in wintering areas were characterized using Google Earth (<http://earth.google.com>).

RESULTS

Ringling data

Analysis of the ringing recoveries from Kazakh birds suggested that central Asian pallid harriers used two alternative migration routes (Fig. 1). A ‘western’ route to the African wintering grounds was evidenced by three ring recoveries in central Kazakhstan from Tunisia and France (Fig. 1). An ‘eastern’ migratory route linking central Kazakhstan to the Indian wintering grounds could be identified by two ring recoveries in south-eastern KZ, and one in northern India (Fig. 1).

Satellite tracking data: migration routes and wintering areas

All of the six migrating birds from north KZ migrated west of the Caspian Sea, through Russia, Georgia, Armenia, then crossed countries of the Middle East (Turkey, Syria and Iraq; Fig. 2a). Two to three of these birds flew a route near the Batumi region, along the

Georgian (east) coast of the Black Sea when crossing the Caucasus area (Fig. 2a). Subsequently, the routes used slightly differed, as some birds used a more “terrestrial” route crossing Israel, Jordan and Egypt, whereas others crossed Saudi Arabia, then the Red sea, Egypt and Eritrea. The wintering areas of these birds were mainly located in sub-Saharan countries from western-central to eastern Africa (Table 1; Fig. 2a), with a large longitudinal spread of c. 4500 km. Two pallid harriers overwintered in Ethiopia, two others in Sudan, and two other birds arrived to Sudan and then shifted to central-western Africa, one settling at the Burkina Faso - Niger border and the other one in Niger (wintering in open, dry savannah habitats). In Sudan, pallid harriers wintered in open and dry grasslands with higher heterogeneity in vegetal cover. Birds wintering in Ethiopia spent the winter in the most anthropized habitats, a mix of pastures and agricultural areas at the vicinity of several villages.

Of the six tagged birds that migrated to Africa, only three transmitted until the following breeding season (two stopped transmitting in Sudan and Ethiopia during winter, and one in Syria during spring migration; Table 1).

Spring migration routes used by pallid harriers differed from the autumn ones by being more direct (Fig. 2b). The two birds wintering in western-central Africa came back via Egypt after a northward trip through the Sahara Desert, thereafter using a similar route to those taken during the autumn. The other two birds that wintered in eastern Africa (Sudan and Ethiopia) crossed the Red Sea, Saudi Arabia, one crossing the Caspian Sea and the other passing east of it (and not west, as did all the birds during the autumn migration).

The bird tagged in winter in India migrated through Pakistan, Afghanistan, Turkmenistan and Uzbekistan, to spend summer in central Kazakh breeding grounds, about 300 km away from the area where the other tagged birds bred in 2007-2008 (Fig. 2b).

Timing and duration of migration and wintering phases

The mean departure date from the breeding grounds was 28 July $\pm$ 6 days and the mean arrival to the wintering areas was 29 October $\pm$ 11 days. Therefore, the mean duration of the autumn migration, was 83 $\pm$ 31 days. Autumn migration was long because most birds (5 of 6) performed one long stop-over (54 $\pm$ 10 days) in an area distant 1695 $\pm$ 610 km from the breeding area (all in Kazakhstan or Russia). During autumn migration, birds travelled a total of 7340 $\pm$ 1378 km. The average migration speed was thus 108 $\pm$ 67 km.day⁻¹ (or 164 $\pm$ 45 km.day⁻¹ when excluding the time spent in the stop-over).

Birds stayed on the wintering grounds an average of 148 $\pm$ 15 days (n = 4). The mean departure date from the African wintering areas was 23 March $\pm$ 6 days (n = 4), and the mean duration of the northward migration was 28 $\pm$ 5 days (n = 3). The mean distance travelled was 6888 $\pm$ 1878 km and average speed during spring migration was thus 258 $\pm$ 105 km.day⁻¹ (n = 3). Overall, the spring migration from Africa was shorter, faster and more direct than the autumn one for all three monitored birds. The first-year male captured in its wintering quarters in India travelled a distance of 3322 km until settlement in the breeding grounds (Table 1, see Verma, in press. for more details).

DISCUSSION

This study documents for the first time the migration timing, routes and wintering areas of pallid harriers of known origin (north-central KZ). A main result was the existence of an intercontinental migratory divide for core breeding populations from central Kazakhstan (tagged birds wintered in sub-Saharan Africa and in north-west India using much differentiated paths). The simultaneous monitoring of several individuals originating from the same breeding area revealed a large longitudinal spread of the African wintering areas, an overall similarity in the initial autumn migration routes of birds going to Africa, but greater inter-individual heterogeneity in spring migration routes, which are shorter (more direct) than the autumn ones.

Migration routes and wintering areas

Analyses of ringing data showed that pallid harriers from Kazakhstan winter either in Africa or in the Indian subcontinent and could therefore use two different migration routes and wintering regions. In addition, the male pallid harrier tagged in India in winter migrated to central Kazakhstan, close to a breeding area where birds wintering in Africa were tagged. Together, these data confirm the existence of an inter-continental migratory divide for pallid harriers breeding in Kazakhstan and suggest that birds breeding in western and north-central KZ would migrate preferentially to east Africa and the Sahelian Belt, whereas those breeding in more eastern locations would have their wintering grounds in the Indian subcontinent. There is no obvious geographical barrier (mountain, desert, sea...) separating the breeding pallid harrier populations wintering in India from those wintering in Africa. Therefore the origins of the evolution of divergent migration routes in this species remain to be investigated, for instance

using molecular approaches (Pérez-Tris et al., 2004). Pallid harriers seem to be more widely distributed in Africa than India (Birdlife International, 2003), but in the absence of reliable wintering population abundance estimates in Africa and India, the relative importance of each wintering area cannot be yet assessed. Further research would be however critical to ascertain this.

Our data indicate at least 4500 km of longitudinal extent in African wintering areas (from Ethiopia to Burkina Faso; Fig. 2) for birds from the same breeding origin (north-central Kazakhstan). This longitudinal variation is much greater than that reported in other raptor species tracked by satellite telemetry (e.g. Limiñana et al., 2007; Gschwend et al., 2009; Strandberg et al., 2009).

Autumn migration routes towards Africa were consistent among individuals, at least during the first part of the migration (Fig. 1), whereas spring migration routes differed substantially from autumn ones, and showed more inter-individual variation, which was likely a consequence of the broad longitudinal range occupied by this species during winter, together with more direct routes used for the return migration. Additionally, climatic conditions (and particularly wind drift) could influence migration patterns (Klaassen et al., 2010; 2011). Migrating raptors are subjected to easterly winds in Africa during both autumn and spring migration, with stronger winds in spring (Meyburg et al., 2003), and there might be individual variation in responses to wind drift (Klaassen et al., 2010; 2011). Finally, spring migration was overall faster than the autumn one. This was a combination of higher flight speed (with an additional 59 km travelled per day even when considering only moving days) and the lack of long stop overs. A faster migration in spring than in autumn has been reported in other raptors (e.g. McGrady et

al., 2002; Alerstam et al., 2006) and could be related to a selection for earlier arrival, with fitness advantages for early breeders (Kokko, 1999).

Risks for migrating and wintering pallid harriers

From the 6 birds that started autumn migration, only 3 apparently survived to the following breeding season. The location of signal losses could indicate where mortality occurred.

Signal from one bird was lost during migration in Syria. Additionally, one of the birds that survived the first year was also lost during migration in Iraq the following year. The convergence of the routes used by tagged birds migrating from central Asia to Africa means a potential vulnerability of birds wintering in Africa to threats like illegal hunting on the migratory pathways or habitat degradation in stopover areas. Widespread capture and shooting of raptors, commonly including harriers, have been reported on migration corridors in Georgia, along the Black sea coast and in other Middle East countries (van Maanen et al., 2001). Our data show that a large proportion of the tagged birds from central Kazakhstan transit through this region, while estimates based on local counts suggest that more than 1500 pallid harriers can be recorded every autumn in the Batumi area (<http://www.batumiraptorcount.org>). These estimates point out the critical importance of this watchsite for monitoring Central Asian pallid harrier populations wintering in Africa. Although comprehensive information is lacking on current shooting and trapping levels on migration routes, a recent report indicates that migratory birds are facing considerable hunting and trapping pressure in Syria (Attar, 2005). Priority should be given to assessing the sensitivity of pallid harrier and other raptors' populations to

illegal hunting along the migration routes, and mitigation effort should be maintained and reinforced meanwhile.

Additionally, signals for two birds were lost in winter, highlighting the potential importance of this stage of the annual cycle for populations. In particular, birds were lost in Sudan and Ethiopia in 2008-09, when a severe drought occurred (NASA: <http://earthobservatory.nasa.gov/NaturalHazards>): low rainfall levels in the Sahel have been associated with reduced survival rates in other migratory raptors and birds (Schaub et al., 2005; Grande et al., 2009; Mihoub et al., 2010).

Overall, different threats may be associated with each of the wintering areas, both intra- and inter-continent. The whole Sahelian and Sudanian biogeographical zones of the pallid harrier wintering range have suffered much habitat degradation (Thiollay, 2006a). Ethiopian grasslands and savannahs have been recently described as threatened by changes in the patterns of extensive seasonal grazing (involving transhumance, use of wet-and dry-season pasturing and the control of encroaching scrub by fire), as sedentarization is associated with a clear pattern of biodiversity loss (Western et al., 2009; Donald et al., 2010). Land degradation seems to be widespread in Ethiopia as a result of rapid population increase, erosion, deforestation and reduced vegetation cover (Tadesse, 2001). Nevertheless, the impact these changes could have on wintering pallid harriers remains largely unknown.

Western Africa has been reported to suffer a huge loss of biodiversity outside the protected areas in recent decades, associated with the dramatic increase in human pressure: deforestation, overgrazing and erosion, increase in pesticide use and

overhunting (Sanderson et al., 2006). These changes were associated with large decline in raptor abundance outside protected areas in west Africa between the 70's and the 2000's and a significant decline in wintering pallid harrier numbers during this period (Thiollay, 2006a, 2006b, 2007).

Finally, results of an ongoing research project in India on the winter ecology indicate that pallid harriers depend on natural grasslands for both roosting and foraging (T. Cornulier et al. unpubl. data; del Hoyo, 1994). Natural grasslands in India have suffered considerable reduction and fragmentation and remnant grasslands are now essentially restricted to protected areas and governmental reclaimed land, with ongoing degradation due to overgrazing and encroachment by invasive trees *Prosopis juliflora* (Joshi et al., 2009).

Further research is needed to better understand winter habitat use and foraging ecology of pallid harriers, as information on the diet in the different wintering areas remains scant. Contrary to the closely related Montagu's harrier, trophic relationships between pallid harriers, grasshoppers or other prey in Africa are still poorly understood. This could have crucial conservation implications as many western European migratory raptors, mainly locust and grasshopper consumers, have suffered severe declines in recent decades, possibly related to climate change and pesticide use to control insects (Newton, 2004; Sánchez-Zapata et al., 2007).

Intercontinental migratory divide and conservation implications

We identified two main migration routes and a wide variety of wintering areas over two continents for the core populations of the pallid harrier breeding in central and north Kazakhstan. Such an intercontinental migratory divide appears to exist in other threatened species from central Asia (including sociable lapwing *Vanellus gregarius*, demoiselle crane *Anthropoides virgo*, steppe eagle *Aquila nipalensis*; del Hoyo et al., 1994; Birdlife International, 2006; Meyburg et al., 2003), although conservation implications are still little understood. Populations with individuals using different migratory pathways and scattered wintering areas are likely vulnerable to contrasted threats. Different wintering areas may be characterised by different levels of environmental and anthropic pressures, such as droughts, habitat destruction or illegal hunting. As a result, the existence of different wintering areas for birds using the same breeding area will impact heterogeneously on the demography of that population, which further complicates the design of effective conservation strategies. While wide wintering ranges have the potential to buffer populations against temporary adverse conditions in one particular wintering site, permanent differences in the quality of migration routes or wintering areas have the potential to constitute source or sink sub-populations. In order to inform global conservation strategies for species like the pallid harrier, it is important for future research to 1) determine the proportion of the global population wintering in India vs. Africa and among the different areas within Africa; 2) identify the specific threats associated with different migratory pathways and wintering areas using a combination of satellite data and detailed field studies, in order to develop appropriate targeted conservation programmes for the species in its wintering areas; 3) estimate the spatial, genetic structuring and exchanges between sub-populations exhibiting different migration and wintering strategies.

ACKNOWLEDGMENTS

We thank Jesús T. García, Bob Stakim, Evgeny and Alexander Bragin, Todd Katzner, Alexei Timoshenko for help during the fieldwork and trapping attempts in Kazakhstan, Mike McGrady and Ewan Weston for technical support, the late Mike Madders for encouragement and inspiration throughout the project, and Natural Research Ltd for funding this research project. We also thank Roine Strandberg, Keith Bildstein and an anonymous referee for constructive comments on the manuscript.

REFERENCES

- Alerstam, T., Hake, M. & Kjellén, N. (2006) Temporal and spatial patterns of repeated migratory journeys by ospreys. *Animal Behaviour*, **71**, 555-566.
- Attar, W. (2005) *National Report on Hunting*. Syrian Arab Republic. Building capacity for sustainable hunting of migratory birds in Mediterranean third countries, Life project.
- BirdLife International. (2000) *Threatened birds of the world*. Lynx Editions. Barcelona and Cambridge.
- Birdlife International. (2003) *International Action plan for the Pallid harrier (Circus macrourus)*. Convention on the Conservation of European Wildlife and Natural habitats. Standing committee. 23rd meeting. Strasbourg, 1-4 December 2003.
- BirdLife International. (2006) *Vanellus gregarius*. 2006. IUCN Red List of Threatened Species. IUCN 2006. www.iucnredlist.org.

- 418 Bragin, EA. (2003) Distribution, number and some features of ecology of the pallid
419 harrier (*Circus macrourus*) in Kustanai region. *Selevinia*, 2003, 145–149. [in
420 Russian].
- 421 Corso, A. & Cardelli, C. (2004) The migration of Pallid Harriers accross the central
422 Mediterranean. *British Birds*, **97**, 238-246.
- 423 Davygora, A.V. & Belik, VP. (1994) The pallid harrier *Circus macrourus* as an
424 endangered species in the Palearctic. *Raptor Conservation Today: Proceedings of*
425 *the IV World Conference on Birds of Prey and Owls* (ed. by B.-U. Meyburg &
426 R.D. Chancellor), pp. 93–96. WWGBP/The Pica Press, Bodmin, Cornwall.
- 427 del Hoyo, J., Elliott, A & Sargatal, J. (1994) *Handbook of the Birds of the World: vol. 2.*
428 *New World Vultures to Guineafowl*. Lynx Editions. Barcelona and Cambridge.
- 429 Donald, P.F., Buchanan, GM., Collar, NJ., Abebe, YD., Gabremichael, MN., Mwangi,
430 MAK., Ndag'ang'a, PK., Spottiswoode, CN. & Wondafrash, M. (2010) Rapid
431 declines in habitat quality and population size of the Liben (Sidamo) Lark
432 *Heteromiraфра sidamoensis* necessitate immediate conservation action. *Bird*
433 *Conservation International*, **20**, 1-12.
- 434 Grande, JM., Serrano, D., Tavecchia, G., Carrete, M., Ceballos, O., Diaz-Delgado, R.,
435 Tella, JL. & Doñazar, JA. (2009) Survival in a long-lived territorial migrant:
436 effects of life-history traits and ecological conditions in wintering and breeding
437 areas. *Oikos*, **118**, 580-590.
- 438 Gschweng, M., Kalko, EKV., Querner, U., Fiedler, W. & Berthold, P. (2009) All across
439 Africa: highly individual migration routes of Eleonora's falcon. *Proceedings of the*
440 *Royal Society-series B, Biological Sciences*, **275**, 2887-2896.
- 441 Guan, HL. & Higuélin, H. (2000) Review of satellite tracking of migratory birds and its
442 prospects. *Zoological Research*, **21**, 412-415.

443 Gunnarsson, TG., Gill, J.A., Newton, J., Potts, PM. & Sutherland, WJ. (2005) Seasonal
444 matching of habitat quality and fitness in a migratory bird. *Proceedings of the*
445 *Royal Society B-Biological Sciences*, **272**, 2319-2323.

446 Hays, GC., Arkesson, S., Godley, B.J., Luschi, P. & Santidrian, P. (2001) The
447 implication of location accuracy for the interpretation of satellite-tracking data.
448 *Animal Behavior*, **61**, 1035-1040.

449 Joshi, P., Kumar, V., Koladiya, M., Patel, Y. & Karthik, T. (2009) Local perceptions of
450 grassland change and priorities for conservation of natural resources of Banni,
451 Gujarat, India. *Frontiers of Biology in China*, **4**, 549-556.

452 Kjellén, N., Hake, M. & Alerstam, T. (2001) Timing and speed of migration in male,
453 female and juvenile Ospreys *Pandion haliaetus* between Sweden and Africa
454 revealed by field observations, radar and satellite tracking. *Journal of Avian*
455 *Biology*, **32**, 57-67.

456 Klaassen, RHG., Strandberg, R., Hake, M., Olofsson, P., Tøttrup, AP. & Alerstam, T.
457 (2010) Loop migration in adult marsh harriers *Circus aeruginosus*, as revealed by
458 satellite telemetry. *Journal of Avian Biology*, **41**, 200-207.

459 Klaassen, RHG., Hake, M., Strandberg, R. & Alerstam, T. (2011) Geographical and
460 temporal flexibility in the response to crosswinds by migrating raptors.
461 *Proceedings of the Royal Society B-Biological Sciences*, **278**, 1339-1346.

462 Kokko, H. (1999) Competition for early arrival in migratory birds. *Journal of Animal*
463 *Ecology*, **68**, 940-950.

464 Kostin, YV. (1983) Ptitsy Kryma [Birds of the Crimea]. M., Nauka. (In Russian).

465 Limiñana, R., Soutullo, A. & Urios, V. (2007) Autumn migration of Montagu's harriers
466 (*Circus pygargus*) tracked by satellite telemetry. *Journal of Ornithology*, **148**,
467 517-523.

468 McGrady, MJ., Maechtle, TL., Vargas, JJ., Seegar, WS. & Pena, PP. (2002) Migration
 469 and ranging of Peregrine falcons wintering on the Gulf of Mexico coast,
 470 Tamaulipas, Mexico. *The Condor*, **104**, 39-48.

471 Meyburg, B-U., Meyburg, C. & Barbraud, JC. (1998) Migration strategies of an adult
 472 short-toed eagle *Circaetus gallicus* tracked by satellite. *Alauda*, **66**, 39-48.

473 Meyburg, B.U, Paillat, P, Meyburg, C. (2003) Migration routes of Steppe Eagles
 474 between Asia and Africa: A study by means of satellite telemetry. *The Condor*,
 475 **105**, 219-227.

476 Meyburg B.-U. & Fuller, MR. (2007) Satellite tracking. Pp. 242-248 in: Bird, D. M. &
 477 K. L. Bildstein (eds.): *Raptor Research and Management Techniques*. Hancock
 478 House Publishers, Surrey, Canada.

479 Mihoub, JB., Gimenez, O., Pilard, P. & Sarrazin, F. (2010) Challenging conservation of
 480 migratory species: sahelian rainfalls drive first-year survival of the vulnerable
 481 Lesser kestrel *Falco naumanni*. *Biological Conservation*, **143**, 839-847.

482 Naidoo, R. & Iwamura, T. (2007) Global-scale mapping of economic benefits from
 483 agricultural lands: Implications for conservation priorities. *Biological*
 484 *Conservation*, **140**, 40-49.

485 Nankinov, D., G. Stoyanov, G. Kouzmanov. & Todorov, R. (1991) The situation of
 486 diurnal birds of prey in Bulgaria. *Birds of prey bulletin n°. 4*. (ed. by R.D.
 487 Chancellor and B.-U. Meyburg). pp. 293-302. World Working Group on Birds of
 488 Prey and Owls, Berlin, Germany.

489 Newton, I. (2004) Population limitation in migrants. *Ibis*, **146**, 197-226.

490 Newton, I. (2008) *Migration ecology of birds*, Academic Press, UK.

491 Norris, DR., Marra, PP., Kyser, TK. & Ratcliffe, KM. (2003) Tropical winter habitat
 492 selection limits reproductive success on the temperate breeding grounds in a

493 migratory bird. *Proceedings of the Royal Society-series B, Biological Sciences*,
494 **271**, 59-64.

495 Panuccio, M. & Agostini, N. (2006) Spring migration of Pallid (*C. macrourus*) and
496 Montagu's harriers (*C. pygargus*) in relation to sex and age classes at two
497 watchsites of the central Mediterranean. *Buteo*, **15**, 3-10.

498 Pérez-Tris, J., Bensch, S., Carbonell, R., Helbig, A. J. & Tellería, J.L. (2004) Historical
499 diversification of migration patterns in a passerine bird. *Evolution*, **58**, 1819–1832.

500 Robinson, W.D., Bowlin, M.S., Bisson, I., Shannon-Barnes, S., Thorup, K., Diehl, R.H.,
501 Keunz, T.H., Mabey, S. & Winkler, D.W. (2010) Integrating concepts and
502 technologies to advance the study of bird migration. *Frontiers in Ecology and*
503 *Environment*, **8**, 354-361.

504 Sánchez-Zapata, J.A., Doñázar, J.A., Delgado, A., Forrero, M.G., Ceballos, O., Hiraldo,
505 F. (2007) Desert locust outbreaks in the Sahel: resource competition, predation
506 and ecological effects of pest control. *Journal of Applied Ecology*, **44**, 323-329.

507 Sanderson, F.J., Donald, P.F., Pain, D.J., Burfield, I.J. & Van Bommel, F.P.J. (2006) Long-
508 term population declines in Afro-Palearctic migrant birds. *Biological Conservation*,
509 **131**, 93-105.

510 Schaub, M., Kania, W. & Köppen, U. (2005) Variation of primary production during
511 winter induces synchrony in survival rates in migratory white storks *Ciconia*
512 *ciconia*. *Journal of Animal Ecology*, **74**, 656-666.

513 Strandberg, R., Klaassen, R.H.G., Hake, M., Olofsson, P. & Alerstam, T. (2009)
514 Converging migration routes of Eurasian hobbies *Falco subbuteo* crossing the
515 African equatorial rain forest. *Proceedings of the Royal Society of London*,
516 *Biological Sciences*, **276**, 727-733.

517 Tadesse, G. (2001) Land degradation: a challenge to Ethiopia. *Journal of Environment*
518 *Management*, **27**, 815-824.

519 Terraube, J., Arroyo, BE., Mougeot, F., Madders, M., Watson, J. & Bragin, EA. (2009)
520 Breeding biology of the pallid harrier *Circus macrourus* in north-central
521 Kazakhstan: implications for the conservation of a Near-Threatened species. *Oryx*,
522 **43**, 104-112.

523 Terraube, J., Arroyo, B., Madders, M. & Mougeot, F (2011) Diet specialisation and
524 foraging efficiency under fluctuating vole abundance: a comparison between
525 generalist and specialist avian predators. *Oikos*, **120**, 234-244.

526 Thiollay, JM. (2006a) The decline of raptors in West Africa: long-term assessment and
527 the role of protected areas. *Ibis*, **148**, 240-254.

528 Thiollay, JM. (2006b) Large bird declines with increasing human pressure in savanna
529 woodlands (Burkina Faso). *Biodiversity and Conservation*, **15**, 2085-2108.

530 Thiollay, JM. (2007) Raptor declines in West Africa: comparison between protected,
531 buffer and cultivated areas. *Oryx*, **41**, 322-329.

532 Trierweiler, C., Koks, B., Drent, R., Exo, KM., Komdeur, J., Dijkstra, C. & Bairlein, F.
533 (2007) Satellite tracking of two Montagu's Harriers (*Circus pygargus*): dual
534 pathways during autumn migration. *Journal of Ornithology*, **148**, 513-516.

535 Trierweiler, C. & Koks, BJ. (2009) Montagu's harrier *Circus pygargus*. *Living on the*
536 *edge: Wetlands and birds in a changing Sahel*. (ed. by Zwarts, L., Bijlsma, RG.,
537 van der Kamp, J & Wymenga, E). KNNV Publishing, Zeist, The Netherlands.

538 Trierweiler, C. (2010) *Travels to feed and food to breed. The annual cycle of a*
539 *migratory raptor, Montagu's harrier, in a modern world*. PhD thesis, University
540 of Groningen, Netherlands.

van Maanen, E., Goradze, I., Gavashelishvili, A. & Goradze, R. (2001) Trapping and hunting of migratory raptors in western Georgia. *Bird Conservation International*, **11**, 77-92.

Verma, A. (in press) Application of telemetry techniques for raptor research in India. *Telemetry in Wildlife Science*. (ed. by Sivakumar, K & Bilal Habib). ENVIS Bulletin: Wildlife and Protected Areas, **13** (1).

Webster, M. S. & Marra PP. (2005) The importance of understanding migratory connectivity and cross-seasonal interactions; pp. 199-209 in *Birds of Two Worlds* (R. Greenberg and P. Marra, eds.), Johns Hopkins University Press.

Western, D., Groom, R. & Worden, J. (2009) The impact of subdivision and sedentarization of pastoral lands on wildlife in an African savannah ecosystem. *Biological Conservation*, **142**, 2538-2546.

BIOSKETCHES

J. Terraube is a post-doctoral researcher in the Ecology group at the National Wildlife Research Institute (IREC), Ciudad Real, Spain. He is mainly interested by approaches integrating animal behaviour into population ecology and conservation biology, particularly in predator-prey interactions and their implications for the conservation of threatened predators in ecosystems facing rapid global changes.

564 Author's contribution: J.T., F.M., T.C. & B.A conceived the ideas; all authors jointly
565 collected the data; J.T., F.M., T.C. & B.A analyzed the data; J.T., F.M., T.C. & B.A led
566 the writing.

567

Table 1: Life cycles and signal transmissions of 10 transmitters applied to pallid harriers, and summarized migratory data for tracked individuals.

Harrier age/sex	Capture/nest site	Start of transmission/end of signals	Start autumn migration (departure breeding site)	Arrival wintering area	Total duration (days)	Total distance (km)	Average autumn migration speed (km.day ⁻¹)	Wintering area location (Lat./long. in decimal degrees)
adult male 1	Naurzum region	07 July 2008 - 19 April 2009	02 Aug.	20 Oct.	79	7301	92,42	Niger (14.68/8.64)
adult female 1	Naurzum region	27 June 2008 - 03 Aug. 2008	signal lost*	-	-	-	-	-
first-year male	Gujarat, India	11 March 2008 - 21 Sept. 2008	-	-	-	-	-	NW India (22.16/71.86)
adult female 2	Naurzum region	27 June 2008 - 24 July 2009	30 July	08 Nov.	101	9773	96,76	Front. Niger/Burk. Faso (14.5/0.21)
adult female 3	Turgai region	09 July 2007 - Oct. 2009	03 Aug.	16 Nov.	105	7691	73,25	Sudan (9.40/29.27)
adult female 4	Naurzum region	27 June 2008 - 10 Dec. 2008	29 July	25 Oct.	88	7184	81,64	Ethiopia (12.51/39.33)
adult male 2	Naurzum region	28 June 2008 - 03 Aug. 2008	signal lost*	-	-	-	-	-
adult male 3	Naurzum region	26 June 2008 - 11 Nov. 2008	22 Sept.	16 Oct.	24	5828	242,83	Sudan (9.126/32.404)
adult male 4	Naurzum region	30 June 2008 - 11 Aug. 2008	signal lost*	-	-	-	-	-
adult female 5	Naurzum region	26- June 2008 - present	18 July	28 Oct.	102	6255	61,32	Ethiopia (7.266/39.985)

*signal lost in the breeding area

Table 1 continued.

Harrier age/sex	Start spring migration	Arrival breeding grounds	Total duration (days)	Total distance (km)	Average spring migration speed (km.day⁻¹)
adult male 1	30 March	signal lost in Syria 19 April	20**	4262**	213**
adult female 1	-	-	-	-	-
first-year male	1 April	26 April	25	3322	133
adult female 2	16 March	7 April	22	8313	378
adult female 3	23 March	24 April	32	6988	218
adult female 4	signal lost in Ethiopia	-	-	-	-
adult male 2	-	-	-	-	-
adult male 3	signal lost in Sudan	-	-	-	-
adult male 4	-	-	-	-	-
adult female 5	23 March	23 April	30	5362	179

**refers to partial data as signal lost in route (data not included for calculation of average duration or distance in text)

Fig. 1: Migratory routes and wintering areas of pallid harriers as evidenced from ringing and recovery data from Kazakhstan (circles, solid lines) and from satellite tracking data (triangles, dashed lines). The western migratory routes (towards Africa) are shown in red, and the eastern routes (towards India) are shown in blue.

Fig. 2: Autumn (a) and spring (b) migration routes of six pallid harriers breeding in north-central Kazakhstan and of one pallid harrier wintering in India.

Fig. 1

Fig. 2