

1 **Geographical variation in inorganic arsenic in paddy field samples and commercial**
2 **rice from the Iberian Peninsula**

3

4 Antonio J. Signes-Pastor*¹, Manus Carey¹, Angel A. Carbonell-Barrachina², Eduardo
5 Moreno-Jiménez³, Andy J. Green⁴, Andrew A. Meharg¹.

6

7 ¹Institute for Global Food Security, Queen's University Belfast, David Keir Building,
8 Malone Road, Belfast, BT9 5BN, Northern Ireland.

9 ² AgroFood Technology Department, Miguel Hernández University, Carretera de
10 Beniel, km 3.2, 03312 Orihuela, Alicante, Spain

11 ³Department of Agricultural Chemistry, Universidad Autónoma de Madrid, 28049
12 Madrid, Spain.

13 ⁴Doñana Biological Station, EBD-CSIC, Amarico Vespucio Rd. 41092, Seville, Spain.

14

15 * Corresponding author, email – a.signes-pastor@qub.ac.uk

16

17 **Abstract**

18 This study investigated total arsenic and arsenic speciation in rice using ion
19 chromatography with mass spectrometric detection (IC-ICP-MS), covering the main
20 rice-growing regions of the Iberian Peninsula in Europe. The main arsenic species found
21 were inorganic and dimethylarsinic acid. Samples surveyed were soil, shoots and field-
22 collected rice grain. From this information soil to plant arsenic transfer was investigated
23 plus the distribution of arsenic in rice across the geographical regions of Spain and
24 Portugal. Commercial polished rice was also obtained from each region and tested for
25 arsenic speciation, showing a positive correlation with field-obtained rice grain.

26 Commercial polished rice had the lowest i-As content in Andalusia, Murcia and
27 Valencia while Extremadura had the highest concentrations. About 26% of commercial
28 rice samples exceeded the permissible concentration^[1] for infant food production as
29 governed by the European Commission. Some cadmium data is also presented, available
30 with ICP-MS analyses, and show low concentration in rice samples.

31

32 **Keywords**

33 Inorganic arsenic; Cadmium; Rice; Soil; Shoots; Iberian Peninsula; Arsenic speciation
34

35 **1. Introduction**

36 The greatest arsenic (As) toxicity is attributed to inorganic arsenic (i-As), a non-
37 threshold class 1 human carcinogen (IARC, 2004). Other than cancers, human exposure
38 to i-As has been associated with diverse health problems, including genotoxic effects
39 (Banerjee et al., 2013). Rice and rice-based products have been identified as prevalent
40 sources of i-As to human diet, particularly in South-East Asia where rice is the staple
41 food (IARC, 2004; Mondal & Polya, 2008; CAC, 2014; EFSA, 2009). A strong
42 correlation between urinary As and rice consumption has been reported (Banerjee et al.,
43 2013; Cascio et al., 2011; Melkonian et al., 2013). Several sub-populations have
44 elevated i-As intakes from rice consumption, notably babies and young children who
45 have 3-fold higher consumption based on body weight. In addition, adults who have
46 gluten allergies or are lactose-intolerant tend to consume more rice, thereby exposing
47 themselves to higher As contamination (EFSA, 2009; IARC, 2004; Meharg, Deacon, et
48 al., 2008; Signes-Pastor, Carey, & Meharg, 2016)

49 The UN FAO/WHO has proposed the maximum level of 0.200 mg/kg i-As in polished

50 rice (CAC, 2014). The European Commission (EC) has recently legislated the maximum
51 limit thresholds of i-As in rice. These are 0.200 mg/kg i-As for polished rice, 0.250
52 mg/kg i-As for parboiled rice and husked rice, 0.300 mg/kg i-As for rice waffles, rice
53 wafers, rice crackers and rice cakes, and 0.100 mg/kg i-As for rice destined for the
54 production of food for infants and young children (EC, 2015). However, there is still
55 much debate that these guidelines and standards are set too high to protect people's
56 health (Schmidt, 2015).

57 Arsenic is ubiquitous in the environment and flooding of soils, as in paddy cultivation,
58 creates anaerobism, which mobilizes i-As into porewater (Lu et al., 2009). This leads to
59 rice being much more efficient at assimilating As into its grain than other cereals
60 (Williams, Villada, et al., 2007). Rice grown under aerobic conditions, such as for
61 upland rice production, has much lower i-As in its grain. However, aerobic cultivation
62 leads to ~10-fold increase in Cadmium (Cd) concentration in rice grain (Xu, McGrath,
63 Meharg, & Zhao, 2008). Cadmium, like i-As is a carcinogen and renal toxicant with a
64 half-life in the human body of ~40 years (Meharg & Zhao, 2012).

65 Unlike other foods such as fish, As species in rice are dominated by i-As and the organic
66 compound dimethylarsinic acid (DMA) (Meharg et al., 2009; Williams et al., 2005;
67 EFSA, 2009). The i-As content in rice and rice-based products have been widely-
68 surveyed and variation in the concentration has been reported between countries
69 (Carbonell-Barrachina et al., 2012; Meharg et al., 2009) and specific differences in i-As
70 concentration in rice have also been reported within countries (Sommella et al., 2013;
71 Williams, Raab, Feldmann, & Meharg, 2007). If low-As rice grain can be sourced and
72 marketed this could potentially be sold as a premium product (Norton et al., 2009).

73 The rice growing regions in Portugal and Spain are distributed throughout the Iberian
74 Peninsula and represent about 36% of the total rice production in Europe (Ferrero &
75 Nguyen, 2004). Belgium, United Kingdom, Germany and France are the main EU
76 countries where Spanish rice is imported (Sainz, Sanz, Aguado, & Martín-Cerdeño,
77 2014).

78 In the current study, total arsenic (t-As) was determined in 40 soils samples from the
79 Iberian Peninsula, and As speciation was determined in 40 shoots and 20 field-collected
80 rice grains. In addition 144 commercial rice samples were categorised for As species,
81 including polished rice, parboiled rice and brown rice. The relationships between t-As in
82 soils and As species in shoot and rice, both field-collected and commercial, were
83 explored. Likewise the differences in As speciation of field-collected and commercial
84 rice were studied. The paddy field regions that produce low i-As were identified, and to
85 assess further potential toxicological risks, Cd concentration in field-collected and
86 commercial rice grain was also analysed.

87

88 **2. Material and Methods**

89 **2.1. Reagents and equipment**

90 Arsenic speciation in shoots and rice, both field-collected and commercial, was
91 performed using a Thermo Scientific IC5000 Ion Chromatography system, with a
92 Thermo AS7, 2 x 250 mm column with a Thermo AG7, 2 x 50 mm guard column
93 interfaced with a Thermo ICAP Q ICP-MS.

94 Total As in soil and Cd analyses in field-collected and commercial rice samples were
95 carried out with the Thermo ICAP Q ICP-MS in direct solution acquisition mode using a

96 Cetac ASX-520 Auto Sampler.
97 Other equipment used included a freeze dryer Christ Alpha 1-4 LD Plus, a Retch PM100
98 rotary ball-mill with a zirconium oxide lined vessel and zirconium oxide grinding balls,
99 a OHAUS- Discovery digital weighing scale with 5 decimals, a CEM MARS 6
100 1800W microwave digester, and a Sorvall Legend RT centrifuge. For As speciation and
101 Cd analyses 50 ml polypropylene centrifuge tubes, conical, with purple HDPE flat screw
102 were used for sample preparation. Teflon pressure vessels (CEM, MARS 6) were used
103 to digest soil samples to determine t-As.
104 Deionised water from a Milli-Q Integral 3 system was used for the preparation of
105 reagents and standards. All chemicals were, at least, pro analysis quality. Commercial
106 DMA and monomethylarsonic acid (MMA) from Supelco Analytical, arsenite and
107 arsenate from Sigma-Aldrich and arsenobetaine BCR n°626 were used to prepare As
108 speciation standards. A commercial Multi-Element Solution 2 in 5% HNO₃ (SPEX
109 CLMS-2) and a Multi-Element Solution 4 in water/Tr-HF (SPEX CLMS-4) were used to
110 prepare the standards for t-As and Cd analyses. In addition, commercial Rhodium from
111 Fluka Analytical was used as internal ICP-MS standard. BDH Prolabo Aristar 69%
112 nitric acid and BDH Prolabo Analar Normapur 30% hydrogen peroxide were used for
113 extraction/digestion and to convert any arsenite to arsenate.

114

115 **2.2. Sample sourcing**

116 Samples of soil, rice shoots and grain were collected in September 2014 from the eight
117 main rice-producing regions of the Iberian Peninsula as shown in **Figure 1**. From each
118 paddy field region 5 replicate samples of topsoil from 0-15 cm deep (n=40), shoots
119 (n=40) and mature rice grain (n=20), when possible, were collected. The mature rice
120 grain samples were collected when available from Andalucía (n=3), Catalunya (n=5),

121 Extremadura (n=1), Murcia (n=5), Portugal (n=1) and Valencia (n=5). The five
122 sampling sites were distributed throughout each region (5 sampling sites per region). At
123 each site, □200 g of topsoil, □200 g of shoots and □200 g of grain sub-samples were
124 gathered within a sampling area of 5 m x 5 m and mixed together in different plastic
125 bags for soil, shoots and rice grain, respectively. A market basket survey was also
126 carried out and commercial rice samples produced in the main rice production regions of
127 the Iberian Peninsula were purchased from supermarkets and local shops (n=144). This
128 set of samples included brown (n=20), parboiled (n=11) and polished (n=113) rice. The
129 rice-growing region was reported for a subset of 107 polished rice samples (Andalucía
130 (n=20), Aragón (n=6), Catalunya (n=14), Extremadura (n=3), Murcia (n=11), Navarra
131 (n=4), Portugal (n=20) and Valencia (n=29)).

132

133 **2.3. Arsenic speciation analysis**

134 *Sample preparation for shoots, dehusked field-collected and commercial rice grain.* All
135 samples were freeze-dried until complete dryness, and then powdered using a rotary
136 ball-mill. The powdered samples were weighed accurately to a weight of 0.1 g into 50
137 ml polypropylene centrifuge tubes. Then, 10 ml of 1% concentrated nitric acid were
138 added and left to stand overnight. Then samples were microwave digested. The
139 temperature was raised to 55°C in 5 min. and held for 10 min. and then to 75°C in 5 min
140 and held for 10 min. Finally the digest was taken up to 95°C in 5 min. and maintained at
141 this temperature for 30 min. Samples were cooled to room temperature. The digestate
142 was centrifuged at 4,500 g for 20 min. and a 1 ml aliquot was transferred to a 2 ml
143 polypropylene vial and 10 µl of analytical grade hydrogen peroxide was added to
144 convert any arsenite to arsenate to facilitate subsequent chromatographic detection.

145

146 *QA/QC procedures:* Each batch of samples included 2-3 blanks and 2-3 replicate
147 samples of rice certified reference material (CRM) were included (NIST 1568b Rice
148 flour). The rice CRM has t-As and the As species DMA, MMA and i-As concentrations
149 certified (0.285 ± 0.014 , 0.182 ± 0.012 , 0.012 ± 0.003 and 0.092 ± 0.010 mg/kg,
150 respectively).

151

152 *Chromatography:* Arsenic speciation was carried out using ion chromatography with
153 mass spectrometric detection (IC-ICP-MS). A gradient mobile phase including A: 20
154 mM ammonium carbonate and B: 200 mM ammonium carbonate, starting at 100% A,
155 changing to 100% B, in a linear gradient over 15 min. was used. The ICP-MS monitored
156 $m/z^+ 75$ using He gas in collision cell mode. The resulting chromatogram was compared
157 with that for authentic standards: arsenobetaine, DMA, MMA, tetramethyl arsonium
158 and i-As. Arsenic present under each chromatographic peak was calibrated using a
159 DMA concentration series. The arsenobetaine, MMA and tetramethyl arsonium
160 concentrations in shoot and field-collected and commercial rice grain samples were
161 below the LOD.

162

163 **2.4. Total arsenic and cadmium analyses**

164 *Sample preparation for dehusked field-collected and commercial rice grain.* Samples
165 were freeze-dried and powdered. The powdered samples were weighed accurately to a
166 weight of 0.1 g into 50 ml polypropylene centrifuge tubes. Then, 2 ml of concentrated
167 nitric acid and 2 ml hydrogen peroxide were added into the 50 ml polypropylene
168 centrifuge tubes and left to stand overnight. Then samples were microwave digested.
169 The temperature was raised to 95°C in 5 min. and held for 10 min. and then to 135°C in
170 5 min. and held for 10 min. Finally the digest was taken up to 180°C in 5 min. and

171 maintained for 30 min. Samples were cooled to room temperature. An internal standard
172 (30 µl of 10 mg/kg Rhodium) was added to the digestate and then accurately diluted to
173 30 ml with deionized distilled water. Lastly samples were analysed for Cd concentration.

174

175 *Sample preparation for soil.* Soil samples were oven-dried and 1.00 mm sieved. Soil
176 samples were weighed accurately to a weight of 0.1 g of soil into Teflon pressure
177 vessels. Then, 2 ml of concentrated nitric acid and 2 ml hydrogen peroxide were added
178 into the Teflon pressure vessels and left to stand overnight. Samples were microwave
179 digested as described earlier. The internal standard was added to the digestate and then
180 accurately diluted to 30 ml with deionized distilled water. Finally samples were analysed
181 for t-As.

182

183 *QA/QC procedures:* Each batch of samples included 2-3 blanks and 2-3 replicate
184 samples of rice CRM (NIST 1568b Rice flour) or 2-3 replicate samples of soil CRMs
185 (ISE 921 and NCS ZC73001). The rice CRM has the Cd concentration certified ($0.022 \pm$
186 0.001 mg/kg) and the soil CRMs have the t-As concentration certified (29.9 ± 1.78 and
187 18.0 ± 2.0 mg/kg for ISE 921 and NCS ZC73001, respectively).

188

189 **2.5. Statistics**

190 The As and Cd concentrations in the Iberian Peninsula samples did not follow a normal
191 distribution. Therefore the rank-based non-parametric test Kruskal-Wallis was used to
192 carry out inference statistical analyses. These statistical analyses and plots were
193 performed using the R Statistical Software (R Core Team, 2014). The limit of detection
194 (LOD) was calculated as the mean of blank concentrations plus three times the standard

195 deviation of the blank concentrations multiplied by the dilution factor. When samples
196 were below the LOD a value of $\frac{1}{2}$ LOD was assigned for statistical analyses of the data.

197

198 **3. Results**

199 *Analytical recoveries, arsenic species analyses.* The mean (\pm SE) recovery of rice CRM
200 flour NIST-1568b As species was $95 \pm 3\%$, $90 \pm 2\%$ and $95 \pm 4\%$ for DMA, MMA and
201 i-As, respectively, based on $n=8$. The LOD for As speciation, calculated from DMA
202 calibration and based on $n=5$ was 0.002 mg/kg.

203 *Analytical recoveries, total arsenic and cadmium analyses.* The mean (\pm SE) recovery of
204 rice CRM flour NIST-1568b Cd was $90 \pm 10\%$ based on $n=3$. The mean (\pm SE) recovery
205 of soil CRM ISE 921 and NCS ZC73001 t-As were $105 \pm 1\%$ and $110 \pm 2\%$ based on
206 $n=3$, respectively. The LOD for t-As and Cd analysis based on $n=5$ was 0.009 mg/kg.

207

208 *Soil.* The t-As concentration in soil ranged from 2.3 to 17 mg/kg with a median
209 concentration of 8.7 mg/kg across all regions. Soil from Portugal had the highest t-As
210 concentration (median of 15 mg/kg). Similar median values were found for Catalunya
211 (11 mg/kg), Andalucía (10 mg/kg), Aragón (9.5 mg/kg), Navarra (8.8 mg/kg) and
212 Valencia (8.2 mg/kg), while lower median t-As concentrations were found for Murcia
213 (5.4 mg/kg) and Extremadura (4.2 mg/kg); $p=0.005$.

214

215 *Shoots.* The i-As was predominant in rice shoots and DMA only represented a small
216 percentage of the sum of As species (Σ As). The i-As concentration ranged from 0.257 to
217 17.1 with a median concentration of 2.7 mg/kg for all shoot samples. It was found that
218 the highest median i-As concentrations were for Extremadura (11.0 mg/kg), Portugal

219 (9.8 mg/kg) and Catalunya (8.5 mg/kg), whereas lower median i-As concentrations were
220 for Valencia (2.9 mg/kg), Andalucía (2.5 mg/kg), Aragón (1.8 mg/kg), Navarra (1.5
221 mg/kg) and Murcia (1.4 mg/kg); $p=0.026$. A positive correlation factor was found
222 between soil t-As concentration and shoots i-As concentration ($R = 0.15$) (**Figure 2**).
223 The DMA concentration ranged from 0.002 to 0.162 with a median concentration of
224 0.006 mg/kg for all shoot samples. The median DMA concentrations in shoots from
225 Portugal (0.042 mg/kg) and Extremadura (0.031 mg/kg) were higher than that from
226 Aragón (0.011 mg/kg), Catalunya (0.008 mg/kg), Andalucía (0.005 mg/kg), Navarra
227 (0.004 mg/kg), Valencia (0.004 mg/kg) and Murcia (0.003 mg/kg); $P=0.014$.

228

229 *Dehusked field-collected rice grain.* The predominant As species in field-collected rice
230 grain was i-As. The i-As percentage, and i-As and DMA concentrations had the
231 following median and range across all the field rice grain samples: 85, 41-97%, 0.088,
232 0.052-0.161 mg/kg and 0.017, 0.003-0.073 mg/kg, respectively. The relationship
233 between soil t-As concentration and field-collected rice grain i-As concentration tended
234 to describe a hyperbolic pattern, approaching a maximum of approximately over 0.100
235 mg/kg (**Figure 2**). A positive correlation was found between field-collected Σ As and
236 commercial rice grain Σ As concentration ($R = 0.98$), and between field-collected i-As
237 and commercial rice grain i-As concentration ($R = 0.76$) (**Figure 3**). The i-As content in
238 the field-collected rice grain samples used to evaluate the relationship with commercial
239 rice had the following median and range concentration according to region: 0.100,
240 0.061-0.130 mg/kg (Andalucía), 0.120, 0.074-0.150 mg/kg (Catalunya), 0.075, 0.063-
241 0.161 mg/kg (Murcia) and 0.093, 0.063-0.097 mg/kg (Valencia) (**Figure 2** and **Figure**
242 **3**).

243

244 *Commercial rice*. Brown rice had the highest i-As concentration, $p < 0.001$; and polished
245 rice had the lowest Σ As concentration; $p = 0.001$ (**Table 1**). The median and range of i-As
246 concentration for the entire polished rice dataset was 0.071 and 0.027 - 0.175 mg/kg,
247 respectively. It was found that the highest median i-As concentration was for
248 Extremadura/Portugal (0.087 mg/kg), whereas the lowest median i-As contents were for
249 Andalucía (0.054 mg/kg), Valencia (0.063 mg/kg) and Murcia (0.057 mg/kg); $p < 0.001$.
250 The percentage of t-As represented by i-As in polished rice had a median of 57% and
251 varied from 14% to 95%. Polished rice from Murcia had the highest median i-As
252 percentage (87%), while similar values were found for Valencia, Catalunya,
253 Aragón/Navarra and Andalucía (ranging from 51% to 62%), and that for
254 Extremadura/Portugal was much lower (41%); $p < 0.001$. The DMA concentration in
255 polished rice had a median concentration of 0.055 mg/kg and ranged from 0.003 to
256 0.291 mg/kg across all regions. It was found that the highest median DMA concentration
257 was for Extremadura/Portugal (0.139 mg/kg), while the lowest median DMA content
258 was for Murcia (0.009 mg/kg); $p < 0.001$ (**Table 1** and **Figure 4**).
259 The Cd concentration was consistently low for all the rice grain samples and was close
260 to the LOD (**Table 1**).
261 The regression analysis of i-As against Σ As concentration in the entire dataset of
262 polished rice had a slope of 0.186 and R^2 equal to 0.38. The regression analysis of i-As
263 against Σ As concentration between regions showed that the slopes for Murcia,
264 Catalunya and Aragón/Navarra, ranging from 0.445 to 0.969, were higher than that for
265 Valencia, Andalucía and Extremadura/Portugal ranging from 0.091 to 0.232. The slope
266 and R^2 of DMA against Σ As concentration for all polished rice was 0.799 and 0.92,
267 respectively. Analysis of DMA against Σ As concentration between regions established
268 that Extremadura/Portugal, Andalucía and Valencia had similar slopes (0.896, 0.787 and

269 0.765) and higher than that for the other regions ranging from 0.150 to 0.539 (**Table 2**).
270 A negative correlation factor was found in polished rice between Σ As concentration and
271 i-As percentage ($R = -0.73$) (**Figure 4**).

272

273 **4. Discussion**

274 The Iberian Peninsula paddy fields soil As has a predominantly geogenic or mining-
275 derived origin (Ramos-Miras et al., 2014). The paddy field soils analysed here had a low
276 to moderate t-As concentration according to previous study (Khan, Stroud, Zhu,
277 McGrath, & Zhao, 2010). Paddy soil in Portugal had 3-fold higher median As content
278 than Murcia and Extremadura, where soil t-As was the lowest. Williams *et al.* (2011)
279 monitored As soil bioavailability to rice by analysing soil porewater dynamics and
280 applying the dynamic sampling technique diffusive gradients in thin films (DGT). They
281 found that paddy soils, even at baseline t-As concentration had As elevated grain due to
282 large labile As reservoirs that resupplied flux from the soil into the porewater (Williams
283 et al., 2011). Khan *et al.* (2010) also evaluated As bioavailability to rice and reported
284 higher As bioavailability in paddy soils irrigated with contaminated water, from which
285 As is likely to bear Fe oxides and absorb to soil minerals, than paddy soil with geogenic
286 or mining-derived As, from which As may be largely present in more insoluble and non-
287 labile forms (Khan et al., 2010).

288 In our Iberian study here i-As in shoots increased with soil t-As, demonstrating a
289 positive correlation. Yet, shoots from Extremadura had high i-As concentration, while
290 low soil t-As content, suggesting that a greater proportion of soil t-As from Extremadura
291 paddy soil was mobilized to be bioavailable for plant uptake compared to the soils from
292 the other regions. As bioavailability strongly depends on environmental factors (Zhao,
293 Ma, Meharg, & McGrath, 2009), suggesting that in Extremadura soil As is likely to be

294 in more labile forms, bearing Fe oxides and absorb to soil minerals (Khan et al., 2010).
295 A high proportion of soil t-As was also transferred to shoots from Catalunya and
296 Portugal soils compared to those from Murcia, Valencia, Aragón and Navarra. In these
297 latter regions, soil As is expected to be present in more insoluble and less labile forms,
298 and probably more influenced by the DOC dynamics as previous studies have suggested
299 (Khan et al., 2010; Williams et al., 2011). The main mechanisms that affect i-As
300 bioavailability in the Iberian Peninsula paddy fields deserve further investigation.

301 The As speciation in shoots here varied geographically and more than 95% of shoot As
302 was in the i-As form, in keeping with other studies (Abedin, Cresser, Meharg,
303 Feldmann, & Cotter-Howells, 2002; Norton et al., 2010). This is understandable since i-
304 As is predominant in porewater (Khan et al., 2010; Williams et al., 2011) and arsenite,
305 which is the dominant species of As in reducing environment, is readily assimilated by
306 rice plant via silicic acid pathway through aquaporin channels (Abedin, Feldmann, &
307 Meharg, 2002; Ma et al., 2008; Zhao et al., 2009). The As levels in rice shoots from
308 Andalucía have been reported previously: median of 2.6 and 1.2 mg/kg in rice shoots
309 from Doñana and Cádiz, respectively (Williams, Villada, et al., 2007). Similar values
310 were found in shoots from Valencia, Andalucía, Aragón, Navarra and Murcia. However,
311 an increase of up to 7-9-fold in the i-As content was found in shoot samples from
312 Extremadura, Portugal and Catalunya compared to that previously reported in Cádiz
313 (Williams, Villada, et al., 2007).

314 In the present study i-As and DMA dominated As speciation in field rice grain, in
315 agreement with earlier studies (Meharg et al., 2009; Williams et al., 2005; Zhao, Zhu, &
316 Meharg, 2013). Yet, DMA showed more efficient above ground translocation than i-As,
317 which may be due to its poor -SH coordination as suggested by previous studies (Norton

318 et al., 2010; Raab, Ferreira, Meharg, & Feldmann, 2007). DMA concentration in both
319 shoots and grain were in the same range, suggesting that DMA was unloaded at a similar
320 rate into these two compartments. In contrast, i-As in rice grain was two orders of
321 magnitude lower than in shoots, suggesting a much less efficient grain unloading of i-
322 As. The relationship between soil t-As and field-collected rice grain i-As suggested a
323 hyperbolic trend, describing a moderation grain i-As concentration at high soil t-As.
324 Similar trend has previously been reported between shoots and grain As concentration
325 due to a decrease translocation efficiency alongside increasing shoot As accumulation
326 (Lu et al., 2009; Norton et al., 2010; Williams, Villada, et al., 2007).

327 Graphical analysis of combined i-As and Σ As concentrations in rice grain from the field
328 and the market basket surveys were in agreement, linking As distribution from field-
329 collected to commercial rice grain samples. The As concentration in commercial rice
330 from the main rice-growing regions of the Iberian Peninsula was strongly influenced by
331 the type of rice and the geographical origin. Commercial brown rice type had twice i-As
332 and Σ As concentration than polished rice. This corroborates earlier studies that reported
333 higher As concentration in brown rice, where As is preferentially localized in the bran,
334 than in polished rice, where As is generally dispersed throughout the grain (Meharg,
335 Lombi, et al., 2008; Sun et al., 2008). An earlier survey including polished and brown
336 rice from Spain found a mean i-As concentration of 0.097 mg/kg (n=11) and of 0.154
337 mg/kg (n=11), respectively (Torres-Escribano, Leal, Vélez, & Montoro, 2008), which
338 compare well to the values reported here. Geographical As speciation variation has been
339 reported previously and environmental conditions seems to play a major role compared
340 to genetic factors (Zhao et al., 2013). Meharg *et al.* (2009) analysed an extensive dataset
341 including 901 polished rice samples from 10 countries and reported significant
342 geographical variation in t-As and i-As concentration. They found similar median i-As

343 concentrations for polished rice from China, Italy and U.S (0.120, 0.120 and 0.100
344 mg/kg, respectively), whereas lower median i-As concentration was for India and
345 Bangladesh (0.030 and 0.070 mg/kg) (Meharg et al., 2009). Geographical variation in i-
346 As content of rice has also been reported within a country. Sommella *et al.* (2013)
347 analysed rice from 4 rice-growing regions in Italy. For Lombardia, Piemonte and Emilia
348 similar i-As concentration was found (mean of ~0.100 mg/kg), while Calabria, located in
349 the south of Italy, had lower i-As content (mean of 0.060 mg/kg)(Sommella et al., 2013).
350 Prior studies have reported t-As and i-As concentration in rice from Spain (Meharg et
351 al., 2009; Williams et al., 2005). However, only a few studies with a limited number of
352 samples have reported i-As concentration in polished rice according to the rice-growing
353 region of the Iberian Peninsula. The mean i-As concentration has been previously
354 reported for polished rice from Valencia (0.075 mg/kg), Catalunya (0.101 mg/kg) and
355 Andalucía (0.101 mg/kg), which are within the range reported in this study (Torres-
356 Escribano et al., 2008). The mean i-As content of 0.180 mg/kg has been reported for
357 Portuguese polished, which is a higher value than that found here (Tattibayeva et al.,
358 2015). A significant regional i-As concentration variation was found with the lowest
359 concentration in Andalucía, Murcia and Valencia, while the highest was in
360 Extremadura/Portugal. This is consistent with the findings of the field study here.
361 Indeed, Portuguese soil had the highest t-As concentration, and although soil from
362 Extremadura had low t-As, the i-As content in shoots from Extremadura and Portugal
363 was much higher than that for Andalucía, Murcia and Valencia, where a less efficient i-
364 As transfer from soil to shoot and grain have been suggested.

365 A wide variation in the relative percentage of i-As and DMA has been reported
366 previously. William *et al.* (2005) compared As speciation in commercial polished rice
367 produced in Bangladesh, India, Europe and the U.S. They found high percentages of i-

368 As (~80%) in Bangladeshi and Indian rice. In comparison, European and U.S. rice had a
369 lower percentage of i-As with a mean of 64% and 42%, respectively, with the
370 corresponding high percentage of DMA (Williams et al., 2005). The mean i-As
371 percentage of 62% have been reported for Spanish rice (Torres-Escribano et al., 2008).
372 Compared to this value, in the study here, polished rice from Murcia had 1.4-fold higher
373 median percentage of i-As concentration, similar to that for Bangladeshi and Indian rice.
374 In contrast, Extremadura/Portugal had 1.5-fold lower median percentage of i-As content,
375 agreeing with that reported for U.S. rice (Williams et al., 2005).

376 The percentage of i-As decreased with Σ As concentration in all regions describing a
377 clear negative correlation, in keeping with earlier studies (Meharg et al., 2009). This
378 may suggest that a physiological switch occurs enhancing methylated As species plant
379 uptake from the soil micro-flora when reaching i-As critical levels, which concurs with
380 recent evidence suggesting the lack of *in planta* methylation ability in rice (Lomax et al.,
381 2012; Zhao et al., 2013).

382 Earlier surveys have carried out regression analyses of t-As against i-As and DMA
383 concentration in rice. Meharg *et al.* (2009) reported that the slope for India and
384 Bangladesh (0.796 and 0.719) were similar, while Chinese and Italian were similar
385 (0.599 and 0.506), and U.S. and Spanish rice much lower (0.275 and 0.193) (Meharg et
386 al., 2009). Zhao *et al.* (2013) reported a strong linear relationship between t-As and i-As
387 for rice from Asia with a slope of 0.78. In contrast, they reported that the U.S. rice
388 showed a hyperbolic pattern in the relationship, approaching a maximum of
389 approximately 0.15 mg/kg. European rice (Italy, Spain and France) samples appear to be
390 more variable and i-As/t-As relationship exhibits a pattern that was intermediate
391 between those of Asian and U.S rice (Zhao et al., 2013). The study here shows that

392 regression analyses of Σ As against i-As for polished rice from the Iberian Peninsula had
393 a low slope and was described with an intermediate pattern between the linear and the
394 hyperbolic trend, which agrees with earlier studies (Meharg et al., 2009; Zhao et al.,
395 2013). However, some differences were identified when carrying out regression analysis
396 of Σ As against i-As between regions. Low slopes and similar to that previously reported
397 for Spain and U.S were found for polished rice from Andalucía, Extremadura/Portugal
398 and Valencia. Aragón/Navarra had a similar slope to that described earlier for Chinese
399 and Italian, whereas Catalunya had a higher slope similar to that reported for India and
400 Bangladesh. Murcia had a much higher regression slope with a high R^2 , meaning that
401 most of the t-As was i-As. Regression analyses of Σ As and DMA described a strong
402 linear regression with a high slope for polished rice from the Iberian Peninsula and
403 similar to that previously reported for the U.S. (0.817) (Meharg et al., 2009). This trend
404 was consistent across all regions but Catalunya and Murcia, due to DMA data showed
405 wide variability.

406 Most of the commercial and field rice grain samples had a Cd concentration below the
407 LOD. This corroborates that rice in the Iberian Peninsula is cultivated under flooded
408 conditions, where the Cd bioavailability is low (Arao, Kawasaki, Baba, Mori, &
409 Matsumoto, 2009; Xu et al., 2008). Therefore, rice Cd concentration did not raise any
410 conflict in the Iberian Peninsula growing regions evaluated in the present study here.

411 The elevated i-As in rice is of concern since 26% of the all our Iberian dataset, which
412 include 144 samples of commercial polished, parboiled and brown rice, and 14% out of
413 113 commercial polished rice samples, would be illegal for the production of food for
414 infants and young children when the EC regulation is enforced in 2016 (EC, 2015). In
415 addition, there is still much debate that the UN WHO guidelines and EC standards are

416 set too high to protect people's health (Schmidt, 2015). Thus, it has been alternatively
417 suggested that the maximum value be 0.100 mg/kg i-As for all types of rice, and 0.05
418 mg/kg i-As for products targeted at young children and babies (Schmidt, 2015). The
419 0.05 mg/kg i-As is a lower value than that obtained for 80% of the polished rice
420 commercial samples included in this study or for even higher percentage when the
421 whole dataset is included in the calculations.

422 **5. Conclusions**

423 In this study it is shown that i-As and DMA are the main arsenic species in shoots and
424 rice, both field-collected and commercial. Paddy field soils had low to moderate t-As
425 concentrations, which was positively correlated with i-As in shoots and described a
426 hyperbolic trend with i-As in field-collected rice grain. The Extremadura paddy soil
427 suggested higher bioavailability for rice plant uptake. However, further studies regarding
428 i-As paddy soil bioavailability in the Iberian Peninsula are required. The As speciation
429 in commercial rice from the Iberian Peninsula compiled here is the largest dataset
430 reported as yet, and highlights that 26% of the rice samples would be illegal for the
431 production of food for infants and young children due to its elevated i-As concentration.
432 On searching for rice with lower i-As concentrations, Andalucía, Murcia and Valencia
433 showed the lowest levels.

434 **Acknowledgements**

435 This work was funded by a Marie Curie Intra-European Fellowship within the 7th
436 European Community Framework Programme (PIEF-GA-2013-622096).

437 **References**

- 438 Abedin, M. J., Cresser, M. S., Meharg, A. A., Feldmann, J., & Cotter-Howells, J. (2002).
439 Arsenic accumulation and metabolism in rice (*Oryza sativa* L.). *Environmental*
440 *Science and Technology*, 36(5), 962–968. doi:10.1021/es0101678
- 441 Abedin, M. J., Feldmann, J., & Meharg, A. A. (2002). Uptake kinetics of arsenic species
442 in rice plants. *Plant Physiology*, 128, 1120–1128. doi:10.1104/pp.010733.0.01
- 443 Arao, T., Kawasaki, A., Baba, K., Mori, S., & Matsumoto, S. (2009). Effects of water
444 management on cadmium and arsenic accumulation and dimethylarsinic acid
445 concentrations in Japanese rice. *Environmental Science and Technology*, 43(24),
446 9361–9367. doi:10.1021/es9022738
- 447 Banerjee, M., Banerjee, N., Bhattacharjee, P., Mondal, D., Lythgoe, P. R., Martínez, et
448 al. (2013). High arsenic in rice is associated with elevated genotoxic effects in
449 humans. *Scientific Reports*, 3, 2195. doi:10.1038/srep02195
- 450 CAC. (2014). Codex Alimentarius Commission. Report of the eighth session of the
451 codex committee on contaminants in food.
452 doi:ftp.fao.org/codex/Reports/Reports_2014/REP14_CFe.pdf. Accessed 02.11.15.
- 453 Carbonell-Barrachina, A. A., Wu, X., Ramírez-Gandolfo, A., Norton, G. J., Burló, F.,
454 Deacon, C., & Meharg, A. A. (2012). Inorganic arsenic contents in rice-based
455 infant foods from Spain, UK, China and USA. *Environmental Pollution*, 163, 77–
456 83. doi:10.1016/j.envpol.2011.12.036
- 457 Cascio, C., Raab, A., Jenkins, R. O., Feldmann, J., Meharg, A. A., & Haris, P. I. (2011).
458 The impact of a rice based diet on urinary arsenic. *Journal of Environmental*
459 *Monitoring : JEM*, 13(2), 257–265. doi:10.1039/c0em00482k

460 EC. (2015). Commission Regulation 2015/1006 of 25 June 2015 amending Regulation
461 (EC) No 1881/2006 as regards maximum levels of inorganic arsenic in foodstuffs.
462 doi:eur-lex.europa.eu/legal-content/EN/TXT/?uri=OJ:JOL_2015_161_R_0006.
463 Accessed 02.11.15.

464 EFSA. (2009). European Food Safety Authority. Scientific opinion on arsenic in food.
465 EFSA panel on contaminants in food chain (CONTAM). (Vol. 7).
466 doi:http://www.efsa.europa.eu/sites/default/files/scientific_output/files/main_docu
467 ments/1351.pdf. Accessed 02.11.15.

468 Ferrero, A., & Nguyen, N. (2004). Proceeding of the FAO Rice Conference. Rice is life.
469 The sustainable development of rice-based production systems in Europe. FAO
470 Rice Conference. doi:www.fao.org/docrep/008/y5682e/y5682e0g.htm. Accessed
471 02.11.15.

472 IARC. (2004). International Agency for Research on Cancer Monographs on the
473 Evaluation of Carcinogenic Risks to Humans. Some drinking-water disinfectants
474 and Contaminants, including Arsenic. (Vol. 84).
475 doi:monographs.iarc.fr/ENG/Monographs/vol84/mono84.pdf. Accessed 02.11.15.

476 Khan, M. A., Stroud, J. L., Zhu, Y. G., McGrath, S. P., & Zhao, F. J. (2010). Arsenic
477 bioavailability to rice is elevated in Bangladeshi paddy soils. *Environmental*
478 *Science and Technology*, 44(22), 8515–8521. doi:10.1021/es101952f

479 Lomax, C., Liu, W., Wu, L., Xue, K., Xiong, J., Zhou, J., & Steve, P. (2012).
480 Methylated arsenic species in plants originate from soil microorganisms. *New*
481 *Phytologist*, 193, 665–672. doi:10.1111/j.1469-8137.2011.03956.x.

482 Lu, Y., Adomako, E. E., Solaiman, A. R. M., Islam, M. R., Deacon, C., Williams, P. N.
483 et al. (2009). Baseline soil variation is a major factor in arsenic accumulation in

484 Bengal delta paddy rice. *Environmental Science and Technology*, 43(6), 1724–
485 1729. doi:10.1021/es802794w

486 Ma, J. F., Yamaji, N., Mitani, N., Xu, X.-Y., Su, Y.-H., McGrath, S. P., & Zhao, F. J.
487 (2008). Transporters of arsenite in rice and their role in arsenic accumulation in rice
488 grain. *Proceedings of the National Academy of Sciences of the United States of*
489 *America*, 105(29), 9931–9935. doi:10.1073/pnas.0802361105

490 Meharg, A. A., Deacon, C., Campbell, R. C. J., Carey, A.-M., Williams, P. N.,
491 Feldmann, J., & Raab, A. (2008). Inorganic arsenic levels in rice milk exceed EU
492 and US drinking water standards. *Journal of Environmental Monitoring : JEM*,
493 10(4), 428–431. doi:10.1039/B800981C

494 Meharg, A. A., Lombi, E., Williams, P. N., Scheckel, K. G., Feldmann, J., Raab, A., et
495 al. (2008). Speciation and localization of arsenic in white and brown rice grains.
496 *Environmental Science & Technology*, 42(4), 1051–7. doi:10.1021/es702212p

497 Meharg, A. A., Williams, P. N., Adomako, E., Lawgali, Y. Y., Deacon, C., Villada, A.
498 et al. (2009). Geographical variation in total and inorganic arsenic content of
499 polished (white) rice. *Environmental Science & Technology*, 43(5), 1612–7.
500 doi:10.1021/es802612a

501 Meharg, A. A., & Zhao, F. J. (2012). *Arsenic & Rice*. Berlin: Springer-Verlag.

502 Melkonian, S., Argos, M., Hall, M. N., Chen, Y., Parvez, F., Pierce, B. et al. (2013).
503 Urinary and dietary analysis of 18,470 Bangladeshis reveal a correlation of rice
504 consumption with arsenic exposure and toxicity. *PLoS ONE*, 8(11), 1–10.
505 doi:10.1371/journal.pone.0080691

506 Mondal, D., & Polya, D. A. (2008). Rice is a major exposure route for arsenic in

507 Chakdaha block, Nadia district, West Bengal, India: A probabilistic risk
508 assessment. *Applied Geochemistry*, 23(11), 2986–2997.
509 doi:10.1016/j.apgeochem.2008.06.025

510 Norton, G. J., Duan, G., Dasgupta, T., Islam, M. R., Lei, M., Zhu, Y. G. et al. (2009).
511 Environmental and genetic control of arsenic accumulation and speciation in rice
512 grain: Comparing a range of common cultivars grown in contaminated sites across
513 Bangladesh, China, and India. *Environmental Science and Technology*, 43(21),
514 8381–8386. doi:10.1021/es901844q

515 Norton, G. J., Islam, M. R., Duan, G., Lei, M., Zhu, Y. G., Deacon, C. M. et al. (2010).
516 Arsenic shoot-grain relationships in field grown rice cultivars. *Environmental*
517 *Science and Technology*, 44(4), 1471–1477. doi:10.1021/es902992d

518 R Core Team. (2014). R: A Language and Environment for Statistical Computing, R
519 Foundation for Statistical Computing. Vienna. doi:www.r-project.org/

520 Raab, A., Ferreira, K., Meharg, A. A., & Feldmann, J. (2007). Can arsenic-phytochelatin
521 complex formation be used as an indicator for toxicity in *Helianthus annuus*?
522 *Journal of Experimental Botany*, 58(6), 1333–1338. doi:10.1093/jxb/erl300

523 Ramos-Miras, J. J., Díaz-Fernández, P., SanJosé-Wery, A., Rodríguez-Martin, J. A.,
524 Roca, N., Bech, J. et al. (2014). Influence of parent material and soil use on arsenic
525 forms in soils: A case study in the Amblés Valley (Castilla-León, Spain). *Journal*
526 *of Geochemical Exploration*, 147, 260–267. doi:10.1016/j.gexplo.2014.09.003

527 Sainz, H., Sanz, A., Aguado, J., & Martín-Cerdeño, V. J. (2014). Alimentación en
528 España. Producción Industria Distribución Consumo. doi:www.mercasa-
529 ediciones.es/alimentacion_2014/AE2014/index.html#4/z. Accessed 02.11.15.

- 530 Schmidt, C. W. (2015). In search of “just right” The Challenge of Regulating Arsenic in
531 Rice, *123*(1), 16–19. doi:10.1289/ehp.123-A16
- 532 Signes-Pastor, A. J., Carey, M., & Meharg, A. A. (2016). Inorganic arsenic in rice-based
533 products for infants and young children. *Food Chemistry*, *191*, 128–134.
534 doi:10.1016/j.foodchem.2014.11.078
- 535 Sommella, A., Deacon, C., Norton, G. J., Pigna, M., Violante, A., & Meharg, A. A.
536 (2013). Total arsenic, inorganic arsenic, and other elements concentrations in
537 Italian rice grain varies with origin and type. *Environmental Pollution*, *181*, 38–43.
538 doi:10.1016/j.envpol.2013.05.045
- 539 Sun, G., Williams, P. N., Carey, A. M., Zhu, Y. G., Deacon, C., Raab, A. et al. A.
540 (2008). Inorganic arsenic in rice bran and its products are an order of magnitude
541 higher than in bulk grain. *Environmental Science and Technology*, *42*(19), 7542–
542 7546. doi:10.1021/es801238p
- 543 Tattibayeva, D., Nebot, C., Miranda, J. M., Cepeda, A., Mateyev, E., Erkebaev, M., &
544 Franco, C. M. (2015). A study on toxic and essential elements in rice from the
545 Republic of Kazakhstan: comparing the level of contamination in rice from the
546 European Community. *Environmental Geochemistry and Health*.
547 doi:10.1007/s10653-015-9687-y
- 548 Torres-Escribano, S., Leal, M., Vélez, D., & Montoro, R. (2008). Total and inorganic
549 arsenic concentrations in rice sold in Spain, effect of cooking, and risk assessments.
550 *Environmental Science and Technology*, *42*(10), 3867–3872.
- 551 Williams, P. N., Price, A. H., Raab, A., Hossain, S. A., Feldmann, J., & Meharg, A. A.
552 (2005). Variation in arsenic speciation and concentration in paddy rice related to
553 dietary exposure. *Environmental Science and Technology*, *39*(15), 5531–5540.

554 doi:10.1021/es0502324

555 Williams, P. N., Raab, A., Feldmann, J., & Meharg, A. A. (2007). Market basket survey
556 shows elevated levels of As in South Central U.S. processed rice compared to
557 California: consequences for human dietary exposure. *Environmental Science &*
558 *Technology*, 41(7), 2178–2183.

559 Williams, P. N., Villada, A., Deacon, C., Raab, A., Figuerola, J., Green, A. J. et al.
560 (2007). Greatly enhanced arsenic shoot assimilation in rice leads to elevated grain
561 levels compared to wheat and barley. *Environmental Science and Technology*,
562 41(19), 6854–6859. doi:10.1021/es070627i

563 Williams, P. N., Zhang, H., Davison, W., Meharg, A. A., Hossain, M., Norton, G. J. et
564 al. (2011). Organic matter-solid phase interactions are critical for predicting arsenic
565 release and plant uptake in Bangladesh paddy soils. *Environmental Science and*
566 *Technology*, 45(14), 6080–6087. doi:10.1021/es2003765

567 Xu, X. Y., McGrath, S. P., Meharg, A. A., & Zhao, F. J. (2008). Growing rice
568 aerobically markedly decreases arsenic accumulation. *Environmental Science and*
569 *Technology*, 42(15), 5574–5579. doi:10.1021/es800324u

570 Zhao, F. J., Ma, J. F., Meharg, A. A., & McGrath, S. P. (2009). Arsenic uptake and
571 metabolism in plants. *The New Phytologist*, 181(4), 777–794. doi:10.1111/j.1469-
572 8137.2008.02716.x

573 Zhao, F. J., Zhu, Y. G., & Meharg, A. A. (2013). Methylated arsenic species in rice:
574 geographical variation, origin, and uptake mechanisms. *Environmental Science &*
575 *Technology*, 47(9), 3957–66. doi:10.1021/es304295n

576

Table 1: Inorganic arsenic (i-As), DMA, ΣAs and Cd concentration (mg/kg dry weight) in commercial rice according to type of rice and region.

Origin	Type of rice	n	i-As (mg/kg)	DMA (mg/kg)	Σ As species (mg/kg)	Cd (mg/kg)
Portugal & Spain	Brown	20	0.157 ^a (0.053-0.247) ^A	0.084 (0.006-0.366)	0.302 ^a (0.083-0.619)	0.004 ^a (0.003-0.035)
	Parboiled	11	0.083 ^b (0.022-0.170)	0.079 (0.015-0.237)	0.201 ^{ab} (0.039-0.407)	0.004 ^a (0.003-0.027)
	Polished	113	0.071 ^b (0.027-0.175)	0.055 (0.003-0.333)	0.143 ^b (0.037-0.433)	0.003 ^b (0.003-0.049)
	<i>p-value</i>		<0.001	0.461	0.001	<0.001
Type of rice	Origin					
Polished	Andalucía	20	0.054 ^b (0.027-0.130)	0.055 ^{ac} (0.019-0.286)	0.107 ^b (0.046-0.397)	0.003 (0.003-0.009)
	Aragón/Navarra	10	0.067 ^{ab} (0.044-0.154)	0.057 ^{ac} (0.029-0.189)	0.128 ^{ab} (0.073-0.343)	0.003 (0.003-0.003)
	Catalunya	14	0.080 ^{ab} (0.046-0.147)	0.057 ^{ac} (0.034-0.237)	0.150 ^{ab} (0.121-0.283)	0.003 (0.003-0.037)
	Extremadura/Portugal	23	0.087 ^a (0.066-0.138)	0.139 ^a (0.045-0.291)	0.224 ^a (0.118-0.421)	0.003 (0.003-0.029)
	Murcia	11	0.057 ^b (0.039-0.121)	0.009 ^b (0.003-0.012)	0.064 ^b (0.043-0.133)	0.004 (0.003-0.005)
	Valencia	29	0.063 ^b (0.028-0.175)	0.044 ^{bc} (0.006-0.259)	0.106 ^b (0.037-0.362)	0.003 (0.003-0.049)
	<i>p-value</i>		<0.001	<0.001	<0.001	0.818

^AMedian (Min-Max); values with the same low case letters were not significantly different at p-value <0.05 for the variable studied (Kruskal Wallis test).

Table 2: Linear regression analysis of ΣA_s versus A_{s_i} and DMA (the intercept is “a” and the slope is “b”).

Type of rice	Origin	n	i-As			DMA		
			a	b	R ²	a	b	R ²
	Iberian Peninsula	113	0.044	0.186	0.38	-0.043	0.799	0.92
	Andalucía	20	0.031	0.188	0.61	-0.029	0.787	0.96
	Aragón/Navarra	10	0.013	0.445	0.80	-0.120	0.539	0.85
Polished	Catalunya	13	-0.022	0.748	0.46	0.021	0.254	0.10
	Extremadura/Portugal	23	0.068	0.091	0.22	-0.068	0.896	0.96
	Murcia	11	-0.006	0.969	0.98	0.007	0.150	0.02
	Valencia	29	0.035	0.232	0.41	-0.035	0.765	0.88

Figure 1: Iberian Peninsula map with the location of the paddy field regions sampled.



Figure 3: Relationship between polished commercial rice and field-collected rice grain i-As and Σ As concentration (mg/kg dry weight). Each value shows the median and standard error according to paddy field region.




Figure 4: Relationship between Σ As and i-As, DMA concentration (mg/kg dry weight) and i-As (%), and i-As, DMA and Σ As concentration (mg/kg dry weight) in commercial polished rice according to paddy field region.

