A lipidomic study on the regulation of inflammation and oxidative stress targeted by Marine ω -3 PUFA and Polyphenols in high- fat high-sucrose diets

Authors

Gabriel Dasilva^{ab*}, Manuel Pazos^a, Eduardo García-Egido^a, José M. Gallardo^a, Sara Ramos-Romero^c, Josep Lluís Torres^c, Marta Romeu^d, María-Rosa Nogués^d, Isabel Medina^a

^a Instituto de Investigaciones Marinas, Consejo Superior de Investigaciones Científicas (IIM-CSIC), E-36208 Vigo, Galicia, Spain. ^bDepartment of Analytical Chemistry, Nutrition and Bromatology and Research Institute for Food Analysis (I.I.A.A.), University of Santiago de Compostela, E-15782 Santiago de Compostela, Galicia, Spain. ^cInstituto de Química Avanzada de Catalunya (IQAC-CSIC), Jordi Girona 18-26, E-08034 Barcelona, Spain. ^dUnidad de Farmacología, Facultad de Medicina, Universidad Rovira i Virgili, Sant Llorenç 21, E-43201 Reus, Spain.

*Corresponding author. Tel. +34986231930; fax: +34986292762. E-mail address: gabrieldasilvaalonso@gmail.com

Abstract

The ability of polyphenols to ameliorate potential oxidative damage of ω -3 PUFAs when they are consumed together and then, to enhance their potentially individual effects on metabolic health are discussed through the modulation of fatty acids profiling and the production of lipid mediators. For that, the effects of the combined consumption of fish oils and grape seed procyanidins on the inflammatory response and redox unbalance triggered by high-fat high-sucrose (HFHS) diets were studied in an animal model of Wistar rats. A standard diet was used as control. Results suggested that fish oils produced a replacement of ω -6 by ω -3 PUFAs in membranes and tissues, and consequently they improved inflammatory and oxidative stress parameters: favored the activity of 12/15-lipoxygenases on ω -3 PUFAs, enhanced glutathione peroxidases activity, modulated pro-inflammatory lipid mediators synthesis through the cyclooxygenase (COX) pathways and down-regulated the synthesis *de novo* of ARA leaded by Δ 5 desaturase. Although polyphenols exerted an anti-oxidative and anti-inflammatory effect in the standard diet, they were less effective to reduce inflammation in the HFHS dietary model. Contrary to the effect observed in the standard diet, polyphenols up-regulated COX pathways towards ω -6 pro-inflammatory eicosanoids as PGE₂ and 11-HETE, and decreased the detoxification of ω -3 hydroperoxides in the HFHS diet. As a result, additive effects between fish oils and polyphenols were found in the standard diet in terms of reducing inflammation and oxidative stress. However, in the HFHS diets, fish oils seem to be the main responsible of the positive effects found in the combined group.

Keywords

Eicosanoids; docosanoids; fish oils; polyphenols; inflammation; oxidative stress.

1. Introduction

Regular consumption of ω -3 PUFAs, mainly EPA and DHA, has been associated to the prevention and palliation of diabetes, cardio vascular diseases (CVD), chronic inflammation and oxidative stress processes closely linked with obesity, insulin resistance, dyslipidemia, and hypertension [1–4]. Although no agreement has been reached on the recommended proportion of these PUFAs in the diet [5], dietary designs with an optimal proportion of EPA and DHA seem to potentiate these health benefits. In previous works, we have suggested that fish oils having a balanced proportion between EPA:DHA (1:1) produce better anti-inflammatory response modulating eicosanoid and docosanoids pathways [6], down-regulate oxidative stress and protein damage [7], and improve biochemical parameters related to metabolic disorders [8,9], compared to oils with unbalanced 1:2 EPA:DHA. Recently, other authors have found that EPA:DHA at 1:1 ratio reduces in higher extent the atherosclerotic plaque area in an animal model of atherosclerosis [10].

Despite the beneficial effects of ω -3 supplements, higher amount of PUFAs in diet leads to greater levels of oxidizable substrate that may deregulate redox balance and increase oxidative stress [11]. Oxidative stress has also been associated with the development of inflammation [12,13]. In this regard, the use of antioxidants is a common strategy to relieve lipid oxidation [14,15]. Polyphenols like grape seed procyanidins (GSP) extracts are known to act as anti-oxidants as well as anti-inflammatory compounds that regulate gene transcription factors and cytokines secretion associated with inflammation [12], reduce C-reactive protein (CRP) [16], and down-regulate the expression of proteins involved in phospholipase A₂ (PLA₂) synthesis, a key enzyme in inflammation [17]. GSP have also prevented oxidative injury by modulating the expression of antioxidant glutathione enzymes [18], regulating glucose homeostasis [19], decreasing plasma levels of triglycerides, low-density lipoprotein (LDL), and circulating free fatty acids (FFA) in plasma [20], and down-regulating the expression of proteins related to *de novo* synthesis of long chain FA [17]. Others have attributed health benefits including vasodilatory, antioxidant, and anti-inflammatory effects by modulating enzyme activities that trigger eicosanoid synthesis [21]. There is a great deal of evidence that antioxidant properties of polyphenols can be exerted by acting at cellular and non-cellular level. Polyphenols act via inhibition of LDL oxidation [22]. At cellular levels, polyphenols can inhibit NADPH oxidase, a major catalyst for the formation of ROS [23], and consequently they down-regulate isoprostanes and thromboxane synthesis [24] in a dose-dependent manner [22]. From a mechanistic point of view, some studies have suggested that the capacity of polyphenols to inhibit dose-dependently protein kinase C (PKC) mediates their effect down-regulating NADPH oxidase activity [22,24]. PKC is an enzyme that activates NADPH oxidase via phosphorylation of p47^{phox}. Lipid mediators from ω -3 PUFAs like resolvins, protectins and lipoxins also reduce NADPH oxidase activity inhibiting the p47^{phox} subunit membrane translocation [25]. The potential benefits of combining ω -3 and polyphenols have been barely described. Maestre *et al.* [26] showed that the inclusion of polyphenol-rich grape seed extracts

1 during the digestion of fish lipids decreased their oxidation. They also related the
2 addition of polyphenols with greater uptake of DHA into a Caco-2 enterocyte model
3 suggesting a protective role on ω -3 PUFAs which favors their intestinal bioavailability.
4 Sekhon *et al.* [27] have found additive properties of both supplements to diminish CRP
5 and cytokine factors in diet-induced inflammation studies.

7 High-fat high-sucrose (HFHS) diets are well-known model diets to intentionally
8 promote inflammation processes and metabolic disorders [16,17]. Zhang *et al.* [28]
9 provided novel evidences that high dietary fat intakes promoted neuronal oxidative
10 stress and inflammation through NADPH oxidase activation and ROS generation, and
11 PGE₂ production in cerebral cortex. The main goal of the anti-inflammatory nutrition
12 has been focused on reducing dietary components that may induce inflammatory
13 responses, such as ω -6 PUFAs, saturated fats, and refined carbohydrates, and increasing
14 ω -3 PUFAs and antioxidants that activate endogenous mechanisms to reduce
15 inflammation [12]. In this context, oxygenated lipid mediators derived from the
16 metabolism of marine ω -3 PUFAs, EPA and DHA, have been suggested as one of the
17 most efficient mechanisms to stop inflammation [29]. Additionally to the effects of ω -3
18 PUFAs on the regulation of gene expression through interacting with nuclear receptors
19 and transcription factors [12], the modulation of the amount and type of mediators
20 derived from PUFAs seems also to be an anti-inflammatory response key achieved by
21 fish oils [30]. For instance, ω -6 ARA eicosanoids are considered strong pro-
22 inflammatory molecules linked to hypercholesterolemia, type-2 diabetes, and obesity
23 [12,25,31], ω -3 eicosanoids and docosanoids from EPA and DHA are weak
24 inflammatory compounds [32,33], and resolvins and protectins from EPA and DHA are
25 potent anti-inflammatory mediators [34,35]. Previous research on genetically obese and
26 hypertensive rats, an animal model for Metabolic Syndrome, demonstrated the
27 effectiveness of fish oils having balanced proportions of EPA and DHA to down-
28 regulate the production of strong pro-inflammatory ω -6 eicosanoids [36].

30 The present study evaluated the role that fish oils, polyphenols, and the potential
31 cooperative effect of both, in the prevention of inflammation and oxidative stress
32 underlying metabolic disorders associated to the consumption of HFHS diets. For this
33 scope, Wistar rats were fed a HFHS diet and compared with controls fed a standard diet.
34 Marine PUFAs in a balanced 1:1 EPA:DHA proportion and GSP were employed as
35 dietary supplements. A lipidomic platform based on SPE-LC-MS/MS was applied to
36 determine the influence of each supplement on the synthesis of ω -3 and ω -6 eicosanoids
37 and docosanoids related to inflammatory and metabolic processes. The regulation of
38 lipid mediators was discussed together to the incorporation of fatty acids into
39 membranes and tissues, fatty acid desaturase activities, and the rise in the plasma
40 concentrations of circulating levels of FFAs. The PUFAs composition of cells has been
41 described to influence inflammation by modifying intracellular signals and altering the
42 pattern of lipid mediators produced [29]. FFAs are signaling molecules participating in
43 oxidative stress and inflammatory processes and have been associated with an increase
44 in fat mass and insulin resistance [37,38].

2. Materials and Methods

2.1 Animals and diets

Eight to ten week old female Wistar-Kyoto rats (n 56, Charles Rivers Lab., Wilmington, MA, USA) weighting 123-168 g were used. All animals were housed in cages (n 2-3/cage) with a 12 h light/dark cycle and constantly regulated humidity (50 ± 10 %) and temperature (22 ± 2 °C). The animals were given *ad libitum* access to feed and water. Animals were randomized in two dietary groups: a standard group (n 28) fed a standard diet (a modification of Teklad Global 2014, Harlan Teklad Inc., IN, USA) used as a control healthy model, and a HFHS group (n 28) fed a high-fat high-sucrose diet (a modification of TD 08811, Harlan Teklad Inc.) used as a risk model to develop chronic inflammation and metabolic disorders. Both groups were also subdivided into four supplemented subgroups (n 7): control, fish oils (ω 3), grape seed proanthocyanidins (GSP), and fish oil with GSP (ω 3&GSP). Each subgroup was fed the corresponding standard or HFHS diet enriched with the assigned supplement as listed in Supplementary Material S1. The experimental diets were pelletized in-house by lyophilization from frozen emulsions in order to incorporate each supplement. EPA:DHA 1:1 oil was obtained by blending adequate proportions of the commercially available fish oils from AFAMPES 121 EPA (AFAMSA, Vigo-Spain), EnerZona Omega 3 RX (ENERZONA, Milano-Italy) and Oligen liquid DHA 80% (IFIGEN-EQUIP 98, Barcelona-Spain). Soybean oil, obtained from cold pressing unrefined organic soybean oil, was from Clearspring Ltd. (London, UK). Grape seed extract Grajfnol (≥ 95 % total oligomeric proanthocyanidins, ≥ 60 % procyanidins dimer B₂) was from JF-Natural product (Tianjin, China). It is important to point out that fatty acids composition of ω 3 and ω 3&GSP supplemented feed had higher proportions of EPA, DPA, DHA, and ARA than control and GSP supplements. In order to compensate the caloric content between the four groups, control and GSP feeds were enriched with soybean oil as explained. Thereby, the fatty acids composition of control and GSP supplements had higher proportions of palmitic and oleic acids, LA, and ALA than ω 3 and ω 3&GSP supplements (Supplementary Material S2). After 25 weeks of nutritional intervention, rats were fasted overnight, anesthetized intraperitoneally with ketamine and xylazine, and killed by exsanguination. Handling and killing of the animals adhered strictly to the European Union guidelines for the management and care of laboratory animals and were approved by the CSIC Bioethical Issues Subcommittee (ref. AGL2009-12 374-C03-03).

2.2 Sample processing

The biological samples for fatty acids and metabolites analysis in plasma and erythrocytes were prepared according to previously described methods [6,36]. Briefly, on collection blood was added to EDTA containing tubes (1 mg/ml). After centrifugation erythrocytes were removed, packed and washed following Sonenberg protocol [39]. Then, PMSF (protease inhibitor) was added to plasma samples (erythrocyte free) and kept at -80 °C until used. Liver and adipose tissue samples were collected, frozen in liquid N₂ and stored at - 80 °C until used.

2.3 Total fatty acids analysis in liver, plasma, adipose tissue, erythrocytes, and circulating FFA analysis in plasma

Methods used for the analysis of total fatty acids (TFA) and FFA in plasma and erythrocytes were widely described in previous works [6,36]. Accordingly, a Bligh and Dyer [40] protocol was applied to plasma, liver, and adipose samples using dichloromethane:methanol:water (2:2:1, v/v) as extraction solvent. To analyze TFA, the organic phase was transesterified as Lepage and Roy [41] and TFA were determined by gas chromatography (GC/FID, Clarus 500, Perkin–Elmer). In the case of FFA in plasma, the organic phase was subjected to solid phase extraction (SPE) using aminopropyl cartridges (500 mg, 6 mL, Biotage, Uppsala-Sweden) as Kaluzny *et al.* [42] and then transesterified and analyzed by GC/FID. Similarly, a Bligh and Dyer modified procedure [43] was applied to erythrocyte samples and TFA were determined as indicated above. Results are summarized in Tables 1-5.

2.4 Fatty acid desaturase (FAD) indexes

FAD indexes were calculated from plasma and liver TFA data [44,45]. Stearoyl-CoA Desaturases SCD-16 and SCD-18 regulate the desaturation from palmitic acid (16:0) into palmitoleic acid (16:1 ω -7) and stearic acid (18:0) into oleic acid (18:1 ω -9), respectively. Desaturases Δ 4, Δ 5, and Δ 6 modulate the formation of ARA, EPA and DHA from LA and ALA. In detail, Δ 4D regulates the desaturation from DPA (22:5 ω -3) into DHA (22:6 ω -3). Δ 5D regulates the desaturation from DGLA (20:3 ω -6) into ARA (20:4 ω -6). Δ 6D regulates the desaturation from LA (18:2 ω -6) into DGLA (20:3 ω -6). The pathway from ALA (18:3 ω -3) into EPA (20:5 ω -3) is mediated by the action of both Δ 5D and Δ 6D, firstly Δ 6D desaturates ALA into SDA (18:4 ω -3) which is elongated and further desaturated by Δ 5D into EPA. Therefore, several FAD indexes were determined as the ratio between product and precursor as follows: SCD-16 = [16:1 ω -7/16:0], SCD-18 = [18:1 ω -9/18:0], Δ 4D = [22:6 ω -3/22:5 ω -3], Δ 5D = [20:4 ω -6/20:3 ω -6], Δ 6D = [20:3 ω -6/18:2 ω -6], and Δ 5/6D = [20:5 ω -3/18:3 ω -3]. Results are shown in Tab. 6.

2.5 Lipid peroxidation, CRP and TNF α levels in liver, protein carbonylation, ORAC and antioxidant enzymes in plasma

Lipid peroxidation levels in liver were measured through conjugated dienes hydroperoxides (intermediate lipid oxidation product) and malondialdehyde (MDA, end lipid oxidation product). For that, liver lipids were extracted [43] and quantified [46]. Then, conjugated dienes were measured following the American Oil Chemists Society method [47]. Total MDA in liver was derivatized with thiobarbituric acid (TBA) after protein hydrolysis [48] and precipitation [49], and determined by HPLC-fluorescence according to Fukunaga *et al.* [50].

Protein oxidation in plasma was measured by determining the level of total protein carbonylation by a fluorescence-based assay as previously described [7]. Briefly,

1 carbonyls groups were labelled with fluorescein-5-thiosemicarbazide and detected on
2 one-dimensional SDS-PAGE gels through a densitometric analysis. Plasma antioxidant
3 capacity was measured as the oxygen radical absorbance capacity (ORAC) [51]. Total
4 superoxide dismutase (SOD) activity was measured in plasma following the method of
5 Misra and Fridovich [52]. Glutathione peroxidase (GPX) activities in plasma were
6 determined according to Wheeler *et al.* [53]. ELISA kits from Cusabio Biotech (Hubei,
7 China) were used to measure CRP and TNF α in liver. Results are shown in Tab. 6.

8 9 2.6 Analysis of lipid metabolites by SPE-LC-ESI-MS/MS

10 The methodology developed by Dasilva *et al.* [54] was applied to extract and determine
11 ω -3 and ω -6 oxidized lipid mediators in plasma and results are shown in Tab. 7. Briefly,
12 plasma samples were subjected to SPE using Oasis-HLB cartridges (60mg, 3mL,
13 Waters, MA-USA) and extracts analyzed by LC-ESI-MS/MS on a Agilent 1260 Series
14 (Agilent, Palo Alto, CA) coupled to a dual-pressure linear ion trap mass spectrometer
15 LTQ Velos Pro with ESI (Thermo Fisher, Rockford, IL, USA). Instrument control and
16 data acquisition were done with Xcalibur software. Retention times for target
17 compounds and individual MS/MS parameters are summarized in Supplementary
18 Material S3.

19 20 2.7 Statistics

21 Statistical analyses were performed by one-way and two-way analysis of variance
22 (ANOVA) with R free software (version 3.2.4). Normal distribution and heterogeneity
23 were evaluated and non-parametric Kruskal Wallis analyses were required when data
24 distribution did not fit a Gaussian model or heterogeneity was found in variances. The
25 means were further compared by the post-hoc test Fisher least square difference (Fisher
26 LSD) and significant differences were set at $p < 0.05$. Data presented are expressed as
27 mean $\pm$ SD.

28 29 3. Results

30 Results are shown according to the two dietary groups. Firstly, the work shows results
31 of animals fed standard diets which were used as a control healthy model. Then, results
32 corresponding to the group fed HFHS diets associated to the development of chronic
33 inflammation and metabolic disorders are presented [55]. Animals from each dietary
34 experiment were divided in four supplemented subgroups: control, fish oils (ω 3), grape
35 seed proanthocyanidins (GSP), and fish oil with GSP (ω 3&GSP). The regulation of
36 plasma, membrane and tissues FAs profile, ω 6/ ω 3 ratios, FADs activity, and several
37 eicosanoid and docosanoids pathways were addressed. Results of
38 eicosanoids/docosanoids and fatty acids profiling were related with lipid peroxidation
39 levels in liver, carbonylation of plasma albumin, and with previous measurements in the
40 same animals for the plasma antioxidant capacity, antioxidant enzymes activities in
41 plasma (SOD and GPX) and inflammation markers in liver (CRP and TNF α).

3.1 Effect of fish oils and polyphenol supplements on the Standard diet

3.1.1 Fatty acids profile in tissues and plasma

3.1.1.1 TFA incorporation into liver

Supplementation with fish oils largely modulated the incorporation of PUFAs in liver in concordance with the FA composition of the supplements. In detail, the proportion of ω -3 EPA, DPA, and DHA and ω -6 DGLA increased in ω 3 and ω 3&GSP groups related to control, and it was observed a specific proportional decrease in the amount of ω -6 ARA, γ -ALA, and ω -9 oleic acid. Supplementation with polyphenols did not influence the incorporation of FA into liver; and thus, TFA profiles in GSP and control groups were similar. As a consequence, the inflammatory index ω -6/ ω -3 in liver was significantly reduced after fish oils treatments compared with control and GSP groups. Animals fed fish oils (EPA:DHA 1:1) accumulated these PUFAs into liver in a proportion of EPA:DHA 1:4. However, animals from control and GSP groups showed EPA:DHA ratios in liver around 1:40 – 1:30, suggesting a higher accumulation and production of DHA synthesized from EPA. Finally, it was observed that all groups accumulated around 50% of PUFAs, 10% of MUFAs, and 40% of SFAs in liver without significant differences due to the supplementation. Results are shown in Tab. 1.

3.1.1.2 TFA profile in plasma

Regarding to the modulation of TFA in plasma due to the supplements (Tab. 2), fish oils increased amounts of ω -3 EPA, DPA, and DHA and ω -6 DGLA with the corresponding proportional decrease of ω -6 ARA and γ -ALA as observed in liver. GSP group showed a similar profile to control group except for a significant increase of DGLA. The inflammatory index ω -6/ ω -3 was also significantly reduced after fish oils treatments compared with control and GSP groups as observed in liver. Percentages of PUFAs, MUFAs, and SFAs in plasma followed the same trend as described in liver. Therefore, the profile of TFA observed in plasma for all groups was similar to the profile of TFA in liver as expected due to plasmatic lipids mostly come from liver secretion in starvation [56]. However, differences were observed in the relative percentage of DHA and ARA between plasma and liver: the amount of DHA in plasma was found decreased for all groups compared with the amount accumulated in liver; similarly, the proportion of ARA was higher in plasma than in liver. In addition, the EPA:DHA ratio found in plasma in animals fed fish oils was 1:1,5, with lower proportion of DHA than ratio found in liver. The same trend was observed in control and GSP groups: EPA:DHA ratio was 1:10 showing a minor proportion of DHA than in liver.

3.1.1.3 TFA incorporation into erythrocyte membranes

The analysis of FA incorporated into erythrocyte membranes (Tab. 3) showed that the proportion of PUFAs was significantly affected by fish oils similarly to the effect observed in liver and TFA of plasma. In detail, the incorporation of EPA, DPA, and DHA was increased in the ω 3 and ω 3&GSP groups accompanied with the specific proportional decrease in the incorporation of ARA. ALA and γ -ALA were not incorporated into membranes in the four groups, and DGLA incorporation was similar

between all groups and not modulated by supplements. Finally, polyphenols did not influence the incorporation of FA into erythrocytes. Therefore, the ω -6/ ω -3 ratio in membranes was significantly reduced in the ω 3 and ω 3&GSP groups compared with control and GSP. The proportion of PUFAs, MUFAs, and SFAs incorporated in membranes was similar than plasma profiles. However, the ratio EPA:DHA showed a higher proportion of DHA vs. EPA in membranes compared with ratio found in plasma. Animals fed fish oils incorporated these PUFAs in a proportion of EPA:DHA 1:3, and animals from control and GSP groups showed ratios around 1:16 – 1:20.

3.1.1.4 TFA accumulation in adipose tissue

The concentration of myristic and palmitic saturated acids, and EPA, DPA, and DHA PUFAs was increased in the ω 3 and ω 3&GSP groups related to control group simultaneously with a decrease of oleic, ARA and LA levels. The accumulation of DGLA was not modulated by supplements. PP group showed a similar fatty acid profile than control. Then, ω -6/ ω -3 ratio found in adipose tissue was significantly reduced in ω 3 and ω 3&GSP groups compared with control and GSP groups. Adipose tissue was characterized by a higher content of oleic and LA acids and a lower content of stearic, ARA and DHA acids in all groups compared with the TFA profile observed in plasma, liver and erythrocytes. Accordingly, the proportion of PUFAs, MUFAs, and SFAs accumulated in adipocytes showed different values than plasma profiles towards a higher accumulation of MUFAs and lower amounts of PUFAs and SFAs. The ratio EPA:DHA 1:3 was similar to that found in erythrocytes in ω 3 and ω 3&GSP groups. Animals from control and GSP groups showed EPA:DHA 1:6. Results are shown in Tab. 4.

3.1.1.5 Circulating FFAs profile in plasma

Circulating FFAs in plasma (Tab. 5) are mostly derived from adipose tissue secretions. Accordingly, fish oils showed the same modulation of FFA profile that had been observed in adipose, and polyphenols did not influence the composition of FFA. In detail, it was found an increase in the proportion of myristic, palmitic, EPA, DPA, and DHA, with the subsequent proportional reduction of oleic, ALA, LA and ARA in the ω 3 and ω 3&GSP groups compared with control. GSP group showed a similar fatty acid profile than control. As a consequence, the ω -6/ ω -3 ratio was significantly reduced in ω 3 and ω 3&GSP groups compared with control and GSP. Although the effect of supplements was similar in both adipose tissue and circulating FFA profiles, the proportion of circulating palmitic acid (SFA) was higher in plasma than adipose, and the contrary was observed for oleic acid (MUFA), suggesting a preference of adipose to accumulate MUFAs instead of SFAs in the standard diet. Regarding to the ratio EPA:DHA found in circulating fatty acids, it was observed a proportion 1:2 EPA:DHA in ω 3 and ω 3&GSP groups, different than that found in adipose tissue which indicated a greater accumulation of DHA vs EPA in adipose (ratio 1:3). Control and GSP groups showed EPA:DHA 1:6 and 1:9, respectively. The total concentration of FFA in plasma showed a trend to lower values for ω 3 and GSP groups, and interestingly this parameter resulted to be significantly lower for the combined ω 3&GSP group.

3.1.2 FADs indexes

FAD indexes were calculated regarding to the TFA data in plasma [44]. The effects of supplementation of fish oils in $\omega 3$ and $\omega 3$ &GSP groups related to control showed the following results: (a) significant lower values of $\Delta 5D$ [ARA/DGLA], (b) and significant higher values of $\Delta 4D$ [DHA/DPA] and $\Delta 5/6D$ [EPA/ALA], and a tendency towards higher values of $\Delta 6D$ [DGLA/LA]. It is remarkable that the increase of $\Delta 5/6D$ [EPA/ALA] was significantly higher than the increase of $\Delta 4D$ [DHA/DPA] after fish oils supplementation.

Interestingly, the supplementation with polyphenols resulted in a modulation of FAD indexes related to control: (a) lower values of $\Delta 5D$ [ARA/DGLA] and $\Delta 4D$ [DHA/DPA] in GSP (b) and higher values of $\Delta 6D$ [DGLA/LA] in GSP and $\omega 3$ &GSP groups. That means a significant accumulation of DGLA related to control and a trend to lower amounts of ARA, EPA and DHA. It is also remarkable the additive effect observed on the reduction of $\Delta 5D$ [ARA/DGLA] for the $\omega 3$ &GSP group which showed the lowest level compared with control, GSP and $\omega 3$ groups. The studied supplements did not exert any significant influence on SCD-16 and SCD-18. Results are shown in Tab. 6.

In high agreement with FAD indexes calculated in plasma, these indexes calculated in liver (Tab. 6) also showed decrease of $\Delta 5D$ [ARA/DGLA], and increase of $\Delta 4D$ [DHA/DPA], $\Delta 5/6D$ [EPA/ALA], and $\Delta 6D$ [DGLA/LA] in $\omega 3$ and $\omega 3$ &GSP groups related to controls. GSP group in liver showed lower values of $\Delta 4D$ [DHA/DPA] and a tendency to decrease $\Delta 5D$ [ARA/DGLA] compared with control.

3.1.3 Eicosanoid and docosanoids levels

A total of nine compounds derived from PUFA oxidative metabolism were identified and quantified in plasma, five EPA eicosanoids, two DHA docosanoids, and two ARA eicosanoids (Tab. 7). The primary oxidation products found were the hydroperoxides 12-HpEPE, 15-HpEPE, and 17-HpDoHE. The secondary metabolites identified are hydroxides (12-HEPE, 15-HEPE, 17-HDoHE, and 11-HETE), a thromboxane (TXB₃), and a prostaglandin (PGE₂). Finally, other searched compounds as 5-HEPE, 8iso-PGF_{3 α} , 8iso-PGF_{2 α} , PGD₃, PGE₃, 11-HDoHE, 4-HDoHE, RvD₁, PD_x, and LTB₄ were under the detection limits of the method.

Data revealed that dietary interventions with fish oils in the $\omega 3$ group compared with control: (a) significantly decreased the concentration of 12-HpEPE and enhanced 15-HpEPE (EPA primary metabolites) coming from 12-LOX and 15-LOX activity respectively, (b) and significantly increased levels of the corresponding secondary metabolites from EPA: 12-HEPE produced from 12-HpEPE, and 15-HEPE from 15-HpEPE; (c) largely decreased the production of 17-HpDoHE (DHA primary metabolite) mediated by 15-LOX, (d) and increased 17HDoHE, secondary metabolite of DHA

produced from 17-HpDoHE; (e) and reduced the synthesis of secondary ARA metabolites (11-HETE and PGE₂) resulting from COX activity. The intervention with polyphenols in the GSP group compared with control: (a) reduced levels of primary metabolites from EPA and DHA produced by 12-LOX and 15-LOX (12-HpEPE, 15-HpEPE, and 17-HpDoHE), (b) did not significantly affect their secondary metabolites (12-HEPE, 15-HEPE, and 17-HDoHE) although it was observed a tendency to decrease their levels; (c) and significantly decreased COX secondary metabolites from ARA. Finally, TBX₃ resulting from COX activity on EPA was not detected in the GSP group, and no differences in its levels were observed between control, ω3, and ω3&GSP groups.

Interestingly, the consumption of fish oils together with polyphenols in the ω3&GSP group produced: (a) a significant decrease in the concentration of primary metabolites, 12-HpEPE and 17-HpDoHE, compared to control group showing an additive effect between both fish oils and polyphenols supplements over 12-LOX and 15-LOX activities, (b) a significant enhancement of 15-HpEPE compared to control as observed in the ω3 group; however, the concentration of 15-HpEPE in the ω3&GSP group was reduced related to the ω3 group in agreement with the reducing effect exerted by polyphenols in this pathway mediated by 15-LOX; (c) a trend to increase levels of 12-HEPE, 15-HEPE, and 17-HDoHE (secondary metabolites) was observed in ω3&GSP related to control although not statistically different, showing again a combined effect of both supplements (up-regulation of the formation of these secondary metabolites due to fish oils and down-regulation produced by polyphenols); (d) and a specially interesting decrease of pro-inflammatory secondary ARA metabolites (11-HETE and PGE₂) synthesis, according to the combined down-regulation effect of individual supplements.

3.1.4 Oxidative stress parameters

Levels of conjugated diene hydroperoxides and TBARS in plasma were under detection limits (data not shown). Lipid peroxidation in liver showed that the level of conjugated diene hydroperoxides was slightly lower in the ω3 and ω3&GSP groups related to control, whereas the concentration of secondary oxidation byproducts measured as TBARS was significantly higher. The GSP group did not modify conjugated dienes levels related to control and showed a trend to reduce the production of TBARS. Consequently, the combination of both supplements (ω3&GSP group) showed lower amount of conjugated diene hydroperoxides than control and slightly lower amount TBARS than ω3 group as a result of polyphenols effect. Data revealed an inhibition of protein carbonylation measured in plasma in the GSP group related to control and no effects on the groups supplemented with fish oils were observed. Finally, plasma antioxidant capacity by ORAC and the activity of some antioxidant enzymes was measured in previous studies executed in the same animals. In detail, no significant differences in ORAC index were recorded between groups, although a tendency towards higher values was observed in the ω3&GSP group compared to other groups. SOD (involved in the detoxification of O₂⁻) and GPX (involved in the detoxification of hydroperoxides to give hydroxides) activities were associated with the synthesis of lipid

mediators. SOD showed a trend to be lower in the ω 3 and GSP groups than in control, and the additive effect was observed when both supplements were combined leading to significant lower values. GPX was found increased in ω 3 and GSP groups, and the additive effect was also observed in ω 3&GSP compared with control. Results are shown in Tab. 6

3.2 Effect of fish oils and polyphenol supplements on the HFHS diet

3.2.1 Fatty acids profile in tissues and plasma

3.2.1.1 TFA incorporation into liver

As observed in the standard diet, supplementation with fish oils largely modulated the incorporation of PUFAs in liver in the HFHS diet. Similarly, the proportion of ω -3 EPA, DPA, and DHA increased in ω 3 and ω 3&GSP groups related to control, and a proportional decrease of ω -6 ARA and γ -ALA was observed. Supplementation with polyphenols did not influence the TFA profile compared with control group. Therefore, the ω -6/ ω -3 index was found significantly reduced in ω 3 and ω 3&GSP compared with control and GSP groups as in the standard diet. Contrary to results observed in the standard diet, the accumulation of DGLA and oleic acid in liver was not significantly modulated by fish oils supplements in the HFHS dietary context. In addition, all groups with HFHS diet significantly accumulated higher proportions of oleic acid in liver and lower proportions of DHA than corresponding groups fed standard diet. EPA:DHA ratios found in liver were similar to that found in animals fed with standard diet and fish oils supplements. In detail, ω 3 and ω 3&GSP groups accumulated EPA:DHA 1:4 as in the standard diet, and control and GSP groups showed higher accumulation and production of DHA (EPA:DHA 1:20 – 1:16). Interestingly, control and GSP groups accumulated/produced half of the proportion of DHA under the HFHS diet under the standard diet (EPA:DHA 1:40 – 1:30). Finally, all groups accumulated similar percentages of PUFAs (45%), MUFAs (15%), and SFAs (40%) as in the standard diet experiment, without significant differences due to the supplementation. Results are shown in Tab.1.

3.2.1.2 TFA profile in plasma

TFA profile in plasma (Tab. 2) in the HFHS diet was similar to that found in the standard experiment and in agreement with the profile found in liver. Fish oils increased amounts of ω -3 EPA, DPA, and DHA and ω -6 DGLA with the corresponding proportional decrease of ω -6 ARA and γ -ALA as observed in liver. As in the standard diet, GSP group did not show significant differences in the proportion of TFA in plasma related to control group with the exception of DGLA which showed a significant enrichment compared to control. The inflammatory index ω -6/ ω -3 was significantly reduced in ω 3 and ω 3&GSP groups compared with control and GSP groups. The profile of TFA observed in plasma and percentages of PUFAs, MUFAs, and SFAs for all groups in the HFHS diet were quite similar to the profiles found in liver, with the exception of the relative amount of ARA and DHA. The relative proportion of DHA in plasma was minor than accumulated in liver; and similarly, the proportion of ARA was

higher in plasma, as it had been observed in the standard diet. Finally, the EPA:DHA ratio was modulated by fish oils supplements similarly to the standard diet. In detail, the ω 3 and ω 3&GSP groups showed 1:1.5 EPA:DHA in plasma. And thus, the proportion of DHA was lower in plasma than the ratio found in liver. Regarding to control and GSP groups, they showed EPA:DHA 1:5, and again a minor proportion of DHA than in liver. Similarly to results observed in liver, control and GSP groups accumulated/produced half of the proportion of DHA in the HFHS diet than these groups in the standard diet (EPA:DHA 1:10).

3.2.1.3 TFA incorporation into erythrocyte membranes

The incorporation of FA into erythrocyte membranes (Tab. 3) in the HFHS diet followed the same trend as in the standard diet. Only the incorporation of PUFAs was significantly affected by fish oils. In the ω 3 and ω 3&GSP groups, the incorporation of EPA, DPA, and DHA increased, and ARA proportionally decreased related to control. ALA and γ -ALA were not incorporated and DGLA proportions were not modulated by the supplements. Polyphenols did not modify the profiles of FA in membranes. Consequently, erythrocyte membranes showed decreased ω -6/ ω -3 ratios in the ω 3 and ω 3&GSP groups compared with control and GSP. The proportion of PUFAs, MUFAs, and SFAs between membranes and plasma were similar as observed in the standard diet. In the same trend as in plasma and liver, the ratio EPA:DHA in erythrocytes showed minor proportion of DHA in control and GSP groups in the HFHS diet (EPA:DHA 1:10) compared with the standard one (EPA:DHA 1:16-1:20). The same was observed in ω 3 and ω 3&GSP groups in the HFHS diet (EPA:DHA 1:2) and standard diet (EPA:DHA 1:3).

3.2.1.4 TFA accumulation in adipose tissue

Regarding to the influence of the supplements in the accumulation of fatty acids in the adipose tissue, results were similar to the standard diet. The proportion of myristic, EPA, DPA, and DHA was increased in the ω 3 and ω 3&GSP group related to control group with the subsequent decrease of oleic, ARA and LA. The accumulation of DGLA was not modulated by the supplements and GSP group showed a similar fatty acid profile than control. Therefore, the inflammatory index ω -6/ ω -3 was significantly reduced in the ω 3 and ω 3&GSP groups compared with control and GSP ones. The profile of TFA in the adipose tissue was richer in oleic acid and less rich in stearic, ARA and DHA acids in all groups compared with plasma, liver and erythrocytes as in the standard diet. The ratio EPA:DHA found in ω 3 and ω 3&GSP groups was around 1:3; meanwhile control showed EPA:DHA 1:6 and GSP did not accumulate EPA. Results are shown in Tab. 4.

3.2.1.5 Circulating FFAs profile in plasma

The influence of fish oils in the proportion of circulating FFA in plasma (Tab. 5) was similar to that found in adipose, as it had been observed in the standard diet. The proportion of stearic, EPA, DPA, and DHA increased proportionally to the reduction of oleic, LA and ARA in ω 3 and ω 3&GSP groups compared with control. Polyphenols did

not influence the profile of circulating FFA. Therefore, fish oils treatments showed the lowest values for the inflammatory index ω -6/ ω -3. Similarly to the standard diet, the proportion of circulating oleic acid (MUFA) was lower in plasma than adipose, and the contrary was observed for palmitic and stearic acids (SFAs). In the ω 3 and ω 3&GSP groups the proportion EPA:DHA was 1:2 as in the standard experiment, and control and GSP groups showed a proportion of EPA:DHA 1:3.5. In a different way to the standard diet, the total concentration of FFA in plasma showed larger values for the GSP group compared with control, and a trend to lower values for both ω 3 and ω 3&GSP groups (this last trend was also observed in the standard diet).

3.2.2 FADs indexes

Comparative results between HFHS and standard diets showed a similar influence of supplements on FAD indexes measured in plasma (Tab. 6). In ω 3 and ω 3&GSP groups related to control it was observed: (a) a significant decrease of Δ 5D [ARA/DGLA], (b) a significant increase of Δ 4D [DHA/DPA], Δ 5/6D [EPA/ALA], and SCD-18 [oleic/stearic], (c) and a tendency to higher values of Δ 6D [DGLA/LA]. As observed in the standard diet, the increase of Δ 5/6D [EPA/ALA] in fish oils groups was significantly higher than the increase of Δ 4D [DHA/DPA]. The effect of polyphenols related to control group showed: (a) lower values of Δ 5D [ARA/DGLA] (b) and higher SCD-18 [oleic/stearic]. Interestingly, the ω 3&GSP group showed an additive effect on the reduction of Δ 5D [ARA/DGLA]. The studied supplements did not exert a different influence on SCD-16. As occurs in the standard diet, these FAD indexes calculated in liver (Tab. 6) mostly agree with FAD indexes in plasma for the ω 3 and ω 3&GSP groups. In detail, liver indexes also showed the decrease of Δ 5D [ARA/DGLA], and increase of Δ 4D [DHA/DPA] and Δ 5/6D [EPA/ALA] in ω 3 and ω 3&GSP groups related to controls. However, FADs in liver did not show differences between GSP and control groups.

3.2.3 Eicosanoid and docosanoids levels

The lipidomic analysis revealed the production of the same compounds in almost all groups as in the standard diet experiment (Tab. 7). Data showed similar tendencies in the formation of eicosanoid and docosanoids in both standard and HFHS experiments when diets were supplemented with fish oils. Briefly, in the ω 3 group compared with control, results showed: (a) a significant decrease of 12-LOX EPA product (12-HpEPE), and enhance of 15-LOX EPA product (15-HpEPE) and secondary metabolites of EPA and DHA (12-HEPE, 15-HEPE, and 17HDoHE), (b) a tendency to decrease the production of 15-LOX product of DHA (17-HpDoHE); (c) and a slightly reduction of the synthesis of secondary ARA metabolites (11-HETE and PGE₂). It is remarkable that the COX product of EPA (TBX₃) was only detected in fish oils groups.

Interestingly, the influence that polyphenols exerted on metabolites synthesis showed some differences between both standard and HFHS experiments. Specifically, the GSP group in the HFHS diet (a) did not significantly affect secondary metabolites of EPA and DHA in concordance with the standard diet. However, contrary to the standard diet

(b) polyphenols did not reduce levels of primary metabolites produced by 12-LOX and 15-LOX from EPA and DHA, and a tendency was even observed to produce higher values of 12-HpEPE and 17-HpDoHE in the HFHS diet; (c) and polyphenols significantly increased the synthesis of ARA metabolites derived from COX activity in the HFHS diet. The different influence of polyphenols between the HFHS and the standard experiment was also observed when polyphenols and fish oils were combined. (a) Levels of primary metabolites from EPA and DHA slightly increased in the ω 3&GSP group related to the ω 3 one (significant for 15-HpEPE) in the HFHS diet in agreement with the effect observed in the GSP group and suggesting a tendency to up-regulate 12-LOX and 15-LOX, contrary to results observed in the standard diet. (b) Secondary metabolites from EPA and DHA also tended to increase in the ω 3&GSP group related to the ω 3 one, contrary to the standard diet; (c) and secondary metabolites from ARA did not decrease in the ω 3&GSP group related to ω 3 as it had been found in the standard diet. Therefore, it seems that polyphenols down-regulated the activity of LOXs and COXs involved in the synthesis of these lipid mediators in the standard diet; however, they turned to up-regulate or slightly increase the activity of these enzymes in the HFHS diet. The two-ways ANOVA analysis showed significant interaction between diet (HFHS and Standard) and supplement group for several compounds due to the different influence exerted by polyphenols depending on the diet.

3.2.4 Oxidative stress parameters

As in the standard diet, levels of conjugated diene hydroperoxides and TBARS in plasma were under detection limits (data not shown). No differences were observed in conjugated diene hydroperoxides in liver between groups, and the concentration of TBARS in liver followed the same trend as in the standard diet: increased in ω 3 and ω 3&GSP groups related to control, and GSP group reduced the production of TBARS compared with control. Consequently, the combination of both supplements (ω 3&GSP group) resulted in a lower production of TBARS than ω 3 group. Data of protein carbonylation in plasma revealed an inhibition of protein oxidation in the ω 3 and GSP groups that was clearly additive in the combined ω 3&GSP group. Finally, regarding the antioxidant parameters: no significant differences were observed in ORAC although a tendency towards higher values in the ω 3&GSP group was recorded as in the standard diet; SOD showed a trend to be lower in the ω 3 group and significantly lower in GSP and ω 3&GSP groups than in control; GPX was found significantly increased in ω 3 and GSP groups, and the combined effect was also observed in ω 3&GSP compared with control. Results are shown in Tab. 6.

4. Discussion

Both ω -3 PUFAs and polyphenols are well established for their individual health benefits. However, their combined effects are not well documented [27]. In previous works, we have suggested that diets supplemented with fish oils were able to modulate inflammation, oxidative stress and CVD risk markers, especially in a balanced proportion 1:1 EPA:DHA [6,36]. Considering that an increase of oxidizable substrates in the diet like PUFAs supplement may deregulate the redox balance and increase

oxidative stress [11], polyphenols have been used in this work in combination with fish oils having a proportion 1:1 EPA:DHA to potentially protect lipids from oxidation processes [14,15] and thus, enhance their health benefits. The potential mechanisms related to the influence on the regulation of lipid profiles in plasma, tissues and membranes, and eicosanoid and docosanoids synthesis have been suggested. Lipid profiles and further metabolites from fatty acids have been closely associated with inflammation, oxidative stress and the endogenous antioxidant system. Interestingly, some differences in the functionality of studied supplements were observed depending on the diet: standard vs. HFHS.

4.1 Fish oils influence

Outcomes suggested that supplementation with fish oils lead to a general improvement of fatty acids and lipid metabolites profiling in both, standard and HFHS diets. Interestingly, the influence of fish oils in the diet was similar in both caloric contexts, standard and HFHS. The replacement of ARA with EPA and DHA in membranes, plasma and tissues in fish oil diets, with the consequent enrichment on ω -3 long chain PUFAs brings a change on the fatty acid profiling accompanied by a modulation on the activity of the LOX and COX enzymes probably due to a competence mechanism. As a result, the production of lipid mediators is being affected and the uptake of ω -3 PUFAs modulates the synthesis *de novo* of ARA through elongases and desaturases.

After digestion, some fatty acids are transported in chylomicrons and directly accumulated in the adipose tissue or incorporated into liver. *Synthesis de novo* of fatty acids occurs in the liver and fatty acids are transported again as chylomicrons in blood to be used in different processes, incorporated into membranes or stored into the adipose tissue [56]. Accordingly, it was observed in this work that the modulation of the hepatic fatty acid profile by the fish oils supplement was also reflected in the TFA fraction of plasma and then in the incorporation into erythrocyte membranes, adipose tissue and circulating FFA in plasma. Fatty acids released from the adipose tissue become circulating FFA in plasma which are reflecting the influence of the diet and synthesis *de novo* in the accumulation of fat in adipocytes. These circulating FFAs in blood are considered signaling molecules participating in oxidative stress and inflammatory processes [38].

In this work, it was observed in both dietary backgrounds that the addition of fish oils (ω 3 and ω 3&GSP groups), with the subsequent increase of EPA, DPA, and DHA proportion in the diet, was reflected in a higher accumulation of these PUFAs in liver, adipose tissue and plasma compared with control and GSP groups. Interestingly, although ARA intake was higher in the diet supplemented with fish oils, the proportion of ARA detected in all samples was reduced related to control and GSP groups. Therefore, these results confirmed the preferential substrate competition of Δ 5D, which modulates *de novo* synthesis of EPA and ARA for ω -3 PUFAs over ω -6 PUFAs [45]. In this regard, the inhibition of ARA synthesis in fish oils groups was generally accompanied by an increment of the concentration of its precursor DGLA.

1
2 In agreement with these findings, Sears *et al* [12] have also found a direct association
3 between ω -3 dietary interventions and FADs modulation towards down-production of
4 ARA and its inflammatory derivatives. Others have also found that interventions in the
5 diet with fish oils enriched FFA fraction with ω -3 PUFAs [57]. Here, it was noteworthy
6 that the analysis of the FAD indexes in plasma [44] was highly consistent with these
7 indexes calculated in liver. Both ω 3 and ω 3&GSP groups showed lower indexes of Δ 5D
8 modulating the synthesis of ARA, higher values of Δ 4D related to DHA formation, and
9 higher Δ 5/6D to produce EPA, compared with control and GSP groups. As a final
10 consequence, the ω 6/ ω 3 ratio in plasma, circulating FFAs, membranes and tissues was
11 lower in both standard and HFHS dietary contexts in the ω 3 and ω 3&GSP groups
12 compared with control groups. The ω 6/ ω 3 ratio is considered an excellent clinical
13 marker for cellular inflammation [58], and higher levels have been associated with
14 increasing prevalence of chronic inflammatory diseases [3].

15
16 Interestingly, adipose tissue showed a high enrichment in oleic acid compared with
17 liver, TFA in plasma and erythrocytes in both dietary contexts regardless of the specific
18 supplement. Accordingly, circulating fatty acids showed a high proportion of oleic acid
19 as well. However, this accumulation of oleic acid in the adipose tissue was significantly
20 reduced when fish oils were uptake. As a consequence, circulating oleic acid in plasma
21 was minor in ω 3 and ω 3&GSP groups than in control and GSP groups in both dietary
22 contexts. The increase of oleic acid in plasma and tissues has been associated with
23 obesity [59], hypertriacylglycerolemia, and the risk of developing insulin resistance
24 [37]. These conditions are usually accompanied by low-grade chronic inflammation
25 [60] and increased oxidative stress [61]. It is remarkable that HFHS diet tended to
26 accumulate higher proportions of oleic acid and lower PUFAs than the standard diet in
27 all groups. In this regard, in the HFHS context more prone to increase oleic acid in
28 adipose, fish oils were effective not only to reduce the accumulation in the adipose
29 tissue as pointed above, but ω 3 and ω 3&GSP groups also showed a decrease of the
30 desaturase index SCD-18 measured in plasma indicating a possible decrease of the
31 synthesis *de novo* mediated by the desaturase. It has been published that increased SCD-
32 18 activity was associated with insulin resistance, fatty liver and metabolic disorders
33 [62].

34
35 The replacement of ARA by EPA, DPA and DHA in erythrocyte membranes in groups
36 supplemented with fish oils was evident in both dietary experiments. Therefore
37 according with literature [45], the synthesis of membrane phospholipids demonstrated
38 to be highly modulated by a substrate competition between ω -3 EPA, DPA and DHA
39 and ω -6 ARA. Consequently, the inflammatory ω 6/ ω 3 ratio in erythrocytes composition
40 was also lower in both standard and HFHS diets supplemented with fish oils, and this
41 fact can lead the production of lipid inflammatory mediators as observed in this work.

42
43 Consequently, the analysis of lipid metabolites produced through enzymatic pathways
44 from EPA, DHA and ARA showed a strong concordance with the fatty acid

composition of erythrocytes. These PUFAs are released from membrane phospholipids into cytosol by the PLA₂ and further oxidized by enzymatic processes [63]. Lipoxygenase (LOX) enzymes regulate the oxidation of EPA, DHA, and ARA to the primary and unstable metabolites (hydroperoxides) [64–66]; glutathione peroxidases (GPX) are responsible for the detoxification of deleterious of oxygen radicals [67] and reduce hydroperoxides into secondary metabolites like hydroxides [68]; cyclooxygenases (COX) produce secondary metabolites like TXB₃ from EPA and strong pro-inflammatory compounds from ARA, 11-HETE and PGE₂ [30]. The partial replacement of ARA by EPA and DHA in membranes in ω 3 and ω 3&GSP groups was consistent with a higher production of derived metabolites from EPA and DHA than derived from ARA. *In vitro* studies showed that the specificity of 12-LOX [69] and PLA₂ [70] was comparable between EPA and ARA; however interventions in the diet with fish oils have shown to favor ω -3 PUFAs, specially EPA, over ω -6 ARA for the same phospholipase enzymes by competence mechanisms [70]. Consequently, in groups supplemented with fish oils, and with lower ω 6/ ω 3 ratios in membranes, the release of PUFAs by PLA₂ was favored towards EPA over ARA. Therefore, the production of eicosanoids was modulated towards the ω -3 hydroxides from EPA instead of the particularly pro-inflammatory ω -6 ARA metabolites, leading to a less inflammatory background compared with controls, probably by substrate competence mechanisms instead of a higher specificity of the enzymes for ω -3 PUFAs. In detail, the ω -3 metabolites have been considered weak-inflammatory compounds [32–35]; meanwhile, the corresponding ω -6 ones have been widely associated with strong pro-inflammatory activities [13,30,31,71], even 10 times more inflammatory than ω -3 derivatives [72]. Then, although both fish oils enriched diets showed higher levels of secondary ω -3 metabolites which are considered weak inflammatory compounds, at the same time there was a significant decrease of corresponding strong pro-inflammatory ω -6.

It is remarkable that the proportion EPA:DHA found in control and GSP groups in erythrocytes membranes, as in other tissues and plasma, was highly moved towards greater amounts of DHA vs. EPA. Accordingly, metabolites from DHA produced by 15-LOX (17-HpDoHE) were significantly more concentrated than EPA metabolites mediated by 15-LOX (15-HpEPE) in control groups. However, supplementation with fish oils modulated EPA:DHA towards more balanced ratios in both standard and HFHS diets. In agreement, the higher increase of EPA than DHA in membranes produced a competence for 15-LOX that preferably increased the pathway of formation of EPA metabolites instead of DHA. Additionally to the substrate competence, *in vitro* studies have shown that the production of LOX and COX metabolites from EPA is favored over DHA which hardly fits into the active catalytic site of the enzyme due to its larger spatial size [70]. It was also proved that the production of 15-LOX metabolites derived from EPA (15-HpEPE) in fish oils supplemented groups was much favored than the production of 12-LOX metabolites (12-HpEPE). Finally, COX activity on EPA was less favored than LOX activity. In fact, the production of COX derived TXB₃ was not increased related to control in the standard diet as could be expected due to the higher concentration of EPA in the membranes. To the best of our knowledge, the catalysis of

1 ARA by LOXs and COXs has been studied [73,74], but little is known about kinetic
2 rates in ω -3 substrates or comparative results *in vivo* related with the specificity of these
3 enzymes between ω -6 vs. ω -3 PUFAs, and ω -3 EPA vs. ω -3 DHA.

4
5 Interestingly, higher levels of 17HDoHE were found after supplementation with fish
6 oils compared with controls in both dietary contexts. 17HDoHE is the precursor of
7 powerful anti-inflammatory eicosanoids as protectins (e.g. PD_x) and resolvins (e.g.
8 RvD₁). These protectins and resolvins have been described as pro resolution targets in
9 several inflammatory conditions as cardiovascular diseases, dry eyes or even Alzheimer
10 [34]. Although searched RvD₁, and PD_x were not detected in plasma or were under the
11 limit of detection, other tissues like brain or adipocytes ought to be analyzed as they
12 have been described as potential staging areas of systemic cellular inflammation
13 [75,76]. Additionally, the production of 17HDoHE from DHA and 18-HEPE from EPA
14 has been associated with the suppression of TNF α , and then, with the down-regulation
15 of pro-inflammatory interleukins synthesis [77].

16
17 This balance towards a lower inflammatory situation described by fatty acid profiles and
18 lipid mediator synthesis in fish oils supplemented groups compared with controls was in
19 highly agreement to the lower values of TNF α and CRP found in liver (Tab. 6). Lower
20 values of CRP levels have been associated with general decrease of inflammation linked
21 to CVD [78]. It has been also suggested a strong association between the increase of
22 CRP and the synthesis of ARA derivates like PGE₂ and 11-HETE as biomarkers of
23 inflammation [36] as it was found in this work. These results were consistent in the
24 standard diet and in the HFHS diet targeting metabolic disorders.

25
26 A set of mechanisms might be implicated in this anti-inflammatory response that may
27 explain data obtained in this work. Firstly, at high dietary intakes, the ω -3 PUFAs
28 substrate can compete vs ω -6 PUFAs substrate for the same enzymes, as well as dilute
29 out the concentration of ω -6 in the cell membrane [25,29], thereby decreasing its
30 potential of being converted into pro-inflammatory eicosanoids as it was observed in
31 this work. Secondly, dietary nutrients like ω -3 EPA and DHA can alter the innate
32 immune system towards the enhancement of anti-inflammatory pathways, modulating
33 toll-like receptors like peroxisome proliferator-activated receptor- γ (PPAR γ) and gene
34 transcription factors like nuclear factor kB (NF-kB) [12]. In detail, ω -3 EPA and DHA
35 can inhibit the binding of saturated fats with toll-like receptors of membranes by direct
36 competition for these sites. The binding with saturated fats would activate NF-kB gene
37 transcription factor, which induces inflammatory responses through COX and cytokine
38 synthesis [79]. In concurrence, it was observed in this work that fish oils
39 supplementation down-regulated the synthesis of strong pro-inflammatory COX
40 products [3] like PGE₂, 11-HETE, and TXB₃ [80–86] compared with controls.

41
42 Finally, the anti-inflammatory response described in ω 3 and ω 3&GSP groups was
43 accompanied by an improvement of the antioxidant system in these groups. Although
44 MDA is a secondary oxidation product from PUFAs, the enhancement of TBARS in

liver in fish oils supplements was not accompanied by a deleterious of oxidative stress markers. In detail, data of conjugated dienes produced in liver did not show an increase in the $\omega 3$ and $\omega 3$ &GSP groups, on the contrary, they revealed a slightly lower production in the standard diet and no differences in the HFHS one. Additionally, data obtained for plasma antioxidant enzymes (Tab. 6) showed lower levels of SOD, and higher GPX in the $\omega 3$ and $\omega 3$ &GSP groups compared with controls in both standard and HFHS diets. Although not statistically different, plasma antioxidant capacity measured by ORAC showed higher levels in $\omega 3$ &GSP groups in both standard and HFHS diets. Carbonylation scores of plasma proteins showed a general decrease in the $\omega 3$ and $\omega 3$ &GSP groups in the HFHS diet. Decreasing protein carbonylation levels [7] and SOD activity [87], and increasing ORAC index [88] and GPX activity [89,90] have been associated with the enhancement of the antioxidant system. Accordingly, GPX is the enzyme involved in the detoxification of hydroperoxides to give hydroxides. The up-production of secondary EPA and DHA metabolites mediated by GPX in fish oils groups is in complete agreement with the significant increase observed in the activity of this enzyme in $\omega 3$ and $\omega 3$ &GSP groups related to controls. The increment of TBARS in $\omega 3$ and $\omega 3$ &GSP groups is in agreement with the high dose of PUFAs incorporated with fish oils which are considered oxidizable substrates [11]. MDA is a side PUFA product by enzymatic processes during the biosynthesis of thromboxanes and hydroxyheptadecatrienoic acid catalyzed by COX and prostacyclin hydroperoxidase, respectively [91]. COX activity on ARA was down-regulated in $\omega 3$ and $\omega 3$ &GSP groups compared with control; however TXB₃ pathway produced by COX on EPA was not reduced in the standard diet and even up-regulated in the HFHS diet. Therefore, the increasing EPA substrate of COX in fish oils may be in agreement with higher MDA levels detected. Therefore, the final outcome was a higher endogenous antioxidant activity, especially on GPX, in $\omega 3$ and $\omega 3$ &GSP groups, which influenced the transformation of primary into secondary eicosanoid and docosanoids [67] and decreased post-translational oxidative damages of proteins indicating lower oxidative stress.

4.2 Polyphenols influence

Fish oils exerted the same anti-inflammatory and anti-oxidant effects in both dietary contexts; however, polyphenols showed differences in the efficacy to reduce inflammation depending on the caloric context of the diet. On one hand, it was observed an anti-inflammatory and anti-oxidant bioactivity of polyphenols compared with the control group in the standard diet as commonly described *in vivo* [92,93]. This effect was similar to that found after intervention with fish oils and the combination of both supplements produced an additive response in most of the parameters studied. On the other hand, in the HFHS diet, polyphenols were less effective to reduce inflammation compared with control and even lead to an up-regulation of some pro-inflammatory biomarkers.

The inclusion of polyphenols did not modify composition of feeds compared with controls in both standard and HFHS diets. Accordingly, profiles of liver, adipose tissue,

erythrocyte membranes, and circulating FFA in plasma were similar between control and GSP groups as expected. However, polyphenols exerted an effect modulating desaturase activities in plasma which was not observed in controls. Such effect was consistent in both, standard and HFHS diets. $\Delta 5D$ and $\Delta 4D$ indexes showed lower values in GSP group than in controls related to the synthesis of ARA and DHA, respectively. As a consequence, the concentration of DGLA (precursor of ARA) was increased related to controls and the index $\Delta 6D$ (DGLA/LA) too. The synthesis of EPA regulated by $\Delta 5D$ and $\Delta 6D$ tended to decrease as well. Therefore, the FAD indexes suggested a decrease of $\Delta 5D$ and $\Delta 4D$ activities due to polyphenols. In this regard, Matsui *et al.* [94] have found that high-fat fed rats supplemented with polyphenols decreased the expression of genes for enzymes involved in fatty acid synthesis in liver and adipose tissue. Although the synthesis *de novo* modulated by these FAD could be affected by the presence of polyphenols, the effect was not reflected in profiles of fatty acids incorporated into phospholipid membranes, adipocytes, or circulating in plasma.

The composition of erythrocyte membranes was not affected by polyphenols, and thus, the substrate of LOX and COX in erythrocytes either. However, polyphenols seemed to modulate the pathways of formation of eicosanoid and docosanoids. Interestingly, their influence was dependent on the background of the diet. In the standard diet the GSP group generally down-regulated the production of ω -3 and ω -6 eicosanoid and docosanoids mediated by LOX, COX, and GPX activities compared with the control group. However, in the HFHS diet, polyphenols showed a tendency to increase products of LOX and GPX, and significantly enhanced the production of ARA eicosanoids mediated by COX.

Several mechanisms could be implicated in the general decrease of PUFAs metabolites production observed in the standard diet. Firstly, flavonoids and other polyphenols have been found to inhibit PLA₂ activity [17,95] which cleaves phospholipids to produce fatty acids, the substrate of LOX and COX. Other studies have suggested a direct influence of grape seed procyanidins inhibiting the activity of LOX [21] and COX [96] at transcriptional and enzyme level; and thus, polyphenols may indirectly block the synthesis of derived eicosanoids as it was observed in this work. Secondly, it was also described the influence of polyphenols up-regulating the nuclear factor-erythroid 2-related factor 2 (Nrf₂) which can activate the expression of antioxidant enzymes like GPX [97,98]. Accordingly, GPX levels were significantly increased after polyphenols interventions in this experiment (Tab. 6) which was directly associated with hydroperoxides detoxification. Polyphenols were also found to down-regulate Nf-kB [99] which activates the expression of COX-2 as explained before; and then, it would explain the decrease of COX products and the total inhibition of TXB₃ synthesis. Finally, polyphenols are well-known for their antioxidant activity [100] as scavengers of ROS [101,102], inhibitors of LDL oxidation and NADPH oxidase activation [22,88], although some outcomes are inconsistent [103,104]. These ROS are initiators of free radical oxidation mechanisms that also lead to the non-enzymatic formation of lipid eicosanoids [105]. Therefore, polyphenols could inhibit lipid peroxidation by

1 scavenging of ROS. In this regard, the possible mechanisms implicated in the down-
2 regulation of oxidative processes due to polyphenols, agree with the significant
3 enhancement of the antioxidant system (increase of GPX and decrease of SOD), the
4 amelioration of protein carbonylation and reduction of TBARS in liver in GSP group
5 compared with control.

6
7 As discussed above, the decrease of most eicosanoid and docosanoids studied is usually
8 associated with the improvement of the inflammatory conditions. Polyphenols have
9 been found to activate PPAR γ enhancing anti-inflammatory pathways [101]. The
10 combination of fish oils and polyphenols in the standard diet produced an additive effect
11 giving rise to the minor levels of pro-inflammatory ω -6 metabolites. Accordingly, GSP
12 and ω 3&GSP groups also showed lower levels of pro-inflammatory markers (TNF α and
13 CRP) (Tab. 6) compared with control group in the standard diet. In this regard, both
14 polyphenols and ω -3 PUFAs through different mechanisms can inhibit NADPH oxidase
15 activity and ROS generation, which activate NF-kB and the production of several
16 inflammatory molecules. ORAC index (Tab. 6) showed a trend to higher values when
17 fish oils and polyphenols were combined. This index measures the oxidative
18 degradation of fluorescein due to free radical generators and ROS, and higher values
19 may be correlated with lower amounts of ROS in plasma.

20
21 Curiously, in the HFHS diet, the presence of procyanidins significantly increased
22 pathways of formation of ω -6 pro-inflammatory products of COX, and tended to
23 increase the production of ω -3 eicosanoids and docosanoids from LOX and GPX,
24 contrary to the effect exerted in the standard dietary context. Thus it suggests an
25 induction to inflammatory conditions compared with controls. Taking into consideration
26 that the Δ 5D activity to rend ARA is diminished and COX activity seems to be
27 increased related to control, both findings bring a situation where the production of
28 ARA is down regulated by polyphenols together to an up-regulated specificity for ARA
29 derivatives oxidation mediated by COX. As a consequence, the HFHS context might
30 reduce the effectiveness of procyanidins to improve inflammation observed in the
31 standard diet. In concordance, the pro-inflammatory markers TNF α and CRP in the
32 HFHS experiment (Tab. 6) were not significantly improved in the GSP group as
33 observed in the standard diet. Consequently, the cooperative effect observed in the
34 ω 3&GSP group in the standard diet decreasing COX activity on ARA and inflammatory
35 parameters was not found in the HFHS diet due to the different modulation exerted by
36 polyphenols.

37
38 Other important outcome was reflected in the amount of circulating FFA. Increasing
39 values have been associated to potentiate and develop an inflammatory status [38] and
40 metabolic disorders associated with lipid overload in non-adipose tissues [106]. Levels
41 of FFA in the standard diet, decreased in GSP group compared with control, and the
42 combined action with ω 3 produced an additive decreasing effect. However, in the

1 HFHS context, values of FFA increased in the GSP group compared with control, and
2 this effect was corrected when they were combined with ω -3.

3
4 Bioactivity and mechanism related to inflammation and polyphenols were reviewed by
5 Yeyi Gu *et al* [21]. A number of *in vitro* studies have reported an inhibitory effect of
6 polyphenols on COXs and LOXs mediated eicosanoid production as we described in the
7 standard diet. The combined effect of ω -3 PUFAs and procyanidins inhibiting COX-1
8 and COX-2 activity *in vitro* was also observed [107]. In addition, animal models fed
9 standard or high fat diets demonstrated a general preventive role of polyphenols in
10 metabolic disorders. However, results on inflammatory markers in the context of
11 Metabolic Syndrome are limited, and the effects on plasma total cholesterol levels are
12 ambiguous. Moreover, recent works have described a strong activation *in vitro* [108]
13 and *in vivo* [109] of COXs catalytic activity by flavonoids. They found a dose-
14 dependent up-regulation *in vivo* of PGE₂ formation, increasing the production at lower
15 doses of polyphenols. A molecular modeling study shed light about the mechanism of
16 activation, and some flavonoids seem to act as cofactors binding tightly to the active site
17 of the COX enzymes facilitating their re-activation [110]. Jenny *et al.* [111] have also
18 found that polyphenols from white tea extracts up-regulated 15-LOX and PPAR- γ
19 signaling pathways, and correlated these findings with the induction of apoptosis in cell
20 lung culture studies. And finally, several works showed that the bioavailability of
21 polyphenols depends on the degree of polymerization, food processing, the presence of
22 other food components, and doses administrated; and then, the bioavailability of
23 different species of polyphenols may influence their bioactivity [112]. In fact, Martinez-
24 Micaelo *et al.* [107] showed specific bioactivities of procyanidins depending on the
25 molecule.

26
27 Questions remain underlying mechanisms of action of polyphenols related to
28 inflammation, and conclusions on the bioavailability and bioactivity of phenolic
29 compounds may depend on the synergistic effects between polyphenols and food
30 matrix, as fat or sugar. Therefore, we hypothesized a lower bioavailability of
31 polyphenols in the HFHS diet compared with the standard context. And then, the lower
32 uptake of some polyphenols may be associated with the up-regulation of COX-mediated
33 eicosanoids from ARA and the lower anti-inflammatory effects observed compared with
34 the standard diet.

35
36 As regards to the anti-oxidative properties of polyphenols, they seem to work in a
37 similar way in both, standard and HFHS, diets. Carbonylation scores of plasma proteins
38 highlighted the efficiency of dietary polyphenols, and especially when combined with
39 ω -3, to reduce *in vivo* damage of proteins due to oxidative stress. These data also agree
40 with the significant down-regulation of plasma SOD activity, enhancement of plasma
41 GPX and ORAC, and reduction of liver TBARS in the GSP and ω 3&GSP groups
42 compared with control (Tab. 6).

5. Conclusion

Fish oils were effective to enhance the anti-inflammatory response and exert an anti-oxidant activity regardless the fat and sucrose composition and content of the diet. Mechanisms related with substrate competition for the same enzymes were suggested to explain the modulation of lipid profiles in tissues, membranes, and plasma, and regulate the synthesis of eicosanoids and docosanoids. As a result, supplementation with fish oils led to an anti-inflammatory situation clearly associated to a lower ω -6/ ω -3 index in plasma and membranes, a lower production of ARA pro-inflammatory lipid mediators, an up-regulation of desaturases related to EPA and DHA synthesis, and a down-regulation of these desaturases to synthesize ARA. Such situation matches with a decrease of biomarkers of inflammation as TNF α and CRP. The anti-oxidant action exerted by fish oils seemed to be clearly associated to an activation of the antioxidant system, noteworthy detected on the major production of lipid mediators derived from GPX activity, and the decrease of SOD activity and protein carbonylation. However, polyphenols bioactivity resulted to be tentatively influenced by the fat and sugar content of the diet. In a standard diet they seemed to modulate enzymes towards an anti-inflammatory and anti-oxidant response, and the combination with fish oils exerted an interesting cooperative influence: down-regulating Δ 5D related with ARA synthesis, decreasing COX activity on ARA, enhancing the antioxidant enzymes, and decreasing total FFA in plasma. Similarly, the combination of both supplements also produced a significant improvement in the antioxidant balance and oxidative stress in HFHS diets. However, the efficacy of polyphenols to reduce inflammation was lower in the HFHS diet and some pro-inflammatory pathways were found even up-regulated. Therefore, fish oils seem to be the main responsible of the anti-inflammatory effects observed in the combined ω 3&GSP group in the HFHS diet. As a final outcome, the combination of both bioactive supplements can result in a significant improvement of metabolic health in both standard and HFHS diets by acting on inflammation and oxidative stress pathways.

Acknowledgements

This work was supported by the Spanish Ministry of Science and Innovation (grants AGL2013-49079-C2-1-R). The Consejo Superior de Investigaciones Científicas (CSIC) and the University of Santiago de Compostela (USC) are gratefully acknowledged for the doctoral fellowship to G. D. and postdoctoral contract to E. G. Xunta de Galicia and European Social Fund are also thankfully recognized by the financial support of the postdoctoral “Isidro Parga Pondal” contract to M. P. The authors thank Lucía Méndez for her excellent assistance and discussion of results, Pilar Comesaña and María Jesús González for her excellent technical assistance, and Eunice Molinar for her contribution to the animal experiment.

Table captions

Tab. 1: Liver TFA composition from animals supplemented with standard (ST) and high-fat high-sucrose (HFHS) diets divided in supplemented subgroups: control (C), fish oils (ω 3), grape seed proanthocyanidins (GSP), and fish oil with GSP (ω 3&GSP)..

Results are expressed as percentage of total fatty acids (mg/100mg of TFA). Values are shown as means $\pm$ SD. Values with different superscript letters in the same row indicate significant difference at $p < 0.05$ between dietary groups.

Tab. 2: Plasma TFA composition from animals supplemented with ST and HFHS diets divided in supplemented subgroups: C, $\omega 3$, GSP, and $\omega 3$ &GSP. Results are expressed as percentage of total fatty acids (mg/100mg of TFA). Values are shown as means $\pm$ SD. Values with different superscript letters in the same row indicate significant difference at $p < 0.05$ between dietary groups.

Tab. 3: FA composition of erythrocyte membranes from animals supplemented with ST and HFHS diets divided in supplemented subgroups: C, $\omega 3$, GSP, and $\omega 3$ &GSP. Results are expressed as percentage of total fatty acids (mg/100mg of TFA). Values are shown as means $\pm$ SD. Values with different superscript letters in the same row indicate significant difference at $p < 0.05$ between dietary groups.

Tab. 4: TFA composition of the adipose tissue from animals supplemented with ST and HFHS diets divided in supplemented subgroups: C, $\omega 3$, GSP, and $\omega 3$ &GSP. Results are expressed as percentage of total fatty acids (mg/100mg of TFA). Values are shown as means $\pm$ SD. Values with different superscript letters in the same row indicate significant difference at $p < 0.05$ between dietary groups.

Tab. 5: Circulating FFA composition of plasma from animals supplemented with ST and HFHS diets divided in supplemented subgroups: C, $\omega 3$, GSP, and $\omega 3$ &GSP. Results are expressed as percentage of total fatty acids (mg/100mg of TFA). Values are shown as means $\pm$ SD. Values with different superscript letters in the same row indicate significant difference at $p < 0.05$ between dietary groups.

Tab. 6: FAD indexes from TFA data of plasma and liver calculated as product/precursor ratio. Conjugate dienes, TBARS, CRP*, and TNF α * calculated in liver. Percentage of total albumin carbonylation index, ORAC*, SOD*, and GPX* activities calculated in plasma. Values are shown as means $\pm$ SD. Values with different superscript letters in the same row indicate significant difference at $p < 0.05$ between dietary groups. *These parameters have already been studied in the same animals.

Tab. 7: Levels of lipid mediators in plasma derived from ARA, EPA and DHA obtained from the dietary interventions. Results are expressed as ng/mL. Values are shown as means $\pm$ SD. One-way and two-way ANOVA analyses were done. Different superscript letters between the subgroups of the same diet indicate significant differences at $p < 0.05$ among them. Superscript * indicates significant differences between ST and HFHS diets at $p < 0.05$. Superscript # indicates significant interaction between diet (ST and HFHS) and supplement (C, $\omega 3$, GSP, $\omega 3$ &GSP) at $p < 0.05$.

Supplementary electronic material captions

S1: Diet composition and experimental design. ST (standard diet), HFHS (high-fat high-sucrose diet), C (control group); ω -3 (fish oils group); GSP (proanthocyanidins group); ω 3&GSP (fish oils and proanthocyanidins group). ^aStandard flour (Tekland Global 2014). ^bHigh fat-high sucrose flour (TD 08811). ^cGrape seed proanthocyanidin extract. ^d27% from sucrose.

S2: Fatty acid composition of supplemented groups in the standard and HFHS diets: C (control group); ω -3 (fish oils group); GSP (polyphenols group); ω 3&GSP (fish oils and polyphenols group). Results are expressed as a percentage of total fatty acids (mg/100mg of Total FA).

S3: Retention times, collision energies and MS/MS transitions for ESI-LC-LIT/MS/MS

References

- [1] Pérez-Martínez P, Pérez-Jiménez F, López-Miranda J. n-3 PUFA and lipotoxicity. *Biochim Biophys Acta-Molecular Cell Biol Lipids* 2010;1801:362–6. doi:10.1016/j.bbalip.2009.09.010.
- [2] Neilson AP, Djuric Z, Ren J, Hong YH, Sen A, Lager C, et al. Effect of cyclooxygenase genotype and dietary fish oil on colonic eicosanoids in mice. *J Nutr Biochem* 2012;23:966–76. doi:10.1016/j.jnutbio.2011.05.003.
- [3] McDaniel JC, Massey K, Nicolaou A. Fish oil supplementation alters levels of lipid mediators of inflammation in microenvironment of acute human wounds. *Wound Repair Regen* 2011;19:189–200. doi:10.1111/j.1524-475X.2010.00659.x.
- [4] Brahmabhatt V, Oliveira M, Briand M, Perrisseau G, Schmid VB, Destailats F, et al. Protective effects of dietary EPA and DHA on ischemia-reperfusion-induced intestinal stress. *J Nutr Biochem* 2013;24:104–11. doi:10.1016/j.jnutbio.2012.02.014.
- [5] Harris WS, Mozaffarian D, Lefevre M, Toner CD, Colombo J, Cunnane SC, et al. Towards Establishing Dietary Reference Intakes for Eicosapentaenoic and Docosahexaenoic Acids. *J Nutr* 2009;139:804S–819S. doi:10.3945/jn.108.101329.
- [6] Dasilva G, Pazos M, García-Egido E, Gallardo JM, Rodríguez I, Cela R, et al. Healthy effect of different proportions of marine ω -3 PUFAs EPA and DHA supplementation in Wistar rats: lipidomic biomarkers of oxidative stress and inflammation. *J Nutr Biochem* 2015;26:1385–92. doi:10.1016/j.jnutbio.2015.07.007.
- [7] Méndez L, Pazos M, Gallardo JM, Torres JL, Pérez-Jiménez J, Nogués R, et al. Reduced protein oxidation in Wistar rats supplemented with marine ω 3 PUFAs. *Free Radic Biol Med* 2013;55:8–20. doi:10.1016/j.freeradbiomed.2012.11.004.
- [8] Lluís L, Taltavull N, Muñoz-Cortés M, Sánchez-Martos V, Romeu M, Giralt M, et al. Protective effect of the omega-3 polyunsaturated fatty acids: Eicosapentaenoic acid/Docosahexaenoic acid 1:1 ratio on cardiovascular disease risk markers in rats. *Lipids Health Dis* 2013;12:1–8. doi:10.1186/1476-511X-12-140.
- [9] Molinar-Toribio E, Pérez-Jiménez J, Ramos-Romero S, Romeu M, Giralt M, Taltavull N, et al. Effect of n-3 PUFA supplementation at different EPA:DHA ratios on the spontaneously hypertensive obese rat model of the metabolic

- 1 syndrome. *Br J Nutr* 2015;113:878–87. doi:10.1017/S0007114514004437.
- 2 [10] Liu L, Hu Q, Wu H, Xue Y, Cai L, Fang M, et al. Protective role of n6/n3 PUFA
3 supplementation with varying DHA/EPA ratios against atherosclerosis in mice. *J*
4 *Nutr Biochem* 2016;32:171–80.
5 doi:http://dx.doi.org/10.1016/j.jnutbio.2016.02.010.
- 6 [11] Liu JK, Yeo HC, Overvik-Douki E, Hagen T, Doniger SJ, Chu DW, et al.
7 Chronically and acutely exercised rats: biomarkers of oxidative stress and
8 endogenous antioxidants. *J Appl Physiol* 2000;89:21–8.
- 9 [12] Sears B, Ricordi C. Role of fatty acids and polyphenols in inflammatory gene
10 transcription and their impact on obesity, metabolic syndrome and diabetes. *Eur*
11 *Rev Med Pharmacol Sci* 2012;16:1137–54.
- 12 [13] Greene ER, Huang S, Serhan CN, Panigrahy D. Regulation of inflammation in
13 cancer by eicosanoids. *Prostaglandins Other Lipid Mediat* 2011;96:27–36.
14 doi:10.1016/j.prostaglandins.2011.08.004.
- 15 [14] Gorelik S, Ligumsky M, Kohen R, Kanner J. The stomach as a “Bioreactor”:
16 When red meat meets red wine. *J Agric Food Chem* 2008;56:5002–7.
17 doi:10.1021/jf703700d.
- 18 [15] Pazos M, Lois S, Torres JL, Medina I. Inhibition of hemoglobin- and iron-
19 promoted oxidation in fish microsomes by natural phenolics. *J Agric Food Chem*
20 2006;54:4417–23. doi:10.1021/jf0530300.
- 21 [16] Terra X, Montagut G, Bustos M, Llopiz N, Ardevol A, Blade C, et al. Grape-seed
22 procyanidins prevent low-grade inflammation by modulating cytokine expression
23 in rats fed a high-fat diet. *J Nutr Biochem* 2009;20:210–8.
24 doi:10.1016/j.jnutbio.2008.02.005.
- 25 [17] Baiges I, Palmfeldt J, Blade C, Gregersen N, Arola L. Lipogenesis Is Decreased
26 by Grape Seed Proanthocyanidins According to Liver Proteomics of Rats Fed a
27 High Fat Diet. *Mol Cell Proteomics* 2010;9:1499–513.
28 doi:10.1074/mcp.M000055-MCP201.
- 29 [18] Puiggros F, Llopiz N, Ardevol A, Blade C, Arola L, Salvado MJ. Grape seed
30 procyanidins prevent oxidative injury by modulating the expression of
31 antioxidant enzyme systems. *J Agric Food Chem* 2005;53:6080–6.
32 doi:10.1021/jf050343m.
- 33 [19] Montagut G, Blade C, Blay M, Fernandez-Larrea J, Pujadas G, Josepa Salvado
34 M, et al. Effects of a grape seed procyanidin extract (GSPE) on insulin resistance.
35 *J Nutr Biochem* 2010;21:961–7. doi:10.1016/j.jnutbio.2009.08.001.
- 36 [20] Del Bas JM, Fernandez-Larrea J, Blay M, Ardevol A, Salvado MJ, Arola L, et al.
37 Grape seed procyanidins improve atherosclerotic risk index and induce liver
38 CYP7A1 and SHP expression in healthy rats. *Faseb J* 2005;19:479–+.
39 doi:10.1096/fj.04-3095fje.
- 40 [21] Yeyi G, Lambert JD. Modulation of metabolic syndrome-related inflammation by
41 cocoa. *Mol Nutr Food Res* 2013;57:948–61.
- 42 [22] Pignatelli P, Ghiselli A, Buchetti B, Carnevale R, Natella F, Germanò G, et al.
43 Polyphenols synergistically inhibit oxidative stress in subjects given red and
44 white wine. *Atherosclerosis* 2006;188:77–83.
45 doi:10.1016/j.atherosclerosis.2005.10.025.
- 46 [23] Cave AC, Brewer AC, Narayanapanicker A, Ray R, Grieve DJ, Walker S, et al.
47 NADPH oxidases in cardiovascular health and disease. *Antioxid Redox Signal*
48 2006;8:691–728.
- 49 [24] Carnevale R, Loffredo L, Pignatelli P, Nocella C, Bartimoccia S, Di Santo S, et
50 al. Dark chocolate inhibits platelet isoprostanes via NOX2 down-regulation in

- 1 smokers. *J Thromb Haemost* 2012;10:125–32. doi:10.1111/j.1538-
- 2 7836.2011.04558.x.
- 3 [25] Massaro M, Scoditti E, Carluccio MA, De Caterina R. Basic mechanisms behind
- 4 the effects of n-3 fatty acids on cardiovascular disease. *Prostaglandins Leukot*
- 5 *Essent Fat Acids* 2008;79:109–15. doi:10.1016/j.plefa.2008.09.009.
- 6 [26] Maestre R, Douglass JD, Kodukula S, Medina I, Storch J. Alterations in the
- 7 Intestinal Assimilation of Oxidized PUFAs Are Ameliorated by a Polyphenol-
- 8 Rich Grape Seed Extract in an In Vitro Model and Caco-2 Cells. *J Nutr*
- 9 2013;143:295–301. doi:10.3945/jn.112.160101.
- 10 [27] Sekhon-Loodu S, Catalli A, Kulka M, Wang Y, Shahidi F, Rupasinghe HPV.
- 11 Apple flavonols and n-3 polyunsaturated fatty acid-rich fish oil lowers blood C-
- 12 reactive protein in rats with hypercholesterolemia and acute inflammation. *Nutr*
- 13 *Res* 2014;34:535–43. doi:10.1016/j.nutres.2014.05.002.
- 14 [28] Zhang XC, Dong F, Ren J, Driscoll MJ, Culver B. High dietary fat induces
- 15 NADPH oxidase-associated oxidative stress and inflammation in rat cerebral
- 16 cortex. *Exp Neurol* 2005;191:318–25. doi:10.1016/j.expneurol.2004.10.011.
- 17 [29] Calder PC. The role of marine omega-3 (n-3) fatty acids in inflammatory
- 18 processes, atherosclerosis and plaque stability. *Mol Nutr Food Res*
- 19 2012;56:1073–80.
- 20 [30] Massey KA, Nicolaou A. Lipidomics of oxidized polyunsaturated fatty acids.
- 21 *Free Radic Bio Med* 2013;59:45–55. doi:10.1016/j.freeradbiomed.2012.08.565.
- 22 [31] Gao L, Yin HY, Milne GL, Porter NA, Morrow JD. Formation of F-ring
- 23 isoprostane-like compounds (F-3-isoprostanes) in vivo from eicosapentaenoic
- 24 acid. *J Biol Chem* 2006;281:14092–9. doi:10.1074/jbc.M601035200.
- 25 [32] Waddington E, Sienuaraine K, Puddey I, Croft K. Identification and quantitation
- 26 of unique fatty acid oxidation products in human atherosclerotic plaque using
- 27 high-performance liquid chromatography. *Anal Biochem* 2001;292:234–44.
- 28 doi:10.1006/abio.2001.5075.
- 29 [33] Kelly L, Grehan B, Della Chiesa A, O'Mara SM, Downer E, Sahyoun G, et al.
- 30 The polyunsaturated fatty acids, EPA and DPA exert a protective effect in the
- 31 hippocampus of the aged rat. *Neurobiol Aging* 2011;32:2318–33.
- 32 doi:10.1016/j.neurobiolaging.2010.04.001.
- 33 [34] Lee CH. Resolvins as New Fascinating Drug Candidates for Inflammatory
- 34 Diseases. *Arch Pharm Res* 2012;35:3–7. doi:10.1007/s12272-012-0121-z.
- 35 [35] Serhan CN, Gotlinger K, Hong S, Lu Y, Siegelman J, Baer T, et al. Anti-
- 36 inflammatory actions of neuroprotectin D1/protectin D1 and its natural
- 37 stereoisomers: Assignments of dihydroxy-containing docosatrienes. *J Immunol*
- 38 2006;176:1848–59.
- 39 [36] Dasilva G, Pazos M, García-Egido E, Pérez-Jiménez J, Torres JL, Giralt M, et al.
- 40 Lipidomics to analyze the influence of diets with different EPA:DHA ratios in
- 41 the progression of Metabolic Syndrome using SHROB rats as a model. *Food*
- 42 *Chem* 2016;205:196–203. doi:http://dx.doi.org/10.1016/j.foodchem.2016.03.020.
- 43 [37] Vessby B, Gustafsson I-B, Tengblad S, Berglund L. Indices of fatty acid
- 44 desaturase activity in healthy human subjects: effects of different types of dietary
- 45 fat. *Br J Nutr* 2013;110:871–9. doi:10.1017/S0007114512005934.
- 46 [38] Martins AR, Nachbar RT, Gorjao R, Vinolo MA, Festuccia WT, Lambertucci
- 47 RH, et al. Mechanisms underlying skeletal muscle insulin resistance induced by
- 48 fatty acids: importance of the mitochondrial function. *Lipids Health Dis*
- 49 2012;11:1–11. doi:10.1186/1476-511X-11-30.
- 50 [39] Tsukamoto T, Sonenberg M. Catecholamine Regulation of Human-Erythrocyte

- 1 Membrane-Protein Kinase. *J Clin Invest* 1979;64:534–40.
2 doi:10.1172/JCI109491.
- 3 [40] Puttmann M, Krug H, Vonochsenstein E, Kattermann R. Fast Hplc
4 Determination of Serum-Free Fatty-Acids in the Picomole Range. *Clin Chem*
5 1993;39:825–32.
- 6 [41] Lepage G, Roy CC. Direct Transesterification of all Classes of Lipids in a One-
7 Step Reaction. *J Lipid Res* 1986;27:114–20.
- 8 [42] Kaluzny MA, Duncan LA, Merritt M V, Epps DE. Rapid Separation of Lipid
9 Classes in High-Yield and Purity using Bonded Phase Columns. *J Lipid Res*
10 1985;26:135–40.
- 11 [43] Bligh EG, Dyer WJ. A Rapid Method of Total Lipid Extraction and Purification.
12 *Can J Biochem Physiol* 1959;37:911–7.
- 13 [44] Warensjo E, Rosell M, Hellenius M-L, Vessby B, De Faire U, Riserus U.
14 Associations between estimated fatty acid desaturase activities in serum lipids
15 and adipose tissue in humans: links to obesity and insulin resistance. *Lipids*
16 *Health Dis* 2009;8:37. doi:10.1186/1476-511X-8-37.
- 17 [45] Ferguson LR. Nutrigenomics and nutrigenetics in functional foods and
18 personalized nutrition. 2014.
- 19 [46] Herbes SE, Allen CP. Lipid Quantification of Fresh-Water Invertebrates -
20 Method Modification for Microquantitation. *Can J Fish Aquat Sci* 1983;40:1315–
21 7.
- 22 [47] AOCS. Official and Tentative Methods of the American-Oil-Chemists-Society.
23 *Analyst* 1947;72:157.
- 24 [48] Mendes R, Cardoso C, Pestana C. Measurement of malondialdehyde in fish: A
25 comparison study between HPLC methods and the traditional spectrophotometric
26 test. *Food Chem* 2009;112:1038–45.
27 doi:http://dx.doi.org/10.1016/j.foodchem.2008.06.052.
- 28 [49] Mateos R, Lecumberri E, Ramos S, Goya L, Bravo L. Determination of
29 malondialdehyde (MDA) by high-performance liquid chromatography in serum
30 and liver as a biomarker for oxidative stress: Application to a rat model for
31 hypercholesterolemia and evaluation of the effect of diets rich in phenolic
32 antioxidant. *Anal Antioxidants Biomarkers Oxidative Stress* 2005;827:76–82.
33 doi:http://dx.doi.org/10.1016/j.jchromb.2005.06.035.
- 34 [50] Fukunaga K, Suzuki T, Takama K. Highly sensitive high-performance liquid
35 chromatography for the measurement of malondialdehyde in biological samples.
36 *J Chromatogr B Biomed Sci Appl* 1993;621:77–81.
37 doi:http://dx.doi.org/10.1016/0378-4347(93)80078-I.
- 38 [51] Cao GH, Booth SL, Sadowski JA, Prior RL. Increases in human plasma
39 antioxidant capacity after consumption of controlled diets high in fruit and
40 vegetables. *Am J Clin Nutr* 1998;68:1081–7.
- 41 [52] Misra HP, Fridovic.i. Role of Superoxide Anion in Autoxidation of Epinephrine
42 and a Simple Assay for Superoxide Dismutase. *J Biol Chem* 1972;247:3170-.
- 43 [53] Wheeler CR, Salzman JA, Elsayed NM, Omaye ST, Korte DW. Automated
44 Assays for Superoxide-Dismutase, Catalase, Glutathione-Peroxidase, and
45 Glutathione-Reductase Activity. *Anal Biochem* 1990;184:193–9.
46 doi:10.1016/0003-2697(90)90668-Y.
- 47 [54] Dasilva G, Pazos M, Gallardo JM, Rodríguez I, Cela R, Medina I. Lipidomic
48 analysis of polyunsaturated fatty acids and their oxygenated metabolites in
49 plasma by solid-phase extraction followed by LC-MS. *Anal Bioanal Chem*
50 2014;406:2827–39. doi:10.1007/s00216-014-7701-3.

- [55] Méndez L, Pazos M, Molinar-Toribio E, Sanchez-Martos V, Gallardo JM, Rosa Nogués M, et al. Protein carbonylation associated to high-fat, high-sucrose diet and its metabolic effects. *J Nutr Biochem* 2014;25:1243–53. doi:10.1016/j.jnutbio.2014.06.014.
- [56] Nelson DL, Michael MC. *Principles of Biochemistry*. vol. 4. New York: W. H. Freeman,; 2004.
- [57] Oh DY, Talukdar S, Bae EJ, Imamura T, Morinaga H, Fan W, et al. GPR120 Is an Omega-3 Fatty Acid Receptor Mediating Potent Anti-inflammatory and Insulin-Sensitizing Effects. *Cell* 2010;142:687–98. doi:10.1016/j.cell.2010.07.041.
- [58] Sears B. *The Anti-inflammation Zone*. New York: Regan Books; 2005.
- [59] Mutch DM, O'Maille G, Wikoﬀ WR, Wiedmer T, Sims PJ, Siuzdak G. Mobilization of pro-inflammatory lipids in obese Plscr3-deficient mice. *Genome Biol* 2007;8:R38–R38. doi:10.1186/gb-2007-8-3-r38.
- [60] Festa A, D'Agostino R, Howard G, Mykkanen L, Tracy RP, Haffner SM. Chronic subclinical inflammation as part of the insulin resistance syndrome - The Insulin Resistance Atherosclerosis Study (IRAS). *Circulation* 2000;102:42–7.
- [61] Armutcu F, Ataymen M, Atmaca H, Gurel A. Oxidative stress markers, C-reactive protein and heat shock protein 70 levels in subjects with metabolic syndrome. *Clin Chem Lab Med* 2008;46:785–90. doi:10.1515/CCLM.2008.166.
- [62] Kazdova L. Relationship of Circulating Fatty Acid Profile to Metabolic Disorders Associated with Insulin Resistance. *Ebiomedicine* 2015;2:1308–9. doi:10.1016/j.ebiom.2015.09.052.
- [63] Balvers MGJ, Verhoeckx KCM, Meijerink J, Bijlsma S, Rubingh CM, Wortelboer HM, et al. Time-dependent effect of in vivo inflammation on eicosanoid and endocannabinoid levels in plasma, liver, ileum and adipose tissue in C57BL/6 mice fed a fish-oil diet. *Int Immunopharmacol* 2012;13:204–14. doi:10.1016/j.intimp.2012.03.022.
- [64] Yin H, Brooks JD, Gao L, Porter NA, Morrow JD. Identification of novel autoxidation products of the omega-3 fatty acid eicosapentaenoic acid in vitro and in vivo. *J Biol Chem* 2007;282:29890–901. doi:10.1074/jbc.M703108200.
- [65] Masoodi M, Mir AA, Petasis NA, Serhan CN, Nicolaou A. Simultaneous lipidomic analysis of three families of bioactive lipid mediators leukotrienes, resolvins, protectins and related hydroxy-fatty acids by liquid chromatography/electrospray ionisation tandem mass spectrometry. *Rapid Commun Mass Spectrom* 2008;22:75–83. doi:10.1002/rcm.3331.
- [66] Serhan CN. Lipoxin Biosynthesis and its Impact in Inflammatory and Vascular Events. *Biochim Biophys Acta-Lipids Lipid Metab* 1994;1212:1–25. doi:10.1016/0005-2760(94)90185-6.
- [67] Biesalski HK. The role of antioxidants in nutritional support. *Nutrition* 2000;16:593–6. doi:10.1016/S0899-9007(00)00316-6.
- [68] Hunt J V, Smith CCT, Wolff SP. Autoxidative Glycosylation and Possible Involvement of Peroxides and Free-Radicals in Ldl Modification by Glucose. *Diabetes* 1990;39:1420–4. doi:10.2337/diabetes.39.11.1420.
- [69] Ikei KN, Yeung J, Apopa PL, Ceja J, Vesci J, Holman TR, et al. Investigations of human platelet-type 12-lipoxygenase: role of lipoxygenase products in platelet activation. *J Lipid Res* 2012;53:2546–59. doi:10.1194/jlr.M026385.
- [70] Swetland LF. *The Zone* - Sears,b, Lawren,b. *Libr J* 1995;120:113.
- [71] Ferreiro-Vera C, Maria Mata-Granados J, Priego-Capote F, Manuel Quesada-Gomez J, Dolores Luque de Castro M. Automated targeting analysis of

- eicosanoid inflammation biomarkers in human serum and in the exometabolome of stem cells by SPE-LC-MS/MS. *Anal Bioanal Chem* 2011;399:1093–103. doi:10.1007/s00216-010-4400-6.
- [72] Mathews CK, Van Holde KE, Ahern KG. Chapter 19: Lipidic metabolism II: membrane lipids, steroids, isoprenoids and eicosanoids. In: Capella I, editor. *Biochemistry*, vol. 3^a, Oregon: Adison Wesley; 2003.
- [73] Segraves EN, Holman TR. Kinetic investigations of the rate-limiting step in human 12-and 15-lipoxygenase. *Biochemistry* 2003;42:5236–43. doi:10.1021/bi0273462.
- [74] Wecksler AT, Jacquot C, van der Donk WA, Holman TR. Mechanistic Investigations of Human Reticulocyte 15-and Platelet 12-Lipoxygenases with Arachidonic Acid. *Biochemistry* 2009;48:6259–67. doi:10.1021/bi802332j.
- [75] Filep JG. Resolution pathways in inflammation: the devil in the adipose tissues and in the details. Focus on “Diversity of lipid mediators in human adipose tissue depots.” *Am J Physiol Physiol* 2013;304:C1127–8. doi:10.1152/ajpcell.00063.2013.
- [76] Claria J, Nguyen BT, Madenci AL, Ozaki CK, Serhan CN. Diversity of lipid mediators in human adipose tissue depots. *Am J Physiol Physiol* 2013;304:C1141–9. doi:10.1152/ajpcell.00351.2012.
- [77] Weylandt KH, Krause LF, Gomolka B, Chiu C-Y, Bilal S, Nadolny A, et al. Suppressed liver tumorigenesis in fat-1 mice with elevated omega-3 fatty acids is associated with increased omega-3 derived lipid mediators and reduced TNF-alpha. *Carcinogenesis* 2011;32:897–903.
- [78] Casas JP, Shah T, Hingorani AD, Danesh J, Pepys MB. C-reactive protein and coronary heart disease: a critical review. *J Intern Med* 2008;264:295–314. doi:10.1111/j.1365-2796.2008.02015.x.
- [79] Bouwens M, van de Rest O, Dellschaft N, Bromhaar MG, de Groot LCPGM, Geleijnse JM, et al. Fish-oil supplementation induces antiinflammatory gene expression profiles in human blood mononuclear cells. *Am J Clin Nutr* 2009;90:415–24. doi:10.3945/ajcn.2009.27680.
- [80] Birklein F. Complex regional pain syndrome. *J Neurol* 2005;252:131–8. doi:10.1007/s00415-005-0737-8.
- [81] Davi G, Alessandrini P, Mezzetti A, Minotti G, Bucciarelli T, Costantini F, et al. In vivo formation of 8-epi-prostaglandin F-2 alpha is increased in hypercholesterolemia. *Arterioscler Thromb Vasc Biol* 1997;17:3230–5.
- [82] Pratico D, Iuliano L, Basili S, Ferro D, Camastra C, Cordova C, et al. Enhanced lipid peroxidation in hepatic cirrhosis. *J Investig Med* 1998;46:51–7.
- [83] Gopaul NK, Anggard EE, Mallet AI, Betteridge DJ, Wolff SP, Nourooz-Zadeh J. Plasma 8-Epi-Pgf(2-Alpha) Levels are Elevated in Individuals with Non-Insulin-Dependent Diabetes-Mellitus. *FEBS Lett* 1995;368:225–9. doi:10.1016/0014-5793(95)00649-T.
- [84] Halliwell B. Establishing the significance and optimal intake of dietary antioxidants: The biomarker concept. *Nutr Rev* 1999;57:104–13.
- [85] Ziboh VA. Special issue - Essential fatty acids/prostanoids/lipoxygenase-derived monohydroxy fatty acids in skin health and disease - Foreword. *Prostaglandins Other Lipid Mediat* 2000;63:1–2. doi:10.1016/S0090-6980(00)00092-7.
- [86] Masoodi M, Nicolaou A. Lipidomic analysis of twenty-seven prostanoids and isoprostanes by liquid chromatography/electrospray tandem mass spectrometry. *Rapid Commun Mass Spectrom* 2006;20:3023–9. doi:10.1002/rcm.2697.
- [87] Garrel C, Alessandri J-M, Guesnet P, Al-Gubory KH. Omega-3 fatty acids

- 1 enhance mitochondrial superoxide dismutase activity in rat organs during post-
2 natal development. *Int J Biochem Cell Biol* 2012;44:123–31.
3 doi:10.1016/j.biocel.2011.10.007.
- 4 [88] Gresele P, Pignatelli P, Guglielmini G, Carnevale R, Mezzasoma AM, Ghiselli
5 A, et al. Resveratrol, at concentrations attainable with moderate wine
6 consumption, stimulates human platelet nitric oxide production. *J Nutr*
7 2008;138:1602–8.
- 8 [89] Chapman C, Morgan LM, Murphy MC. Maternal and early dietary fatty acid
9 intake: Changes in lipid metabolism and liver enzymes in adult rats. *J Nutr*
10 2000;130:146–51.
- 11 [90] Yessoufou A, Soulaïmann N, Merzouk SA, Moutairou K, Ahissou H, Prost J, et
12 al. N-3 Fatty acids modulate antioxidant status in diabetic rats and their
13 macrosomic offspring. *Int J Obes* 2006;30:739–50. doi:10.1038/sj.ijo.0803211.
- 14 [91] Hornstra G. Dietary fats, prostanoids and arterial thrombosis. vol. Martinus N.
15 Maastricht, Netherlands.: 1983. doi:90-247-2667-0.
- 16 [92] Fiesel A, Gessner DK, Most E, Eder K. Effects of dietary polyphenol-rich plant
17 products from grape or hop on pro-inflammatory gene expression in the intestine,
18 nutrient digestibility and faecal microbiota of weaned pigs. *Bmc Vet Res*
19 2014;10:196. doi:10.1186/s12917-014-0196-5.
- 20 [93] Niu L, He X, Wang Q, Fu M, Xu F, Xue Y, et al. Polyphenols in Regulation of
21 Redox Signaling and Inflammation During Cardiovascular Diseases. *Cell*
22 *Biochem Biophys* 2015;72:485–94. doi:10.1007/s12013-014-0492-5.
- 23 [94] Matsui N, Ito R, Nishimura E, Yoshikawa M, Kato M, Kamei M, et al. Ingested
24 cocoa can prevent high-fat diet-induced obesity by regulating the expression of
25 genes for fatty acid metabolism. *Nutrition* 2005;21:594–601.
26 doi:10.1016/j.nut.2004.10.008.
- 27 [95] Lambert JD, Rice JE, Hong JG, Hou Z, Yang CS. Synthesis and biological
28 activity of the tea catechin metabolites, M4 and M6 and their methoxy-
29 derivatives. *Bioorg Med Chem Lett* 2005;15:873–6.
30 doi:10.1016/j.bmcl.2004.12.070.
- 31 [96] Yoon JH, Baek SJ. Molecular targets of dietary polyphenols with anti-
32 inflammatory properties. *Yonsei Med J* 2005;46:585–96.
- 33 [97] Kluth D, Banning A, Paur I, Blomhoff R, Brigelius-Flohe R. Modulation of
34 pregnane X receptor-and electrophile responsive element-mediated gene
35 expression by dietary polyphenolic compounds. *Free Radic Biol Med*
36 2007;42:315–25. doi:10.1016/j.freeradbiomed.2006.09.028.
- 37 [98] Patel R, Maru G. Polymeric black tea polyphenols induce phase II enzymes via
38 Nrf2 in mouse liver and lungs. *Free Radic Biol Med* 2008;44:1897–911.
39 doi:10.1016/j.freeradbiomed.2008.02.006.
- 40 [99] Banerjee S, Bueso-Ramos C, Aggarwal BB. Suppression of 7,12-
41 dimethylbenz(a)anthracene-induced mammary carcinogenesis in rats by
42 resveratrol: Role of nuclear factor-kappa B, cyclooxygenase 2, and matrix
43 metalloprotease 9. *Cancer Res* 2002;62:4945–54.
- 44 [100] Cazzola R, Cestaro B. Red wine polyphenols protect n-3 more than n-6
45 polyunsaturated fatty acid from lipid peroxidation. *Food Res Int* 2011;44:3065–
46 71. doi:10.1016/j.foodres.2011.07.029.
- 47 [101] Mitjavila MT, Moreno JJ. The effects of polyphenols on oxidative stress and the
48 arachidonic acid cascade. Implications for the prevention/treatment of high
49 prevalence diseases. *Biochem Pharmacol* 2012;84:1113–22.
50 doi:10.1016/j.bcp.2012.07.017.

- 1 [102] Koutelidakis AE, Argiri K, Serafini M, Proestos C, Komaitis M, Pecorari M, et
2 al. Green tea, white tea, and Pelargonium purpureum increase the antioxidant
3 capacity of plasma and some organs in mice. *Nutrition* 2009;25:453–8.
4 doi:10.1016/j.nut.2008.10.007.
- 5 [103] Chu KO, Chan S-O, Pang CP, Wang CC. Pro-oxidative and Antioxidative
6 Controls and Signaling Modification of Polyphenolic Phytochemicals:
7 Contribution to Health Promotion and Disease Prevention? *J Agric Food Chem*
8 2014;62:4026–38. doi:10.1021/jf500080z.
- 9 [104] Lambert JD, Elias RJ. The antioxidant and pro-oxidant activities of green tea
10 polyphenols: A role in cancer prevention. *Arch Biochem Biophys* 2010;501:65–
11 72. doi:10.1016/j.abb.2010.06.013.
- 12 [105] Frankel EN. *Lipid Oxidation*. West Ferry, Dundee, Scotland: The Oily Press
13 LTD; 1998.
- 14 [106] Lewis GF, Carpentier A, Adeli K, Giacca A. Disordered fat storage and
15 mobilization in the pathogenesis of insulin resistance and type 2 diabetes. *Endocr*
16 *Rev* 2002;23:201–29. doi:10.1210/er.23.2.201.
- 17 [107] Martinez-Micaelo N, Gonzalez-Abuin N, Terra X, Richart C, Ardevol A, Pinent
18 M, et al. Omega-3 docosaehaenoic acid and procyanidins inhibit cyclo-
19 oxygenase activity and attenuate NF-kappa B activation through a p105/p50
20 regulatory mechanism in macrophage inflammation. *Biochem J* 2012;441:653–
21 63. doi:10.1042/bj20110967.
- 22 [108] Bai HW, Zhu BT. Strong activation of cyclooxygenase I and II catalytic activity
23 by dietary bioflavonoids. *J Lipid Res* 2008;49:2557–70.
24 doi:10.1194/jlr.M800358-JLR200.
- 25 [109] Bai H-W, Zhu BT. Myricetin and quercetin are naturally occurring co-substrates
26 of cyclooxygenases in vivo. *Prostaglandins Leukot Essent Fat Acids* 2010;82:45–
27 50. doi:10.1016/j.plefa.2009.10.006.
- 28 [110] Pan W, Hyoun-woo B, Bao Ting Z. Structural basis for certain naturally
29 occurring bioflavonoids to function as reducing co-substrates of cyclooxygenase
30 I and II. *PLoS One* 2010;5:e12316 [11pp.]-e12316 [11pp.].
- 31 [111] Mao JT, Nie W-X, Tsu I-H, Jin Y-S, Rao JY, Lu Q-Y, et al. White Tea Extract
32 Induces Apoptosis in Non-Small Cell Lung Cancer Cells: the Role of Peroxisome
33 Proliferator-Activated Receptor-gamma and 15-Lipoxygenases. *Cancer Prev Res*
34 2010;3:1132–40. doi:10.1158/1940-6207.CAPR-09-0264.
- 35 [112] D'Archivio M, Filesi C, Vari R, Scazzocchio B, Masella R. Bioavailability of the
36 Polyphenols: Status and Controversies. *Int J Mol Sci* 2010;11:1321–42.
37 doi:10.3390/ijms11041321.
- 38

	ST DIET								HFHS DIET							
	C		$\omega 3$		GSP		$\omega 3$ &GSP		C		$\omega 3$		GSP		$\omega 3$ &GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
14:0	0,2 ^a	0,02	0,2 ^{ab}	0,02	0,2 ^{ab}	0,01	0,2 ^b	0,01	0,5 ^a	0,03	0,7 ^b	0,004	0,5 ^a	0,10	0,7 ^{ab}	0,14
15:0	0,2 ^a	0,03	0,1 ^a	0,02	0,1 ^a	0,03	0,1 ^a	0,01	0,3 ^a	0,02	0,3 ^b	0,005	0,3 ^a	0,02	0,4 ^b	0,01
16:0	18,5 ^a	0,30	19,4 ^a	0,35	19,3 ^a	0,43	18,3 ^a	0,80	20,4 ^a	0,75	22,6 ^b	0,32	19,9 ^a	0,68	22,8 ^b	0,32
16:1 ω 7	0,6 ^a	0,08	0,7 ^a	0,07	0,8 ^a	0,23	0,7 ^a	0,04	1,0 ^a	0,15	1,1 ^a	0,12	0,8 ^a	0,14	1,0 ^a	0,18
17:0	0,6 ^a	0,06	0,5 ^a	0,07	0,5 ^a	0,10	0,5 ^a	0,10	0,6 ^a	0,03	0,5 ^a	0,02	0,6 ^a	0,03	0,5 ^a	0,01
17:1 ω 7	0,2 ^a	0,02	0,1 ^b	0,00	0,2 ^a	0,01	0,1 ^b	0,02	0,2 ^a	0,02	0,2 ^a	0,01	0,2 ^a	0,01	0,1 ^b	0,01
18:0	18,8 ^a	0,34	19,6 ^a	1,59	17,6 ^a	0,51	19,3 ^a	0,57	16,6 ^a	1,13	13,6 ^b	0,11	17,9 ^a	0,93	12,6 ^b	0,79
18:1 ω 9	6,7 ^a	0,38	4,9 ^b	0,29	7,2 ^a	1,09	5,1 ^b	0,70	14,3 ^a	1,10	13,0 ^{ab}	0,02	12,6 ^{ab}	1,51	12,9 ^b	0,11
18:1 ω 7	1,9 ^a	0,02	1,7 ^a	0,05	1,9 ^a	0,09	1,6 ^a	0,11	1,5 ^a	0,03	1,4 ^a	0,03	1,6 ^a	0,09	1,3 ^a	0,08
18:2 ω 6	16,5 ^a	0,27	17,1 ^a	0,06	17,1 ^a	2,14	17,4 ^a	0,67	15,4 ^a	0,79	13,7 ^a	0,08	14,7 ^a	0,37	14,3 ^a	0,46
18:3 ω 6	0,3 ^a	0,06	0,1 ^b	0,01	0,3 ^a	0,01	0,1 ^b	0,02	0,3 ^a	0,04	0,1 ^b	0,02	0,2 ^a	0,01	0,1 ^b	0,01
18:3 ω 3	0,4 ^a	0,02	0,3 ^b	0,01	0,5 ^a	0,12	0,3 ^b	0,02	0,6 ^{ab}	0,06	0,7 ^a	0,04	0,5 ^b	0,05	0,7 ^a	0,07
20:1 ω 9	0,1 ^a	0,003	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01	0,3 ^a	0,05	0,4 ^a	0,01	0,3 ^a	0,03	0,4 ^a	0,01
20:2 ω 6	0,3 ^a	0,01	0,2 ^b	0,02	0,3 ^a	0,03	0,2 ^b	0,01	0,1 ^a	0,01	0,1 ^b	0,02	0,2 ^a	0,02	0,1 ^b	0,01
20:3 ω 6	0,2 ^a	0,01	0,4 ^{bc}	0,06	0,2 ^{ab}	0,03	0,5 ^c	0,15	0,3 ^a	0,06	0,3 ^a	0,02	0,3 ^a	0,003	0,2 ^a	0,03
20:4 ω 6	21,3 ^a	0,29	11,8 ^b	0,69	20,4 ^a	0,68	12,1 ^b	0,21	17,9 ^a	1,08	7,9 ^b	0,15	19,6 ^a	1,33	7,6 ^b	0,26
20:5 ω 3	0,3 ^a	0,04	4,2 ^b	0,49	0,3 ^a	0,04	3,8 ^b	0,23	0,5 ^a	0,08	4,3 ^b	0,51	0,4 ^a	0,04	4,0 ^b	0,36
24:0	0,1 ^a	0,02	0,1 ^a	0,01	0,2 ^b	0,02	0,2 ^b	0,01	0,1 ^a	0,01	0,1 ^a	0,01	0,2 ^b	0,03	0,2 ^b	0,005
24:1 ω 11	0,5 ^a	0,05	0,1 ^b	0,01	0,6 ^a	0,11	0,1 ^b	0,01	0,3 ^a	0,03	0,2 ^b	0,03	0,3 ^a	0,04	0,2 ^b	0,02
22:5 ω 3	1,5 ^a	0,18	1,7 ^a	0,04	1,8 ^a	0,23	1,7 ^a	0,13	1,8 ^a	0,06	2,6 ^b	0,19	1,8 ^a	0,16	2,7 ^b	0,21
22:6 ω 3	10,8 ^a	0,05	18,1 ^b	0,48	9,0 ^a	0,21	17,2 ^b	1,33	6,9 ^a	0,30	16,1 ^b	0,45	7,1 ^a	0,36	17,0 ^b	0,93
$\omega 6/\omega 3$	3,0 ^a	0,1	1,2 ^b	0,03	3,4 ^a	0,03	1,3 ^b	0,1	3,5 ^a	0,1	0,9 ^b	0,01	3,5 ^a	0,1	0,9 ^b	0,01
$\Sigma \omega 3$	12,9 ^a	0,3	24,3 ^b	0,1	11,6 ^a	0,2	23,1 ^b	1,6	9,7 ^a	0,3	23,7 ^b	0,3	9,9 ^a	0,5	24,5 ^b	0,3
$\Sigma \omega 6$	38,6 ^a	0,5	29,6 ^b	0,8	39,3 ^a	0,3	30,3 ^b	0,9	34,1 ^a	0,7	22,0 ^b	0,1	35,0 ^a	1,1	22,3 ^b	0,2
Σ SFA	38,4 ^a	0,1	40,0 ^a	1,4	38,3 ^a	0,5	38,8 ^a	0,2	38,4 ^a	0,4	37,9 ^a	0,5	39,5 ^a	0,3	37,2 ^a	0,4
Σ MUFA	10,0 ^a	0,5	7,6 ^b	0,4	10,9 ^a	1,3	7,8 ^b	0,9	17,8 ^a	1,3	16,4 ^b	0,1	15,7 ^{ab}	1,7	16,0 ^b	0,3
Σ PUFA	51,5 ^a	0,6	53,9 ^b	0,8	50,9 ^a	0,5	53,5 ^b	0,8	43,8 ^a	1,0	45,8 ^b	0,4	44,8 ^{ab}	1,6	46,8 ^b	0,6

	ST DIET								HFHS DIET							
	C		ω3		GSP		ω3&GSP		C		ω3		GSP		ω3&GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
14:0	0,4 ^a	0,03	0,5 ^a	0,04	0,4 ^a	0,04	0,4 ^a	0,03	0,9 ^a	0,14	1,1 ^a	0,10	1,0 ^a	0,14	1,1 ^a	0,12
15:0	0,3 ^a	0,03	0,3 ^a	0,07	0,3 ^a	0,03	0,3 ^a	0,05	0,4 ^a	0,03	0,5 ^b	0,05	0,5 ^{ab}	0,04	0,5 ^a	0,04
16:0	21,2 ^a	0,5	23,7 ^b	0,83	21,7 ^{ab}	0,62	21,6 ^{ab}	1,12	21,6 ^a	0,93	25,3 ^b	1,74	23,2 ^{ab}	0,72	22,8 ^{ab}	0,65
16:1ω7	1,0 ^a	0,1	1,1 ^a	0,18	1,0 ^a	0,26	1,2 ^a	0,23	1,0 ^a	0,18	1,3 ^a	0,11	1,2 ^a	0,08	1,2 ^a	0,20
17:0	0,5 ^a	0,04	0,4 ^a	0,05	0,4 ^a	0,04	0,4 ^a	0,03	0,5 ^a	0,01	0,5 ^a	0,03	0,4 ^a	0,05	0,4 ^a	0,04
18:0	15,8 ^a	1,2	14,5 ^a	1,14	13,3 ^a	0,98	13,5 ^a	1,37	17,3 ^a	1,17	14,0 ^b	0,93	13,3 ^b	0,76	12,8 ^b	0,71
18:1ω9	6,6 ^a	0,8	6,4 ^a	0,96	7,8 ^a	1,26	6,4 ^a	1,35	9,3 ^a	1,07	10,3 ^{ab}	1,20	14,0 ^b	0,59	10,1 ^a	0,67
18:1ω7	1,2 ^a	0,1	1,1 ^a	0,05	1,3 ^a	0,12	1,1 ^a	0,13	0,9 ^a	0,05	0,8 ^a	0,06	1,1 ^a	0,02	0,9 ^a	0,04
18:2ω6	18,3 ^a	1,0	20,4 ^a	1,07	19,9 ^a	1,67	21,2 ^a	1,77	14,0 ^a	0,92	15,1 ^a	0,44	16,5 ^b	0,15	16,2 ^{ab}	0,55
18:3ω6	0,4 ^a	0,1	0,2 ^b	0,07	0,4 ^a	0,04	0,2 ^b	0,03	0,4 ^a	0,04	n.d.		0,4 ^a	0,03	n.d.	
18:3ω3	0,4 ^a	0,02	0,4 ^a	0,08	0,5 ^a	0,11	0,4 ^a	0,08	0,4 ^a	0,08	0,4 ^a	0,06	0,6 ^b	0,04	0,4 ^a	0,05
20:3ω6	0,4 ^a	0,03	0,6 ^b	0,04	1,0 ^c	0,11	1,4 ^d	0,16	0,6 ^a	0,08	0,9 ^{abc}	0,44	1,0 ^b	0,11	1,4 ^c	0,14
20:4ω6	27,5 ^a	0,8	13,1 ^b	1,25	26,4 ^a	2,36	14,4 ^b	2,02	27,6 ^a	1,72	12,4 ^b	0,77	22,3 ^a	0,82	13,4 ^b	1,38
20:5ω3	0,5 ^a	0,1	6,5 ^b	0,70	0,4 ^a	0,05	6,6 ^b	0,76	0,7 ^a	0,10	6,1 ^b	1,13	0,6 ^a	0,02	6,5 ^b	0,78
22:5ω3	0,8 ^a	0,05	1,3 ^b	0,07	1,0 ^a	0,05	1,5 ^b	0,11	1,1 ^a	0,04	1,6 ^b	0,14	1,1 ^a	0,09	1,7 ^b	0,20
22:6ω3	4,7 ^a	0,4	9,3 ^b	0,35	3,9 ^a	0,49	9,4 ^b	0,48	3,2 ^a	0,16	9,7 ^b	0,74	3,0 ^a	0,23	10,6 ^b	0,83
ω6/ω3	7,3 ^a	0,6	2,0 ^b	0,1	8,3 ^a	0,5	2,1 ^b	0,2	8,1 ^a	0,4	1,6 ^b	0,1	7,7 ^a	0,3	1,6 ^b	0,1
Σω3	6,4 ^a	0,4	17,5 ^b	0,8	5,8 ^a	0,4	17,9 ^b	1,1	5,3 ^a	0,2	17,8 ^b	1,8	5,2 ^a	0,3	19,2 ^b	0,7
Σω6	46,6 ^a	0,5	34,4 ^b	0,6	47,7 ^a	1,2	37,1 ^c	1,4	42,7 ^a	1,4	28,4 ^b	1,2	40,1 ^a	0,7	31,0 ^b	1,2
ΣSFA	37,6 ^{ab}	1,0	39,5 ^b	0,7	36,1 ^a	0,5	36,2 ^a	1,3	40,6 ^{ab}	1,4	41,4 ^a	2,3	38,3 ^{ab}	1,5	37,6 ^b	0,5
ΣMUFA	9,4 ^a	1,0	8,6 ^a	1,2	10,4 ^a	1,6	8,7 ^a	1,7	11,4 ^a	1,1	12,4 ^a	1,3	16,4 ^b	0,7	12,2 ^a	0,7
ΣPUFA	53,0 ^{ab}	0,3	51,9 ^a	1,0	53,5 ^{ab}	1,3	55,0 ^b	1,5	48,0 ^{ab}	1,5	46,2 ^b	2,6	45,4 ^b	1,0	50,2 ^a	0,8
TFA ug/mL	3242,9 ^a	380,9	2873,8 ^a	343,7	3426,7 ^a	338,5	2770,9 ^a	438,6	2442,2 ^a	245,0	2111,7 ^a	305,7	3638,4 ^b	183,4	2101,4 ^a	231,1

	ST DIET								HFHS DIET							
	C		ω3		GSP		ω3&GSP		C		ω3		GSP		ω3&GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
14:0	0,3 ^a	0,1	0,6 ^b	0,1	0,5 ^{ab}	0,1	0,4 ^a	0,02	0,7 ^{bc}	0,1	0,8 ^c	0,1	0,7 ^{bc}	0,1	0,7 ^{bc}	0,1
15:0	0,3 ^a	0,02	0,3 ^a	0,1	0,3 ^a	0,04	0,3 ^a	0,05	0,4 ^b	0,03	0,5 ^b	0,02	0,4 ^b	0,1	0,5 ^b	0,03
16:0	28,1 ^a	0,9	29,2 ^a	1,6	28,5 ^a	0,4	28,2 ^a	0,4	25,5 ^b	0,7	27,1 ^{ab}	0,6	25,4 ^b	0,6	27,0 ^{ab}	0,6
16:1 ω7	0,4 ^a	0,1	0,5 ^a	0,05	0,4 ^a	0,04	0,4 ^a	0,02	0,5 ^a	0,1	0,5 ^a	0,1	0,4 ^a	0,04	0,5 ^a	0,1
17:0	0,7 ^a	0,1	0,7 ^a	0,1	0,6 ^a	0,1	0,6 ^a	0,04	0,7 ^a	0,1	0,7 ^a	0,03	0,7 ^a	0,1	0,7 ^a	0,03
18:0	18,5 ^a	1,3	17,8 ^a	0,8	19,0 ^a	0,5	18,0 ^a	0,4	19,2 ^a	0,3	16,8 ^a	0,2	18,8 ^a	0,5	17,1 ^a	0,4
18:1 ω9	6,7 ^a	0,4	6,2 ^{ab}	0,3	6,6 ^a	0,1	6,1 ^b	0,1	8,6 ^c	0,2	8,7 ^c	0,1	8,4 ^c	0,3	8,6 ^c	0,2
18:1 ω7	2,2 ^{ab}	0,1	2,0 ^a	0,1	2,2 ^b	0,0	1,9 ^a	0,05	1,7 ^c	0,1	1,6 ^c	0,1	1,7 ^c	0,04	1,6 ^c	0,1
18:2 ω6	10,0 ^a	1,1	10,5 ^a	0,4	9,5 ^a	0,4	10,5 ^a	0,2	10,1 ^a	0,5	9,8 ^a	0,2	10,1 ^a	0,4	10,1 ^a	0,3
18:3 ω6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
18:3 ω3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
20:3 ω6	0,7 ^{ab}	0,1	0,7 ^{ab}	0,03	0,6 ^a	0,05	0,8 ^b	0,02	0,9 ^c	0,1	0,9 ^c	0,01	0,9 ^c	0,05	0,9 ^c	0,1
20:4 ω6	24,2 ^a	0,9	15,4 ^{bc}	0,7	24,7 ^a	0,3	16,1 ^b	0,4	24,0 ^a	0,5	14,2 ^c	0,5	24,8 ^a	0,3	14,6 ^c	0,2
20:5 ω3	0,3 ^{ac}	0,1	3,0 ^b	0,3	0,2 ^a	0,03	3,1 ^b	0,2	0,4 ^c	0,02	4,0 ^d	0,3	0,4 ^c	0,02	3,9 ^d	0,4
22:5 ω3	2,5 ^a	0,2	4,2 ^b	0,2	2,2 ^a	0,2	4,5 ^b	0,1	3,3 ^c	0,2	5,1 ^d	0,1	3,3 ^c	0,1	5,1 ^d	0,3
22:6 ω3	4,9 ^a	0,7	8,7 ^b	0,4	4,4 ^a	0,4	8,9 ^b	0,2	3,8 ^c	0,2	9,2 ^b	0,2	3,6 ^c	0,3	8,7 ^b	0,4
ω6/ω3	4,6 ^a	0,7	1,7 ^b	0,1	5,1 ^a	0,2	1,7 ^b	0,04	4,7 ^a	0,1	1,4 ^b	0,1	4,9 ^a	0,2	1,4 ^b	0,03
Σω3	7,7 ^a	0,9	15,9 ^b	0,8	6,9 ^a	0,3	16,4 ^b	0,3	7,4 ^a	0,2	18,3 ^c	0,3	7,4 ^a	0,3	17,7 ^c	0,3
Σω6	35,2 ^a	1,1	26,8 ^{bc}	0,9	35,1 ^a	0,6	27,6 ^b	0,4	35,3 ^a	0,7	25,1 ^c	0,7	36,1 ^a	0,4	25,7 ^c	0,4
ΣSFA	47,9 ^{ab}	0,7	48,7 ^a	1,6	48,9 ^a	0,6	47,5 ^{ab}	0,4	46,4 ^{bc}	0,8	45,8 ^c	0,7	46,0 ^b	0,6	45,9 ^c	0,4
ΣMUFA	9,3 ^a	0,5	8,6 ^{ab}	0,4	9,2 ^a	0,1	8,4 ^b	0,2	10,8 ^c	0,2	10,8 ^c	0,2	10,5 ^c	0,4	10,6 ^c	0,2
ΣPUFA	42,9 ^a	0,5	42,7 ^a	1,4	42,0 ^a	0,6	44,1 ^a	0,5	42,8 ^a	0,8	43,4 ^a	0,7	43,4 ^a	0,5	43,4 ^a	0,5

	ST DIET								HFHS DIET							
	C		ω3		GSP		ω3&GSP		C		ω3		GSP		ω3&GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
14:0	0,8 ^a	0,01	1,2 ^b	0,02	0,7 ^a	0,02	1,3 ^b	0,03	4,5 ^a	0,06	5,3 ^b	0,24	4,4 ^a	0,28	5,4 ^b	0,26
14:1ω5	0,1 ^a	0,01	0,1 ^b	0,01	0,1 ^a	0,01	0,1 ^{ab}	0,01	0,4 ^{ab}	0,02	0,5 ^b	0,05	0,4 ^a	0,04	0,5 ^b	0,04
15:0	0,3 ^a	0,02	0,3 ^a	0,07	0,2 ^a	0,05	0,3 ^a	0,05	0,7 ^a	0,02	0,8 ^a	0,03	0,7 ^a	0,02	0,8 ^a	0,04
16:0	19,4 ^a	0,02	21,4 ^b	0,65	19,6 ^a	0,52	22,4 ^b	0,81	23,9 ^a	0,55	26,2 ^b	0,77	24,7 ^{ab}	0,63	25,3 ^{ab}	1,57
16:1ω7	2,1 ^a	0,22	3,2 ^b	0,10	2,3 ^a	0,21	3,4 ^b	0,30	3,2 ^{ab}	0,24	4,0 ^a	0,65	2,7 ^b	0,29	3,5 ^{ab}	0,66
17:0	0,4 ^{ab}	0,06	0,5 ^a	0,05	0,3 ^b	0,03	0,5 ^a	0,04	0,6 ^a	0,01	0,7 ^b	0,04	0,6 ^{ab}	0,05	0,7 ^b	0,03
17:1ω7	0,2 ^a	0,01	0,3 ^a	0,02	0,2 ^a	0,01	0,3 ^a	0,03	0,4 ^a	0,01	0,4 ^a	0,04	0,4 ^a	0,01	0,4 ^a	0,04
18:0	3,2 ^a	0,15	3,2 ^a	0,08	2,9 ^a	0,21	3,3 ^a	0,21	5,4 ^a	0,38	5,9 ^a	0,63	5,7 ^a	0,21	6,0 ^a	0,39
18:1ω9	28,4 ^a	0,57	25,0 ^b	0,69	29,2 ^a	0,81	24,9 ^b	0,54	37,5 ^a	0,25	35,2 ^b	1,03	37,6 ^a	0,38	35,2 ^b	2,02
18:1ω7	3,1 ^a	0,04	3,0 ^a	0,10	3,2 ^a	0,08	2,9 ^a	0,13	2,3 ^a	0,02	2,1 ^a	0,06	2,3 ^a	0,11	2,2 ^a	0,18
18:2ω6	38,0 ^a	0,47	33,3 ^b	0,71	37,5 ^a	1,07	32,7 ^b	1,08	18,3 ^a	0,27	14,5 ^b	0,48	17,8 ^a	0,22	14,9 ^b	0,36
18:3ω6	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01	n.d.		n.d.		n.d.		n.d.	
20:0	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,02	0,1 ^a	0,01	0,1 ^a	0,005	0,1 ^a	0,003	0,1 ^a	0,03
18:3ω3	1,6 ^a	0,05	1,5 ^a	0,01	1,6 ^a	0,02	1,5 ^a	0,03	1,2 ^a	0,004	1,1 ^a	0,03	1,1 ^a	0,03	1,1 ^a	0,19
20:1ω9	0,4 ^a	0,03	0,5 ^a	0,03	0,4 ^a	0,07	0,5 ^a	0,07	0,8 ^a	0,01	0,9 ^a	0,03	0,8 ^a	0,02	0,9 ^a	0,04
20:2ω6	0,3 ^a	0,02	0,2 ^a	0,02	0,2 ^a	0,02	0,2 ^a	0,03	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01
20:3ω6	0,1 ^{ab}	0,01	0,2 ^b	0,01	0,1 ^a	0,003	0,2 ^b	0,002	0,1 ^a	0,01	0,1 ^a	0,004	0,1 ^a	0,01	0,1 ^b	0,01
20:4ω6	0,6 ^a	0,09	0,4 ^b	0,03	0,6 ^a	0,07	0,4 ^b	0,02	0,3 ^a	0,02	0,2 ^b	0,02	0,3 ^a	0,03	0,2 ^{ab}	0,01
20:5ω3	0,1 ^a	0,01	0,7 ^a	0,06	n.d.		0,8 ^b	0,07	n.d.		0,3 ^a	0,02	n.d.		0,3 ^a	0,06
24:1ω11	0,05 ^a	0,01	0,2 ^b	0,01	0,2 ^b	0,03	0,2 ^b	0,02	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,00
22:5ω3	0,3 ^a	0,03	1,5 ^b	0,14	0,2 ^a	0,02	1,3 ^b	0,16	0,1 ^a	0,01	0,6 ^b	0,07	0,1 ^a	0,02	0,8 ^b	0,07
22:6ω3	0,4 ^a	0,04	2,6 ^b	0,20	0,2 ^a	0,02	2,5 ^b	0,24	0,1 ^a	0,003	0,8 ^b	0,12	0,1 ^a	0,01	1,2 ^b	0,13
ω6/ω3	17,1 ^a	0,8	5,3 ^b	0,2	20,2 ^a	0,2	5,3 ^b	0,5	14,3 ^a	0,3	5,3 ^b	0,3	14,6 ^a	0,6	4,5 ^b	0,6
Σω3	2,3 ^a	0,1	6,5 ^b	0,3	1,9 ^a	0,1	6,3 ^b	0,5	1,3 ^a	0,04	2,8 ^b	0,2	1,2 ^a	0,1	3,5 ^b	0,3
Σω6	39,1 ^a	0,6	34,3 ^b	0,8	38,6 ^a	1,2	33,5 ^b	1,1	18,7 ^a	0,3	14,9 ^b	0,5	18,2 ^a	0,2	15,4 ^b	0,4
ΣSFA	24,3 ^a	0,2	26,9 ^b	0,5	24,0 ^a	0,3	27,9 ^b	0,6	35,3 ^a	0,7	39,1 ^b	1,3	36,3 ^a	0,4	38,4 ^b	1,9
ΣMUFA	34,4 ^a	0,7	32,4 ^b	0,6	35,6 ^a	1,0	32,3 ^b	0,7	44,7 ^a	0,5	43,2 ^a	1,7	44,2 ^a	0,6	42,8 ^a	1,8
ΣPUFA	41,4 ^a	0,6	40,7 ^a	1,1	40,5 ^a	1,2	39,8 ^a	1,2	20,0 ^a	0,3	17,7 ^b	0,7	19,5 ^a	0,3	18,8 ^b	0,2

Table 5
Circulating plasma
FFA

	ST DIET								HFHS DIET							
	C		ω3		GSP		ω3&GSP		C		ω3		GSP		ω3&GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
14:0	1,6 ^a	0,04	2,0 ^b	0,2	1,9 ^b	0,1	2,3 ^b	0,3	4,6 ^c	0,3	5,1 ^c	0,2	3,8 ^d	0,2	5,8 ^e	0,3
15:0	0,6 ^a	0,05	0,7 ^a	0,1	0,4 ^a	0,4	0,7 ^a	0,2	1,0 ^b	0,1	1,1 ^b	0,1	0,9 ^a	0,00	1,1 ^b	0,1
16:0	29,5 ^a	0,2	32,2 ^{bc}	0,6	31,2 ^b	0,6	32,1 ^{bc}	1,7	33,0 ^c	0,8	33,9 ^c	0,5	33,1 ^c	0,1	33,7 ^c	1,0
16:1 ω7	3,6 ^a	0,2	3,4 ^a	0,6	3,5 ^a	0,5	3,5 ^a	0,9	3,7 ^a	0,6	4,2 ^a	0,4	3,3 ^a	0,2	4,5 ^a	0,7
17:0	1,6 ^a	1,0	1,6 ^a	1,2	1,2 ^a	1,2	1,6 ^a	1,4	1,5 ^a	1,2	2,0 ^a	1,4	1,3 ^a	1,0	1,8 ^a	1,5
18:0	7,3 ^a	0,3	9,2 ^{ab}	1,3	8,5 ^{ab}	0,9	10,2 ^{bc}	1,4	10,7 ^{bc}	1,3	12,7 ^c	0,9	9,6 ^b	0,2	12,5 ^c	1,6
18:1 ω9	18,7 ^a	0,4	13,5 ^b	1,0	17,2 ^a	0,7	12,8 ^b	1,7	21,6 ^{cd}	1,9	18,4 ^{ac}	2,5	22,8 ^d	1,0	17,6 ^a	1,4
18:1 ω7	2,3 ^a	0,2	1,7 ^{bc}	0,1	2,2 ^{ab}	0,3	1,7 ^{bcd}	0,4	1,4 ^{cd}	0,2	1,2 ^d	0,2	1,5 ^c	0,04	1,2 ^d	0,1
18:2 ω6	27,0 ^a	0,6	22,1 ^b	1,4	26,4 ^a	1,0	21,9 ^b	1,1	15,1 ^c	0,6	10,8 ^d	0,2	16,7 ^c	0,1	11,2 ^d	1,3
18:3 ω6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
18:3 ω3	1,4 ^a	0,1	1,2 ^b	0,1	1,4 ^{ab}	0,1	1,2 ^{bc}	0,1	1,2 ^b	0,1	1,0 ^c	0,1	1,3 ^{ab}	0,1	1,0 ^c	0,1
20:3 ω6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
20:4 ω6	4,1 ^a	0,4	2,6 ^b	0,5	4,3 ^a	0,7	2,7 ^b	0,4	4,2 ^a	0,6	1,9 ^b	0,2	4,1 ^a	0,1	2,0 ^b	0,2
20:5 ω3	0,2 ^a	0,1	2,3 ^b	0,3	0,1 ^a	0,01	2,3 ^b	0,2	0,2 ^a	0,1	2,1 ^b	0,6	0,2 ^a	0,1	2,0 ^b	0,2
22:5 ω3	0,5 ^a	0,1	1,8 ^b	0,3	0,5 ^a	0,2	1,8 ^b	0,3	0,6 ^a	0,1	1,5 ^b	0,5	0,7 ^a	0,1	1,4 ^b	0,3
22:6 ω3	1,2 ^a	0,1	5,3 ^b	0,3	0,9 ^c	0,1	4,8 ^b	1,0	0,7 ^c	0,1	4,0 ^b	0,9	0,7 ^c	0,03	3,9 ^b	0,7
ω6/ω3	9,5 ^{ac}	1,2	2,3 ^b	0,1	10,8 ^a	1,2	2,5 ^b	0,4	7,3 ^c	0,7	1,6 ^d	0,4	7,3 ^c	0,7	1,6 ^d	0,2
Σω3	3,3 ^a	0,4	10,7 ^b	0,4	2,9 ^a	0,2	10,1 ^b	1,4	2,7 ^a	0,3	8,6 ^b	2,1	2,9 ^a	0,3	8,4 ^b	1,0
Σω6	31,1 ^a	0,9	24,7 ^b	1,0	30,8 ^a	1,2	24,5 ^b	1,0	19,2 ^c	1,2	12,8 ^d	0,2	20,8 ^c	0,2	13,2 ^d	1,2
ΣSFA	40,7 ^a	0,9	45,7 ^{bc}	1,2	43,3 ^{ac}	2,3	47,0 ^{bc}	2,2	50,9 ^{bd}	1,6	54,7 ^d	1,0	48,7 ^b	0,8	54,8 ^d	2,7
ΣMUFA	24,9 ^a	0,3	18,9 ^b	1,1	23,1 ^a	1,4	18,3 ^b	3,2	27,2 ^c	2,8	24,0 ^a	2,5	27,6 ^c	0,9	23,6 ^a	2,1
ΣPUFA	34,4 ^a	0,7	35,4 ^a	1,1	33,6 ^a	1,0	34,7 ^a	1,5	21,9 ^b	1,3	21,3 ^b	2,1	23,7 ^b	0,2	21,6 ^b	1,8
Total FFA (μg/mL)	212,0 ^a	22,1	164,2 ^{ab}	32,0	166,6 ^{ab}	24,2	147,5 ^b	30,0	169,4 ^{ab}	30,0	140,2 ^{ac}	15,9	187,8 ^b	4,5	136,6 ^c	25,2

Table 6

	ST DIET								HFHS DIET							
	C		ω3		GSP		ω3&GSP		C		ω3		GSP		ω3&GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
FAD indexes in plasma																
SCD-16 [palmitoleic/palmitic]	0,05 ^a	0,005	0,05 ^a	0,006	0,05 ^a	0,01	0,05 ^a	0,009	0,05 ^a	0,01	0,1 ^a	0,01	0,1 ^a	0,005	0,1 ^a	0,01
SCD-18 [oleic/stearic]	0,4 ^a	0,09	0,4 ^a	0,1	0,6 ^a	0,1	0,5 ^a	0,1	0,5 ^a	0,10	0,7 ^{ab}	0,1	1,1 ^c	0,1	0,8 ^b	0,1
Δ5D [ARA/DGLA]	62,9 ^a	4,1	20,8 ^b	1,8	26,9 ^c	2,9	10,7 ^d	0,8	43,0 ^a	4,9	15,6 ^b	4,5	23,7 ^c	3,2	9,7 ^d	1,3
Δ6D [DGLA/LA]	0,02 ^a	0,002	0,03 ^a	0,004	0,1 ^b	0,008	0,1 ^b	0,01	0,05 ^a	0,01	0,1 ^{ab}	0,03	0,1 ^a	0,01	0,1 ^b	0,01
Δ6/5D [EPA/ALA]	1,2 ^a	0,3	17,2 ^b	4,3	0,8 ^a	0,2	17,3 ^b	6,4	2,0 ^a	0,5	14,3 ^b	3,8	1,0 ^a	0,06	15,3 ^b	1,9
Δ4D [DHA/DPA]	5,7 ^a	0,7	7,0 ^b	0,3	4,1 ^c	0,6	6,5 ^{ab}	0,8	3,1 ^a	0,1	6,3 ^b	0,5	2,7 ^a	0,3	6,3 ^b	0,9
FAD indexes in liver																
SCD-16 [palmitoleic/palmitic]	0,03 ^a	0,004	0,04 ^a	0,003	0,04 ^a	0,01	0,04 ^a	0,003	0,05 ^a	0,01	0,05 ^a	0,01	0,04 ^a	0,01	0,04 ^a	0,01
SCD-18 [oleic/stearic]	0,4 ^a	0,01	0,3 ^b	0,01	0,4 ^a	0,1	0,3 ^b	0,03	0,9 ^a	0,1	1,0 ^a	0,01	0,7 ^a	0,1	1,0 ^a	0,1
Δ5D [ARA/DGLA]	107,0 ^a	4,8	31,9 ^b	2,9	92,5 ^a	12,1	26,5 ^b	8,5	64,3 ^a	8,7	31,4 ^b	1,5	71,2 ^a	4,3	31,4 ^b	4,2
Δ6D [DGLA/LA]	0,01 ^a	0,0004	0,02 ^b	0,003	0,01 ^a	0,002	0,03 ^b	0,007	0,02 ^a	0,004	0,02 ^a	0,001	0,02 ^a	0,001	0,02 ^a	0,002
Δ6/5D [EPA/ALA]	0,6 ^a	0,07	13,7 ^b	1,5	0,6 ^a	0,08	12,1 ^b	1,3	0,8 ^a	0,2	6,3 ^b	0,4	1,0 ^a	0,2	5,9 ^b	0,3
Δ4D [DHA/DPA]	7,4 ^a	0,9	10,6 ^b	0,5	4,9 ^c	0,6	9,9 ^b	0,8	3,9 ^a	0,1	6,3 ^b	0,7	3,9 ^a	0,2	6,3 ^b	0,8
Lipid oxidation parameters in liver																
Conjugated dienes (mmol hydroperoxides/kg of lipid)	19,6 ^a	0,9	17,2 ^b	1,1	20,3 ^a	1,8	16,2 ^b	2,1	27,5 ^a	1,5	29,8 ^a	2,9	27,7 ^a	2,2	27,5 ^a	2,8
TBARS (mg MDA/kg of liver)	4,5 ^{ac}	1,4	10,0 ^b	2,4	3,2 ^c	0,2	7,6 ^{ab}	1,7	3,6 ^a	0,7	10,7 ^b	2,0	2,7 ^a	0,3	6,6 ^c	1,4
Oxidative stress parameters in plasma																
Total albumin carbonylation index (%)	36,8 ^a	2,6	43,2 ^a	4,1	28,1 ^b	1,9	34,3 ^{ab}	9,1	41,8 ^a	3,3	37,3 ^a	3,2	30,3 ^b	2,8	24,3 ^b	4,3
ORAC (μmol Trolox/mL plasma)	17,6 ^a	8,3	18,8 ^a	4,4	16,6 ^a	5,1	22,1 ^a	2,9	17,7 ^{ab}	7,9	10,7 ^a	5,7	19,3 ^{ab}	5,0	24,0 ^b	3,5
SOD (U/g Hb)*	3698 ^a	512	3277 ^{ab}	395	3033 ^{ab}	274	2285 ^b	1083	4688 ^a	1344	3856 ^{ab}	689	2619 ^b	1338	2787 ^{ab}	1462
GPX (U/g Hb)*	0,4 ^a	0,1	1,2 ^a	0,7	6,1 ^b	0,5	19,2 ^c	3,01	0,4 ^a	0,04	3,8 ^a	0,9	13,4 ^b	1,9	36,7 ^c	8,5
Inflammation parameters in liver																
CRP (mg/mL)*	91,4 ^a	21,6	45,4 ^b	15,7	49,7 ^b	7,9	56,1 ^b	26,5	93,9 ^a	53,7	64,3 ^a	18,9	76,7 ^a	41,7	60,3 ^a	46,9
TNFα (mg/mL)*	116,0 ^a	72,5	63,8 ^b	26,3	73,1 ^b	66,7	62,6 ^b	47,6	91,5 ^a	47,8	55,2 ^b	49,3	85,2 ^{ab}	59,5	54,5 ^b	65,4

Table 7	ST DIET								HFHS DIET							
	C		ω 3		GSP		ω 3&GSP		C		ω 3		GSP		ω 3&GSP	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
Eicosanoids from EPA (ng/mL)																
5HEPE	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
12HEPE	80 ^a	44	468 ^b	433	34 ^a	15	195 ^{ab}	191	35 ^a	18	312 ^b	176	19 ^a	10	370 ^b	96
15HEPE [*]	3,4 ^{ac}	1,2	7,5 ^b	3,5	1,6 ^a	0,1	5,6 ^{bc}	1,5	1,6 ^a	0,2	4,6 ^b	1,7	1,6 ^a	0,2	5,0 ^b	1,7
12HpEPE ^{**}	593 ^a	188	208 ^b	66	200 ^b	95	117 ^b	81	270 ^a	178	46 ^b	23	387 ^a	181	99 ^b	32
15HpEPE ^{**}	219 ^a	131	2752 ^b	594	86 ^a	36	2269 ^b	539	86 ^a	43	376 ^b	121	24 ^a	6	511 ^c	127
8isoPGF _{3α}	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
TXB ₃	3,4 ^a	0,2	4,4 ^a	1,5	n.d.	n.d.	3,6 ^a	0,6	n.d.	n.d.	3,7 ^a	0,6	n.d.	n.d.	4,0 ^a	1,0
PGD ₃ /E ₃	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Docosanoids from DHA (ng/mL)																
17HDoHE [#]	55 ^a	20	148 ^b	104	35 ^a	12	77 ^a	51	23 ^a	11	101 ^b	45	13 ^a	5	170 ^c	73
11HDoHE	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
4HDoHE	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
17HpDoHE ^{**}	2062 ^a	434	1437 ^b	367	1466 ^b	333	1269 ^b	195	1752 ^a	476	1677 ^a	313	1857 ^a	298	1856 ^a	598
RvD ₁	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
PD _x	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Eicosanoids from ARA (ng/mL)																
11HETE [#]	7,7 ^a	4,5	2,5 ^b	2,4	3,2 ^b	1,6	0,5 ^b	0,3	2,4 ^a	0,6	1,0 ^a	0,7	8,2 ^b	4,8	1,4 ^a	1,0
PGE ₂ [#]	16 ^a	12	7 ^b	4	7 ^b	3	4 ^b	1	6 ^a	2	4 ^a	2	10 ^b	6	4 ^a	1
LTB ₄	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
8isoPGF _{2α}	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

S1

Dietary groups	ST-C	ST- ω 3	ST-GSP	ST- ω 3&GSP	HFHS-C	HFHS- ω 3	HFHS-GSP	HFHS- ω 3&GSP
Diet composition								
Flour (g)	1000 ^a	1000 ^a	1000 ^a	1000 ^a	1000 ^b	1000 ^b	1000 ^b	1000 ^b
Porcine gelatine (g)	25	25	25	25	25	25	25	25
Soybean lecithin (g)	6	6	6	6	22	22	22	22
Polyphenol extract ^c (mg)	-	-	880	880	-	-	1090	1090
Oil (mL)	19 mL soybean	19 mL EPA:DHA 1:1	19 mL soybean	19 mL EPA:DHA 1:1	24 mL soybean	24 mL EPA:DHA 1:1	24 mL soybean	24 mL EPA:DHA 1:1
Macronutrients (% weight)								
Protein	16.4	16.4	16.4	16.4	21.7	21.7	21.7	21.7
Fat	6.2	6.2	6.2	6.2	24.1	24.1	24.1	24.1
Carbohydrates	46.6	46.6	46.5	46.5	45.0	45.0	44.9	44.9
Macronutrients (% caloric value)								
Protein	21,3	21.3	21.3	21.3	17,9	17.9	17.9	17.9
Fat	18,2	18.2	18.2	18.2	44,9	44.9	44.9	44.9
Carbohydrates	60,5	60.5	60.5	60.5	37,2 ^d	37.2 ^d	37.2 ^d	37.2 ^d
Total energy density (kcal/g)	3.1	3.1	3.1	3.1	4.8	4.8	4.8	4.8

	C and GSP groups		ω 3 and ω 3&GSP groups	
	Standard diet	HFHS diet	Standard diet	HFHS diet
14:0	0,017	11,5	0,075	11,6
16:0	14,8	32,2	14,6	32,0
16:1 ω 7	0,015	1,7	0,08	1,7
18:0	2,9	13,7	2,9	13,7
18:1 ω 9	20,5	26,4	20,3	26,2
18:1 ω 7	0,026	0,67	0,033	0,68
18:2 ω 6	58,6	9,7	57,8	8,7
20:0	n.d.	n.d.	0,0055	0,0069
18:3 ω 3	2,9	1,1	2,9	1,1
20:1 ω 9	0,025	0,031	0,017	0,021
18:4 ω 3	0,0026	0,0032	0,026	0,033
20:2 ω 6	0,0034	0,0043	0,0036	0,0045
20:3 ω 6	n.d.	n.d.	0,0038	0,0048
20:4 ω 6	0,0069	0,0086	0,029	0,036
22:1 ω 11	0,019	0,023	0,02	0,024
22:1 ω 9	0,0043	0,0054	0,0048	0,006
20:4 ω 3	0,0034	0,0043	0,018	0,022
20:5 ω 3	0,012	0,015	0,43	0,54
24:1 ω 9	0,0048	0,0060	0,0065	0,0082
22:5 ω 3	0,0045	0,0056	0,074	0,093
22:6 ω 3	0,02	0,024	0,44	0,56
Total SFAs	17,5	59,4	18,3	60,5
Total MUFAs	21,1	29,3	20,3	28,4
Total PUFAs	61,1	10,8	61,0	10,7
Total ω 3	3,3	1,6	3,2	1,4
Total ω 6	58,8	10,4	58,9	10,6

Compound	Retention Time (min)	LIT parameters	
		Collision energy (eV)	Quantification transition (m/z)
8iso-PGF _{3α}	6.79	30	351>253
TXB ₃	6.97	19	367>195
PGD ₃ /PGE ₃	7.67	19	349>313
8iso-PGF _{2α}	8.28	28	353>299
PGE ₂	9.09	20	351>315
RvD ₁	9.80	25	375>141
PD _x	12.89	30	359>153
LTB ₄	13.79	27	335>195
15HpEPE	17.36	20	333>315
12HpEPE	17.80	25	333>315
15HEPE	17.92	27	317>219
12HEPE	18.72	27	317>179
5HEPE	20.47	25	317>255
17HpDoHE	21.90	26	359>341
17HDoHE	21.94	27	343>245
11HETE	22.09	30	319>167
12HETEd ₈	22.66	30	325>307
11HDoHE	23.20	27	343>149
4HDoHE	23.64	27	343>281
EPA	24.43	27	301>257
DHA	24.93	30	327>283
ARA	25.10	30	303>259