Influence of the honeybee and trait similarity on the effect of a non-native plant on pollination and network rewiring

Ana Montero-Castaño^{* a} and Montserrat Vilà^a

^aEstación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio s/n, Isla
de la Cartuja, 41092 Sevilla, Spain

*Corresponding author: ana.montero.castano@gmail.com

Running headline: Plant invasions: the role of the honeybee and trait similarity

Summary

1. Introduced entomophilous non-native plants usually become well integrated into the diet of generalist pollinators. This integration can affect the entire recipient plant-pollinator network. Effects vary from facilitative to competitive, and understanding the factors that govern such variability is one of the fundamental goals in invasion ecology.

Species traits determine the linking patterns between plant and pollinator species. Therefore, trait similarity among plants or among pollinators might modulate how they affect each other.

2. We conducted a flower removal experiment to investigate the effects of the non-native entomophilous legume *Hedysarum coronarium* on the pollination patterns of a Mediterranean shrubland plant-pollinator network. Specifically, we explored whether effects were influenced by similarity with the resident plant species in flower morphology (papilionate vs. non-papilionate), and whether effects on the pollinator community were influenced by similarity in functional group with its main visitor species (bees vs. non-bees). In addition, we explored whether *Hedysarum* had an effect on the identity of interactions. For this purpose, we calculated the interaction rewiring; i.e., the number of plant-pollinator interactions that were gained or lost after invasion.

3. *Hedysarum* was well integrated into the diet of 15 generalist pollinators having the honeybee as its main visitor species. Such integration did not affect visitation rates, normalized degree (i.e., proportion of pollinators they are visited by) and niche overlap (i.e., proportion of plant species they share pollinators with) of plants, irrespective of their flower morphology. Only the proportion of honeybee visits to resident plants decreased with invasion. On the other hand, *Hedysarum* reduced

visitation rates and niche overlap of pollinators, mainly those of bee species. Finally, we observed that changes in the foraging behaviour of the honeybee were positively associated with the interaction rewiring involving the rest (92 taxa) of pollinators.

4. In conclusion, pollinators show a plastic use of floral resources, responding to the presence of non-native plants. When the non-native attracts highly competitive pollinators such as the honeybee, plasticity is especially significant in pollinators that are functionally close to that competitive pollinator. The result is an interaction rewiring due to pollinators avoiding competition with the honeybee. Though this plasticity might not quantitatively affect the pollination of plants, consequences on their reproduction and the functioning of the network can derive from the interaction rewiring.

Key-words: *Apis mellifera*, flower morphology, *Hedysarum coronarium*, plant invasion, plant-pollinator communities, pollinator functional group

Introduction

Many entomophilous and obligate out-crossing non-native plants become well integrated into the diet of resident pollinators (Vilà et al. 2009). Many of these pollinators are super-generalist species such as the honeybee (*Apis mellifera* L.) or bumblebees (*Bombus* spp.) (Olesen, Eskildsen & Venkatasamy 2002; Stout, Kells & Goulson 2002; Gross et al. 2010). The new plant-pollinator interactions established do not occur in isolation but rather are embedded in complex interaction networks (Montoya, Pimm & Solé 2006), and thus the effect of non-natives can expand to the

entire recipient plant-pollinator network (Lopezaraiza-Mikel et al. 2007, Bartomeus, Vilà & Santamaría 2008, Padrón et al. 2009, Ferrero et al. 2013).

The effects of non-native entomophilous plants on both pollinators and the pollination of plants (Bjerknes et al. 2007; Montero-Castaño & Vilà 2012) vary from facilitative to competitive, being highly context dependent. Understanding the factors that govern such variability would allow us to predict the impact of non-native plant species on recipient communities, which is one of the fundamental goals of invasion ecology (Simberloff et al. 2013).

The linking patterns of plant-pollinator interactions are determined by several factors. First of all, a match between plant traits (e.g. corolla size, shape and colour, type of reward offered) and pollinator traits (e.g. body size, tongue length) is required for interactions to be established (Stang, Klinkhamer & van der Meijden 2006; Santamaría & Rodríguez-Gironés 2007; Olesen et al. 2011; Encinas-Viso, Revilla & Etienne 2012; Bartomeus 2013). Consequently, the similarity among species in some traits can determine how plant and pollinator species affect each other (Lázaro, Hegland & Totland 2008; Morales & Traveset 2009; Campbell et al. 2010; Gibson, Richardson & Pauw 2012; Carvalheiro et al. 2014). For instance, observations on pairs of co-flowering non-native and native species have found that similarity in flower morphology can reduce pollinator visitation rates to native species (Morales & Traveset 2009). However, this trend has not been found when considering the entire native plant community (Morales & Aizen 2006; Vilà et al. 2009). In the case of non-native plants with flower morphologies not accessible to all pollinators, such as papilionate flowers (Córdoba & Cocucci 2011), we would expect them to compete more strongly for pollinators with resident species which share similar flower morphology.

77 Second, pollinators adapt their foraging behaviour to the abundance and quality of
78 available floral resources (Mustajarvi et al. 2001) in order to optimize their food
79 intake (Armbruster & Herzig 1984). Therefore, the arrival of a highly rewarding non-
80 native plant can alter the foraging behaviour of pollinators by modifying the floral
81 environment (Memmott & Waser 2002; Vilà et al. 2009). Pollinators would respond in
82 accordance with their body size, sociability, preferred food resource, flying distances,
83 etc. (Greenleaf et al. 2007; Bommarco et al. 2010). Pollinator taxa sharing some of
84 these ecological characteristics might behave similarly and might thus belong to the
85 same functional group due to their functional redundancy (Hagen *et al.* 2012). We
86 would expect the effect of non-native plants to be higher in those pollinators
87 belonging to the same functional group than those visiting them.

88 Finally, the linking patterns of plant-pollinator interactions are also influenced by
89 interspecific competition among taxa (Carstensen et al. 2014). Some highly
90 competitive pollinator species are able to displace other pollinators by depleting floral
91 resources (e.g. Roubik 1980, Paine 2004) and/or by physical disturbance (Gross &
92 Mackay 1998). Therefore, the arrival of a highly rewarding non-native plant can also
93 alter the foraging behaviour of pollinators by altering the behaviour of one or more
94 highly competitive species able to displace other pollinators.

95 Through the above mentioned mechanisms, invasion not only alters the frequency of
96 plant-pollinator interactions, but can also entail their reshuffling, with gains or losses
97 of exclusive pair-wise interactions in recipient communities (Bartomeus *et al.* 2008).
98 The turnover of interactions when co-occurring species interact differently over time
99 or space is known as interaction rewiring (Burkle & Alarcón 2011; Poisot et al. 2012;
100 Trojelsgaard et al. 2015). Interaction rewiring can have important consequences on
101 networks. The way interactions are shaped; i.e., their topology, affects networks'
102 functioning and persistence to species loss (Kaiser-Bunbury et al. 2010; Tylianakis et

al. 2010; Ramos-Jiliberto et al. 2012). Even if topological properties are conserved after invasion (e.g. Vilà et al. 2009, but see Aizen, Morales & Morales 2008), the functional role of particular species may change (Campos-Navarrete et al. 2013; Nielsen & Totland 2014).

We present a flower removal field experiment, conducted during two consecutive years, to investigate the effect of a highly rewarding non-native legume species on the pollination patterns of the entire plant-pollinator network. Specifically, we address the following questions: (a) how is a non-native entomophilous plant species with a floral morphology that restricts pollinator visitation integrated into the diet of the resident pollinator community? (b) is the effect of the non-native plant on the visitation of the resident plants dependent on the similarity in flower morphology between the non-native and resident species? (c) is the effect of the non-native plant on the foraging behaviour of pollinators dependent on the functional similarity between the non-native's main visitor and the pollinators?; and finally (d) is there an interaction rewiring and is it influenced by the foraging behaviour of the non-native's main visitor and its functional closeness to the rest of pollinators?

Materials and methods

Non-native species and study area

Hedysarum coronarium L. (Leguminosae) is a short-lived N-fixing perennial (Sulas et al. 2000) with either erect (0.8 m average height) or prostrate growth (Bustamante et al. 1998). Its inflorescences are racemes with up to 30 pink flowers rich in pollen and nectar (Rodríguez-Riaño, Ortega-Olivencia & Devesa 1999) that bloom during April

and May. Its papilionate and restrictive flowers (Córdoba & Cocucci 2011) are self-compatible but present high out-crossing rates (Louati-Namouchi, Louati & Chriki 2000, Yagoubi & Chriki 2000). *Hedysarum* is mainly pollinated by bees, and in particular the honeybee which is its main pollinator in the study area and in other areas as well (Satta et al. 2000; Montero-Castaño, Vilà & Ortiz-Sánchez 2014) (Fig. 1).

Hedysarum is native of the southwestern part of the Mediterranean basin (Talavera et al. 1988), where it grows from sea level to low frost-free altitudes (Gutiérrez 1982). It has been introduced as a forage plant into other semiarid regions of the Mediterranean basin due to its high palatability and feeding value to cattle (Yagoubi & Chriki 2000). It is also used for erosion control, re-vegetation and high quality honey production (Flores et al. 1997; Satta et al. 2000). Currently it grows in many Mediterranean basin countries; from Turkey to Spain (Flores et al. 1997). We conducted our study in Menorca (the northernmost of the Balearic Islands, Spain). *Hedysarum* was introduced to this island between the end of the 18th century and the beginning of the 19th century (Ortells & Campos 1983). Since 1860 it has been used in a traditional cyclical agro-farming system (Bustamante, Allés & Espadas 2007). In addition, *Hedysarum* has escaped from cultivated fields and has become naturalised (*sensu* Pyšek et al. 2004) in natural and semi-natural areas such as ditches, old-fields, field edges and ruderal areas (Fraga et al. 2004).

Experimental design and pollination censuses

We located three sites with early successional shrublands (Carreras, Pons & Canals 2007) where we established three pairs (one in each site) of invaded 20x20 m² plots. Paired plots were located at an average distance of 138.3 m (ranging from 95 to 164

m), so that they could potentially share the same pollinator community because most of pollinator flying distances fall within this range (Gathmann & Tschardt 2002; Osborne *et al.* 2008; Mawdsley & Sithole 2009; Bommarco *et al.* 2010). Meanwhile, the average distance between pairs was 11.5 km (ranging from 600 m to 17.2 km) to assure their independence.

Despite *Hedysarum* being one of the most dominant species in the shrublands (cover ranging from 26.4% to 48.6% across plots), in each plot it coexisted with 8.33 ± 0.33 (mean $\pm$ SE, hereafter) native co-flowering species. Overall, ten native plant species also belonged to the family Leguminosae and represented on average $36.4 \pm 7.3\%$ of the plant species in each plot. The rest of the native plant species (17) belonged to seven different families and had open and accessible flowers (Appendix S1 in Supporting Information).

To investigate the effect of *Hedysarum* on recipient plant-pollinator networks, we manually removed all *Hedysarum* inflorescences from one randomly selected plot of each pair (removal plot, hereafter), while the other plot was not manipulated (invaded plot, hereafter).

We conducted pollination censuses in the same study plots in the springs of 2009 and 2010 throughout the entire flowering period of *Hedysarum* (April-May). In both years weather conditions fell within the average ranges for these months in the study area (AEMET).

Pollination censuses were conducted on sunny, warm (≥ 17 °C) and non-windy days, from 10 a.m. to 6 p.m. During each observation period (15 min), we counted the number of floral units (hereafter flowers, according to Dicks, Corbet & Pywell 2002) under observation, the number and identity of pollinators and the number of visits of each pollinator species. A visitor was considered a pollinator if it entered a flower and touched its sexual parts. The pollinator species that could not be identified in the

field were sorted into distinct morphospecies and caught for later identification by specialists. Voucher specimens are deposited at EBD-CSIC.

The observation schedule for each plant species and individual under observation was randomly established. We considered a plot to be properly surveyed when, according to its rarefaction curve, we found no new plant-pollinator interaction after six or more censuses (Appendix S2). In total, we conducted 1252 censuses (313 h). On average each plant species was observed for 5.79 ± 0.60 h per plot, ranging from 1 h to 23.25 h.

Statistical analyses

For each study plot we built plant-pollinator networks with the data gathered during the two study years (i.e. six networks: three invaded and three removal plots). A network is defined as a two dimensional matrix (i^*j) describing the interaction between the flowering plant species (i) and the pollinator species (j) in the community. Each cell in the matrix (a_{ij}) can be 1 or 0 indicating whether the interaction between the plant species i and the visitor species j is observed or not, respectively. Quantitative networks were built following the same criteria, except that each a_{ij} value is the weight of the interaction between the plant species i and the pollinator j measured as the visitation rate (n^0 visits/flower/hour) (Jordano, Bascompte & Olesen 2003).

In order to explore both quantitative and qualitative alterations in pollination patterns, for each plant and pollinator species we calculated visitation rate, normalised degree and niche overlap in each network (Table 1). These response variables, respectively, inform about the intensity and number of interactions established by each species, and about how these interactions are distributed. The proportion of honeybee visits

(the main pollinator of *Hedysarum* in the study area) was also calculated for each plant species. Interaction rewiring was calculated for plant species shared between paired invaded and removal plots. Based on our experimental design, all pollinator species were assumed to be shared between invaded and removal paired plots (Table 1). We estimated two values of interaction rewiring: one considering all the interactions between shared species and the other excluding singletons (i.e., interactions that were only observed once), which represent interactions that are rare and difficult to detect and, thus, could potentially overestimate rewiring (Chacoff et al. 2012).

To explore the effect of *Hedysarum* on the pollination of resident plant species and whether their similarity in flower morphology with *Hedysarum* influenced such an effect, we built linear and generalised mixed models with the effect of treatment (invaded vs. removal), flower morphology (papilionate vs. non-papilionate) and their interaction as fixed effects. Site was included as a random factor in the models. The response variable visits (V_p), standardised by the number of flowers and hours of observation, was log-transformed and analysed through a linear mixed model. For the response variables normalised degree (D_p) and niche overlap (NO_p) we built generalised mixed models with binomial as error distribution family. The proportion of honeybee visits was logit-transformed according to Warton & Hui (2011) and analysed through a linear mixed model.

To explore the effect of *Hedysarum* on the foraging behaviour of pollinator species and whether their functional similarity to the main visitor of *Hedysarum* (i.e. the honeybee) influenced such an effect, we built linear and generalised models with the effect of treatment (invaded vs. removal), functional group (bees vs. non-bees) and their interaction as fixed effects. The functional group of bees included all bee

species observed and the short tongued bumblebee *Bombus terrestris*. The functional group of non-bees included wasps, dipterans and coleopterans. Though the functional group of bees includes a wide variety of taxa, they all have larger flying ranges, visitation rates and capacities to reach low accessible floral resources than the non-bees, and are functionally closer to the honeybee. In addition, this functional classification is a compromise between the information available about the ecology of the observed taxa and an acceptable representation of functional groups across invaded and removal plots for allowing robust statistical analyses.

For the log-transformed response variable visits (V_a), we built a linear model with the logarithms of the number of flowers and hours of observation included as offsets. For the response variables normalised degree (D_a) and niche overlap (NO_a) we built generalised models with binomial as error distribution family. We also calculated the three response variables for the honeybee and analysed them through paired Wilcoxon tests.

Finally, we explored whether interaction rewiring of pollinators excluding the honeybee was influenced by changes in the foraging behaviour of the honeybee or by their functional similarity with the honeybee. We defined the changes in the foraging behaviour of the honeybee as the difference in the proportion of visits to a given plant species that the honeybee achieved in invaded and removal paired networks. For those plant species present in more than one site, data were pooled for all invaded and all removal plots where present. We built a generalised linear model with the change in foraging behaviour of the honeybee, the functional group of the pollinators involved (bees vs. non-bees) and their interaction as fixed factors. The binomial was the error distribution family. The analysis was repeated for the response variable interaction rewiring calculated excluding singletons.

The calculation of the network parameters and the analyses were performed in R (R Development Core Team 2014). Network parameters were calculated with the library *bipartite*. Linear and generalised mixed models were conducted with the libraries *nlme* and *lme4*, respectively. *Post hoc* multiple comparisons were conducted by building our own contrast matrices and analysing them with the library *multcomp*.

Results

We observed a total of 28 flowering plant species from eight different families. The eleven papilionate species, including *Hedysarum*, represented $41.8 \pm 7.6\%$ of the species in all study plots. Plants were visited by 93 pollinator species belonging to 38 families of Coleoptera (19.4%), Diptera (38.7%) and Hymenoptera (41.9%) (Appendix S1). All pollinator species are considered native to the study site.

Integration of Hedysarum into the diet of resident pollinators

Hedysarum was visited by a total of 15 pollinator species: 11 hymenopterans (including 7 Apidae) and four coleopterans, which represented 16.1% of the total community of pollinators. With the exception of the bees *Andrena ovatula* (Kirby, 1802) and *Synhalonia hungarica* (Fries, 1895), which represented only the 0.7% of its visits, all pollinators that visited *Hedysarum* were also observed visiting other plant species. Pollinators visiting *Hedysarum* were on average more generalised than pollinators visiting only natives ($D_a = 0.24 \pm 0.07$ and 0.09 ± 0.01 , respectively; $Z = -5.081$, $p\text{-value} < 0.001$).

Although *Hedysarum* received 54% of the visits observed in invaded plots, when standardised by the number of flowers, its visitation rate was low and similar to that of the resident papilionate species and lower than that of non-papilionate species (Fig. 2a). On average, *Hedysarum* had a normalised degree higher than papilionate species but similar to that of non-papilionate species (Fig. 2b). *Hedysarum* also had an averaged niche overlap higher than papilionate species but lower than non-papilionate species (Fig. 2c). *Hedysarum* was mostly visited ($92.7 \pm 4.2\%$) by the honeybee. On average, the proportion of honeybee visits to *Hedysarum* was higher than that to resident plant species, whether papilionate or not (Fig. 2d).

Effect of Hedysarum on the pollination of resident plants and the influence of flower morphology similarity

There were no significant differences in pollinator visits, normalised degree or niche overlap of resident plants between invaded and removal plots. Flower morphology had a significant effect on these variables, with papilionate species showing lower values for the three response variables than the non-papilionate ones. The interaction between treatment and flower morphology was not significant for any of the response variables, indicating that *Hedysarum* did not affect the pollination of resident plants, irrespective of their similarity in flower morphology (Table 2, Figs 2a, b and c).

The abovementioned results obtained for the entire pool of pollinators contrasted with those considering only the honeybee. The proportion of honeybee visits was lower in invaded than in removal plots, regardless of flower morphology. Once again, the interaction between treatment and flower morphology was not significant (Table 2, Fig. 2d).

Effect of Hedysarum on the foraging behaviour of pollinators and the influence of functional similarity

Pollinator visitation rates differed between invaded and removal plots but did not differ between functional groups. The interaction between the two variables was not significant, indicating that *Hedysarum* did not more strongly affect the visits conducted by bees, i.e., those taxa functionally closer to the honeybee, than those conducted by non-bee pollinators. Bees conducted less visits in invaded than in removal plots; while a similar but not significant trend was observed for the non-bees functional group (Table 2, Fig. 3c). No significant trend was observed for honeybee visits ($N = 6$, $V = 6$, $p\text{-value} = 0.250$) (Fig. 3c).

The normalised degree of pollinators did not differ between invaded and removal plots, neither between functional groups. The interaction between the two variables was not significant (Table 2, Fig. 3a). The honeybee also showed a similar normalised degree in invaded and removal plots ($N = 6$, $V = 1$, $p\text{-value} = 0.500$) (Fig. 3b).

The niche overlap of pollinators differed between invaded and removal plots and between functional groups. In addition, the interaction between the two variables was also significant: while both functional groups had lower niche overlap in invaded than in removal plots, bees were more strongly affected than non-bee pollinators (Table 2, Fig. 3b). No significant trend was observed for honeybee niche overlap ($N = 6$, $V = 0$, $p\text{-value} = 0.250$) (Fig. 3b).

Effect of Hedysarum on interaction rewiring and the influence of honeybee foraging behaviour

In the experiment, $29.7 \pm 4.4\%$ of the interactions among shared species were observed both in invaded and removal paired plots. Meanwhile, $34.3 \pm 8.8\%$ and $36.7 \pm 4.4\%$ of the interactions were exclusive to invaded and removal plots, respectively. The contribution to exclusive interactions was similar across plant species, as for all of them at least half of their interactions were exclusive to either invaded or removal plots. In the case of pollinators, their contribution to exclusive interactions was highly diverse across species. Most pollinators were only involved in exclusive interactions; few of them showed high fidelity and were only involved in interactions shared between invaded plots. All the intermediate contributions were also observed.

Interaction rewiring was influenced by changes in foraging behaviour of the honeybee. That is, for a given plant species, the greater the difference between invaded and removal plots in honeybee visits, the higher the proportion of rewired interactions ($\text{Chisq} = 5.185$, $p\text{-value} = 0.023$) (Fig. 4). We observed this relationship irrespective of the functional group of the pollinators involved ($\text{Chisq} = 0.001$, $p\text{-value} = 0.983$). The interaction between the two explicative variables (i.e. functional group and changes in the foraging behaviour of the honeybee) was also not significant ($F = 0.203$, $p\text{-value} = 0.652$). Results did not qualitatively differ when singletons were excluded from the analysis.

Discussion

The important role of the honeybee in the integration of Hedysarum into the recipient plant-pollinator network

361

362 *Hedysarum* was well integrated into recipient plant-pollinator networks, being visited
363 by more species than the average for the resident plants. As pollinators seek to
364 optimize their foraging behaviour (Olesen et al. 2008; Petanidou et al. 2008; Lázaro
365 & Totland 2010), it might be advantageous for them to include abundant and high-
366 rewarding species like *Hedysarum* in their diet. Particularly, the honeybee played an
367 important role in such integration by performing most of the *Hedysarum* visits. The
368 honeybee, like other generalised, abundant and ubiquitous pollinators (Goulson
369 2003), is able to include many plant species in its diet, even non-natives (Memmott &
370 Waser 2002; Olesen et al. 2002; Morales & Aizen 2006; Padrón et al. 2009). In
371 addition, plants with flowers arranged in inflorescences like *Hedysarum*, might be
372 particularly attractive to the honeybee due to its flower constancy (Grüeter et al.
373 2011) and intense foraging behaviour with short flying distances between
374 consecutive visits (Gross 2001). This behaviour would also explain the high
375 percentage of honeybee visits observed in *Hedysarum*.

376 Despite *Hedysarum* integration, pollinator species visiting *Hedysarum* represented a
377 low percentage of the total pool of pollinator species (16.1%) in the community
378 compared to other non-native plant species invading other systems. Vilà et al. (2009)
379 studied five non-native plant species and found that they were visited by 31 to 50%
380 of the pollinator species in the community. However, the five non-native species
381 studied had open and non-restrictive flower morphologies, allowing a wider range of
382 pollinators to visit them. Meanwhile, non-natives with more restrictive flower
383 morphologies like legumes, filter pollinators according to their ability to access
384 rewards (Córdoba & Cocucci 2011). Non-native plants with restrictive flower
385 morphologies might face similar limitations in introduced areas than in their native

ranges. In fact, *Hedysarum* has a normalised degree in its native range comparable to that in the introduced communities on Menorca (Montero-Castaño et al. 2014).

The effect of Hedysarum on resident plant pollination was not influenced by similarity in flower morphology

The integration of *Hedysarum* into the recipient plant-pollinator networks did not have a larger effect on the pollination of plants exhibiting similar flower morphology to *Hedysarum* (i.e., those with papilionate flowers). Papilionate flowers are not accessible to all types of pollinators (Córdoba & Cocucci 2011). In fact, plants with papilionate flowers showed low visitation rates and normalised degree in both invaded and removal plots, obscuring the detection of an influence of *Hedysarum* presence. The expected influence of similarity in flower morphology on pollination between non-native and resident plants might be more easily observed for non-native plant species with minimally restrictive flower morphologies such as Composites, as reported by Morales & Traveset (2009).

Though *Hedysarum* did not have an overall effect on the average number or frequency of interactions involving resident plants (i.e. no quantitative effect), it affected the identity of some of those interactions (i.e. qualitative effect) as demonstrated by the lower proportion of honeybee visits observed in invaded networks. As has already been mentioned, *Hedysarum* was highly attractive to the honeybee, which reduced its presence on resident plants. Consequently, in invaded plots more resident floral resources were available to other pollinators. Honeybees can outcompete other pollinator species by depleting floral resources (Roubik 1983; Paini 2004; Valido, Rodríguez-Rodríguez & Jordano 2014) due to their abundance, generalised diet, communication skills, wide activity periods and systematic foraging

behaviour (Huryñ 1997, Gathmann & Tschardtke 2002, Steffan-Dewenter et al. 2002, Steffan-Dewenter & Kuhn 2003). Besides, honeybees have been shown to displace smaller species from flowers by physical disturbance (Gross & Mackay 1998).

Hedysarum affected the foraging behaviour of bees

As expected, *Hedysarum* affected the behaviour of pollinators and more noticeably those functionally closer to the honeybee. *Hedysarum* decreased the frequency (visitation rate), though not the number (normalised degree), of the interactions established by bees. Besides, *Hedysarum* altered the distribution of such interactions and more noticeably decreased the niche overlap of bees. The decrease in the visitation rate of bees in invaded plots seemed to be the result of an indirect effect of *Hedysarum* through the alteration of the foraging behaviour of the honeybee, as suggested by the opposite trends observed for both groups. As we have previously discussed, the honeybee can be a strong competitor for other pollinator species; especially for functionally redundant species. Resource partitioning in time and space reduces competition and allows species coexistence (Westphal, Steffan-Dewenter & Tschardtke 2006). However, such partitioning might be blurred among species belonging to the same functional group due to shared phenotypic and behavioural traits (e.g. tongue length, preferred resources, periods of maximum activity, etc.).

Despite the fact that the normalised degree was not altered, niche overlap of bees decreased in invaded plots. That is, in invaded plots there was a reorganisation of plant-pollinator interactions. In invaded plots, bees able to visit papilionate species (including *Hedysarum*) were more interconnected among them than with those bees

visiting non-papilionate species, and vice versa. Meanwhile, in removal plots, bees able to visit papilionate species also visited non-papilionate plants in order to fulfil their requirements, and consequently increased their niche overlap. The same trend was observed for non-bees. However, as only a few non-bee species are able to access papilionate flowers, the effect of *Hedysarum* was less significant.

The honeybee leads the interaction rewiring between invaded and removal networks

We observed that plant-pollinator interactions are highly plastic, suggesting a high lability of pollinators in resource use. Due to our experimental set up, we assume that paired invaded and removal networks share the same pollinator community. However, we cannot disregard potential differences in the abundance of some pollinators, mainly of the less mobile ones. Though that could slightly overestimate the lability of pollinators in resource use, the observed percentage of exclusive interactions to invaded or removal networks is consistent with the 30% found by other authors (Petanidou et al. 2008).

Plasticity of plant-pollinator interactions can have several non-exclusive explanations. First, it can be determined by the local floral environment. On the one hand, the abundance of floral rewards affects the probability of interactions (Vázquez et al. 2007), as stated by neutral theory (Hubbell 2001). On the other hand, neutrality can be diluted by magnet effects (Johnson et al. 2003; Molina-Montenegro, Badano & Cavieres 2008; Montero-Castaño & Vilà 2015) or conversely, by the monopolisation of visits by particular neighbours (Chittka, Gumbert & Kunze 1997; Kandori et al. 2009; Morales & Traveset 2009). Second, coexisting pollinator species can influence each other's foraging behaviour; especially, when abundant and/or highly competitive species are involved, such as the honeybee (Valido et al. 2014).

464 Though both drivers may be acting in this study case, we have evidence for the
465 second explanation, as the interaction rewiring was associated with the changes in
466 honeybee foraging behavior between invaded and removal networks.

467 Floral resources seem to be a limiting factor in the study system. Therefore, for a
468 given resident plant, the greater the differences in honeybee visits, the greater the
469 differences in floral resources available to other pollinators and the higher the
470 chance of finding different interactions between invaded and removal plots. This
471 result sheds some light on the conditions under which rewiring occurs, which is an
472 important topic ripe for empirical and theoretical consideration (Bourke & Alarcón
473 2011).

474 For instance, for a more realistic projection of the long-term response of plant
475 pollinator networks to the arrival or removal of species, modellers are incorporating
476 information on rewiring (e.g. Kaiser-Bunbury et al. 2010, Ramos-Jiliberto et al. 2012,
477 Valdovinos et al. 2013). In fact, the incorporation of topological plasticity based on
478 interaction rewiring, seems to increase network robustness to secondary extinctions
479 (Ramos-Jiliberto et al. 2012).

480 Even if network topological properties are conserved after invasion, the effects that
481 we have observed at the species level can have consequences on the reproduction
482 of resident plant and pollinator species.

483 Pollination visitation patterns of resident plants were not quantitatively altered, but
484 the identity of visitor species differed between invaded and removal plots: the
485 honeybee accounted for a higher proportion of visits to plants in removal than in
486 invaded plots. Visitor species differ in their pollination effectiveness in terms of pollen
487 removal, transport and deposition (Ne'eman *et al.* 2010) thus, implications on the
488 reproductive success of resident plants would be expected. In terms of pollen quality,
489 the honeybee is considered a low efficient pollinator as it usually increases

geitonogamy (Westerkamp 1991; Garibaldi *et al.* 2013). Therefore, higher reproductive success of resident plants would be expected in invaded plots. Regarding to pollinator species, their pollination patterns were quantitatively and qualitatively altered, what would also have consequences on their fitness. However, the impacts and underlying mechanisms of changes on food resources on pollinators' fitness are poorly understood, preventing us to advance any predictions. In fact, the literature on the effect of invasions on pollinators show diverse and even contradictory impacts on pollinators (Montero-Castaño & Vilà 2012; Litt *et al.* 2014).

Acknowledgements

We are very grateful to C. Coll, M. López, R. Molina, C. Montero, R. Ruiz, V. Sánchez and M. Vallés for field work assistance. We thank the staff of the Parc Natural Albufera des Grau and Observatori de la Sostenibilitat de Menorca (OBSAM) for their advising and logistic suport. P. Beuk, R. Castro, P. Grootaert, K. Heller, J.J. López Pérez, M.A. Marcos, A. Martínez, B. Merz, V. Michelsen, E. Nartshuk, J.L. Nieves Aldrey, F.J. Ortiz-Sánchez, A. Ricarte, M.C. Tolrá and H.P. Tschorsnig for insect indentification. We also thank J.M. Olesen for comments on the manuscript and J. Macaluso for English revision. EBD recieves financial support from the Spanish Ministerio de Economía y Competitividad, through the Severo Ochoa Program for Centres of Excellence in R+D+I (SEV-2012-0262). Funding was provided by the Spanish Ministerio de Ciencia e Innovación projects REDESIN (CGL 2007-61165-BOS), FLORMAS (CGL 2012-33801) and MONTES (CSD2008-00040), and by the EU 7FP project STEP

(244090-STEP-CP-FP). A. Montero-Castaño was supported by a JAE-Predoc fellowship.

Data Accessibility

DData deposited in the Dryad Digital Repository:
<http://dx.doi.org/10.5061/dryad.77rn2> (Montero-Castaño & Vilà, 2016)

References

- AEMET, Agencia Estatal de Meteorología. <http://www.aemet.es/es/elclima/datosclimatologicos/valoresclimatologicos> (accessed 27.09.15).
- Aizen, M.A., Morales, C.L. & Morales, J.M. (2008) Invasive mutualists erode native pollination webs. *PLoS BIOLOGY*, **6**, e31 doi:10.1371/journal.pbio.0060031.
- Armbruster, W.S. & Herzig, A.L. (1984) Partitioning and sharing of pollinators by four sympatric species of *Dalechampia* (Euphorbiaceae) in Panama. *Annals of the Missouri Botanical Garden*, **71**, 1–16.
- Bartomeus, I. (2013) Understanding linkage rules in plant-pollinator networks by using hierarchical models that incorporate pollinator detectability and plant traits. *PloS ONE*, **8**, e69200.
- Bartomeus, I., Vilà, M. & Santamaría, L. (2008) Contrasting effects of invasive plants in plant-pollinator networks. *Oecologia*, **155**, 761–770.

539 Bjerknes, A.L., Totland, Ø., Hegland, S.J. & Nielsen, A. (2007) Do alien plant
540 invasions really affect pollination success in native plant species? *Biological*
541 *Conservation*, **138**, 1–12.

542 Bommarco, R., Biesmeijer, J.C., Meyer, B., Potts, S.G., Poyry, J., Roberts, S.P.M.,
543 Steffan-Dewenter, I. & Ockinger, E. (2010) Dispersal capacity and diet breadth
544 modify the response of wild bees to habitat loss. *Proceedings of the Royal*
545 *Society B-Biological Sciences*, **277**, 2075–2082.

546 Burkle, L.A. & Alarcón, R. (2011) The future of plant-pollinator diversity:
547 Understanding interaction networks across time, space, and global change.
548 *American Journal of Botany*, **98**, 528–538.

549 Bustamante, J., Allés, A. & Espadas, M. (2007) *Alternativa de Tres Hojas O*
550 *Sementers*. Consell Insular de Menorca, Maó, Menorca.

551 Bustamante, J., Allés, A., Espadas, M. & Muñoz, J. (1998) *El Cultivo de La Zulla En*
552 *Menorca (IA): La Siembra*. Consell Insular de Menorca, Maó, Menorca.

553 Campbell, D.R., Bischoff, M., Lord, J.M. & Robertson, A.W. (2010) Flower color
554 influences insect visitation in alpine New Zealand. *Ecology*, **91**, 2638–2649.

555 Campos-Navarrete, M.J., Parra-Tabla, V., Ramos-Zapata, V., Díaz-Castelazo, C. &
556 Reyes-Novelo, E. (2013) Structure of plant-Hymenoptera networks in two
557 coastal shrub sites in Mexico. *Arthropod-Plant Interactions*, **7**, 607–617.

558 Carreras, D., Pons, C. & Canals, A. (2007) *Cartografia Digital de L'ocupació Del*
559 *Territori de Menorca - 2002*. Maó, Menorca.

560 Carstensen, D.W., Sabatino, M., Trøjelsgaard, K. & Morellato, L.P.C. (2014) Beta
561 diversity of plant-pollinator networks and the spatial turnover of pairwise
562 interactions. *PLoS ONE*, **9**.

563 Carvalheiro, L.G., Biesmeijer, J.C., Benadi, G., Fründ, J., Stang, M., Bartomeus, I.,
564 Kaiser-Bunbury, C.N., Baude, M., Gomes, S.I.F., Merckx, V., Baldock, K.C.R.,

565 Bennett, A.T.D., Boada, R., Bommarco, R., Cartar, R., Chacoff, N., Dänhardt, J.,
 566 Dicks, L. V., Dormann, C.F., Ekroos, J., Henson, K.S.E., Holzschuh, A., Junker,
 567 R.R., Lopezaraiza-Mikel, M., Memmott, J., Montero-Castaño, A., Nelson, I.L.,
 568 Petanidou, T., Power, E.F., Rundlöf, M., Smith, H.G., Stout, J.C., Temitope, K.,
 569 Tscharntke, T., Tscheulin, T., Vilà, M. & Kunin, W.E. (2014) The potential for
 570 indirect effects between co-flowering plants via shared pollinators depends on
 571 resource abundance, accessibility and relatedness. *Ecology Letters*, **17**, 1389–
 572 1399.

573 Chacoff, N.P., Vázquez, D.P., Lomascolo, S.B., Stevani, E.L., Dorado, J. & Padrón,
 574 B. (2012) Evaluating sampling completeness in a desert plant-pollinator
 575 network. *Journal of Animal Ecology*, **81**, 190–200.

576 Chittka, L., Gumbert, A. & Kunze, J. (1997) Foraging dynamics of bumble bees:
 577 Correlates of movements within and between plant species. *Behavioral Ecology*,
 578 **8**, 239–249.

579 Córdoba, S.A. & Cocucci, A.A. (2011) Flower power: its association with bee power
 580 and floral functional morphology in papilionate legumes. *Annals of Botany*, **108**,
 581 919–931.

582 Dicks, L. V, Corbet, S.A. & Pywell, R.F. (2002) Compartmentalization in plant-insect
 583 flower visitor webs. *Journal of Animal Ecology*, **71**, 32–43.

584 Encinas-Viso, F., Revilla, T. a. & Etienne, R.S. (2012) Phenology drives mutualistic
 585 network structure and diversity. *Ecology Letters*, **15**, 198–208.

586 Ferrero, V., Castro, S., Costa, J., Acuña, P., Navarro, L. & Loureiro, J. (2013) Effect
 587 of invader removal: pollinators stay but some native plants miss their new friend.
 588 *Biological Invasions*, **15**, 2347–2358.

589 Flores, F., Gutiérrez, J.C., López, J., Moreno, M.T. & Cubero, J.I. (1997) Multivariate
 590 analysis approach to evaluate a germplasm collection of *Hedysarum coronarium*
 591 L. *Genetic Resources and Crop Evolution*, **44**, 545–555.

592 Fraga, P., Mascaró, C., Carreras, D., García, O., Pallicer, X., Pons, M., Seoane, M.
 593 & Truyol, M. (2004) *Catàleg de La Flora Vascular de Menorca*. Institut Menorquí
 594 d'Estudis, Menorca.

595 Garibaldi, L.A., Steffan-Dewenter, I., Winfree, R., Aizen, M.A., Bommarco, R.,
 596 Cunningham, S. a., Kremen, C., Carvalheiro, L.G., Harder, L.D., Afik, O.,
 597 Bartomeus, I., Benjamin, F., Boreux, V., Cariveau, D., Chacoff, N., Dudenhöffer,
 598 J.H., Freitas, B.M., Ghazoul, J., Greenleaf, S., Hipólito, J., Holzschuh, A.,
 599 Howlett, B., Isaacs, R., Javorek, S.K., Kennedy, C.M., Krewenka, K., Krishnan,
 600 S., Mandelik, Y., Mayfield, M.M., Motzke, I., Munyuli, T., Nault, B.A., Otieno, M.,
 601 Petersen, J., Pisanty, G., Potts, S.G., Rader, R., Ricketts, T.H., Rundlöf, M.,
 602 Seymour, C.L., Schüepp, C., Szentgyörgyi, H., Taki, H., Tscharntke, T.,
 603 Vergara, C.H., Viana, B.F., Wanger, T.C., Westphal, C., Williams, N.M. & Klein,
 604 A.M. (2013) Wild pollinators enhance fruit set of crops regardless of honey bee
 605 abundance. *Science*, **339**, 1608–1611.

606 Gathmann, A. & Tscharntke, T. (2002) Foraging ranges of solitary bees. *Journal of*
 607 *Animal Ecology*, **71**, 757–764.

608 Gibson, M.R., Richardson, D.M. & Pauw, A. (2012) Can floral traits predict an
 609 invasive plant's impact on native plant-pollinator communities? *Journal of*
 610 *Ecology*, **100**, 1216–1223.

611 Goulson, D. (2003) Effects of introduced bees on native ecosystems. *Annual Review*
 612 *of Ecology Evolution and Systematics*, **34**, 1–26.

613 Greenleaf, S.S., Williams, N.M., Winfree, R. & Kremen, C. (2007) Bee foraging
 614 ranges and their relationship to body size. *Oecologia*, **153**, 589–596.

615 Gross, C.L. (2001) The effect of introduced honeybees on native bee visitation and
 616 fruit-set in *Dillwynia juniperina* (Fabaceae) in a fragmented ecosystem.
 617 *Biological Conservation*, **102**, 89–95.

618 Gross, C.L., Gorrell, L., Macdonald, M.J. & Fatemi, M. (2010) Honeybees facilitate
 619 the invasion of *Phyla canescens* (Verbenaceae) in Australia - no bees, no seed!
 620 *Weed Research*, **50**, 364–372.

621 Gross, C.L. & Mackay, D. (1998) Honeybees reduce fitness in the pioneer shrub
 622 *Melastoma affine* (Melastomataceae). *Biological Conservation*, **86**, 169–178.

623 Grüeter, C., Moore, H., Firmin, N., Helantera, H. & Ratnieks, F.L.W. (2011) Flower
 624 constancy in honey bee workers (*Apis mellifera*) depends on ecologically
 625 realistic rewards. *Journal of Experimental Biology*, **214**, 1397–1402.

626 Gutiérrez, J.C. (1982) *Hedysarum Coronarium L.: Estudio de Su Variabilidad Y*
 627 *Mejora Genética*. Universidad de Córdoba.

628 Hagen, M., Kissling, W.D., Rasmussen, C., De Aguiar, M. a M., Brown, L.E.,
 629 Carstensen, D.W., Alves-Dos-Santos, I., Dupont, Y.L., Edwards, F.K., Genini, J.,
 630 Guimarães, P.R., Jenkins, G.B., Jordano, P., Kaiser-Bunbury, C.N., Ledger,
 631 M.E., Maia, K.P., Marquitti, F.M.D., Mclaughlin, Ó., Morellato, L.P.C., O’Gorman,
 632 E.J., Trøjelsgaard, K., Tylianakis, J.M., Vidal, M.M., Woodward, G. & Olesen,
 633 J.M. (2012) *Biodiversity, Species Interactions and Ecological Networks in a*
 634 *Fragmented World*, 1st ed. Elsevier Ltd.

635 Hubbell, S. (2001) *The Unified Neutral Theory of Biodiversity and Biogeography*.
 636 Princeton University Press.

637 Huryn, V.M.B. (1997) Ecological impacts of introduced honey bees. *Quarterly*
 638 *Review of Biology*, **72**, 275–297.

639 Johnson, S.D., Peter, C.I., Nilsson, L.A. & Agren, J. (2003) Pollination success in a
 640 deceptive orchid is enhanced by co-occurring rewarding magnet plants.
 641 *Ecology*, **84**, 2919–2927.

642 Jordano, P., Bascompte, J. & Olesen, J.M. (2003) Invariant properties in
 643 coevolutionary networks of plant-animal interactions. *Ecology Letters*, **6**, 69–81.

644 Kaiser-Bunbury, C.N., Muff, S., Memmott, J., Muller, C.B. & Caflisch, A. (2010) The
 645 robustness of pollination networks to the loss of species and interactions: a
 646 quantitative approach incorporating pollinator behaviour. *Ecology Letters*, **13**,
 647 442–452.

648 Kandori, I., Hirao, T., Matsunaga, S. & Kurosaki, T. (2009) An invasive dandelion
 649 unilaterally reduces the reproduction of a native congener through competition
 650 for pollination. *Oecologia*, **159**, 559–569.

651 Lázaro, A., Hegland, S.J. & Totland, Ø. (2008) The relationships between floral traits
 652 and specificity of pollination systems in three Scandinavian plant communities.
 653 *Oecologia*, **157**, 249–257.

654 Lázaro, A. & Totland, Ø. (2010) Local floral composition and the behaviour of
 655 pollinators: attraction to and foraging within experimental patches. *Ecological*
 656 *Entomology*, **35**, 652–661.

657 Litt, A.R., Cord, E.E., Fulbright, T.E. & Schuster, G.L. (2014) Effects of Invasive
 658 Plants on Arthropods. *Conservation Biology*, **28**, 1532–1549.

659 Lopezaraiza-Mikel, M.E., Hayes, R.B., Whalley, M.R. & Memmott, J. (2007) The
 660 impact of an alien plant on a native plant-pollinator network: an experimental
 661 approach. *Ecology Letters*, **10**, 539–550.

662 Louati-Namouchi, I., Louati, M. & Chriki, A. (2000) Mating system and multiple
 663 paternity in *Hedysarum coronarium* L. (Fabaceae). *Agronomie*, **20**, 655–663.

664 Mawdsley, J.R. & Sithole, H. (2009) Diversity and abundance of insect visitors to
 665 flowers of trees and shrubs in a South African savannah. *African Journal of*
 666 *Ecology*, **48**, 691–698.

667 Memmott, J. & Waser, N.M. (2002) Integration of alien plants into a native flower-
 668 pollinator visitation web. *Proceedings of the Royal Society of London Series B-*
 669 *Biological Sciences*, **269**, 2395–2399.

670 Molina-Montenegro, M.A., Badano, E.I. & Cavieres, L.A. (2008) Positive interactions
 671 among plant species for pollinator service: assessing the “magnet species”
 672 concept with invasive species. *Oikos*, **117**, 1833–1839.

673 Montero-Castaño, A. & Vilà, M. (2012) Impact of landscape alteration and invasions
 674 on pollinators: a meta-analysis. *Journal of Ecology*, **100**, 884–893.

675 Montero-Castaño, A. & Vilà, M. (2015) Direct and indirect influence of non-native
 676 neighbours on pollination and fruit production of a native plant. *Plos One*, **10**,
 677 e0128595.

678 Montero-Castaño, A. & Vilà, M. Data from: Influence of the honeybee and trait
 679 similarity on the effect of a non-native plant on pollination and network rewiring.
 680 Dryad Digital Repository. <http://dx.doi.org/10.5061/dryad.77rn2>

681 Montero-Castaño, A., Vilà, M. & Ortiz-Sánchez, F.J. (2014) Pollination ecology of a
 682 plant in its native and introduced areas. *Acta Oecologica*, **56**, 1–9.

683 Montoya, J.M., Pimm, S.L. & Solé, R. (2006) Ecological networks and their fragility.
 684 *Nature*, **442**, 259–264.

685 Morales, C.L. & Aizen, M.A. (2006) Invasive mutualisms and the structure of plant-
 686 pollinator interactions in the temperate forests of north-west Patagonia,
 687 Argentina. *Journal of Ecology*, **94**, 171–180.

688 Morales, C.L. & Traveset, A. (2009) A meta-analysis of impacts of alien vs. native
689 plants on pollinator visitation and reproductive success of co-flowering native
690 plants. *Ecology Letters*, **12**, 716–728.

691 Mustajarvi, K., Siikamaki, P., Rytönen, S. & Lammi, A. (2001) Consequences of
692 plant population size and density for plant-pollinator interactions and plant
693 performance. *Journal of Ecology*, **89**, 80–87.

694 Ne'eman, G., Jurgens, A., Newstrom-Lloyd, L., Potts, S.G. & Dafni, A. (2010) A
695 framework for comparing pollinator performance: effectiveness and efficiency.
696 *Biological Reviews*, **85**, 435–451.

697 Nielsen, A. & Totland, Ø. (2014) Structural properties of mutualistic networks
698 withstand habitat degradation while species functional roles might change.
699 *Oikos*, **123**, 323–333.

700 Olesen, J.M., Bascompte, J., Dupont, Y.L., Elberling, H., Rasmussen, C. & Jordano,
701 P. (2011) Missing and forbidden links in mutualistic networks. *Proceedings*.
702 *Biological sciences / The Royal Society*, **278**, 725–32.

703 Olesen, J.M., Bascompte, J., Elberling, H. & Jordano, P. (2008) Temporal dynamics
704 in a pollination network. *Ecology*, **89**, 1573–1582.

705 Olesen, J.M., Eskildsen, L.I. & Venkatasamy, S. (2002) Invasion of pollination
706 networks on oceanic islands: importance of invader complexes and endemic
707 super generalists. *Diversity and Distributions*, **8**, 181–192.

708 Ortells, V. & Campos, X. (1983) *Els Anglicismes de Menorca*. Editorial Moll, Palma
709 de Mallorca.

710 Osborne, J.L., Martin, A.P., Carreck, N.L., Swain, J.L., Knight, M.E., Goulson, D.,
711 Hale, R.J. & Sanderson, R.A. (2008) Bumblebee flight distances in relation to
712 the forage landscape. *Journal of Animal Ecology*, **77**, 406–415.

713 Padrón, B., Traveset, A., Biedenweg, T., Díaz, D., Nogales, M. & Olesen, J.M.
 714 (2009) Impact of alien plants invaders on pollination networks in two
 715 archipelagos. *Plos One*, **4**, e6275.

716 Paini, D.R. (2004) Impact of the introduced honey bee (*Apis mellifera*)
 717 (Hymenoptera : Apidae) on native bees: A review. *Austral Ecology*, **29**, 399–
 718 407.

719 Petanidou, T., Kallimanis, A.S., Tzanopoulos, J., Sgardelis, S.P. & Pantis, J.D.
 720 (2008) Long-term observation of a pollination network: fluctuation in species and
 721 interactions, relative invariance of network structure and implications for
 722 estimates of specialization. *Ecology Letters*, **11**, 564–575.

723 Poisot, T., Canard, E., Mouillot, D., Mouquet, N., Gravel, D. & Jordan, F. (2012) The
 724 dissimilarity of species interaction networks. *Ecology letters*, **15**, 1353–61.

725 Pyšek, P., Richardson, D.M., Rejmanek, M., Webster, G.L., Williamson, M. &
 726 Kirschner, J. (2004) Alien plants in checklists and floras: towards better
 727 communication between taxonomists and ecologists. *Taxon*, **53**, 131–143.

728 R Development Core Team. (2014) R: A language and environment for statistical
 729 computing.

730 Ramos-Jiliberto, R., Valdovinos, F.S., Moisset de Espanés, P. & Flores, J.D. (2012)
 731 Topological plasticity increases robustness of mutualistic networks. *Journal of*
 732 *Animal Ecology*, **81**, 896–904.

733 Rodríguez-Riño, T., Ortega-Olivencia, A. & Devesa, J.A. (1999) *Biología Floral En*
 734 *Fabaceae* (ed S Castroviejo). Consejo Superior de INvestigaciones Científicas,
 735 Madrid.

736 Roubik, D.W. (1980) Foraging behavior of competing Africanized honeybees and
 737 stingless bees. *Ecology*, **61**, 836–845.

738 Roubik, D.W. (1983) Experimental community studies: time-series tests of
739 competition between African and Neotropical bees. *Ecology*, **64**, 971–978.

740 Santamaría, L. & Rodríguez-Gironés, M.A. (2007) Linkage rules for plant – pollinator
741 networks : Trait complementarity or exploitation barriers ? *PLoS BIOLOGY*, **5**,
742 e31. doi:101371/journal.pbio.0050031.

743 Satta, A., Acciaro, M., Floris, I., Lentini, A. & Sulas, L. (2000) Insect pollination of
744 sulla (*Hedysarum coronarium* L.) and its effect on seed production in a
745 Mediterranean environment. *Legumes for Mediterranean forage crops, pastures*
746 *and alternative uses* (ed L. Sulas), pp. 373–377. CIHEAM-IAMZ, Zaragoza.

747 Simberloff, D., Martin, J.L., Genovesi, P., Maris, V., Wardle, D. a., Aronson, J.,
748 Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R.,
749 Tabacchi, E. & Vilà, M. (2013) Impacts of biological invasions: What's what and
750 the way forward. *Trends in Ecology and Evolution*, **28**, 58–66.

751 Stang, M., Klinkhamer, P.G.L. & van der Meijden, E. (2006) Size constraints and
752 flower abundance determine the number of interactions in a plant-flower visitor
753 web. *Oikos*, **112**, 111–121.

754 Steffan-Dewenter, I. & Kuhn, A. (2003) Honeybee foraging in differentially structured
755 landscapes. *Proceedings. Biological sciences / The Royal Society*, **270**, 569–
756 575.

757 Steffan-Dewenter, I., Munzenberg, U., Burger, C., Thies, C. & Tscharntke, T. (2002)
758 Scale-dependent effects of landscape context on three pollinator guilds.
759 *Ecology*, **83**, 1421–1432.

760 Stout, J.C., Kells, A.R. & Goulson, D. (2002) Pollination of the invasive exotic shrub
761 *Lupinus arboreus* (Fabaceae) by introduced bees in Tasmania. *Biological*
762 *Conservation*, **106**, 425–434.

763 Sulas, L., Stangoni, A.P., Re, G.A. & Ledda, L. (2000) Growing cycle of *Hedysarum*
764 *coronarium* L. (sulla): relationship between plant density, stem length, forage
765 yield and phytomass partitioning. *Legumes for Mediterranean forage crops,*
766 *pastures and alternative uses* (ed L. Sulas), pp. 147–151. CIHEAM-IAMZ,
767 Zaragoza.

768 Talavera, S., Herrera, J., Arroyo, J., Ortiz, P.L. & Devesa, J.A. (1988) Estudio de la
769 flora apícola de Andalucía occidental. *Lagascalia*, **15**, 567–591.

770 Trojelsgaard, K., Jordano, P., Carstensen, D.W. & Olesen, J.M. (2015) Geographical
771 variation in mutualistic networks: similarity, turnover and partner fidelity.
772 *Proceedings of the Royal Society B: Biological Sciences*, **282**, 20142925–
773 20142925.

774 Tylianakis, J.M., Laliberte, E., Nielsen, A. & Bascompte, J. (2010) Conservation of
775 species interaction networks. *Biological Conservation*, **143**, 2270–2279.

776 Valdovinos, F.S., Moisset de Espanés, P., Flores, J.D. & Ramos-Jiliberto, R. (2013)
777 Adaptative foraging allows the maintenance of biodiversity of pollination
778 networks. *Oikos*, **118**, 907–917.

779 Valido, A., Rodríguez-Rodríguez, M.C. & Jordano, P. (2014) Impacto de la
780 introducción de la abeja doméstica (*Apis mellifera*, Apidae) en el Parque
781 Nacional del Teide (Tenerife, Islas Canarias). *Ecosistemas*, **23**, 58–66.

782 Vázquez, D.P., Melian, C.J., Williams, N.M., Bluthgen, N., Krasnov, B.R. & Poulin, R.
783 (2007) Species abundance and asymmetric interaction strength in ecological
784 networks. *Oikos*, **116**, 1120–1127.

785 Vilà, M., Bartomeus, I., Dietzsch, A.C., Petanidou, T., Steffan-Dewenter, I., Stout,
786 J.C. & Tscheulin, T. (2009) Invasive plant integration into native plant-pollinator
787 networks across Europe. *Proceedings of the Royal Society B-Biological*
788 *Sciences*, **276**, 3887–3893.

- Warton, D.I. & Hui, F.K.C. (2011) The arcsine is asinine: the analysis of proportions in ecology. *Ecology*, **92**, 3–10.
- Westerkamp, C. (1991) Honeybees are poor pollinators - why? *Plant Systematics and Evolution*, **177**, 71–75.
- Westphal, C., Steffan-Dewenter, I. & Tscharntke, T. (2006) Bumblebees experience landscapes at different spatial scales: possible implications for coexistence. *Oecologia*, **149**, 289–300.
- Yagoubi, N. & Chriki, A. (2000) Estimation of mating system parameters in *Hedysarum coronarium* L. (Leguminosae, Fabaceae). *Agronomie*, **20**, 933–942.

Supporting Information

Additional supporting information may be found in the online version of this article.

Appendix S1. Plant-pollinator networks.

Table S1.1. Flowering plant species present in the study plots.

Table S1.2. Pollinator species observed visiting flowering plants

Appendix S2. Rarefaction curves.

Figure legends

Figure 1. Non-native *Hedysarum* (left) and detail of an inflorescence being visited by a honeybee (right). Photographs by A. Montero-Castaño.

Figure 2. Effect of *Hedysarum* on the pollination patterns of resident plants. Mean $\pm$ 95%CI (a) visits (standardised per flower and hour and log-transformed), (b) normalised degree, (c) niche overlap and (d) proportion of honeybee visits to *Hedysarum* and resident papilionate and non-papilionate species in invaded and removal plots. Upper case symbols represent the significance levels for differences between invaded and removal plots according to the *post hoc* multilevel comparisons conducted for the models: ** $p < 0.01$, n.s = no significant.

Figure 3. Effect of *Hedysarum* on the foraging behaviour of pollinators. Mean $\pm$ 95%CI (a) visits (standardised per flower and hour and log-transformed), (b) normalised degree, (b) and (c) niche overlap of the honeybee, bees and non-bees in invaded and removal plots. Upper case symbols represent the significance levels for differences between invaded and removal plots; in the case of bees and non-bees functional groups, according to the *post hoc* multilevel comparisons conducted for the models: * $p\text{-value} = 0.05$, ** $p < 0.01$, *** $p < 0.001$, n.s = no significant.

Figure 4. Relationship between interaction rewiring and changes in the foraging behavior of the honeybee. (a) Positive relationship between these two variables and (b) its bipartite network representation. Changes in the foraging behaviour of the honeybee are defined as the difference in the proportion of visits to a given plant species that the honeybee achieved in invaded and removal paired networks.

835 Shared species of the three pairs of networks studied are pooled and represented
836 together. In (b) the size of the circles representing plant species indicates differences
837 in the proportion of honeybee visits between invaded and removal plots. Grey lines
838 represent rewired interactions (whether exclusive of invaded or removal plots) and
839 black lines represent non-rewired interactions (i.e. those observed in both invaded
840 and removal plots).

841 **Tables**

842

843 **Table 1.** Response variables estimated for both plant (p) and animal pollinator (a) species.

Parameter	Symbol	Definition
Visits	V_p	Number of visits a plant species receives
	V_a	Number of visits a pollinator species makes
Normalised degree	D_p	Proportion of the total number of pollinator species a particular plant species is visited by
	D_a	Proportion of the total number of plant species a particular pollinator species visits
Niche overlap	NO_p	Proportion of the total number of plant species a particular plant species shares pollinators with
	NO_a	Proportion of the total number of pollinator species a particular pollinator species shares visited plants with
Interaction rewiring		Proportion of the interactions involving plant species shared between paired invaded and removal plots that are exclusive to invaded or to removal plots

Table 2. Likelihood Ratio Tests (LRT) of the effects of treatment and flower morphology and their interaction on visitation rates, normalised degree and niche overlap for resident plant and pollinator communities. Significance levels:

* p-value ≈ 0.05 , * < 0.05, ** < 0.01, *** < 0.001.

Resident community	N	Response variable	Explicative variable	p-value
Plants	54	Visits (V_p)	Treatment	0.732
			Flower morphology	< 0.001 ***
			Treatment*Flower morphology	0.854
	54	Normalised degree (D_p)	Treatment	0.892
			Flower morphology	< 0.001 ***
			Treatment*Flower morphology	0.973
	54	Niche overlap (NO_p)	Treatment	0.187
			Flower morphology	< 0.001 ***
			Treatment*Flower morphology	0.562
	54	Proportion honeybee visits	Treatment	< 0.001 ***
			Flower morphology	0.845
			Treatment*Flower morphology	0.403
Pollinators	178	Visits (V_a)	Treatment	0.005 **
			Functional group	0.988
			Treatment*Functional group	0.405
	191	Normalised degree (D_a)	Treatment	0.828
			Functional group	0.140
			Treatment*Functional group	0.929
	191	Niche overlap (NO_a)	Treatment	< 0.001 ***
			Functional group	< 0.001 ***
			Treatment*Functional group	0.002 **