Elsevier Editorial System(tm) for Peptides
Manuscript Draft

Manuscript Number: PEPTIDES-D-07-00332R1

Title: Identification of a tachykinin-related peptide with orexigenic properties in the German cockroach

Article Type: Research Paper

Keywords: Tachykinin; *Blattella germanica*; food consumption; myotropic peptide.

Corresponding Author: Dr. José L. Maestro, Ph. D.

Corresponding Author's Institution: IBMB-CSIC

First Author: Núria Pascual

Order of Authors: Núria Pascual; José L. Maestro; Cristina Chiva; David Andreu; Xavier Bellés

Abstract: A number of evidences suggest that tachykinin-related peptides (TRPs) of insects can stimulate food consumption after being released from the midgut to the haemolymph. The idea of the present work has been to test this hypothesis in the anautogenous cockroach *Blattella germanica*. Firstly, we have identified the peptide LemTRP-1 (APSGFLGVR-NH₂) from brain extracts, by means of an ELISA developed with a polyclonal antibody against this peptide. ELISA studies have also shown that, whereas brain LemTRP-1 levels were fairly constant, midgut levels increase to a maximum on day 3 after adult emergence, falling thereafter until the end of the gonadotrophic cycle. Interestingly, maximum values of food consumption are concomitant with the decrease of LemTRP-1 immunoreactivity in the midgut. Furthermore, starvation decreases LemTRP-1 immunoreactivity in midgut, whereas in the haemolymph it increases. Finally, injection of synthetic LemTRP-1 to adult females significantly stimulates food consumption. The whole observations suggest that LemTRP-1 is released from the midgut to the haemolymph when sustained food consumption is required to maintain vitellogenesis at the highest levels, and that LemTRP-1 in the haemolymph stimulates food consumption in these days.

Suggested Reviewers:

Opposed Reviewers:

**Identification of a tachykinin-related peptide with orexigenic properties in the
German cockroach**

Núria Pascual^a, José L. Maestro^{a*}, Cristina Chiva^b, David Andreu^b, Xavier Bellés^{a,*}

^aDepartment of Physiology and Molecular Biodiversity, Institut de Biologia Molecular
de Barcelona (CSIC), Jordi Girona 18-26, 08034 Barcelona, Spain.

^bDepartment of Experimental and Health Sciences, Universitat Pompeu Fabra, Doctor
Aiguader 80, 08003 Barcelona, Spain.

*Co-corresponding authors. Tel.: +34-934006124; fax: +34-932045904. (X. Bellés);
Tel.: +34-934006135; fax: +34-932045904. (J.L. Maestro).

E-mail addresses: xbragr@cid.csic.es (X. Bellés); jmgagr@ibmb.csic.es (J.L. Maestro)

Abstract

A number of evidences suggest that tachykinin-related peptides (TRPs) of insects can stimulate food consumption after being released from the midgut to the haemolymph. The idea of the present work has been to test this hypothesis in the anautogenous cockroach *Blattella germanica*. Firstly, we have identified the peptide LemTRP-1 (APSGFLGVR-NH₂) from brain extracts, by means of an ELISA developed with a polyclonal antibody against this peptide. ELISA studies have also shown that, whereas brain LemTRP-1 levels were fairly constant, midgut levels increase to a maximum on day 3 after adult emergence, falling thereafter until the end of the gonadotrophic cycle. Interestingly, maximum values of food consumption are concomitant with the decrease of LemTRP-1 immunoreactivity in the midgut. Furthermore, starvation decreases LemTRP-1 immunoreactivity in midgut, whereas in the haemolymph it increases. Finally, injection of synthetic LemTRP-1 to adult females significantly stimulates food consumption. The whole observations suggest that LemTRP-1 is released from the midgut to the haemolymph when sustained food consumption is required to maintain vitellogenesis at the highest levels, and that LemTRP-1 in the haemolymph stimulates food consumption in these days.

Keywords: Tachykinin, *Blattella germanica*, food consumption, myotropic peptide.

1. Introduction

Tachykinin-Related Peptides (TRPs) constitute a family of invertebrate peptides with a characteristic carboxy-terminal sequence $\text{FX}_1\text{GX}_2\text{R-NH}_2$. The name comes from their relative sequence similarity with vertebrate tachykinins, which show a conserved C-terminal sequence FXGLM-NH_2 . In addition, TRPs and vertebrate tachykinins share other characteristics, as their occurrence in both nervous system and gut tissues, and their stimulatory activity of gut musculature contractions [20].

The first peptides belonging to the TRP family (Lom-TK I and II) were purified from brain-corpora cardiaca-corpora allata-suboesophageal ganglion extracts of the locust, *Locusta migratoria*, by monitoring their myotropic activity on cockroach hindgut [21]. Since then, peptides belonging to TRP family have been identified in insects belonging to Orthoptera, Diptera, Dictyoptera and Hymenoptera orders [20,23]. Furthermore, cDNAs encoding TRP precursors have been cloned and sequenced in the fruit fly *Drosophila melanogaster* [22], the mosquito *Anopheles gambiae* [18], the honeybee *Apis mellifera* [23] and the cockroaches *Leucophaea maderae* and *Periplaneta americana* [17]. In all cases, the cDNA sequence confirmed the identity of the previously reported peptides.

Immunocytochemical studies in cockroaches have revealed the occurrence of TRP immunoreactivity in interneurons of the central nervous system, in the stomatogastric nervous system, in processes to the corpora cardiaca glandular lobe, in nerves innervating different gut areas and in midgut endocrine cells [10,13]. This wide distribution has been observed in all the species tested [14], and suggests that TRPs have multiple functions. Indeed, a remarkable variety of activities has been reported for these peptides in insects, including stimulation of visceral and skeletal muscle contractions, induction of adipokinetic hormone release by corpora cardiaca, neuronal depolarization, stimulation of urine production, and induction of pheromone biosynthesis [6,14,20].

Furthermore, it has been demonstrated that the midgut of *L. maderae* and *L. migratoria* incubated in vitro release TRPs in response to an increase of K^+ levels in the bathing solution [25]. In *L. migratoria*, starvation increases the concentration of TRP-immunoreactive material in the haemolymph, concomitantly with a decrease in the midgut, which suggested that TRPs are released as hormones from the midgut, and that this release could be linked to the nutritional status [25]. These results make TRPs good

1
2
3
4 candidates for being tested as orexigenic factors, under the hypothesis that its release in
5 starved specimens might stimulate food consumption.

6
7 A good model to test this hypothesis would be the German cockroach, *Blattella*
8 *germanica*, given that it is anautogenous and show a well defined feeding cycle
9 paralleling that of vitellogenesis [16]. The feeding cycle suggests that food intake is
10 finely regulated, and possible regulatory mechanisms have been already reported. Thus,
11 the peptide perisulfakinin has been identified as a putative satiety factor [8], and a
12 number of YXFGL-NH₂ allatostatins, W²W⁹-amide myoinhibitory peptides and
13 leucomyosuppressin have been shown to inhibit food intake in this cockroach [1-3].
14 Nevertheless, no information about factors stimulating food intake has been reported.

15
16 The aim of the present work has been to identify TRPs in *B. germanica*, and to
17 study whether they may play a stimulatory role in the regulation of food intake. The
18 reference peptide to search native TRPs in *B. germanica* was LemTRP-1
19 (APSGFLGVR-NH₂), which had been already identified in the cockroaches *L. maderae*
20 and *P. americana* [11,12,17].

21 22 23 24 25 26 27 28 29 30 31 **2. Material and methods**

32 33 34 35 *2.1. Insect rearing*

36
37 Adult females of *B. germanica* (L.) were obtained from a colony reared on dog
38 chow and water, at 30 ± 1°C and 60-70 % r.h. Freshly moulted adult virgin females
39 were isolated and used at the appropriate physiological ages within the first
40 gonadotrophic cycle. Physiological age was assessed by measuring the basal oocyte
41 length [4]. For starvation experiments, animals were supplied only with water since the
42 imaginal moult.
43
44
45
46
47

48 49 50 *2.2. Synthesis of peptides and conjugates*

51
52 Peptides LemTRP-1: APSGFLGVR-NH₂, LemTRP-2:
53 APEESPKRAPSGFLGVR-NH₂, LemTRP-4: APSGFMGMR-NH₂ and LemTRP-5:
54 APAMGFQGVR-NH₂ [11] were synthesized using standard Fmoc solid phase methods
55 [5]. The identity and purity (ca. 90%) of each peptide were assessed by amino acid
56 analysis, matrix-assisted laser desorption ionization time of flight (MALDI-TOF) mass
57
58
59
60
61
62
63
64
65

spectra and HPLC. To raise antibodies against LemTRP-1, the peptide was synthesized with its N-terminus extended by two residues of 2-aminohexanoic acid and conjugated to keyhole-limpet-haemocyanin (KLH) (Sigma, St. Louis, MO, USA) according to [19]. For ELISA plate coating, LemTRP-1 was conjugated to bovine serum albumin (BSA) (Sigma), using glutaraldehyde [24]. The conjugates were dialyzed, lyophilized and stored at -20°C.

2.3. Antibody production and titre test

Three male white New Zealand rabbits were used to raise antibodies using LemTRP-1-KLH as immunogen. Rabbits were injected subcutaneously with 100 µg of peptide, in the conjugated form, diluted in 500 µl of water emulsified with 500 µl of Freund's complete adjuvant (Sigma) on days 0 and 7, and with incomplete adjuvant on day 14. Blood samples were obtained on day 21. Rabbits were boosted again once a month using the same dose and incomplete adjuvant, and serum was obtained one week after each booster injection, during 6 months. Serum was added with 0.1% thimerosal (Serva, Heidelberg, Germany) and stored at -20°C. The titre of serum from each rabbit was determined by measuring the binding of serial dilutions to microtiter plates coated with 1 µg/ml of LemTRP1-BSA. A two-dimensional titration protocol was used for the screening and determination of the optimum concentration of both coating antigens and antisera to be used later in the competitive experiments [7].

2.4. ELISA method

LemTRP-1-BSA at a concentration of 0.15 µg/ml in 0.1 M carbonate-bicarbonate buffer (pH 9.6) was used for coating polystyrene 96 wells microtiter plates (Nunc Maxisorp, Roskilde, Denmark), in a volume of 100 µl/well, and incubated overnight at 4°C. The plates were washed five times with PBST buffer (0.2 M, phosphate-buffered saline solution containing 0.05% Tween 20, pH 7.4). The plate was blocked with 1% polyvinylpyrrolidone (Sigma) in PBST buffer. After 1h, plates were washed again as described above. The immunological reaction was initiated by adding dilutions of the samples or standard peptide analyte in PBST buffer (from 10^{-6} M to 10^{-10} M) in volume of 50 µl/well followed by 50 µl/well of the antibody previously diluted 1/30,000 in PBST buffer (final dilution in the well: 1/60,000). After incubation at room

temperature for 2 h, the plates were washed as described above, and 100 μ l/well of a 1/6,000 diluted goat antirabbit IgG peroxidase conjugated (Sigma) solution were added. After 1 h incubation and a washing step, 100 μ l/well of substrate solution were added and incubated in the dark with gentle shaking. Substrate solution was prepared with 12.5 ml of citrate buffer (pH=5), 200 μ l of 0.6% 3,3',5,5'-tetramethylbenzidine in dimethyl sulfoxide and 50 μ l of 1% H₂O₂. Reaction was stopped by adding 50 μ l/well of 2N H₂SO₄. Absorbance was read at 450 nm with a Titertek Multiscan Plus MKII spectrophotometer (Labsystems, Helsinki, Finland). The calibration curves were analyzed using a four parameter logistic equation.

2.5. HPLC procedures

A total of 1440 brains from 5- to 7-day-old females were dissected out under saline solution (NaCl 9 g/l; KCl 0.2 g/l; NaHCO₃ 0.2 g/l; CaCl₂ 0.2 g/l), and homogenized in methanol/water/acetic acid (87/8/5, v/v/v), using a mechanical homogenizer Eurostar digital (Ika labortechnik, Staufen, Germany) designed for 1.5 ml tubes. After centrifugation (8000xg for 10 min at 4°C), the supernatant was collected and stored at -20 °C until use. Brain extract was processed in five consecutive HPLC steps. LemTRP-1-immunoreactive fractions were detected using the above described ELISA and used for further purification. Steps 1, 2 and 3 were carried out with a Merck-Hitachi (Darmstadt, Germany) low-pressure system, L-6200A pump with a L-4200 UV-VIS detector. Steps 4 and 5 were carried out with a Waters (Milford, MA, USA) low-pressure system, 626 pump with a 600S controller and 996 PDA detector.

Step 1: Waters DeltaPak semi-preparative C₁₈ column (300x7.8 mm, 15 μ m, 300 Å). Linear gradient of CH₃CN/0.1% TFA. The gradient change was 1.67%/min and the flow rate 1.5 ml/min. HPLC fractions were analysed with the ELISA described above, which revealed immunoreactive material in fractions eluting between 20.0% and 23.3% CH₃CN.

Step 2: Column and solvents as in step 1. The gradient change was 0.5%/min and the flow rate 1.5 ml/min. Immunoreactive material was detected in fractions eluting between 21.0% and 22.5% CH₃CN.

Step 3: Merck LiChroCART C₁₈ column (125x4 mm, 5 μ m, 100 Å). Linear gradient of CH₃CN/0.1% TFA. The gradient change was 0.25%/min and the flow rate

1
2
3
4 1.5 ml/min. Immunoreactive material was detected in fractions eluting between 18.5%
5 and 19.2% CH₃CN.
6

7 Step 4: Waters DeltaPak C₁₈ column (150x2 mm, 5 µm, 300 Å). Linear gradient
8 of CH₃CN/0.05% TFA. The gradient change was 0.25%/min and the flow rate 0.2
9 ml/min. Immunoreactive material was detected in fractions eluting between 16.4% and
10 17.0% CH₃CN.
11
12
13

14 Step 5: Column and solvents as in step 4. The gradient change was 0.16%/min
15 and the flow rate 0.2 ml/min. A single peak of immunoreactive material eluted at 21.1%
16 CH₃CN.
17
18

19 For the chromatographic separation of haemolymph from 3-day-old fed and
20 starved females, the HPLC system and column described for steps 1 and 2 of the brain
21 extract separation were used. Samples of 155 µl and 229 µl of haemolymph from fed
22 and starved females, respectively, were used in the HPLC separation. A linear gradient
23 of CH₃CN/0.1% TFA with a linear change of 1%/min and a flow rate of 1.5 ml/min was
24 carried out, and HPLC fractions were analysed using the ELISA described above.
25
26
27
28
29
30

31 2.6. MS and sequencing 32 33

34 An aliquot of the purified LemTRP-1-immunoreactive peak was analysed using
35 an Applied Biosystems (Foster City, CA, USA) Voyager DE-RP MALDI-TOF mass
36 spectrometer. The amino-acid sequence of the purified factor was determined by Edman
37 degradation using an Applied Biosystems Procise instrument.
38
39
40
41
42

43 2.7. Myotropic assay 44 45

46 LemTRP-1 was tested on foregut and hindgut of *B. germanica* females prepared
47 in a standard organ bath as previously described [8]. The composition of the bath was:
48 154 mM NaCl, 2.7 mM KCl, 1.8 mM CaCl₂, 22mM glucose and 5 mM HEPES, pH 6.8.
49 An FSG-01 transducer (Experimetria, Budapest, Hungary) was used for isometric
50 recording. Myotropic activity was calculated as the difference of the mean of the force
51 produced by the tissue one min after and one min before the treatment.
52
53
54
55
56
57

58 2.8. Sampling for physiological observations 59 60 61 62 63 64 65

1
2
3
4 Brains and midguts were dissected under saline solution and homogenized in
5 PBS. Then, samples were boiled for 5 min and centrifuged at 16000xg for 10 min.
6 Supernatants were collected and the pellets were re-extracted, centrifuged again and the
7 supernatants were pooled together with the previous ones, lyophilized and stored at -
8 20°C until use. Haemolymph samples to determine peptide concentration by HPLC
9 separation and ELISA were obtained by cutting off one leg of the animal and applying
10 gentle pressure to the abdomen. A pool from 110 and 150 fed and starved females
11 respectively was used for the determination. The haemolymph was collected and diluted
12 in methanol/water/trifluoroacetic acid (90/10/0.1, v/v/v). After evaporation of organic
13 solvent and lyophilization, samples were stored at -20°C until use.
14
15
16
17
18
19
20
21

22 *2.9. Measurement of food consumption*

23
24
25
26 Food consumption was measured as reported previously [16], with some
27 modifications. Briefly, individual specimens were provided with a weighed food (dog
28 chow) portion, and 24 h later the remaining food was dried out in an oven and weighed
29 again. The water lost by evaporation of food placed in a control box, containing only the
30 water vial, was used as a correction factor. In this case, instead of using the oviposition
31 time for realigning the results to a 8 days scale as in [16], food consumption was
32 quantified in chronological periods of 24 h throughout the gonadotrophic cycle.
33
34
35
36
37
38
39

40 *2.10. Feeding bioassay*

41
42
43 The feeding bioassay was carried out as previously reported [8], although with
44 some modifications. Freshly ecdysed adult females were fed with carrot for 24 h, after
45 which they were injected with saline or with the testing peptide and provided again
46 carrot ad libitum. After 5 h, the gut was dissected out and the foregut, midgut and
47 hindgut regions were separated by sectioning just before the gastric caeca and at the
48 level of Malpighian tubules evagination. Each gut region together with its content was
49 extracted with methanol, and carotenoid concentration was estimated by
50 spectrophotometric measurement of the absorbance at 450 nm. Total weight of carrot
51 ingested was quantified by interpolation on a standard curve constructed using
52 methanolic extracts of increasing amounts of lyophilized carrot.
53
54
55
56
57
58
59
60
61
62
63
64
65

3. Results

3.1. ELISA characterization

Figure 1 shows the sensitivity curve obtained with different concentrations of LemTRP-1, and the antiserum used at a dilution of 1:60,000. The ED₅₀ (concentration required for 50% displacement of binding to conjugate) was 1.15×10^{-7} μ mol.

In order to study the selectivity of the antibody, cross-reactivity assays using anti-LemTRP-1 serum with synthetic LemTRP-2, -4 and -5 were carried out (sequences indicated in 2.2). Results indicate that antiserum shows high cross-reactivity with LemTRP-2, comparable to that with LemTRP-1. Conversely, cross-reaction with LemTRP-4 and -5 is very low (Fig. 1). The limit of detection of standard LemTRP-1 was around 10^{-10} M.

3.2. Purification and identification of LemTRP-1

A crude extract of 1440 brains from adult females of *B. germanica* was processed through five consecutive HPLC steps, using LemTRP-1 immunoreactivity for monitoring the fractions of interest. In the last step, an immunoreactive homogeneous peak resulted in the nonapeptide APSGFLGVR-NH₂, identical to LemTRP-1 previously identified from the cockroaches *L. maderae* and *P. americana*. The final yield was approximately 110 pmol. We presume that the C-terminus was amidated on the basis of the MS analysis, which showed a molecular mass of 902.52 (MH⁺). The identification was further confirmed by the coelution in HPLC of the native peptide with synthetic APSGFLGVR-NH₂.

3.3. Myostimulatory activities on gut tissues

Given that tachykinins and TRPs have been described as myotropic peptides in both vertebrates and insects, we first assessed this biological effect in our insect model. Thus, we studied the effects of LemTRP-1 on foregut and hindgut motility in *B. germanica*, using a standard organ bath system. Results (Fig. 2) showed that LemTRP-1 elicits myostimulatory activity on both gut regions, with an ED₅₀ for the force increase within the nanomolar range.

3.4. *LemTRP-1* levels and feeding rhythm

LemTRP-1 levels were quantified by ELISA in extracts of individual brains and midguts of virgin females during the first gonadotrophic cycle. The profile of LemTRP-1 immunoreactivity in the brain (Fig. 3) shows fairly constant levels (between 40 and 60 pg/brain equivalent), throughout the whole period. In contrast, LemTRP-1 levels in the midgut starts to rise from day 1, reaches the maximal value on day 3 (ca. 70 pg/midgut equivalent), and decreases thereafter, showing relative low levels until day 7, when chorionation and oviposition occurs (Fig. 3).

Quantification of food ingestion during the first gonadotrophic cycle indicates that *B. germanica* females undergo a food consumption cycle, with maximum values (ca. 10 mg/24 h) observed between days 3 and 6, decreasing thereafter until the time of oviposition and ootheca formation (Fig. 3).

3.5. *Effect of starvation and feeding on LemTRP-1* levels

To study the effect of starvation, brain, midgut and haemolymph from fed and starved 3-day-old *B. germanica* females were extracted, and LemTRP-1 immunoreactivity was measured by ELISA. Results show no differences in LemTRP-1 immunoreactivity between brains from fed and starved females (Fig. 4A). Conversely, 57% significant reduction in LemTRP-1 levels was observed in midgut from starved females when compared to the midgut of fed controls (Fig. 4A).

Haemolymph samples were separated by HPLC before being submitted to ELISA quantification. A single peak of LemTRP-1 immunoreactive material, coeluting with the synthetic LemTRP-1, was detected in the HPLC separation of haemolymph from both fed and starved 3-day-old females. The amounts of LemTRP-1 equivalents per μ l of haemolymph were 0.266 pg and 0.496 pg for fed and starved females, respectively (Fig. 4, inset).

In further experiments, we compared LemTRP-1 immunoreactivity in midguts from 3-day-old fed control females with midguts from 3-day-old females which have fed for the first 2 days, and starved during the third day. Results showed 64% reduction on LemTRP-1 levels in midguts from the group that had been starved the third day (Fig. 4B).

3.6. Effects of LemTRP-1 on food intake

The effects of synthetic LemTRP-1 on food intake were tested on adult females at doses of 25 and 50 µg per specimen on food intake in *B. germanica*, using the carrot feeding bioassay. Both doses resulted in ca. 120% significant increase of food content in the foregut. This asymptotic level of response suggests that the increase observed is the maximum that can be reached in our experimental conditions. Conversely, no significant differences were observed either in the midgut or in the hindgut (Fig. 5), as expected, given the short duration (5 h) of the assay.

4. Discussion

In order to identify TRPs in the cockroach *B. germanica*, we developed an ELISA using a polyclonal antibody raised against the peptide LemTRP-1 (APSGFLGVR-NH₂). The limit of detection of LemTRP-1 was around 10⁻¹⁰M (Fig. 1), which allowed the quantification of LemTRP-1 immunoreactivity in crude extracts of individual brains and midguts. Cross-reactivity tests show that the LemTRP-1 antiserum is specific to the C-terminal tetrapeptide -LGVR-NH₂, given that it cross-reacts with LemTRP-2, which is a N-terminal extended version of LemTRP-1, whereas cross-reaction with LemTRP-4 or -5, which have a different N-terminal sequence (sequences indicated in 2.2), is very low (Fig. 1). The ELISA helped to isolate the peptide LemTRP-1 from brain extracts of the cockroach *B. germanica*. The peptide LemTRP-1 had previously been identified in midgut and brain of the cockroaches *L. maderae* and *P. americana* [11,12,17].

In the adult female of *B. germanica*, LemTRP-1 stimulates foregut and hindgut contractions, with ED₅₀ values within the nanomolar range for both gut regions (Fig. 2). This is not surprising given that TRPs were originally detected in the locust *L. migratoria* using *L. maderae* hindgut myotropic assay [21]. Nevertheless, although ED₅₀s for LemTRP-1 were similar to the values found for proctolin and sulfakinins tested in *B. germanica* tissues [8], the absolute values of force increase are lower for LemTRP-1 than for those peptides [8]. The stimulatory activity of LemTRPs on *L. maderae* hindgut contractions gave an ED₅₀ around 10 nM [27]. In the cockroach *P. americana*, TRPs stimulate foregut muscle contractions, with an ED₅₀ around 5 nM, but

1
2
3
4 TRPs failed to activate foregut contractions in the cockroaches *L. maderae* and
5
6 *Nauphoeta cinerea* [15]. These results point to foregut sensitivity differences between
7
8 the more primitive cockroach families Blattidae (*P. americana*) and Blattellidae (*B.*
9 *germanica*) and the more modified Blaberidae (*L. maderae* and *N. cinerea*).

10
11 ELISA studies showed that LemTRP-1 levels in the brain of *B. germanica*
12
13 females are quite constant throughout the first gonadotrophic cycle (Fig. 3). In *L.*
14 *maderae*, a large number of TRP-immunoreactive interneurons in the proto-, deuto- and
15
16 tritocerebrum, supplying processes to most part of the brain, in addition to a few
17
18 number of protocerebral neurons which send immunoreactive processes to the glandular
19
20 lobe of the corpora cardiaca, have been reported [13]. The distribution of TRP
21
22 immunoreactivity in the cockroach brain suggests that these peptides may play
23
24 neuromodulatory roles [13], which seems compatible with the fairly constant expression
25
26 observed in *B. germanica* brain.

27
28 In contrast, LemTRP-1 immunoreactivity in the midgut shows marked changes
29
30 during the gonadotrophic cycle. In the midgut, LemTRP-1 levels increase from the day
31
32 of emergence to a maximum on day 3, and then, they steadily decrease until day 6 (Fig.
33
34 3). In *L. maderae*, TRP-immunoreactivity has been detected in a fairly dense supply of
35
36 varicose fibres in the wall of the midgut, which originated in the stomatogastric nervous
37
38 system, and in numerous midgut endocrine cells [13].

39
40 *B. germanica* females show a cyclic pattern of feeding throughout the
41
42 gonadotrophic cycle (Fig. 3). Maximum values are observed between days 3 and 6 (with
43
44 food consumption values of ca. 10 mg/day, Fig. 3), a period coincident with the
45
46 maximum of vitellogenin production [9], which is one of the most energy-consuming
47
48 process in the adult female. Interestingly, maximum values of food consumption are
49
50 concomitant with TRP-like immunoreactivity decrease in the midgut. A dual hypothesis
51
52 arises from these observations: 1) TRPs are released from the midgut to the
53
54 haemolymph when sustained food consumption is required to maintain vitellogenesis at
55
56 the highest levels (between days 3 and 6 of the gonadotrophic cycle), and 2) TRPs in the
57
58 haemolymph stimulate food consumption in these days.

59
60 In support of that hypothesis we found that while LemTRP-1 immunoreactive
61
62 brain levels do not change in starved animals (Fig. 4A), starvation decreases LemTRP-1
63
64 immunoreactivity in midgut, whereas increases it in the haemolymph (Fig. 4A and
65
inset), and that 24 h of starvation are enough to produce that effect (Fig. 4B). In *L.*
maderae, TRPs have not been detected in typical neurohemal organs, and, therefore, the

1
2
3
4 source of TRPs detected in the haemolymph seems to be the midgut endocrine cells
5 [13,25]. Congruently with that, it has been demonstrated that increasing levels of K^+ in
6 the bathing saline, elicited the release of TRPs from isolated midguts of *L. migratoria*
7 and *L. maderae* [25]. In *L. migratoria* starvation also resulted in a decrease of TRPs in
8 the midgut and an increase in the haemolymph [25]. In the locust, although the
9 concentration increase in the haemolymph might be a consequence from the reduced
10 haemolymph volume in starved specimens, it was concluded that TRPs were released
11 from midgut in relation with nutritional stress [25]. In the present work, starved *B.*
12 *germanica* females were provided with water ad libitum, and the haemolymph volumes
13 obtained when sampling fed and starved animals were similar, which indicates that
14 haemolymph volumes were comparable in fed and starved specimens.

15
16 Also in support of the above dual hypothesis are the stimulatory effects of
17 LemTRP-1 on food consumption observed in our quantitative carrot feeding assays.
18 Food content in the foregut was significantly higher in LemTRP-1-treated animals,
19 which indicates a stimulation of food intake during the 5 h treatment period (Fig. 5).
20 The similar results between treated and controls observed in midgut and hindgut
21 contents (Fig. 5) are explained by the short duration of the assay. Carrot contents in
22 these compartments correspond to food ingested before the treatment. The fact that
23 LemTRP-1 immunoreactivity in the haemolymph of *B. germanica* females corresponds
24 to the peptide LemTRP-1 is suggested by the occurrence of a single immunoreactive
25 peak in the haemolymph HPLC separations, and the coincidence of the retention time of
26 this peak with the retention time of synthetic LemTRP-1.

27
28 Thus, our observations in *B. germanica* strongly suggest that TRPs are released
29 from midgut to the haemolymph, and that circulating TRPs contribute to maintain high
30 levels of food consumption, especially in periods of high energetic demand, like during
31 full vitellogenesis. This does not rule out, of course, the possibility that TRPs are
32 involved in regulating other processes. For example, in *D. melanogaster*, the use of a
33 RNAi construct to silence TRP gene expression specifically in the nervous system has
34 demonstrated that these peptides modulate odour perception and locomotory activity
35 [26]. Although odour perception may be closely related to food intake, knocking-down
36 of TRPs gene expression on midgut would provide a more specific approach to
37 demonstrate orexigenic roles for TRPs in insects.

38
39 Finally, the stimulation of food consumption induced by LemTRP-1 in *B.*
40 *germanica* is mirrored by the inhibitory effects induced by other peptides, like
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 leucomyosuppressin [1] and perisulkakinin [8] in the same cockroach. This suggests
5 that food consumption in *B. germanica* is finely tuned by the concerted regulatory
6 actions of stimulatory and inhibitory factors.
7
8
9

10 **Acknowledgements**

11
12
13
14 Financial support from the Spanish Ministry of Science and Technology
15 (projects AGL2002-01169, AGL2005-00773 and BFU2006-01090) and the Generalitat
16 de Catalunya (2005 SGR 00053) are gratefully acknowledged.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- [1] Aguilar R, Maestro JL, Vilaplana L, Chiva C, Andreu D, Bellés X. Identification of leucomyosuppressin in the German cockroach, *Blattella germanica*, as an inhibitor of food intake. Regul Pept 2004;119:105-12.
- [2] Aguilar R, Maestro JL, Vilaplana L, Pascual N, Piulachs MD, Bellés X. Allatostatin gene expression in brain and midgut, and activity of synthetic allatostatins on feeding-related processes in the cockroach *Blattella germanica*. Regul Pept 2003;115:171-7.
- [3] Aguilar R, Maestro JL, Bellés X. Effects of myoinhibitory peptides on food intake in the German cockroach. Physiol Entomol 2006;31:257-61.
- [4] Bellés X, Casas J, Messeguer A, Piulachs MD. In vitro biosynthesis of JH III by the corpora allata of adult females of *Blattella germanica* (L.). Insect Biochem 1987;17:1007-10.
- [5] Fields GB, Noble RL. Solid phase peptide synthesis utilizing 9-fluorenylmethoxycarbonyl amino acids. Int. J. Pept. Protein Res 1990;35:161-214
- [6] Johard HA, Muren JE, Nichols R, Larhammar DS, Nässel DR. A putative tachykinin receptor in the cockroach brain: molecular cloning and analysis of expression by means of antisera to portions of the receptor protein. Brain Res 2001;919:94-105.
- [7] Jung F, Meyer HHD, Hamm RT. Development of a sensitive enzyme-linked immunosorbent assay for the fungicide fenpropimorph. J Agric Food Chem 1989;37:1183-7.
- [8] Maestro JL, Aguilar R, Pascual N, Valero ML, Piulachs MD, Andreu D, Navarro I, Bellés X. Screening of antifeedant activity in brain extracts led to the identification of sulfakinin as a satiety promoter in the German cockroach. Are arthropod

- sulfakinins homologous to vertebrate gastrins-cholecystokinins? Eur J Biochem
2001 ;268 :5824-30.
- [9] Martín D, Piulachs MD, Bellés X. Patterns of haemolymph vitellogenin and ovarian
vitellin in the German cockroach, and the role of juvenile hormone. Physiol
Entomol 1995;20:59-65.
- [10] Muren JE, Nässel DR. Radioimmunoassay determination of tachykinin-related
peptide in different portions of the central nervous system and intestine of the
cockroach *Leucophaea maderae*. Brain Res 1996a;739:314-21.
- [11] Muren JE, Nässel DR. Isolation of five tachykinin-related peptides from the midgut
of the cockroach *Leucophaea maderae*: existence of N-terminally extended
isoforms. Regul Pept 1996b;65:185-96.
- [12] Muren JE, Nässel DR. Seven tachykinin-related peptides isolated from the brain of
the Madeira cockroach: evidence for tissue-specific expression of isoforms. Peptides
1997;18:7-15.
- [13] Muren JE, Lundquist CT, Nässel DR. Abundant distribution of locustatachykinin-
like peptide in the nervous system and intestine of the cockroach *Leucophaea*
maderae. Philos Trans R Soc Lond B Biol Sci 1995;348:423-44.
- [14] Nässel DR. Tachykinin-related peptides in invertebrates: a review. Peptides
1999;20:141-58.
- [15] Nässel DR, Eckert M, Muren JE, Penzlin H. Species-specific action and
distribution of tachykinin-related peptides in the foregut of the cockroaches
Leucophaea maderae and *Periplaneta americana*. J Exp Biol 1998;201:1615-26.
- [16] Osorio S, Piulachs MD, Bellés X. Feeding and activation of corpora allata in the
cockroach *Blattella germanica* (L.) (Dictyoptera, Blattellidae). J Insect Physiol
1998;44:31-8.

- [17] Predel R, Neupert S, Roth S, Derst C, Nässel DR. Tachykinin-related peptide precursors in two cockroach species. *Febs J* 2005;272:3365-75.
- [18] Riehle MA, Garczynski SF, Crim JW, Hill CA, Brown MR. Neuropeptides and peptide hormones in *Anopheles gambiae*. *Science* 2002;298:172-5.
- [19] Sambrook J, Fritsch EF, Maniatis T. Molecular cloning. A laboratory manual. New York: Cold Spring Harbor Laboratory Press;1989.
- [20] Satake H, Kawada T, Nomoto K, Minakata H. Insight into tachykinin-related peptides, their receptors, and invertebrate tachykinins: a review. *Zoolog Sci* 2003;20:533-49.
- [21] Schoofs L, Holman GM, Hayes TK, Nachman RJ, De Loof A. Locustatachykinin I and II, two novel insect neuropeptides with homology to peptides of the vertebrate tachykinin family. *FEBS Lett* 1990;261:397-401.
- [22] Siviter RJ, Coast GM, Winther AM, Nachman RJ, Taylor CA, Shirras AD, Coates D, Isaac RE, Nässel DR. Expression and functional characterization of a *Drosophila* neuropeptide precursor with homology to mammalian preprotachykinin A. *J Biol Chem* 2000;275:23273-80.
- [23] Takeuchi H, Yasuda A, Yasuda-Kamatani Y, Kubo T, Nakajima T. Identification of a tachykinin-related neuropeptide from the honeybee brain using direct MALDI-TOF MS and its gene expression in worker, queen and drone heads. *Insect Mol Biol* 2003;12:291-8.
- [24] Van Regenmortel MHV, Briand JP, Muller S, Plaué S. Synthetic polypeptides as antigens. In *Laboratory techniques in biochemistry and molecular biology*, vol. 19 (ed. V. K. P. Burdon RH), pp. 95-129. Amsterdam: Elsevier Science;1988.

- 1
2
3
4 [25] Winther AM, Nässel DR. Intestinal peptides as circulating hormones: release of
5
6 tachykinin-related peptide from the locust and cockroach midgut. J Exp Biol
7
8 2001;204:1269-80.
9
- 10 [26] Winther AM, Acebes A, Ferrus A. Tachykinin-related peptides modulate odor
11
12 perception and locomotor activity in *Drosophila*. Mol Cell Neurosci 2006 ;31 :399-
13
14 406.
15
16
- 17 [27] Winther AM, Muren JE, Lundquist CT, Osborne RH, Nässel DR. Characterization
18
19 of actions of *Leucophaea* tachykinin-related peptides (LemTRPs) and proctolin on
20
21 cockroach hindgut contractions. Peptides 1998;19:445-58.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legends

Fig. 1. Test of the cross-reactivity of LemTRP-1 antiserum to synthetic peptides: LemTRP-1, 2, 4 and 5. Each measurement was in duplicate.

Fig. 2. Stimulatory effect of LemTRP-1 on foregut (left) and hindgut (right) motility in *B. germanica* females. Results (mean $\pm$ SE; $n = 4-7$) are expressed as the difference of the mean of the force produced by the tissue during one minute after and before the treatment. The ID₅₀ for each tissue is indicated.

Fig. 3. LemTRP-1 equivalents in brain ($n = 7-14$) and midgut ($n = 8$) from *B. germanica* females throughout the first gonadotrophic cycle. Food consumption during the same period ($n = 11$). Results are expressed as the mean $\pm$ SE.

Fig. 4. A) LemTRP-1 equivalents in brain and midgut from 3-day-old fed and starved *B. germanica* females ($n = 15-17$). Inset: LemTRP-1 equivalents/ μ l of haemolymph, in immunoreactive fractions from HPLC separations of 155 μ l of haemolymph from 3-day-old fed, and 229 μ l from 3-day-old starved *B. germanica* females. The retention time of the immunoreactive fractions coincided with that of synthetic LemTRP-1. B) LemTRP-1 equivalents in midgut from 3 days fed and 2 days fed + 1 day starved *B. germanica* females ($n = 8$). For A and C results are expressed as the mean $\pm$ SE. Asterisks indicate significant differences (Student's *t*-test) (** $P < 0.005$; *** $P < 0.0005$).

Fig. 5. Food (carrot) content within the foregut (FG), midgut (MD) and hindgut (HD) in control and LemTRP-1-treated *B. germanica* adult females. Results are expressed as the mean $\pm$ SE ($n = 24$). The asterisk indicates significant differences (Student's *t*-test) ($P < 0.05$).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5