**From classical to nonparametric growth models: towards comprehensive
modelling of mussel growth patterns**

Isabel Fuentes-Santos, Uxío Labarta*, Kristina Arranz, M^a José Fernández-Reiriz

Consejo Superior de Investigaciones Científicas (CSIC), Instituto de Investigaciones Marinas (IIM),
C/Eduardo Cabello 6, 36208 Vigo, Spain

*Corresponding author: labarta@iim.csic.es

Telephone: +34 986 231 930 Ext: 214.

FAX:(+34) 986 292 762

11 **Abstract**

12 Understanding biological processes, such as growth, is crucial to development management and
13 sustainability plans for bivalve populations. von Bertalanffy and Gompertz models have been commonly
14 used to fit bivalve growth. These models assume that individual growth is only determined by size,
15 overlooking the effects of environmental and intrinsic conditions on growth patterns. The comparison
16 between classical models and nonparametric GAM (generalized additive models) fits conducted in this
17 work shows that the latter provide a more realistic approach of mussel growth measured in terms of shell
18 length, and dry weight of hard and soft tissues. GAM fits detected a reduction in growth during the cold
19 season, under unfavourable nutritional conditions. These fits also captured the decoupling between hard
20 and soft tissue growth, widely addressed in the literature but not incorporated in growth models. In
21 addition a GAM fit of condition index allowed us to explain annual changes in resources allocation,
22 identifying the asymptotic growth of shell and the effects of the reproductive cycle on soft tissue
23 fluctuations.

24 **Keywords:** condition index, GAM, Gompertz, shell, soft tissue, von Bertalanffy

Introduction

Mussels are dominant organisms on many rocky shores worldwide, where they play an important ecological role as habitat or prey for a multitude of organisms (Rilov et al., 2008), and in the pelagic-benthic coupling (Alonso-Pérez et al., 2010; Dame, 1993; Zúñiga et al., 2014). In addition, mussel aquaculture is a sustainable food production system with important commercial (Díaz et al., 2014; Labarta et al., 2004) and environmental value, since mussel farming has been proposed as a tool to mitigation of eutrophic coastal areas (Lindahl et al., 2005). The growing importance of aquaculture in food production along the past decades has resulted on an increasing demand for management and sustainability plans (Bergström et al., 2015). Understanding how biological processes, such as reproduction or growth, respond to environmental changes is crucial to improve management strategies.

The most commonly applied growth models are those proposed by Gompertz (1825) and von Bertalanffy (1938). The latter is thought to be a better describer of growth in fish and bivalves (Gosling, 2003). These models focus on characterizing the mean growth pattern of a given species along its lifespan. For this purpose both von Bertalanffy and Gompertz models assume that individual growth is asymptotic and that the growth rate is only determined by size. However development of management and sustainability plans requires understanding the growth dynamics and seeks for models able to detect seasonal or short-term changes in bivalve growth.

It is well known that bivalve growth is driven by intrinsic physiological processes and influenced by environmental conditions. During the past decades several works have highlighted the effect of factors such as physiological dynamics, ecological memory, and environmental conditions on bivalve growth (Babarro et al., 2000; Blicher et al., 2010; Borrero and Hilbish, 1988; Hilbish, 1986; Kautsky, 1982; Lewis and Cerrato, 1997; Mallet et al., 1987; Okumuş and Stirling, 1994; Pérez-Camacho et al., 2014, 1995). Two different strategies have been adopted to account for these effects. Some works have fitted von Bertalanffy (Bagur et al., 2013; Connor and Robles, 2015; Nedoncelle et al., 2013; Ozernyuk and Zotin, 2006) and Gompertz (Cubillo et al., 2012; Irisarri et al., 2015; Peteiro et al., 2008, 2006) models to populations growing under different conditions and compared the fitted parameters to test whether the factors under study affected bivalve growth. Other works have developed extended von Bertalanffy models incorporating the effects of food availability (Marambio et al., 2012) or maturity status (Ohnishi et al., 2012) on bivalve growth.

An important feature of bivalve growth, which deserves special attention to schedule aquaculture production, is the decoupling between hard and soft tissue (Blicher et al., 2010; Borrero and Hilbish, 1988; Hilbish, 1986; Lewis and Cerrato, 1997; Witbaard et al., 2015). This decoupling can be attributed to a mismatch between the favourable conditions to shell growth and those leading to increases of soft tissue (Borrero and Hilbish, 1988), and to seasonal changes in the allocation of resources between growth and reproduction (Peterson and Fegley, 1986). Despite these evidences, changes in shell length and soft tissue have been wrongly used as equivalent measures of growth and consequently classical and new parametric growth models have overlooked this uncoupling. For instance the extended von Bertalanffy model introduced by Ohnishi and Akamine (2006) to incorporate hard and soft tissue growth patterns assumes an allometric relationship between hard and soft tissue growth rates, in contrast with the observed uncoupling.

The common practice in fish and bivalve growth modelling is to select a model, usually von Bertalanffy or Gompertz, and fit this model to the data. This procedure can lead to biased estimators, and consequently wrong decisions, when the observed data are inconsistent with the selected model. Recent studies (Katsanevakis and Maravelias, 2008; Rabaoui et al., 2007; Rogers-Bennett, 2003) have pointed out this problem and proposed the Akaike Information Criterion (AIC) as goodness-of-fit measure to select the best growth model among a set of candidates. Once again this procedure relies on the subjective selection of some candidate models leading to unrealistic estimates if none of them is consistent with the real data.

Recent works have applied semiparametric approaches, such as generalized additive models (GAM, Hastie and Tibshirani, 1990) to include environmental and endogenous variables in fish growth curves (Ligas et al., 2015; Olsen et al., 2006; Smith et al., 2005). For bivalves, some authors have computed instantaneous growth rates as the ratio between shell length increase and time between sequential samplings and applied GAM to fit the corresponding temporal pattern (Katsanevakis, 2007; Witbaard et al., 2015). To our knowledge any comparison between the performance of classical and semiparametric approaches, which would check the goodness-of-fit of the former, has not been conducted up to date.

To address the management of both wild and cultured bivalve populations at ecosystem level we need procedures able to detect short-term or seasonal changes in the growth pattern of bivalves. This

work is a first attempt to test whether the commonly used von Bertalanffy and Gompertz provide accurate fits for the growth patterns of hard and soft tissues in the mussel *Mytilus galloprovincialis* (Lamarck, 1819) or more flexible procedures are required. For this purpose, we have considered our dataset used previously in Babarro et al. (2003), which analyzed mussel growth in suspended culture using the classical models. We tested the goodness-of-fit of the von Bertalanffy and Gompertz models through comparison with GAM fits. Instantaneous growth rates were also estimated as the first derivatives of the fitted models, in contrast with previous works that estimated instantaneous growth rates as the average change in size between samplings. Finally we tested whether GAM models can detect differences between the growth patterns of hard and soft tissues.

Materials and methods

Experimental design

Seed of *Mytilus galloprovincialis* with a mean shell length of 21.2 mm (sd =8.5) was gathered from collector ropes on a raft located in the mid-to-outer area of the Ría de Arousa (Galicia, NW Iberian Coast, see Figure A.1 in appendix A), and were socked on culture ropes deployed in an adjacent commercial raft in January 1998. Culture ropes were kept in the water up to July 1999 (526 days) covering both pre-fattening, up to November 1998, and fattening, from November 1998 to harvest in July 1999, phases of mussel culture (see details in Babarro et al. (2003)).

Sampling procedure

Sequential samplings were conducted on 28 January (seeding), 11 March, 6 May, 3 June, 1 July, 24 September and 11 November during 1998; and on 24 February, 28 April, 26 May and 7 July (harvest) during 1999. Duplicated samples of 200-300 individuals were taken from adjacent ropes in each sampling date. Individual mussel length (L, mm) was measured to the nearest 1 mm using callipers to obtain the mean shell length of each sample. Subsamples of 5-15 mussels each from 5-6 length classes covering the whole size range were used to determine total (TDW, g), shell (DWs, g) and tissue dry weight (DWt, g) by dissection and after drying at 100° C until constant weight. Condition index was calculated as the ratio between tissue and shell dry weights, $CI=DWt/DWs$ (Lucas and Beninger, 1985).

Data analysis

Classical growth models

Specific von Bertalanffy (vB) and Gompertz (G) models were used to fit growth curves in terms of shell length (mm), total, shell and tissue weight (g). The von Bertalanffy and Gompertz models for shell length are defined as follows:

$$vB: L_t = L_{\infty} \left(1 - e^{-k(t-t_0)} \right) \quad (1)$$

$$G: L_t = L_{\infty} \exp \left(-e^{-k(t-t^*)} \right) \quad (2)$$

where L_{∞} (mm) represents the asymptotic length, L_t (mm) is the shell length at time t (months), k is the growth parameter indicating the speed at which asymptotic growth is attained, t_0 (1) represents the theoretical time at which length is 0, and t^* (2) denotes the time of growth inflexion. The same models were used to fit growth curves in terms of weight.

The first derivatives of the models under comparison were computed to estimate growth rates along the culture period. For the von Bertalanffy and Gompertz model we used the expressions:

$$vB: \frac{\partial L}{\partial t} = kL_{\infty} e^{-k(t-t_0)} \quad (3)$$

$$G: \frac{\partial L}{\partial t} = ke^{-k(t-t_0)} L_{\infty} \exp \left(-e^{-k(t-t_0)} \right) = kL_t e^{-k(t-t_0)} \quad (4)$$

GAM fit of growth curves

Despite von Bertalanffy (eq. 1) and Gompertz (eq. 3) models have been widely used to estimate mussel growth in terms of shell length (mm) and dry weight (g), they can be very restrictive and lead to biased estimations and consequently incorrect conclusion when the data do not support the relationships imposed by these models. Nonparametric regression techniques provide an approach to growth curves without specifying in advance any function for the effect of time on size. Therefore mussel growth can be expressed as follows

$$Y = E[Y|t] + e = m(t) + e \quad (5)$$

where Y is the size measure, m is an unknown smooth function and e is the error term. In this work, m was fitted using regression splines, i.e. we applied a generalized additive model (GAM, Hastie & Tibshirani, 1990; Wood, 2006), which can be defined as follows:

$$E[Y|t] = H(\alpha + f(t)) \quad (6)$$

where α is the intercept, f_j is an unknown smooth function, which in this case was fitted by thin plate regression splines, and H is a fixed, known, monotone link function, selected according to the distribution of the response variable, Y . In this work we fitted GAM with Gaussian family and “identity” link function for shell length and dry weight, which were normally distributed (Shapiro test, p-value > 0.05). A GAM (1) with Gamma family and logarithmic link was applied to fit the temporal pattern of condition index. We also computed the first derivative of the GAM fits to obtain instantaneous growth rates.

Testing the goodness-of-fit of the classical growth models

Graphical comparison of the growth curves provided by the classical and GAM fits can be used to check the accuracy of von Bertalanffy and Gompertz models. Classical models are correct when the corresponding growth curves lay within the 95% confidence interval of the GAM growth curves. Nevertheless we also conducted a formal goodness-of-fit test for this purpose.

We applied a Generalized likelihood ratio (GLR) test (Fan and Jiang, 2007) to check the goodness-of-fit of the classical von Bertalanffy and Gompertz models, that is, to test whether any of these models provides a correct specification of the growth curves. Thus, considering the general model in equation (3), we test the null hypothesis $H_0: m(\text{time}) = m_0(\text{time})$, where m_0 is given by expression (1) for von Bertalanffy and expression (2) for Gompertz, respectively, versus the general alternative hypothesis H_1 where m is an unknown smooth function estimated by GAM. Let $\{t_i, Y_i\}_{i=1}^n$ be the observed data, the likelihood ratio test is

$$T = \frac{n}{2} \log \frac{RSS0}{RSS1} = \frac{n}{2} \log \frac{\sum_{i=1}^n (Y_i - m_0(t_i))^2}{n^{-1} \sum_{i=1}^n (Y_i - m(t_i))^2} \quad (7)$$

where $RSS0$ and $RSS1$ are the maximum likelihood estimators for the error variance under H_0 and H_1 , respectively. For a fixed significance level α , we reject the null hypothesis when $T > T^{1-\alpha}$, where $T^{1-\alpha}$ is the $100(1-\alpha)$ -percentile of T under H_0 . As asymptotic theory for nonparametric tests is not closed yet, we can apply resampling methods such as bootstrap (Efron and Tibshirani, 1994) to calibrate the test. Wild bootstrap (Hardle and Mammen, 1993) was used in this case to generate the resamples. Thus the empirical p-value of the test is the proportion of simulated T values larger than that obtained for the observed data. The algorithm performed to conduct this goodness-of-fit test is detailed in Appendix A.2.

Comparison of shell and soft tissues growth patterns

In order to test whether hard and soft tissue growths are coupled, we used a GAM fit with interaction factor (type of tissue) by curve (time) defined as follows:

$$E[Y|t, Z] = H \left(\alpha_j I(Z = Z_j) + \sum_{j=1}^2 f_j(t) I(Z = Z_j) \right) \quad (8)$$

where, $j=1,2$ identifies the group defined by hard (shell) and soft tissue, α_j , is the intercept for the j th type of tissue. And f_j is the smooth function that describes the effect of time on the response for the j th type of tissue. As well as in model (5) we used Gaussian family with “identity” link function (H).

We used the shrinkage variable selection procedure proposed by Marra and Wood (2011) to test whether the effect of time on each response variable depends on type of tissue, i.e. to select between model (8) and the model without interactions:

$$E[Y|time, Z] = H \left(\alpha_j I(Z = Z_j) + f(time) \right) \quad (9)$$

Once model selection was conducted, normality and independence of residuals were tested by the Shapiro-Wilk and Ljung–Box tests, respectively. In addition to the growth curves, we fitted their first derivatives in order to estimate growth rates along the culture period.

Software

Data analysis was performed with the statistical software R.3.1.3 (R Development Core Team, 2015). In particular, the `nls` function in the `stats` package was used to fit Gompertz and von Bertalanffy

models by nonlinear least squares (Bates and Watts, 1988), and the *mgcv* package (Wood, 2006) was used to fit the GAM models.

Results

Figure 1 shows that the GAM (eq. 5) and classical growth models -von Bertalanffy (eq. 1) and Gompertz (eq. 2) provide similar growth curves up to September 1998. Although the GLR tests accept the goodness-of-fit of both classical models ($p\text{-value} > 0.1$, Table B1 in Appendix B), we observe differences between models from autumn onwards when the GAM detects stagnation in shell length growth, which was not identified by the classical models. In Appendix B (Figure B1, Table B1) we report the result for growth curves in terms of weight. The GLR tests (Table B1) point out the lack of fit of the classical growth models, mainly for tissue dry weight. In view of these results GAM fits shall be used to estimate the different growth curves considered along this work.

Figure 1: Comparison of growth models. Observed shell lengths (points), GAM (black), von Bertalanffy (vB, red), Gompertz (G, green) fits of growth curves (left) and their first derivative (right). Fitted values (solid line) and 95% confidence intervals (dashed lines) for GAM fit.

The model selection conducted to test for differences between the growth patterns of shell and soft tissue detected a significant effect of type of tissues on the intercept ($p\text{-value} \approx 0$) and smooth terms of the growth curves in terms of dry weight ($p\text{-value} \approx 0$), indicating a mismatch between the growth of hard and soft tissues. Indeed the AIC for model (8), which assumed different growth patterns for hard and soft tissue (AIC = 16.33) was smaller than that obtained overlooking the differences between tissues

(model (9), AIC=152.29). Therefore, model (8) was used to estimate the growth patterns of shell (DWs) and tissue (DWt) dry weight.

A summary of the fitted models, goodness-of-fit measures and tests conducted for model checking is shown in Table 1. Shell length increased up to October; stagnated during autumn-winter, and increased again from February onwards, but at a lower rate than during the first spring (Figure 1). As observed for shell length (Figure 1), total (Figure 2, top) and shell dry weight (Figure 2, bottom, black lines) stopped growing from October to March. Nevertheless, soft tissue (Figure 2, bottom, grey lines) increased up to September, decreased from November to February and increased again from March onwards.

Table 1: GAM fits of mussel growth in terms of shell length (L, mm) total dry weight (TDW, g), shell dry weight (DWs, g), tissue dry weight (DWt, g), and condition index (CI = DWt/DWs). Normality of residuals was tested through the Shapiro–Wilk (SW) while independence of residuals was checked through the Ljung–Box (LB) tests.

L (mm)	Parametric coefficients						Goodness of fit		Residuals	
	Estimate	Std. Error	t value	Pr(> t)		Adj R ²	%Dev Exp	SW	LB	
	Intercept	54.903	0.169	325.5	<2e-16	***	0.998	99.90%	0.0969	0.4327
	Smoth terms									
	edf	Ref.df	F	p-value						
	s(time)	6.673	9	1194	<2e-16	***				
TDW (g)	Parametric coefficients						Goodness of fit		Residuals	
	Estimate	Std. Error	t value	Pr(> t)		Adj R ²	%Dev Exp	SW	LB	
	Intercept	6.027	0.103	58.62	8.7e-16	***	0.997	99.20%	0.1462	0.1760
	Smoth terms									
	edf	Ref.df	F	p-value						
	s(time)	7.327	9	162.2	<2e-16	***				
DWs vs DWt (g)	Parametric coefficients						Goodness of fit		Residuals	
	Estimate	Std. Error	t value	Pr(> t)		Adj R ²	%Dev Exp	SW	LB	
	(Intercept)	4.981	0.054	91.77	<2e-16	***	0.994	99.60%	0.0495	0.0179
	Tissue	-3.936	0.077	-51.27	<2e-16	***				
	Smoth terms									
	edf	Ref.df	F	p-value						
	s(time):Shell	8.594	9	417.4	<2e-16	***				
	s(time):Tissue	4.872	9	13.98	2.5e-14	***				
CI	Parametric coefficients						Goodness of fit		Residuals	
	Estimate	Std. Error	t value	Pr(> t)		Adj R ²	%Dev Exp	SW	LB	
	Intercept	-1.525	0.026	-59.53	3.2e-15	***	0.803	91.90%	0.5895	0.2529
	Smoth terms									
	edf	Ref.df	F	p-value						
	s(time)	7.936	8.684	14.02	1.6e-06	***				

(***) p-value < 0.001, (**) p-value < 0.01, (*) p-value < 0.05, (.) p-value < 0.1.

Figure 2: Mussel growth. Top: total dry weight (TDW, top). Bottom: shell (black) and tissue dry weight (grey). Observed values (points), and GAM fits of growth curves (left) and their first derivative (right). Fitted values (solid line) and 95% confidence intervals (dashed lines).

The differences between hard and soft tissue growth curves and the significant effect of time on the condition index (Table 1) indicate a temporal mismatch between the investment of mussels on shell and soft tissue. The decrease in CI during the first spring and winter (Figure 3) reflects two different situations: during spring both tissue and shell grew, but the latter grew at a higher rate; while the decrease in winter is caused by soft tissue losses (Figure 2, bottom). We also observe opposite behaviours between the first and second spring, as the latter reported an increase in the condition index (Figure 3), i.e. higher growth rates in soft tissue than in shell.

Figure 3: Condition index ($CI = DW_t/DW_s$). Observed values (points), and GAM fit of the temporal pattern of mussel CI (left) and its first derivative (right). Fitted values (solid line) and 95% confidence intervals (dashed lines).

Discussion

The important ecological and commercial role of bivalve aquaculture have motivated an increasing demand for management and sustainability plans (Bergström et al., 2015). Modelling individual growth is a key factor to understand the dynamics of these species and, consequently in the development of management and sustainability plans for aquaculture. Up to date, the majority of works have focused on spatial planning through the comparison of bivalve growth patterns at different locations (Bergström et al., 2015; Brigolin et al., 2009; Pérez-Camacho et al., 2014, 1995). Short term variability on bivalve growth in response to changes on the environmental conditions, which is crucial for site specific management plans, have received less attention.

Bivalve growth has been modelled by the classical von Bertalanffy and Gompertz models, which aim to estimate the mean growth along the individual lifespan and assume that growth is only determined by size. However a proper understanding of the growth dynamics requires detecting changes in the growth patterns along the culture period, as well as differences between hard and soft tissue growth. For this reason this work tests whether the classical growth models provide realistic fits of bivalve growth or more flexible techniques, such as GAM, are required

Although the small sample size ($n = 20$) can limit the power of the goodness-of-fit test, we have confirmed that von Bertalanffy and Gompertz models miss-specify mussel growth in terms of dry weight.

The GAM fit detected temporal variability in mussel growth, extensively described (Borrero and Hilbish, 1988; Gangnery et al., 2004; Hilbish, 1986; Kautsky, 1982; Lewis and Cerrato, 1997; Pernet et al., 2012; Urrutia et al., 1999; Witbaard et al., 2015). The large amount of endogenous and exogenous variables involved in a complex process such as bivalve growth, which hampers the development of accurate parametric models for age-size relationships, and the goodness-of-fit of GAMs, which do not need to assume any restrictive relationship between the response and the explanatory variables, support the use of the latter to fit bivalve growth. The first order derivatives of the fitted growth curves provide accurate estimates of the instantaneous growth rates, which up to date have been obtained as the ratio between size increases and time between samplings (Katsanevakis, 2007; Witbaard et al., 2015), GAM fits also detected a decoupling between hard and soft tissues, widely addressed in the literature (Blicher et al., 2010; Borrero and Hilbish, 1988; Hilbish, 1986; Lewis and Cerrato, 1997; Witbaard et al., 2015) but not incorporated to growth models.

The temporal variability on mussel growth detected by the GAM fits is in agreement with important features such as the effect of the environmental conditions and the reproductive cycle on bivalve growth, which have been described along several decades. The fitted growth curves and rates in terms of shell length and shell weight reflect the common pattern at temperate latitudes, where shell growth from spring to early autumn is favoured by optimal nutritional conditions and increasing temperatures, in contrast with the low food availability in colder months (Kautsky, 1982; Loo and Rosenberg, 1983; Mallet et al., 1987). Soft tissue losses during winter can be attributed to an early spawning event. *Mytilus* species are thought to synchronize their reproductive cycle with favourable environmental conditions (Edwards and Richardson, 2004; Newell et al., 1982; Philippart et al., 2012). In coastal upwelling systems spawning of many invertebrate and fish species occurs during the upwelling season (Otero et al., 2009; Snodden and Roberts, 1997; Suárez et al., 2005), which in the Galician Rías extends between spring and autumn (Torres et al., 2003; Wooster et al., 1976).

The temporal pattern of the condition index suggest a higher investment in shell up to summer followed by a period of isometric growth up to mid-autumn, i.e. the end of the upwelling season. In agreement with these findings, Hilbish (1986) argued that in *Mytilus edulis* shell growth preceded soft tissue growth. The decrease in CI during winter reflects soft tissue losses, which may be attributed to energy investment on gametogenesis coupled with unfavourable nutritional and environmental conditions

(Witbaard et al., 2015) and the subsequent spawning event in late winter. The increase in CI during the second spring indicates a higher investment on tissue than on shell, in contrast with the first spring. This interannual shift may be attributed to the progressive reduction in shell growth as asymptotic size is approached, and the post-spawning recovery of soft tissue. Accordingly, Peterson and Fegley (1986) found that shell growth in juvenile clams was much higher than in adult clams. These results are in agreement with the different hard and soft tissue growth patterns of bivalves. While shell growth is asymptotic (Sebens, 1987), soft tissue of adult mussels undergo substantial annual changes mainly driven by their reproductive cycle (Peterson and Fegley, 1986).

The decoupling between shell and soft tissues points out the need for testing whether mussels fulfil the allometric length-weight relationship, widely used to describe relative bivalve growth (Cubillo et al., 2012; Irisarri et al., 2015; Peteiro et al., 2008, 2006, Rabaoui et al., 2011, 2007). In this line Sestelo and Roca-Pardinas (2011) and Martínez-Silva et al. (2014) show that nonparametric regression provides a better fit for the relative growth of barnacles and sea urchins, which condition index also exhibit seasonal variation, than allometric models.

The productivity of mussel farming, depend on individual size and meat yield, defined as the ratio between meat and total fresh weight, since market prices are based on these parameters (Fuentes-Santos et al., 2015; Pérez-Camacho et al., 2013). More exhaustive sampling procedures than those required by the classical models, and the flexibility of nonparametric approaches, such as GAM, shall capture short term variability of bivalve growth and meat yield. These approaches shall contribute to the development of site specific management strategies through a proper schedule of seeding and harvesting.

Appendix A: Supporting information for Materials and Methods

A.1 Experimental design

Figure A 1: Mussel culture polygons in the Ría de Arousa (Galicia, NW Spain). A: raft where the experimental culture was conducted. B: raft where mussel seed was collected.

A.2 Details of the goodness-of- fit test.

We applied a Generalized likelihood ratio test (Fan and Jiang, 2007) to check the goodness-of-fit of the classical von Bertalanffy and Gompertz models, that is, to test whether any of these models provides a correct specification of the growth curves. Thus, considering the general model in equation (3), we test the null hypothesis $H_0: m(t) = m_0(t)$, where m_0 is given by expression (1) for von Bertalanffy and expression (1) for Gompertz, respectively, versus the general alternative hypothesis H_1 where m is an unknown smooth function estimated by GAM. Let $\{t_i, Y_i\}_{i=1}^n$ the observed data, the likelihood ratio test is

$$T = \frac{n}{2} \log \frac{RSS0}{RSS1} = \frac{n}{2} \log \frac{\sum_{i=1}^n (Y_i - m_0(t_i))^2}{n^{-1} \sum_{i=1}^n (Y_i - m(t_i))^2} \quad (10)$$

where $RSS0$ and $RSS1$ are the maximum likelihood estimators for the error variance under H_0 and H_1 , respectively. For a fixed significance level α , we reject the null hypothesis when $T > T^{1-\alpha}$, where $T^{1-\alpha}$ is the $100(1-\alpha)$ -percentile of the T under H_0 . As asymptotic theory for nonparametric tests is not closed yet, we can apply resampling methods such as bootstrap (Efron and Tibshirani, 1994) to calibrate the test. Wild bootstrap (Hardle and Mammen, 1993) was used in this case to generate the resamples. Thus, we applied the following algorithm to test the goodness-of-fit of the classical growth model.

1. Obtain the null, $m_0(t)$, and alternative, $m(t)$, regression functions for the sample data $\{t_i, Y_i\}_{i=1}^n$ and compute the test statistic T_{obs} as defined in expression (7).
2. For $b = 1, \dots, B$ ($B = 10000$ in this work)
 - 2.1. generate bootstrap resamples $\{t_i, Y_i^*\}_{i=1}^n$, with $Y_i^* = m_0(t_i) + e_i^*$ being

$$e_i^* = \begin{cases} \frac{(1-\sqrt{5})e_i}{2} & \text{with probability } p = \frac{5+\sqrt{5}}{10} \\ \frac{(1+\sqrt{5})e_i}{2} & \text{with probability } p = \frac{5-\sqrt{5}}{10} \end{cases}$$

where $e_i = Y_i - m_0(t_i)$ are the errors of the parametric model in H_0 .

- 2.2. Given $\{t_i, Y_i^*\}_{i=1}^n$, compute T^{*b} as in step 1.
3. The empirical p-value of the test is the proportion of simulated T-statistics larger than that obtained for the observed data.

Appendix B: Supplementary data for the Results Section

This Appendix provides the parameters and the p-values of the likelihood ratio tests for the fitted von Bertalanffy and Gompertz models (Table B1). Figure B1 provides graphical comparison between classical and GAM fits of mussel growth in terms of total, shell and soft tissue weight .

Table B1: Parameters of the von Bertalanffy and Gompertz fits for shell length (L), total (TDW), shell (DWs), and tissue (DWt) dry weight. GoF test: p-value of the Generalized likelihood ratio goodness-of fit test calibrated by wild bootstrap (B = 10000).

		von Bertalanffy						Gompertz					
		Estimate	Std.Error	t-value	Pr(> t)		GoF test	Estimate	Std.Error	t-value	Pr(> t)		GoF test
L (mm)	L_{∞}	75.83	1.423	53.28	<2.e-16	***	0.314	72.96	1.043	69.94	<2.e-16	***	0.313
	K	0.175	0.014	12.30	7e-10	***		0.274	0.012	22.18	5e-14	***	
	t_0	-1.760	0.230	-7.661	7e-07	***		0.831	0.077	10.730	5e-09	***	
TDW (g)	w_{∞}	16.44	3.334	4.93	0.0001	***	0.013	11.19	0.469	23.89	2e-14	***	0.065
	K	0.065	0.023	2.87	0.0106	*		0.246	0.012	21.13	1e-13	***	
	t_0	0.163	0.437	0.372	0.7142			4.907	0.242	20.290	2e-13	***	
DWs (g)	w_{∞}	14.57	3.146	4.63	0.0002	***	0.012	9.74	0.423	23.01	3e-14	***	0.056
	K	0.060	0.021	2.83	0.0116	*		0.226	0.011	21.26	1e-13	***	
	t_0	0.232	0.413	0.562	0.5814			5.362	0.266	20.200	2e-13	***	
DWt (g)	w_{∞}	2.118	0.375	5.64	3e-05	***	0.007	1.584	0.122	13.02	3e-10	***	0.01
	K	0.103	0.041	2.48	0.0237	*		0.383	0.038	9.97	2e-08	***	
	t_0	-0.048	0.657	-0.073	0.9426			3.269	0.332	9.858	2e-08	***	

(***) p-value < 0.001, (**) p-value < 0.01, (*) p-value < 0.05, (.) p-value < 0.1.

Figure B2: Observed dry weights (points), GAM (black), von Bertalanffy (vB, red) and Gompertz (green) fits of growth curves (left) and their first derivative (right) for mussels gathered from collector ropes (subtidal mussels) . Fitted values (solid line) and 95% confident intervals (dashed lines) for GAM fit. Top: total dry weight (TDW), centre: shell dry weight (DWs) , bottom: tissue dry weight (DWt).

Acknowledgements

We are grateful to Lourdes Nieto and Beatriz Gonzalez for their technical support. We acknowledge Dr. X.A. Álvarez-Salgado for his critical comments that contributed to improve this manuscript. This study was funded by PIE project (CSIC 201540E107), EU H2020 project ClimeFish (EU 677039), and PROINSA-CSIC contract-project (CSIC0704101100001)

References

- Alonso-Pérez, F., Ysebaert, T., Castro, C., 2010. Effects of suspended mussel culture on benthic–pelagic coupling in a coastal upwelling system (Ría de Vigo, NW Iberian Peninsula). *J. Exp. Mar. Biol. Ecol.* 382, 96–107.
- Babarro, J.M., Fernández-Reiriz, M.J., Labarta, U., 2000. Growth of seed mussel (*Mytilus galloprovincialis* Lmk): effects of environmental parameters and seed origin. *J. Shellfish Res.* 19, 187–193.
- Babarro, J.M., Labarta, U., Fernández-Reiriz, M.J., 2003. Growth patterns in biomass and size structure of *Mytilus galloprovincialis* cultivated in the Ría de Arousa (north-west Spain). *J. Mar. Biol. Assoc. UK* 83, 151–158.
- Bagur, M., Richardson, C.A., Gutiérrez, J.L., Arribas, L.P., Doldan, M.S., Palomo, M.G., 2013. Age, growth and mortality in four populations of the boring bivalve *Lithophaga patagonica* from Argentina. *J. Sea Res.* 81, 49–56.
- Bates, D.M., Watts, D.G., 1988. Nonlinear regression: iterative estimation and linear approximations, in: *Nonlinear Regression Analysis and Its Applications*. Wiley, New York, pp. 32–66.
- Bergström, P., Lindegarth, S., Lindegarth, M., 2015. Modeling and predicting the growth of the mussel, *Mytilus edulis*: implications for planning of aquaculture and eutrophication mitigation. *Ecol. Evol.* 5, 5920–5933.
- Blicher, M.E., Rysgaard, S., Sejr, M.K., 2010. Seasonal growth variation in *Chlamys islandica* (Bivalvia) from sub-Arctic Greenland is linked to food availability and temperature. *Mar Ecol Prog Ser* 407, 71–86.
- Borrero, F.J., Hilbish, T.J., 1988. Temporal variation in shell and soft tissue growth of the mussel *Geukensia demissa*. *Mar. Ecol. Prog. Ser.* Oldendorf 42, 9–15.

370 Brigolin, D., Dal Maschio, G., Rampazzo, F., Giani, M., Pastres, R., 2009. An individual-based
 371 population dynamic model for estimating biomass yield and nutrient fluxes through an off-shore
 372 mussel (*Mytilus galloprovincialis*) farm. *Estuar. Coast. Shelf Sci.* 82, 365–376.

373 Connor, K.M., Robles, C.D., 2015. Within-Site Variation of Growth Rates and Terminal Sizes in *Mytilus*
 374 *californianus* Along Wave Exposure and Tidal Gradients. *Biol. Bull.* 228, 39–51.

375 Cubillo, A.M., Peteiro, L.G., Fernández-Reiriz, M.J., Labarta, U., 2012. Influence of stocking density on
 376 growth of mussels (*Mytilus galloprovincialis*) in suspended culture. *Aquaculture* 342–343, 103–
 377 111. doi:10.1016/j.aquaculture.2012.02.017

378 Dame, R.F., 1993. The role of bivalve filter feeder material fluxes in estuarine ecosystems, in: *Bivalve*
 379 *Filter Feeders*. Springer, pp. 245–269.

380 Díaz, C., Figueroa, Y., Sobenes, C., 2014. Seasonal effects of the seeding on the growth of Chilean
 381 mussel (*Mytilus edulis platensis*, d’Orbigny 1846) cultivated in central Chile. *Aquaculture* 428–
 382 429, 215–222. doi:10.1016/j.aquaculture.2014.03.013

383 Edwards, M., Richardson, A.J., 2004. Impact of climate change on marine pelagic phenology and trophic
 384 mismatch. *Nature* 430, 881–884.

385 Efron, B., Tibshirani, R.J., 1994. *An introduction to the bootstrap*. CRC press, London.

386 Fan, J., Jiang, J., 2007. Nonparametric inference with generalized likelihood ratio tests. *Test* 16, 409–444.

387 Fuentes-Santos, I., Cubillo, A.M., Labarta, U., 2015. A bioeconomic approach to optimize mussel culture
 388 production. *Rev. Aquac.* doi:10.1111/raq.12108

389 Gangnery, A., Bacher, C., Buestel, D., 2004. Application of a population dynamics model to the
 390 Mediterranean mussel, *Mytilus galloprovincialis*, reared in Thau Lagoon (France). *Aquaculture*
 391 229, 289–313.

392 Gompertz, B., 1825. On the nature of the function expressive of the law of human mortality, and on a new
 393 mode of determining the value of life contingencies. *Philos. Trans. R. Soc. Lond.* 513–583.

394 Gosling, E., 2003. *Bivalve molluscs: biology, ecology and culture*, Blackwell Publishing. ed. Elsevier,
 395 Oxford, UK.

396 Hardle, W., Mammen, E., 1993. Comparing nonparametric versus parametric regression fits. *Ann. Stat.*
 397 1926–1947.

398 Hastie, T.J., Tibshirani, R.J., 1990. *Generalized Additive Models*. Chapman and Hall/CRC, London.

399 Hilbish, T.J., 1986. Growth trajectories of shell and soft tissue in bivalves: seasonal variation in *Mytilus*
400 *edulis* L. J. Exp. Mar. Biol. Ecol. 96, 103–113.

401 Irisarri, J., Cubillo, A.M., Fernández-Reiriz, M.J., Labarta, U., 2015. Growth variations within a farm of
402 mussel (*Mytilus galloprovincialis*) held near fish cages: importance for the implementation of
403 integrated aquaculture. Aquac. Res. 46, 1988–2002. doi:10.1111/are.12356

404 Katsanevakis, S., 2007. Growth and mortality rates of the fan mussel *Pinna nobilis* in Lake Vouliagmeni
405 (Korinthiakos Gulf, Greece): a generalized additive modelling approach. Mar. Biol. 152, 1319–
406 1331.

407 Katsanevakis, S., Maravelias, C.D., 2008. Modelling fish growth: multi-model inference as a better
408 alternative to a priori using von Bertalanffy equation. Fish Fish. 9, 178–187.

409 Kautsky, N., 1982. Growth and size structure in a baltic *Mytilus edulis* population. Mar. Biol. 68, 117–
410 133. doi:10.1007/BF00397599

411 Labarta, U., Fernández-Reiriz, Pérez-Camacho, Alejandro, Pérez-Corbacho, E., 2004. Bateeiros, mar,
412 mejillón. Una perspectiva bioeconómica., Serie Estudios Sectoriales. Fundación Caixa Galicia.,
413 A Coruña, Spain.

414 Lewis, D.E., Cerrato, R.M., 1997. Growth uncoupling and the relationship between shell growth and
415 metabolism in the soft shell clam *Mya arenaria*. Mar. Ecol. Prog. Ser. 158, 177–189.

416 Ligas, A., Colloca, F., Lundy, M.G., Mannini, A., Sartor, P., Sbrana, M., Voliani, A., Belcari, P., 2015.
417 Modeling the growth of recruits of European hake (*Merluccius merluccius*) in the northwestern
418 Mediterranean Sea with generalized additive models. Fish. Bull. 113, 69–82.

419 Lindahl, O., Hart, R., Hernroth, B., Kollberg, S., Loo, L.-O., Olrog, L., Rehnstam-Holm, A.-S., Svensson,
420 J., Svensson, S., Syversen, U., 2005. Improving marine water quality by mussel farming: a
421 profitable solution for Swedish society. AMBIO J. Hum. Environ. 34, 131–138.

422 Loo, L.-O., Rosenberg, R., 1983. *Mytilus edulis* culture: Growth and production in western Sweden 35,
423 137–150. doi:10.1016/0044-8486(83)90081-9

424 Lucas, A., Beninger, P.G., 1985. The use of physiological condition indices in marine bivalve
425 aquaculture. Aquaculture 44, 187–200.

426 Mallet, A.L., Carver, C.E.A., Coffen, S.S., Freeman, K.R., 1987. Winter growth of the blue mussel
427 *Mytilus edulis* L.: importance of stock and site. J. Exp. Mar. Biol. Ecol. 108, 217–228.

428 Marambio, J., Maturana, S., Campos, B., 2012. Modelo dinámico de crecimiento de la biomasa para
 429 *Mytilus chilensis* en sistemas de cultivo en líneas. Rev. Biol. Mar. Oceanogr. 47, 51 – 64.

430 Marra, G., Wood, S.N., 2011. Practical variable selection for generalized additive models. Comput. Stat.
 431 Data Anal. 55, 2372–2387.

432 Martínez-Silva, I., Sestelo, M., Bidegain, G., Lorenzo-Arribas, A., Roca-Pardiñas, J., 2014.
 433 Nonparametric regression applied to sea urchin growth, in: Sea Urchins: Habitat, Embryonic
 434 Development and Importance in the Environment. NOVA, Editors: E. Raymond Banks, pp. 1–
 435 32.

436 Nedoncelle, K., Lartaud, F., De Rafelis, M., Boulila, S., Le Bris, N., 2013. A new method for high-
 437 resolution bivalve growth rate studies in hydrothermal environments. Mar. Biol. 160, 1427–
 438 1439.

439 Newell, R.I., Hilbish, T.J., Koehn, R.K., Newell, C.J., 1982. Temporal variation in the reproductive cycle
 440 of *Mytilus edulis* L.(Bivalvia, Mytilidae) from localities on the east coast of the United States.
 441 Biol. Bull. 162, 299–310.

442 Ohnishi, S., Akamine, T., 2006. Extension of von Bertalanffy growth model incorporating growth
 443 patterns of soft and hard tissues in bivalve molluscs. Fish. Sci. 72, 787–795.

444 Okumuş, İ., Stirling, H.P., 1994. Physiological energetics of cultivated mussel (*Mytilus edulis*)
 445 populations in two Scottish west coast sea lochs. Mar. Biol. 119, 125–131.

446 Olsen, E., Knutsen, H., Simonsen, J., Jonsson, B., Knutsen, J., 2006. Seasonal variation in marine growth
 447 of sea trout, *Salmo trutta*, in coastal Skagerrak. Ecol. Freshw. Fish 15, 446–452.

448 Otero, J., Álvarez-Salgado, X.A., González, Á.F., Gilcoto, M., Guerra, Á., others, 2009. High-frequency
 449 coastal upwelling events influence *Octopus vulgaris* larval dynamics on the NW Iberian shelf.
 450 Mar. Ecol. Prog. Ser. 386, 123–132.

451 Ozernyuk, N., Zotin, A., 2006. Comparative analysis of growth of edible mussel *Mytilus edulis* from
 452 different White Sea regions. Biol. Bull. 33, 149–152.

453 Pérez-Camacho, A., Aguiar, E., Labarta, U., Vinseiro, V., Fernández-Reiriz, M.J., Álvarez-Salgado, X.A.,
 454 2014. Ecosystem-based indicators as a tool for mussel culture management strategies. Ecol.
 455 Indic. 45, 538–548.

456 Pérez-Camacho, A., Labarta, U., Beiras, R., 1995. Growth of mussels (*Mytilus edulis galloprovincialis*)
457 on cultivation rafts: influence of seed source, cultivation site and phytoplankton availability.
458 *Aquaculture* 138, 349–362.

459 Pérez-Camacho, A., Labarta, U., Vinseiro, V., Fernández-Reiriz, M.J., 2013. Mussel production
460 management: Raft culture without thinning-out. *Aquaculture* 406–407, 172–179.
461 doi:10.1016/j.aquaculture.2013.05.019

462 Pernet, F., Malet, N., Pastoureaud, A., Vaquer, A., Quéré, C., Dubroca, L., 2012. Marine diatoms sustain
463 growth of bivalves in a Mediterranean lagoon. *J. Sea Res.* 68, 20–32.

464 Peteiro, L.G., Babarro, J.M., Labarta, U., Fernández-Reiriz, M.J., 2006. Growth of *Mytilus*
465 *galloprovincialis* after the Prestige oil spill. *ICES J. Mar. Sci. J. Cons.* 63, 1005–1013.

466 Peteiro, L.G., Filgueira, R., Labarta, U., Fernández-Reiriz, M.J., 2008. Growth and biochemical responses
467 of the offspring of mussels directly affected by the “Prestige” oil spill. *ICES J. Mar. Sci. J. Cons.*
468 65, 509–513.

469 Peterson, C.H., Fegley, S.R., 1986. Seasonal allocation of resources to growth of shell, soma, and gonads
470 in *Mercenaria mercenaria*. *Biol. Bull.* 171, 597–610.

471 Philippart, C.J., Amaral, A., Asmus, R., van Bleijswijk, J., Bremner, J., Buchholz, F., Cabanellas-
472 Reboredo, M., Catarino, D., Cattrijsse, A., Charles, F., others, 2012. Spatial synchronies in the
473 seasonal occurrence of larvae of oysters (*Crassostrea gigas*) and mussels (*Mytilus*
474 *edulis/galloprovincialis*) in European coastal waters. *Estuar. Coast. Shelf Sci.* 108, 52–63.

475 Rabaoui, L., Zouari, S.T., Katsanevakis, S., Belgacem, W., Hassine, O.K.B., 2011. Differences in
476 absolute and relative growth between two shell forms of *Pinna nobilis* (Mollusca: Bivalvia)
477 along the Tunisian coastline. *J. Sea Res.* 66, 95–103.

478 Rabaoui, L., Zouari, S.T., Katsanevakis, S., Hassine, O.K.B., 2007. Comparison of absolute and relative
479 growth patterns among five *Pinna nobilis* populations along the Tunisian coastline: an
480 information theory approach. *Mar. Biol.* 152, 537–548.

481 Rilov, G., Dudas, S.E., Menge, B.A., Grantham, B.A., Lubchenco, J., Schiel, D.R., 2008. The surf zone: a
482 semi-permeable barrier to onshore recruitment of invertebrate larvae? *J. Exp. Mar. Biol. Ecol.*
483 361, 59–74. doi:10.1016/j.jembe.2008.04.008

484 Rogers-Bennett, L.D., 2003. Modeling red sea urchin growth using six growth functions. *Fish. Bull.* 101,
485 614–626.

- Sebens, K.P., 1987. The ecology of indeterminate growth in animals. *Annu. Rev. Ecol. Syst.* 371–407.
- Sestelo, M., Roca-Pardinas, J., 2011. A New Approach to Estimation of the Length—Weight Relationship of *Pollicipes pollicipes* (Gmelin, 1789) on the Atlantic Coast of Galicia (Northwest Spain): Some Aspects of Its Biology and Management. *J. Shellfish Res.* 30, 939–948.
- Smith, J.M., Pierce, G.J., Zuur, A.F., Boyle, P.R., 2005. Seasonal patterns of investment in reproductive and somatic tissues in the squid *Loligo forbesi*. *Aquat. Living Resour.* 18, 341–351.
- Snodden, L. m., Roberts, D., 1997. Reproductive Patterns and Tidal Effects on Spat Settlement of *Mytilus Edulis* Populations in Dundrum Bay, Northern Ireland 77, 229–243. doi:10.1017/S0025315400033890
- Suárez, M.P., Alvarez, C., Molist, P., Juan, F.S., 2005. Particular aspects of gonadal cycle and seasonal distribution of gametogenic stages of *mytilus galloprovincialis* cultured in the estuary of vigo 24, 531–540. doi:10.2983/0730-8000(2005)24[531:PAOGCA]2.0.CO;2
- Torres, R., Barton, E.D., Miller, P., Fanjul, E., 2003. Spatial patterns of wind and sea surface temperature in the Galician upwelling region. *J. Geophys. Res. Oceans* 108.
- Urrutia, M., Ibarrola, I., Iglesias, J., Navarro, E., 1999. Energetics of growth and reproduction in a high-tidal population of the clam *Ruditapes decussatus* from Urdaibai Estuary (Basque Country, N. Spain). *J. Sea Res.* 42, 35–48.
- von Bertalanffy, L., 1938. A quantitative theory of organic growth (Inquiries on growth laws. II). *Hum. Biol.* 10, 181–213.
- Witbaard, R., Duineveld, G.C., Bergman, M.J., Witte, H.I., Groot, L., Rozemeijer, M.J., 2015. The growth and dynamics of *Ensis directus* in the near-shore Dutch coastal zone of the North Sea. *J. Sea Res.* 95, 95–105.
- Wood, S., 2006. Generalized additive models: an introduction with R. Chapman and Hall/CRC, London.
- Wooster, W.S., Bakun, A., McLain, D.R., 1976. Seasonal upwelling cycle along the eastern boundary of the North Atlantic. *J. Mar. Res.* 34, 131–141.
- Zúñiga, D., Castro, C.G., Aguiar, E., Labarta, U., Figueiras, F., Fernández-Reiriz, M.J., 2014. Biodeposit contribution to natural sedimentation in a suspended *Mytilus galloprovincialis* Lmk mussel farm in a Galician Ría (NW Iberian Peninsula). *Aquaculture* 432, 311–320.