**Paleoflood records from sinkholes. An example from the Ebro River floodplain,
NE Spain**

Francisco Gutiérrez ^{a*}, Mario Zarroca^b, Carmen Castañeda^c, Domingo Carbonel^a, Jesús
Guerrero^a, Rogelio Linares^b, Carles Roqué^d, Pedro Lucha^e

^a Departamento de Ciencias de la Tierra; Universidad de Zaragoza; C/. Pedro Cerbuna 12; 50009
Zaragoza; Spain; Phone: 34 976 761090

^b Departamento de Geología; Universidad Autónoma de Barcelona; E-08193 Barcelona; Spain

^c Estación Experimental de Aula Dei; EEAD-CSIC; Av. Montañana 1001; 50059 Zaragoza; Spain

^d Àrea de Geodinàmica Externa i Geomorfologia; Universitat de Girona; Campus Montilivi; E-17071
Girona; Spain

^e Departamento de Didáctica de las Ciencias Experimentales; Universidad de Zaragoza; C/. Pedro
Cerbuna 12; 50009 Zaragoza; Spain

* Corresponding author: Tel: +34 976 761090; Fax: +34 976 761106; e-mail address: fgutier@unizar.es

18

19 **Abstract**

20 This work introduces for the first time the concept of using sinkholes in fluvial valleys
21 as recorders of past floods. The notion is illustrated through the investigation of a
22 complex sinkhole located in a broad floodplain underlain by salt-bearing Cenozoic
23 evaporites. This active sinkhole comprises a large subsidence depression affecting the
24 floodplain and the edge of a terrace, and a nested collapse sinkhole that used to host a
25 sinkhole pond. A borehole drilled in the buried sinkhole pond revealed a ca. 7.8 m thick
26 fill that records around 2,700 yr of clayey lacustrine deposition interrupted by three
27 types of detrital facies. Two thick pebble gravel beds have been attributed to major
28 high-competence flood; a paleoflood occurred in Visigothic times (1537-1311 cal yr
29 BP) and the 1961 Great Ebro River flood, which is the largest event of the instrumental
30 record. A trench dug in the portion of the terrace affected by subsidence exposed a mid-
31 Holocene slackwater paleoflood deposit. The disadvantages and advantages of sinkholes
32 as archives of past flood histories are discussed.

33

34 **Key words:** paleohydrology, sinkhole lakes, evaporite karst, subsidence rate, trenching

35

36

38 **Introduction**

39 Traditional flood-frequency analyses are based on gauging records that cover short time
40 spans and historical data with limited accuracy. Moreover, the operation of gauging
41 stations frequently fail during the most important events; extremely large and rare
42 floods. Consequently, the hazard estimates for large and long return-period floods,
43 derived from extrapolations, have a questionable reliability. The magnitude and
44 frequency relationships of large floods are essential for flood risk management (e.g.,
45 definition of the flooding zone) and the design or retrofitting of costly engineering
46 structures (e.g., large dams, nuclear facilities). These limitations can be partially
47 overcome by conducting paleoflood hydrological studies, which provide information on
48 the magnitude and chronology of large floods using geological evidence (Kochel and
49 Baker, 1982; Baker, 2003). Paleoflood data allow expanding the temporal length of
50 flood catalogues and reduce the uncertainty in estimates of long return-period floods
51 (e.g., Benito et al., 2004). Past floods may be recorded as erosional and depositional
52 features formed near maximum water levels, notably slackwater deposits. The latter are
53 fine-grained sediments that accumulate from sediment-laden flood waters at high
54 localities, where the flow becomes separated from the main tread of flood flow, and can
55 be used as paleostage indicators (Baker, 1987; Kochel and Baker, 1988; Baker et al.,
56 2002). They are most frequent in deep and narrow bedrock canyons, where floods
57 produce large increases of water stage, and typically form in tributary mouths, caves or
58 alcoves, zones of sharp channel widening, and on high terraces (Baker et al., 2002).
59 Moreover, those bedrock channels, characterised by relatively stable boundaries, are the
60 most appropriate for converting reliably floodwater elevations into paleoflood discharge
61 estimates.

Alluvial channels and broad floodplains are commonly inadequate geomorphic settings for finding long and complete geological records of paleofloods (e.g., Baker, 2003; Benito et al., 2004). However, in valley reaches underlain by soluble bedrock, the sedimentary fill of sinkhole lakes developed within the floodplain and low terraces might include valuable archives of past floods.

Sinkholes are enclosed depressions endemic of karst regions that may result from the differential corrosional lowering of soluble bedrock (solution sinkholes) or from subsurface dissolution and the subsidence of the overlying sediments (subsidence sinkholes) by different mechanisms (i.e., collapse, sagging and suffosion) (Gutiérrez, 2016 and references therein). Sinkhole sediments have been investigated for multiple purposes: (1) Reconstructing paleoenvironmental and paleoclimatic variability (e.g., Laury, 1980; Whitmore et al., 1996; Hyatt and Gilbert, 2004; Morellón et al., 2009; Barreiro-Lostres et al., 2014); (2) Investigating paleontological and archaeological sites (e.g., Carbonell et al., 2008; Calvo et al., 2013; Zaidner et al., 2014; Gutiérrez et al., 2016); (3) Estimating erosion rates and their temporal variability in small catchments (e.g., Turnage et al., 1997; Hart, 2014); (4) Inferring quantitatively the evolution of the subsidence phenomena using the trenching technique in combination with geochronological data (e.g., Gutiérrez et al., 2009, 2011, 2014; Carbonel et al., 2014, 2015; Gutiérrez, 2016). (5) Identifying the sedimentary signature of hurricanes recorded in bedrock collapse sinkholes, both onshore (cenotes) and offshore (blue holes) (Gischler et al., 2008; Lane et al., 2011; Brown et al., 2014). Brown et al. (2014) recognized two historical hurricanes (1967 Hurricane Beulah and 1991 Hurricane Gilbert) in the sediments of a 80 m deep sinkhole lake located 10 km inland of the Caribbean coast on the Yucatan Peninsula, Mexico. The hurricanes were inferred from

prominent fining-upward coarse-grained beds intercalated within fine-grained carbonate mud.

This work explores using sinkhole lakes in alluvial valleys as recorders of floods. The working hypothesis is that large flood events may deposit distinctive coarse-grained sediments in these lacustrine basins that normally have fine-grained background sedimentation. This geomorphic setting has similarities with oxbow lakes, which have been already used as proxies of past floods. Oliva et al. (2016) reconstruct a 1300 year long record of paleofloods using sediment cores from two oxbow lakes along the broad floodplain of the Désert River in temperate-subpolar southwestern Québec, Canada. Here, flood deposits are detected through magnetic susceptibility peaks associated with fine-grained layers, intercalated within fine sediment rich in organic matter and lower values in magnetic susceptibility (Oliva et al., 2016). Sheffer et al. (2003) investigated paleofloods recorded in the deposits accumulated in a perched abandoned meander of the Ardèche River, Southern France. The paleoflood record associated with a buried collapse sinkhole located in the floodplain of the Ebro River, NE Spain, that used to host a permanent pond, is investigated. Sinkholes in this area are related to the karstification of the evaporitic bedrock and are generally affected by high subsidence rates due to rapid dissolution of high-solubility beds (halite and glauberite).

Geological setting

The investigated sinkhole is located in the middle reach of the Ebro River valley, NE Spain. The sinkhole pond, nowadays buried by anthropogenic deposits, is situated in the floodplain, 7 km to the W of Zaragoza city and 1 km to the SW of Monzalbarba village (**Figs. 1 and 2**). From the geological perspective, the site lies in the central sector of the Ebro Cenozoic basin, which is the southern foreland basin of the Pyrenean Alpine

orogen. The valley has been carved in horizontally lying evaporites of the late Oligocene-Miocene Zaragoza Formation, deposited in an extensive high-salinity playalake system (Quirantes, 1978; Ortí and Salvany, 1997). This formation, more than 850 m in thickness, is composed of anhydrite (CaSO_4), halite (NaCl), glauberite ($\text{Na}_2[\text{CaSO}_4]_2$), clay and marl in the subsurface. Secondary gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) is the only evaporitic rock exposed at the surface (Torrescusa and Klimowitz, 1990; Salvany et al., 2007; Salvany, 2009). On the basis of 9 boreholes drilled along a 50 km long stretch of the Ebro Valley, this upper part of the evaporitic sequence has been subdivided into four lithostratigraphic units, in ascending order (Salvany et al., 2007; Salvany, 2009): (1) marl and anhydrite basal unit; (2) halite unit (up to 150 m); (3) glauberite-halite unit (ca. 50 m); and (4) anhydrite unit. In two nearby boreholes (Utebo, and Santa Inés 2), the contact between the anhydrite unit and the underlying glauberite-halite unit occurs at 130-150 m a.s.l. (Salvany et al., 2007). The elevation of the studied sinkhole (205 m a.s.l.) suggests that this subsidence feature is likely related to interstratal karstification of high-solubility salts. The equilibrium solubilities of gypsum, glauberite, and halite in distilled water at normal conditions are 2.4 g/l, 118 g/l, and 360 g/l, respectively (Langer and Offerman, 1982; Ford and Williams, 2007; Gutiérrez and Cooper, 2013).

Geomorphological setting

The NW-SE-trending Ebro Valley displays a markedly asymmetric geometry, with a prominent and linear gypsum escarpment on the NE margin, and a stepped sequence of terraces on the opposite side (**Figs. 1 and 2**). The terrace deposits are locally thickened, filling kilometre-sized basins generated by dissolution-induced synsedimentary subsidence, in which the alluvium may reach more than 50 m in thickness. The terrace

deposits are dominated by channel gravel facies that record broad braided channels, whereas fine-grained floodplain and palustrine-lacustrine facies are abundant in the thickened alluvium and in paleosinkholes. The alluvial cover typically shows numerous gravitational deformations and paleosinkholes, mainly related to sagging and/or collapse subsidence mechanisms. The subsidence structures commonly also affect to the underlying gypsiferous bedrock, indicating that they are largely related to interstratal karstification of salts (e.g., Gutiérrez et al., 2008, 2015; Guerrero et al., 2013).

The Ebro River flows along a low-gradient and broad floodplain around 5 km wide. The channel has a sinuous geometry (sinuosity index of 1.6; Ollero, 2010) with gravelly point-bars and mid-channel bars. The floodplain, mainly occupied by crop fields, is underlain by fine-grained facies and shows multiple cross-cutting abandoned channels, some oxbow lakes (locally known as *galachos*) (**Fig. 1**) and sinkhole ponds (**Fig. 3**).

The Ebro River in its middle reach has experienced frequent changes mainly by migration and also by avulsion and cut-off processes (Regato, 1988; Ollero, 1995, 2010; Gutiérrez et al., 2007; Benito-Calvo et al., 2016). Magdaleno and Fernández-Yuste (2011), in a study on the evolution of the middle reach of Ebro River between 1927 and 2003, document a significant reduction in the mobility of the channel and its average bankfull width. Benito-Calvo et al. (2016) document a displacement of 540 m in the river channel at Alcalá village between 1927 and 1957, resulting in the abandonment of a river passage. These are largely episodic adjustments associated with flood events. For instance, the Juslibol oxbow lake, located 4 km NE of the studied site, was created by the 1961 Great Ebro River flood, which caused the cut-off of a meander (Ollero, 2010) (**Fig. 1**). At the present time, the potential for the river to change its path has been significantly hampered by the construction of channel-bank protection structures (e.g., rip-rap and concrete walls), dykes and flood lamination using the headwater reservoirs

(e.g., López-Moreno et al., 2002; Cabezas et al., 2009; Ollero, 2010). The integrity of the dikes is locally compromised by active sinkholes (Benito-Calvo et al., 2016). The floodplain is affected by numerous active sinkholes (around 8% of the area underlain by evaporites) which can be grouped into three types (Galve et al., 2009): (1) sagging sinkholes hundreds of meters long with diffuse edges; (2) large collapse sinkholes tens of meters across; and (3) small cover collapse sinkholes (**Fig. 1**). The large collapse sinkholes commonly intercept the water table and consequently host permanent lakes that may reach more than 6 m in depth (Gutiérrez et al., 2007, 2011) (**Fig. 3**). These active sinkholes have been largely filled by anthropogenic deposits including a wide variety of materials (**Fig. 1**) and many of them have been used for the construction of human structures (e.g. buildings, roads, conventional and high-speed railways) (Galve et al., 2009, 2015). Consequently, the area is affected by severe subsidence damage, mainly related to the activity of pre-existing sinkholes improperly used for development. Galve et al. (2009), in a large area upstream of Zaragoza city (40.8 km²), including the investigated sinkhole, estimated that around 70% of the sinkholes identifiable in aerial photographs from 1957 have been obliterated by human activity. The studied sinkhole, associated with the southern edge of the floodplain, is a large diffuse-edged depression with a nested collapse sinkhole that used to host a permanent pond (**Figs. 4 and 5**). The buried pond is located at the foot of the lowermost terrace (+8-10 m) and 2.4 km apart from the present-day river channel.

The Ebro River floods

The Ebro River is the largest Mediterranean fluvial system of the Iberian Peninsula, with a watershed area of 85,362 km² and a channel length of 930 km. In its middle reach, around Zaragoza city, it is essentially an allocthonous drainage that flows across

a semiarid area (300-400 mm/yr), conveying the run-off mostly contributed by the Pyrenean headwaters (**Fig. 1**), where the mean annual precipitation reaches more than 2000 mm. At Zaragoza gauging station, with a contributing area of 40,434 km², the mean annual discharge is 216.5 m³/s (5.35 l/s/km²). Here, the river has a pluvial-nival regime, with the highest monthly flow recorded in February and prolonged low flows in summer (Ollero, 2010).

Floods commonly occur in winter and in early spring and are typically related to long-lasting cyclonic rains in the Pyrenees that may induce rapid snow melting. The historical floods of the Ebro River in Zaragoza area compiled by the Comisión Nacional de Protección Civil are shown in **Table 1** (CNPC, 1985). The oldest event dates back to AD 827. Most of these floods caused damage on bridges, crop fields and villages located in the floodplain, and there is no information on the peak discharge. The largest flood of the 19th century occurred in January 1871, which caused several fatalities (CNPC, 1985; Espejo et al., 2008; Mejón, 2011). A smaller and spatially restricted flood in 1880 overturned a boat in Logroño (160 km upstream Zaragoza) killing 90 soldiers. The biggest flood in the 20th century was the Great Ebro flood of January 1961, with a peak flow rate of 4130 m³/s at Zaragoza, and a return period of around 80 years according to the Ebro Basin Water Authority. This event, with higher magnitude than the 1871 event, was produced by a long rainfall episode accompanied by substantial snow melting in the Pyrenees. This is the largest and most damaging flood event within the instrumental record starting in 1943. The flood waters caused severe economic damage and had a major morpho-sedimentary impact, including significant changes in the route of the river channel and the formation of the Juslibol oxbow lake through a meander cut-off (CNPC, 1983; Ollero, 1995; Mejón, 2011) (**Fig. 1**). The February 1952 and the well-documented February 2003 flood were the second and fourth largest events

occurred so far in the 20th and 21st centuries, with peak discharges of 3260 m³/s and 2988 m³/s, respectively (**Table 1**). The flood waters in the 2003 event covered 83% of the floodplain (Losada et al., 2004), but not the analysed sinkhole. The peak discharge of floods that overflow the dykes experience a significant downstream decrease between Castejón and Zaragoza gauging stations, around 85 km apart, and a twofold increase in the propagation time of the hydrograph crest due to the laminating and retarding effect of the floodplain (Espejo et al., 2008). Moreover, some floodplain areas located beyond the dykes are frequently flooded by quasi-static waters related to water table rise. The magnitude and frequency relationships of floods have changed substantially in the second half of the 20th century, involving a significant hazard reduction, mainly due to the construction of reservoirs (Batalla et al., 2004; Cabezas et al., 2009). The storage capacity of reservoirs upstream Zaragoza is around 2700 hm³, mainly related to the Reiosa (540 hm³, 1945), Yesa (447 hm³, 1960) and Itoiz reservoirs (586 hm³, 2002).

Methodology

Before exploring the potential paleoflood record associated with the large depression and the nested pond, we applied several methods in order to characterise this complex sinkhole. Priority was given to the following issues: (1) definition of the edges of the large depression and the nested sinkhole pond; (2) identification of potential paleoflood deposits along the edge of the terrace, which is affected by the large sinkhole, and the buried sinkhole pond; (3) internal geometry of the compound sinkhole; (4) long-term evolution of the subsidence structure and current activity. Initially, we analysed old aerial photographs and topographic maps that show the large depression and the nested sinkhole pond before its filling (**Figs. 4 and 5**). The aerial photographs were orthorectified and georeferenced with a GIS in order to map the edges of the infilled

depressions. Particularly useful were images taken in 1927 (Fotoplano of the Ebro Basin Water Authority), 1946 (American flight A), 1957 (American flight B), and 1970 (**Fig. 4**). The location of the sinkhole pond was also constrained using old detailed topographic maps of the Municipality of Zaragoza produced in 1969 (1:2,000 scale, 1 m contour interval) and 1971-73 (1:1,000 scale, 2 m contour interval) (**Fig. 5**).

A detailed field survey was conducted in the area, paying especial attention to the identification of evidence of recent deformation, both on the ground surface and in the human-built structures (**Fig. 6**). These features were used to refine the limits of the area affected by subsidence and assess its activity. Subsequently, we excavated five trenches, four of them on the terrace next to its riser (T1-T4), and another one in the floodplain, within a crop field in the western sector of the large depression (T5). Trenches T1 to T4 were aimed at defining the southern edge of the large sinkhole and exploring the probable existence of paleoflood deposits on a minor bench associated with the terrace scarp. Trenches were excavated with a rubber-tired backhoe. The vertical walls were cleaned and one of them was logged on graph paper using a reference grid with horizontal and vertical strings spaced 1 m apart. Samples were collected for AMS dating giving priority to the most adequate stratigraphic units for determining the age of flood deposits. The deformation and sedimentation history recorded in trench T1 was reconstructed by performing a retrodeformation analysis (e.g., McCalpin, 2009a; Carbonel et al., 2014, 2015).

Ground deformation data derived from DInSAR (Differential Interferometry Synthetic Aperture Radar) displacement rate maps have been integrated in the sinkhole characterization (**Fig. 6**). Subsidence rates were derived from 29 ENVISAT ASAR images acquired from May 2, 2003 to September 17, 2010. The data were processed using the Stable Point Network (SPN) Persistent Scatters (PS) approach (Arnaud et al.,

2003; Crosetto et al., 2008), through which coherent points (20 m pixels) were identified considering a coherence threshold of >0.46 and a deviation amplitude of <0.50 . In this study we have considered subsidence rates higher than 2 mm/yr, based on the assessed error margin (Galve et al., 2015).

An electrical resistivity section was acquired across the buried sinkhole pond (i.e. Griffiths and Barker, 1993; Loke et al., 2013). The bulk electrical resistivity in alluvial environments is largely governed by the grain size of the different units. Therefore, ERT was a priori an adequate method to image the clayey lacustrine-palustrine facies of the large subsidence depression and the nested sinkhole pond, overlain by coarse and porous anthropogenic deposits and underlain by fluvial gravels. Moreover, this geophysical method is particularly useful for obtaining information on the internal geometry of sinkholes below the investigation depths of other techniques (e.g. ground penetrating radar, trenching; Carbonel et al., 2014; Zarroza et al., 2016). The ERT profile was acquired with a multi-electrode Lund Imaging System (ABEM) and an interelectrode spacing of 2 m. The system is composed of a Terrameter SAS 4000 resistivimeter, electrode selector ES10-64 and 4 signal cable and reels and 64 steel electrodes. We selected a Dipole-Dipole electrode array (Dahlin and Zhou, 2004) and the measured apparent resistivity data were processed using the 2D finite-difference inversion commercial software RES2DINV of Loke and Barker (1996) to produce the resistivity image.

A 20 m long borehole with full-core recovery was drilled in the central sector of the sinkhole pond filled between 1970 and 1984. Some deposits of the core underwent considerable compaction during the drilling operations, especially the clayey lacustrine facies. To construct the stratigraphic log, the original thickness of the different units was estimated by applying a backstripping factor for each section of the core given by the

ratio between the length of the borehole section versus the length of the retrieved compacted core. This factor was generally lower than 1.5. The core was longitudinally halved in the laboratory and the stratigraphic units were differentiated by visual examination, mainly on the basis of their textural characteristics, colour, and internal structure. Three charcoal samples were collected for AMS dating. The obtained conventional radiocarbon ages were calibrated using CALIB 7.1 and the dataset IntCal 13 (**Table 2**). The number of dated samples was constrained by budget limitations, rather than by the availability of datable material.

Sinkhole characterisation

Aerial photographs, topographic maps and field surveying

The aerial photographs from 1927, 1957 and 1970 show a WNW-ESE-oriented shallow depression with palustrine vegetation around 600 m long and 240 m wide (**Fig. 4**). This large sinkhole is apparently confined to the floodplain with its southern edge located at the foot of the lower terrace (Simón et al., 1998; Galve et al., 2009). The northern edge of the topographic depression is defined by rectilinear scarps associated with crop fields, suggesting that its extent has been reduced by anthropogenic filling in order to increase the cultivable land. The three aerial photographs clearly show a nested sinkhole pond around 40 m in diameter on the eastern half. The images from 1970, with higher resolution, allow the recognition of another pond, apparently shallower, nowadays located beneath the A-68 highway (**Fig. 6**). Aerial photographs from 1984 show the depressions completely buried and a large fuel plant built along its northern margin; i.e. the sinkhole was obliterated sometime between 1970 and 1984. The detailed topographic maps produced in 1969 and 1971-73 roughly depict the large depression with dashed contour lines, and the nested sinkhole is represented as a pond (*balsa*) 80

and 40 m across, respectively (**Fig. 5**). Elevation data in the 1971-73 map indicate that the large depression was at least 1.5 m deep. Both maps show an old drainage ditch (*escorredero*) that used to flow into the pond.

The walls of the factory located in the western sector of the depression show numerous curved subhorizontal cracks with apertures of up to 10 mm indicative of basal support loss and differential settlement (**Fig. 6**). Severe subsidence damage was detected in a concrete ditch and buildings located on the terrace, next to its riser. The presence of these recent deformation features on the northern sector of the terrace led us to consider that the sinkhole is not confined to the floodplain, and that its southern edge might be located on the terrace. This interpretation would help to explain a secondary bench associated with the terrace scarp, which might include paleoflood deposits.

Trenching and slackwater paleoflood deposits

In order to test the hypothesis indicated above on the southern boundary of the sinkhole, we excavated four trenches (T1-T4) along the northern edge of the terrace and oriented roughly perpendicular to the scarp (**Fig. 6**). Trenches T2, T3 and T4, up 3.5 m deep and around 41, 22 and 12 m long, respectively, exposed crudely-bedded terrace gravels with no evidence of deformation.

Trench T1 was excavated with a N12E orientation next to a building affected by conspicuous cracks parallel to the adjacent terrace scarp (**Fig. 7**). The excavation was 25 m long and reached a depth of 2.4 m, limited by flooding of the trench bottom by groundwater coming from the gravels exposed in the southern sector (perched aquifer). The terrace gravels occur in two blocks separated by a deformation zone around 1.5 m wide. This deformation band comprises, from N to S: (1) an oversteepened down-to-the north normal fault with an associated shear zone overlain by a colluvial wedge; and (2)

a downward tapering fissure fill opened between the terrace gravels of the footwall and the colluvial wedge underlain by the shear zone of the downthrown block; i.e. the fissure clearly postdates the colluvial wedge. The shear zone, 0.6 m wide, is made up of terrace gravels with reoriented fabrics and its northern edge is defined by a fault plane dipping antithetically 65S. This pseudo-reverse fault is in fact a normal fault rotated towards the sinkhole during the development of the fissure. The colluvial wedge is 0.4 m thick and 2 m long and its base dips around 8° to the north. It is made up of poorly-sorted and matrix-rich rounded gravels shed from the fault scarp underlain by terrace gravels. The orientation of the planar clasts is roughly concordant with the top boundary of the wedge. The fissure fill, 1 m wide in the upper part, consists of a well-cemented deposit including two units separated by a downward-pointing v-shaped contact. The lower unit is gravel with silt matrix and chaotic fabrics and the upper unit massive marly silts. The top of the terrace gravels on both sides of the deformation band (shear zone and fissure) show a vertical separation of 0.7 m. This is consistent with the rule of thumb whereby the thickness of colluvial wedges tend to be half of the height of the fault scarps from which they are derived (McCalpin, 2009b). Most probably the top of the terrace gravels in the footwall has experienced some erosion. The vertical separation of the upper contact of the terrace gravels rises to 1.75 m considering the whole length of its exposure in the trench.

In the downthrown block, there is a fine grained-deposit that overlies the terrace gravels and the colluvial wedge. This unit pinches out just at the northern edge of the fissure. This deposit consists of light brown massive fines, from coarse sand to silt, with normal grading and abundant gastropods. In the northern sector of the trench it includes thin lenticular beds of pebbly-granule gravel (**Fig. 7**). Shells collected 35 cm below the top of this unit have yielded a calibrated age of 7004-6788 cal yr BP (error margin at two

sigma). This unit is interpreted as a mid-Holocene slackwater paleoflood deposit (Baker, 1987) of the Ebro River, most probably occurred when the river channel was located at a higher elevation; the deposit lies at approximately 6 m above the current floodplain. The paleoflood deposit is tilted towards the sinkhole. The top of the unit dips 3-4° and the dip of the intercalated gravel bed reaches 5-6° in the northern sector of the trench. Moreover, the deposit is affected by a small monoclinal fold around the vertical reference lines 5-7, probably related to a secondary buried normal fault (drape fold). The top of the unit shows an elevation difference of 1.2 m along the trench. The deformed units of the footwall, deformation zone and downthrown block are overlain by two apparently non-deformed anthropogenic deposits that thicken towards the north, indicating that the sinkhole at this site used to have geomorphic expression in historical times.

The stratigraphic and structural relationships observed in the trench allow us to infer the following deformational and depositional history, as illustrated in a simplified retrodeformation sequence shown in **figure 8**. A first collapse event, older than 7 ka and controlled by a subvertical normal fault, caused the downdropping of a portion of the terrace and the generation of a lower bench. The displacement along the fault of around 1 m produced a shear zone, and erosion on the oversteepened fault scarp resulted in the development of the colluvial wedge. Subsequently, a flood of the Ebro River, dated at 7004-6788 cal yr BP, deposited a package of fine-grained sediments (slackwater deposits) on the bench onlapping the colluvial wedge. Later, the downthrown block experienced a rotation towards the sinkhole centre, with the consequent opening of a fissure. The two units in the fissure fill with a synformal structure may be related to downward ravelling and compaction and/or two episodes of fissure opening. The total vertical displacement accommodated in this second deformation event in the trench

zone was around 1.2 m. Subsequently, the geomorphic expression of the sinkhole margin was masked by the accumulation of artificial deposits. Although these massive anthropogenic deposits do not show evidence of deformation, the conspicuous damage observed in the adjacent building reveals that this sector of the sinkhole is still active.

The southern edge of the large sinkhole was mapped integrating the distribution of subsidence damage on human structures and the trenching investigation, including a portion of the northern sector of the terrace (**Fig. 6**). The proposed map excludes trenches T2, T3 and T4, adjusts the edge to the deformation zone exposed in trench T1, and embraces the recorded subsidence damage.

Trench 5 was excavated in the NW sector of the large depression, within a crop field located in the floodplain (**Fig. 6**). The excavation, 6 m long and 2 m deep, exposed, from top to bottom: (1) a gravelly anthropogenic deposit 65 cm thick; (2) 115 cm of dark brown, massive and bioturbated mud with scattered granules and abundant gastropods and vegetation remains; and (3) the upper part of a bed of loose rounded gravel. Shells collected from the palustrine facies 20 cm above its base yielded a calendar age of 1705-1562 cal yr BP. The data provided by this trench reveals that this sector, although levelled by anthropogenic fill, forms part of the large depression, and that this part of the sinkhole was occupied by a palustrine environment in the Ebro River floodplain at least since the 3rd century.

DInSAR deformation data

The available DInSAR ground deformation data is essentially restricted to human structures and paved areas, due to lack of coherence in crop fields and wastelands, including the buried sinkhole pond. Deformation data points tend to be distributed in clusters with similar values, from which we have depicted the highest ones in **figure 6**.

Consistently with the geomorphological map, mean subsidence rates along the line-of-sight (LOS) on the western sector of the large depression, including the damaged factory, range from 2.3 to 4.8 mm/yr (**Fig. 6**). The DInSAR displacement rate map also captures stable pixels in buildings located outside the mapped boundary of the large sinkhole, like the large factory situated on the terrace. In apparent contradiction with our geomorphological map, the fuel plant shows numerous deformation points (2.2-4.5 mm/yr), mainly clustered on the SW side of the tanks. This disagreement may be related to artefacts caused by double-bounce reflection. Double-bounce is a type of scattering behaviour frequent in built-up areas, and depends on the geometrical relationship between the sensor flight line and the feature orientation (Henderson and Lewis, 1998). Considering the ascending right-looking pass of the images and the incidence angle of the ENVISAT SAR satellite (20 -30°), clusters of double-bounce PS points may have formed due to the bouncing effect of the tank walls (ca. 16 m high).

Electrical resistivity tomography

The 126 m long electrical resistivity section was acquired across the buried sinkhole pond with an ENE-WSW orientation and approximately centred on its middle point (**Figs. 6 and 9**). The interpretation of the resistivity model has been aided by knowledge on the geomorphic setting and the 20 m deep borehole drilled in the centre of the sinkhole pond, which traversed, from top to bottom, three sedimentary packages: (1) a coarse man-made fill 3.9 m thick; (2) a sinkhole fill 7.85 m thick consisting of clayey facies with some gravel intercalations; and (3) fluvial gravels more than 8.25 m thick (the base of the alluvial cover was not reached). Consistently with the borehole data, the resistivity image shows three electro-layers. The upper layer with medium-high resistivity (50 - 500 Ω m) corresponds to the anthropogenic deposits. In the central

sector of the buried sinkhole pond, between stations 52 and 62, it displays an abrupt thickening (ca. 6 m) that might be related to a collapse structure nested within the sinkhole pond. The intermediate low-resistivity layer ($<10 \Omega\text{m}$) is ascribed to clayey sediments deposited in the palustrine and lacustrine environments developed in the large depression and the inner sinkhole pond, respectively. The thickness of this unit increases from 1-4 m in the margins of the section, to around 8 m in the sinkhole pond, consistently with the borehole log. Its base, especially in the western half of the resistivity image, shows abrupt drops towards the sinkhole centre attributable to collapse faults. The central sector where the intermediate layer reaches the highest thickness is approximately 40 m wide, in agreement with the diameter of the sinkhole pond. The lower medium-high resistivity layer ($>75 \Omega\text{m}$) corresponds to the coarse fluvial gravel logged in the borehole. The stepped geometry of the top of this unit, as well as the heterogeneous distribution of the resistivity, should be related to the aforementioned collapse faults. Overall, the resistivity image indicates that this is a large sagging sinkhole (large depression) with an inner collapse structure (sinkhole pond) controlled by a master ring fault around 40 m in diameter, and other secondary outer and inner collapse faults.

Sinkhole pond fill and paleoflood record

Three main sedimentary packages comprising 35 stratigraphic units were differentiated in the 20 m long borehole core obtained from the central sector of the buried sinkhole pond (**Figs. 6 and 10**). The uppermost package, corresponding to unit U1, is an anthropogenic deposit 3.9 m thick consisting of massive and chaotic matrix-rich gravel and rubble. The thickness of this artificial fill, dumped between 1970 and 1984, indicates that by that time the sinkhole pond was around 3-4 m deep. The lowest

package, designated as unit U35, is composed of more than 8.25 m of relatively well-sorted, rounded, polymict pebble-cobble gravel with sand matrix, ascribed to channel facies. The borehole did not reach the base of the Quaternary cover. The 7.85 m thick intermediate package includes units U2 to U34. This is a natural sinkhole fill, mainly composed of dark clays with intercalations of gravel and silt-sand beds. Four lithofacies types have been differentiated in the intermediate package:

(I) Dark brown-grey clay, commonly massive, but locally vaguely laminated. These soft clays generally contain plant remains, gastropods and traces of bioturbation. The layers of this facies range from 2 to 59 cm thick and have an aggregate thickness of 5.5 m (67% of the natural sinkhole fill). This facies records low-energy lacustrine deposition in the freshwater sinkhole pond with relatively high production of organic matter.

(II) Thick beds of massive and rounded pebble gravel with a high proportion of light brown sand-silt matrix. This gravel facies, intercalated within the clayey lake deposit, is represented by units U9 and U15, with thicknesses of 67 and 76 cm, and clasts up to 8 and 4 cm long, respectively. Facies II represents 17% of the natural sinkhole fill. These beds are ascribed to two major Ebro River floods, during which the flood waters reached flow competence conditions high enough to transport gravels across the low gradient floodplain. Sudden water depth increase at the sinkhole pond and the consequent energy dissipation prompted rapid sediment load deposition.

(III) Thin beds (2-8 cm thick) of rounded pebble gravel with clayey silt matrix (units U3 and U7).

(IV) Silt-sand beds 6-69 cm thick with rounded pebble-sized clasts up to 4 cm long.

Facies III and IV can be attributed to different processes. Our preferred interpretation is that they correspond to flood events with significantly lower competence than those recorded by the thick gravel beds. Unit U14, U10 and U16, in contact with the thick

gravel beds, may have been deposited during the two large floods, before and after peak flow conditions. Other plausible interpretations, considering the geomorphic context, include storm-derived water flows coming from the adjacent terrace scarp (**Fig. 6**) and accumulation of alluvium transported by subaqueous mass movements from the scarped margins of the collapse sinkhole.

Charcoal samples from units U34 (base of intermediate package) and U16 (overlain by gravel unit U15) yielded calibrated ages of 2759-2681 and 1537-1311 cal yr BP, respectively (**Table 2**). The sample from the top of unit U10 (overlain by gravel unit U9) provided a ^{14}C concentration of 131 ± 039 pMC, indicating an age younger than 1950. The numerical age obtained from the top of unit U10 (>1950), situated just below the youngest thick gravel bed (U9), strongly suggests that the latter was deposited during the Great Ebro River flood of January 1961. The age range obtained for unit U16 (1537-1311 cal yr BP, error margin at 2 sigma) should cover the calendar age of the flood recorded by the overlying thick gravel bed U15. This paleoflood, occurred in Visigothic times, is older than the oldest flood documented from historical data, which dates back to AD 827 (CNPC, 1983).

The numerical ages also allow inferring the following information on the evolution of the sinkhole: (1) The development of the collapse sinkhole and the creation of the sinkhole pond occurred at around 2759-2681 cal yr BP. (2) An average long-term subsidence rate of around 4.2-4.6 mm/yr can be computed considering the age of the sinkhole and the cumulative subsidence given by the aggregate thickness of the two upper packages (11.7 m). Although subsidence in the sinkhole is probably characterised by episodic kinematics, this value is consistent with the rates measured and estimated by various methods in active sinkholes of the salt-bearing evaporite karst of the Ebro Valley (e.g., Galve et al., 2009, 2015; Desir et al., 2016a, b). (3) Average aggradation

rates between deposition of units U34-U16 and U16-U10 were approximately 3.4 and 1.6 mm/yr, respectively. The overall vertical accretion rate has decreased more than 50% in the last 1,4000 yr, despite that the two thick gravel beds (143 cm), which represent quasi-instantaneous deposition, were accumulated within this time span.

Discussion

The subsidence depression used to explore the potential of sinkholes in alluvial systems as archives for flood events is in the mantled evaporite karst of the Ebro River valley, upstream of Zaragoza city, NE Spain. The investigation carried out integrating data gathered by geomorphological mapping, surveys of structural damage, trenching, ERT, DInSAR, and a borehole reveal the following features on the selected sinkhole:

(1) The depression is a complex sinkhole comprising a large diffuse-edged basin around 600 m long and 240 m wide associated with the southern edge of the floodplain and the adjacent terrace, and a main nested collapse sinkhole that used to host a permanent lake approximately 40 m across (**Fig. 6**).

(2) The spatial distribution of damage on human structures and a trench dug next to the terrace riser indicates that the southern edge of the large depression is located on the terrace, as revealed by a buried scarp, a colluvial wedge, and a fissure fill exposed in the trench. Episodic dissolution-induced subsidence on the terrace created a bench associated with the terrace riser, on which at least one Ebro River paleoflood, dated at 7004-6788 cal yr BP, accumulated fine-grained slackwater deposits (**Figs. 7 and 8**).

(3) The sinkhole pond, located in the floodplain at the foot of the terrace scarp, was filled by anthropogenic deposits sometime between 1970 and 1984 (**Fig. 4**). By that time, water depth, controlled by the water table of the alluvial aquifer, was around 3-4 m. This pond, as suggests the ERT section, was the geomorphic expression of a collapse

structure, probably comprising a master ring fault around 40 m in diameter, and other outer and inner secondary concentric faults (**Fig. 9**).

(4) Multiple lines of evidence indicate that both the large depression and the nested collapse sinkhole are currently active. DInSAR ground deformation data indicate LOS displacement rates of the order of 2-5 mm/yr in some sectors of the large depression (**Fig. 6**). Consistently, the numerical age of the oldest unit of the sinkhole-pond fill (2759-2494 cal yr BP), located at a depth of 11.7 m, indicate a long-term subsidence rate of 4.2-4.6 mm/yr (Fig. 9). These high subsidence rates strongly suggest that the sinkhole is related to interstratal dissolution of high-solubility salts (halite and or glauberite) (e.g., Castañeda et al., 2009; Galve et al., 2015; Desir et al., 2016).

Although unexpectedly, trench T1 dug next to the terrace scarp exposed a mid-Holocene slackwater paleoflood deposit accumulated in a peculiar geomorphic setting; i.e., an inset bench formed by collapse faulting at the margin of a large karst depression. The age of the slackwater deposits (7004-6788 cal yr BP) is older than the stratigraphic record of the sinkhole pond and falls within a period of increased fluvial activity in Spain situated at 7980-6860 cal yr BP by Thorndycraft and Benito (2006), using a compilation of 74 Holocene radiocarbon dates. These sediments allow obtaining information on the chronology of a pre-historical flood, but they cannot be used as paleostage indicators for paleodischarge estimation due to two reasons: (1) The slackwater deposits have been affected by post-sedimentary subsidence (>1.2 m). (2) The sediments occur in a non-stable alluvial valley reach, whose bottom was probably at higher elevation around 7 kyr ago.

The limited radiocarbon dates obtained from the sinkhole pond deposits provide a fairly week age-depth structure. Nonetheless, the available numerical ages allow inferring relevant features related to the subsidence and sedimentation patterns, including

paleoflood deposits. Sedimentation in the sinkhole pond between 2759-2681 cal yr BP and 1970-1984 was dominated by lacustrine dark clayey facies, eventually interrupted by the accumulation of three types of coarser detrital facies: (1) two thick pebble gravel beds (>0.5 m thick); (2) thin layers of pebble gravel (<10 cm thick); and (3) silt-sand units with scattered pebble-sized clasts. The thick gravel beds, and probably the underlying and overlying sand-silt facies, record two major Ebro River floods with an inter-event time lapse of 1,300-1,550 yr. During these events, fine gravel transported across the floodplain was rapidly accumulated in the sinkhole pond. Local flow energy dissipation at this sediment trap due to sharp increase in water depth caused rapid gravity-controlled deposition of massive gravelly facies with abundant sand-silt matrix. To our knowledge, this is the first paleoflood record from a sinkhole-fill documented in the literature. The lower thick gravel bed records a paleoflood probably occurred in Visigothic times between 1537-1311 cal yr BP, before the oldest flood documented with historical data, which dates back to AD 827 (**Table 1**; CNPC, 1985). The upper one is younger than 1950 as indicated by radiocarbon dating, and older than the anthropogenic infill of the sinkhole pond bracketed at 1970-1984. Consequently, this unit is ascribed to the 1961 Great Ebro River flood, with an estimated peak discharge of 4,130 m³/s. This is by far the biggest flood of the instrumental period starting in 1913, and probably larger than all the historical events (**Table 1**; Mejón 2011). The Ebro River Basin Water Authority, based on the annual maximum discharge values recorded from 1943, estimates a return period of around 80 years for the 1961 flood. Probably, that event is an outlier that biases the magnitude and frequency scaling relationships, leading to overestimated discharge values for high recurrence periods (e.g., Webb et al., 2002). The thin gravel beds and the sand-silt units separated from the thick gravel beds by lacustrine facies may have been generated by lower-competence floods at the site.

Nonetheless, they could also be related to other processes such as storm-derived water flows coming from the adjacent terrace scarp, or subaqueous sediment-gravity flows originated from failures in the scarped margin of the collapse sinkhole. The palustrine facies dated in trench 5 are probably correlative to the fine-grained facies (units 14, 13) that overlie the thick gravelly paleoflood deposit of the sinkhole pond. There is the possibility that the gravels underlying the palustrine facies of trench 5 could correspond to the first major paleoflood identified in the deposits of the sinkhole pond.

The investigated geomorphic settings have significant limitations as paleoflood recorders that affect their usefulness for flood-frequency analysis. Sinkhole sediments may considerably under-represent the paleoflood history. The potential for a flood to be recorded in a sinkhole pond located in a broad floodplain and the sedimentologic signature of the deposits may depend on factors independent of the magnitude of the event, like the distance to the shifting channel, presence of riparian and palustrine vegetation (**Fig. 3**), flood duration or the geometry and size of the sinkhole. Moreover, no flood magnitude estimates can be obtained from these stratigraphic records since they do not allow inferring paleostage data. Paleocompetence methods based on the size of the largest particles may be applied to roughly estimate flow velocity (e.g., Maiziels, 1983; Williams, 1984), but this variable may be also affected by some of the local factors indicated above.

Despite the aforementioned limitations, sinkhole ponds have some advantages: (1) In the case of active sinkholes, long-sustained subsidence higher than the aggradation rate facilitates the accumulation and preservation of long and continuous records with no erosional hiatuses. The long-term subsidence rate estimated for the investigated sinkhole is 4.2-4.6 mm/yr, whereas average aggradation rates are 2.8-3.1 mm/yr and 2-2.2 mm/yr considering and ignoring the thick detrital units intercalated within the

lacustrine facies, respectively. (2) The chronology of the paleofloods deposits identified in borehole cores can be accurately constrained thanks to the generally high dating potential of the associated lacustrine facies rich in organic remains. In some areas, the success of these investigations may be hindered by the recency of the sinkholes, as well as by high sedimentation rates that overwhelms dissolution-induced subsidence. (3) Oxbow lakes could be used to infer paleoflood histories, as satisfactorily illustrated by Oliva et al. (2016) in the Désert River, southwestern Québec, Canada. However, in the studied reach of the Ebro River Valley, most of the lakes related to abandoned meanders date back to the second half of the 20th Century, and consequently they cannot be used as archives of prehistorical floods, in contrast with the sinkhole ponds that may be older than 2,700 yr (**Fig. 10**). (4) Sinkhole ponds in many regions have been commonly filled by anthropogenic deposits in the last few decades (e.g. Gutiérrez, 2016). These buried depressions, in which drilling is particularly easy, are excellent candidates for paleoflood investigations. (5) Sinkholes may be found in a considerable proportion of fluvial systems worldwide, considering that karst rocks underlie around 20% of the Earth's continental surface (Ford and Williams, 2007). By extending the temporal length of the flood catalogues with this information, more reliable frequency estimates for high-competence floods could be calculated. Moreover, temporal clusters of paleofloods may be identified attributable to climatic forcing and/or anthropogenic impacts (e.g., Baker, 2003). Further investigations integrating the information from multiple cores obtained in several sinkhole lakes with variable characteristics and situated in different sectors of the floodplain (**Figs. 1 and 3**) would help to get deeper insight into the potential of these landforms for paleoflood studies and flood hazard analyses.

The cores derived from sinkhole lakes may also provide practical information on the subsidence phenomenon and may be used for paleoenvironmental investigations. As this study illustrates, quantitative data of interest for sinkhole hazard analysis that may be inferred from the numerically dated successions include: (1) cumulative subsidence magnitude; (2) age of the sinkhole; (3) long-term subsidence rates; and (4) variations in aggradation and subsidence rates through time. A great part of the Holocene paleoenvironmental investigations carried out in the central sector of the Ebro Depression have been focused on playa-lakes largely related to wind deflation. These geomorphic settings and the associated stratigraphic records pose important problems, including the presence of long hiati related to wind erosion and the scarcity of datable material (e.g., González-Sampériz et al., 2008; Gutiérrez et al., 2013). However, the infill of sinkhole lakes offers continuous and datable records, whose temporal length can be optimised by integrating information from sinkholes of different ages (e.g., floodplain and terraces).

Conclusions

Sinkhole fills in alluvial systems may be used as archives for paleoflood events. The investigated complex sinkhole illustrates that dissolution-induced subsidence may create two types of geomorphic settings with the potential of preserving stratigraphic records of paleoflood deposits:

- Benches generated by collapse faulting at the edge of low terraces, where fine-grained slackwater paleoflood deposits may be accumulated. These deposits may have limitations as paleostage indicators for paleodischarge estimation due to two main reasons: (1) they may have been affected by post-sedimentary subsidence; (2) occur on non-stable valley reaches.

- Sinkhole ponds located in the floodplain, where the background autochthonous fine-grained sedimentation may be suddenly interrupted by the accumulation of allochthonous coarse-grained deposits during major flood events.

The use of sinkhole ponds as recorders of past floods has a number of advantages: (1) Long-sustained active subsidence contributes to the preservation of continuous stratigraphical successions with no erosional hiatuses. In general, sinkholes related to evaporite dissolution are more favourable since they are typically affected by higher subsidence rates. (2) Generally, paleoflood chronology can be resolved thanks to the high dating potential of the lacustrine facies and the presence of organic remains in the detrital allochthonous facies. (3) The paleoflood investigations could be conducted integrating data gathered from multiple sinkhole ponds located within the same valley reach, provided precise chronologies are available. The investigations could include sinkholes of different ages, whose aggregate record could cover long paleoflood histories. The selected sinkholes could be also located on variable morpho-stratigraphic settings (e.g., floodplain, low terraces) where flood deposits may have different sedimentological signatures and their formation is controlled by variable paleodischarge thresholds.

Nonetheless, they also have significant drawbacks related to the nature of the geomorphic context: (1) paleoflood deposits do not allow inferring paleostage data, although paleocompetence methods could be applied; (2) floods of similar magnitude may produce significantly different sedimentological signatures due to multiple factors such as the variable distance between the sinkhole and the river channel and the distribution and density of riparian and palustrine vegetation; (3) the investigation of these deposits, typically located in the subsurface and beneath the water-table, need to be investigated via boreholes.

685

686 Before using sinkholes for the investigation of past floods, it is highly recommended to
687 characterise the subsidence depressions integrating data gathered by multiple
688 approaches such as geomorphic mapping, detailed filed surveying shallow geophysics
689 and trenching.

690

691 **Acknowledgements**

692 This work has been funded by project CGL2013-40867-P (Ministerio de Economía y
693 Competitividad, Spain). We are also grateful to Mr. Octavio Plumed from the company
694 ENSAYA for his support for the borehole drilling. MZ has a position at the Universidad
695 Autónoma de Barcelona as a Serra Hunter Fellow. The work conducted by CC has been
696 supported by project PCIN-2014-106. We thank Kelsey Budahn (Akron University,
697 Ohio) for improving the English of the manuscript.

698

References

- Arnaud, A., Adam, N., Hanssen, R., Inglada, J., Duro, J., Closa, J., Eineder, M., 2003. ASAR ERS interferometric phase continuity. Proceedings of the International Geoscience and Remote Sensing Symposium, Toulouse, pp. 1133–1135.
- Baker, V.R., 1987. Paleoflood hydrology and extraordinary flood events. *Journal of Hydrology* 96, 79-99.
- Baker, V.R., 2003. Paleofloods and extended discharge records. In: Gregory, K.J., Benito, G. (Eds.), *Paleohydrology. Understanding Global Change*. Wiley, Chichester, pp. 307-323.
- Baker, V.R., Webb, R.H., House, P.K., 2002. The scientific and societal value of paleoflood hydrology. In: House, P.K., Webb, R.H., Baker, V.R., Levish, D.R. (Eds.), *Ancient Floods, Modern Hazards. Principles and Applications of Paleoflood Hydrology*. American Geophysical Union. *Water Science and Applications* 5, pp. 1-19.
- Barreiro-Lostres, F., Moreno, A., Giralt, S., Caballero, M., Valero-Garcés, B., 2014. Climate, palaeohydrology and land use change in the Central Iberian Range over the last 1.6 kyr: The La Parra Lake record. *The Holocene* 24, 1177-1192.
- Batalla, R.J., Gómez, C.M., Kondolf, G.M., 2004. Reservoir-induced hydrological changes in the Ebro River basin (NE Spain). *Journal of Hydrology* 290, 117-136.
- Benito, G., Lang, M., Barriendos, M., Llasat, C., Francés, F., Ouarda, T., Thorndycraft, V.R., Enzel, Y., Bardossy, A., Coeur, D., Bobée, B., 2004. Use of systematic, paleoflood and historical data for the improvement of flood risk estimation. Review of scientific methods. *Natural Hazards* 31, 623-643.
- Benito-Calvo, A., Gutiérrez, F., Carbonel, D., Desir, G., Guerrero, J., Magri, O., Karampaglidis, T., Fabregat, I., 2016. Measuring deformation related to active

724 sinkholes with ground-based 3D laser scanner. A case study in the evaporite karst of
 725 the Ebro Valley, NE Spain. In: Durán, J.J., Montes, M., Robador, A., Salazar, A.
 726 (Eds.), *Comprendiendo el relieve: Del pasado al futuro*. XIV Reunión Nacional de
 727 Geomorfología, Málaga, pp. 599-606.

728 Brown, A.L., Reinhardt, E.G., van Hengstum, P.J., Pilarczyk, J.E., 2014. A coastal
 729 Yucatan sinkhole records intense hurricane events. *Journal of Coastal Research* 30,
 730 418-428.

731 Cabezas, A., Comín, F.A., Beguería, S., Trabucchi, M., 2009. Hydrologic and landscape
 732 changes in the Middel Ebro River (NE Spain): implications for restoration and
 733 management. *Hydrology and Earth System Sciences* 13, 273-284.

734 Calvo, J.P., Pozo, M., Silva, P.G., Morales, J., 2013. Pattern of sedimentary infilling of
 735 fossil mammal traps formed in pseudokarst at Cerro de los Batallones, Madrid Basin,
 736 central Spain. *Sedimentology* 60, 1681-1708.

737 Carbonel, D., Rodríguez, V., Gutiérrez, F., McCalpin, J.P., Linares, R., Roqué, C.,
 738 Zarroca, M., Guerrero, J., 2014. Evaluation of trenching, ground penetrating radar
 739 (GPR) and electrical resistivity tomography (ERT) for sinkhole characterization.
 740 *Earth Surface Processes and Landforms* 39, 214-227.

741 Carbonel, D., Rodríguez-Tribaldos, V., Gutiérrez, F., Galve, J.P., Guerrero, J., Zarroca,
 742 M., Roqué, C., Linares, R., McCalpin, J.P., Acosta, E., 2015. Investigating a
 743 damaging buried sinkhole cluster in an urban area integrating multiple techniques:
 744 geomorphological surveys, DinSAR, GPR, ERT, and trenching. *Geomorphology*
 745 229, 3-16.

746 Carbonell, E., Bermúdez de Castro, J.M., Parés, J.M., Pérez-González, A., Cuenca-
 747 Bescos, G., Ollé, A., Mosquera, M., Huguet, R., van der Made, J., Rosas, A., Sala,
 748 R., Vallverdú, J., García, N., Granger, D.E., Martínón-Toeers, M., Rodríguez, X.P.,

749 Stock, G.M., Vergès, J.M., Allué, E., Burjachs, F., Cáceres, I., Canals, A., Benito,
750 A., Díez, C., Lozano, M., Mateos, A., Navazo, M., Rodríguez, J., Rosell, J., Arsuaga,
751 J.L., 2008. The first hominid of Europe. *Nature* 452, 465-469.

752 Castañeda, C., Gutiérrez, F., Manunta, M., Galve, J.P., 2009. DInSAR measurements of
753 ground deformation by sinkholes, mining subsidence, and landslides, Ebro River,
754 Spain. *Earth Surface Processes and Landforms* 34, 1562–1574.

755 CHE, 2015. Informe sobre las avenidas del primer trimestre de 2015 en la Cuenca del
756 Ebro. Confederación Hidrográfica del Ebro, Zaragoza.
757 <http://www.chebro.es/contenido.visualizar.do?idContenido=43453&idMenu=4840>

758 CNPC, 1985. Estudio de inundaciones históricas. Mapa de riesgos potenciales. Cuenca
759 del Ebro. Comisión Técnica de Inundaciones. Gobierno de España, Madrid.

760 Crosetto, M., Biescas, E., Duro, J., Closa, J., Arnaud, A., 2008. Generation of advanced
761 ERS and Envisat interferometric SAR products using the Stable Point Network
762 technique. *Photogrammetric Engineering and Remote Sensing* 74, 443.

763 Dahlin, T., Zhou, B., 2004. A numerical comparison of 2D resistivity imaging with 10
764 electrode arrays. *Geophysical Prospecting* 52, 379-398.

765 Desir, G., Guerrero, J., Gutiérrez, F., Carbonel, D., Merino, J., Benito, A., Fabregat, I.,
766 Roqué, C., Zarroca, M., Linares, R., 2016a. Monitorización de dolinas activas en el
767 entorno de Zaragoza mediante nivelación geométrica de alta precisión. In: Durán,
768 J.J., Montes, M., Robador, A., Salazar, A. (Eds.), *Comprendiendo el relieve: Del*
769 *pasado al futuro*. XIV Reunión Nacional de Geomorfología, Málaga, pp. 607-613.

770 Desir, G., Gutiérrez, F., Merino, J., Carbonel, D., Benito-Calvo, A., Guerrero, J.,
771 Fabregat, I., 2016b. Monitoring damaging sinkholes using high-precision levelling in
772 the salt karst of the central Ebro Basin, NE Spain. *Earth Surface Processes and*
773 *Landforms*, submitted.

774 Espejo, F., Domenech, S.M, Ollero, A., Sanchez. M., 2008. La crecida del Ebro de
 775 2007: Procesos hidrometeorológicos y perspectivas de gestión del riesgo. Boletín de
 776 la Asociación de Geógrafos Españoles 48, 129-154.

777 Ford, D.C., Williams, P., 2007. Karst Hydrogeology and Geomorphology. Wiley,
 778 Chichester.

779 Galve, J.P., Gutiérrez, F., Lucha, P., Bonachea, J., Cendrero, A., Gimeno, M.J.,
 780 Gutiérrez, M., Pardo, G., Remondo, J., Sánchez, J.A., 2009. Sinkholes in the salt-
 781 bearing evaporite karst of the Ebro River valley upstream of Zaragoza city (NE
 782 Spain). Geomorphological mapping and analysis as a basis for risk management.
 783 Geomorphology 108, 145-158.

784 Galve, J.P., Castañeda, C., Gutiérrez, F., 2015. Railway deformation detected by
 785 DInSAR over active sinkholes in the Ebro Valley evaporite karst, Spain. Natural
 786 Hazards and Earth System Science 3, 3967-3981.

787 Gischler, E., Shinn, E.A., Oschmann, W., Fiebig, J., Buter, N.A., 2008. A 1500-yr
 788 Holocene Caribbean climate archive from the Blue Hole, Lighthouse Reef, Belize.
 789 Journal of Coastal Research 24, 1495-1505.

790 González-Sampériz, P., Valero-Garcés, B.L., Moreno, A., Morellón, M., Navas, A.,
 791 Machín, J., Delgado-Huertas, A., 2008. Vegetation changes and hydrological
 792 fluctuations in the Central Ebro Basin (NE Spain) since the Late Glacial period:
 793 Saline lake records. Paleogeography, Paleoclimatology, Paleoecology 259, 157-181.

794 Griffiths, D.H., Barker, R.D., 1993. Two-dimensional resistivity imaging and modelling
 795 in areas of complex geology. Journal of Applied Geophysics 29, 211-226.

796 Guerrero, J., Gutiérrez, F., Galve, J.P., 2013. Large depressions, thickened terraces and
 797 gravitational deformation in the Ebro River valley (Zaragoza area, NE Spain).

798 Evidence of glauberite and halite interstratal karstification. *Geomorphology* 196,
799 162-176.

800 Gutiérrez, F., 2016. Sinkhole hazards. *Oxford Research Encyclopedia of Natural Hazard*
801 *Science*. Oxford University Press, Oxford.

802 Gutiérrez, F., Cooper, A.H., 2013. Surface morphology of gypsum karst. In: Frumkin,
803 A. (Ed.), *Treatise on Geomorphology. Karst Geomorphology*. Elsevier, vol. 6, pp.
804 425-437.

805 Gutiérrez, F., Galve, J.P., Guerrero, J., Lucha, P., Cendrero, A., Remondo, J., Bonachea,
806 J., Gutiérrez, M., Sánchez, J.A., 2007. Typology, spatial distribution and detrimental
807 effects of the sinkholes developed in the alluvial evaporite karst of the Ebro River
808 valley downstream Zaragoza city. *Earth Surface Processes and Landforms* 32, 912-
809 928.

810 Gutiérrez, F., Guerrero, J., Lucha, P., 2008. A genetic classification of sinkholes
811 illustrated from evaporite paleokarst exposures in Spain. *Environmental Geology* 53,
812 993-1006.

813 Gutiérrez, F., Galve, J.P., Lucha, P., Bonachea, J., Jordá, L., Jordá, R., 2009.
814 Investigation of a large collapse sinkhole affecting a multi-storey building by means
815 of geophysics and the trenching technique, Zaragoza city, NE Spain. *Environmental*
816 *Geology* 58, 1107-1122.

817 Gutiérrez F., Galve, J.P., Lucha, P., Castañeda, C., Bonachea, J., Guerrero, J., 2011.
818 Integrating geomorphological mapping, trenching, InSAR and GPR for the
819 identification and characterization of sinkholes in the mantled evaporite karst of the
820 Ebro Valley (NE Spain). *Geomorphology* 134. 144–156.

821 Gutiérrez, F., Valero-Garcés, B., Desir, G., González-Sampériz, P., Gutiérrez, M.,
822 Linares, R., Zarroca, M., Moreno, A., Guerrero, J., Roqué, C., Arnold, L.J., Demuro,

823 M., 2013. Late Holocene evolution of playa lakes in the central sector of the Ebro
824 Depression based on geophysical surveys and morpho-stratigraphic analysis of
825 lacustrine terraces. *Geomorphology* 196, 177-197.

826 Gutiérrez, F., Parise, M., De Waele, J., Jourde, H., 2014. A review on natural and
827 human induced geohazards and impacts in karst. *Earth Science Reviews* 138, 61-88.

828 Gutiérrez, F., Mozafari, M., Carbonel, D., Gómez, R., Raeisi, E., 2015. Leakage
829 problems in dams built on evaporites. The case of La Loteta Dam (NE Spain), a
830 reservoir in a large karstic depression generated by interstratal salt dissolution.
831 *Engineering Geology* 185, 139-154.

832 Gutiérrez, F., Fabregat, I., Roqué, C., Carbonel, D., Guerrero, J., García-Hermoso, F.,
833 Zarroca, M., Linares, R., 2016. Sinkholes and caves related to evaporite dissolution
834 in a stratigraphically and structurally complex setting, Fluvia Valley, eastern Spanish
835 Pyrenees. Geological, geomorphological and environmental implications.
836 *Geomorphology* 267, 76–97.

837 Hart, E.A., 2014. Legacy sediment stored in sinkholes: a case study of three urban
838 watersheds in Tennessee, United States of America. *Physical Geography* 35, 514-
839 531.

840 Henderson, F.M., Lewis, A.J. 1998. *Manual of remote sensing: principles and*
841 *applications of imaging radars*. John Wiley and Sons, New York.

842 Hyatt, J.A., Gilbert, R., 2004. Subbottom acoustic and sedimentary records of past
843 surface water-groundwater exchange through sinkhole lakes in south Georgia, USA.
844 *Environmental Geology* 46, 32-46.

845 Kochel, R.C., Baker, V.R., 1982. Paleoflood hydrology. *Science* 215, 353-361.

846 Kochel, R.C., Baker, V.R., 1988. Paleoflood analysis using slackwater deposits. In:
847 Baker, V.R., Kochel, R.C., Patton, P.C. (Eds.), Flood Geomorphology. Wiley,
848 NewYork, pp. 357-402.

849 Lane, P., Donnelly, J.P., Woodruff, J.D., Hawkes, A.D., 2011. A decadal-resolved
850 paleohurricane record archived in the late Holocene sediments of a Florida sinkhole.
851 Marine Geology 287, 14-30.

852 Langer, H., Offermann, H., 1982. On the solubility of sodium chloride in water. Journal
853 of Crystal Growth 60, 389-392.

854 Laury, R. L., 1980. Paleoenvironment of a late Quaternary mammoth-bearing sinkhole
855 deposit, Hot Springs, South Dakota. Geological Society of America Bulletin 91, 465-
856 475.

857 Loke, M.H., Barker, R.D., 1996. Rapid least-squares inversion of apparent resistivity
858 pseudosections by a quasi-Newton method. Geophysical Prospecting 44, 131-152.

859 Loke, M.H., Chambers, J.E., Rucker, D.F., Kuras, O., Wilkinson, P.B., 2013. Recent
860 developments in the direct-current geoelectrical imaging method. Journal of Applied
861 Geophysics 95, 135–156.

862 López-Moreno, J.I., Beguería, S., García-Ruiz, J.M., 2002. Influence of the Yesa
863 reservoir on floods of the Aragón River, central Spanish Pyrenees. Hydrology and
864 Earth System Sciences 6, 753-762.

865 Losada, J.A., Montesinos, S., Omedas, M., García-Vera, M.A., Galván, R., 2004.
866 Cartografía de las inundaciones del río Ebro en febrero de 2003: trabajos de
867 fotointerpretación, teledetección y análisis SIG en el GIS-Ebro. In: Conesa, C.,
868 Álvarez, Y., Martínez, J.B. (Eds.), Medio ambiente, recursos y riesgos naturales:
869 análisis mediante tecnología SIG y teledetección. Asociación de Geógrafos
870 Españoles – Universidad de Murcia. Murcia, vol. 2, pp. 207-218.

871 Magdaleno, F., Fernández-Yuste, J.A., 2011. Meander dynamics in a changing corridor.
872 Geomorphology 130, 197-207.

873 Maiziels, J.K., 1983. Paleovelocity and paleodischarge determination for coarse gravel
874 deposits. In: Gregory, K.J. (Ed.), Background to Paleohydrology. Wiley, Chichester,
875 pp. 101-139.

876 McCalpin, J.P., 2009a. Field techniques in Paleoseismology. Terrestrial environments.
877 In: Paleoseismology, McCalpin, J.P. (Ed), Academic Press, San Diego, pp. 29-118.

878 McCalpin, J.P., 2009b, Paleoseismology in extensional tectonic environments. In:
879 McCalpin, J.P. (Ed.), Paleoseismology. San Diego, Academic Press, p. 171-269.

880 Mejón, M. (2011). El Ebro desbordado. Ayuntamiento de Zaragoza. Zaragoza, 59 p.

881 Morellón, M., Valero-Garcés, B., Anselmetti, F., Ariztegui, D., Schnellmann, M.,
882 Moreno, A.N.A., Mata, P., Rico, M., Corella, J.P., 2009. Late Quaternary deposition
883 and facies model for karstic Lake Estanya (North-eastern Spain). Sedimentology 56,
884 1505-1534.

885 Oliva, F., Viau, A.E., Bjornson, J., Desrochers, N., Bonneau, M.-A., 2016. A 1300 year
886 reconstruction of paleofloods using oxbow lake sediments in temperate southwestern
887 Québec, Canada. Canadian Journal of Earth Sciences 53, 378-386.

888 Ollero A., 1995. Dinámica reciente del cauce del Ebro en la Reserva Natural de los
889 Galachos (Zaragoza). Cuaternario y Geomorfología 9, 85-93.

890 Ollero, A., 2010. Channel changes and floodplain management in the meandering
891 middle Ebro River, Spain. Geomorphology 117, 247-260.

892 Ortí, F., Salvany, J.M., 1997. Continental evaporitic sedimentation in the Ebro Basin
893 during the Miocene. In: Busson, G., Schreiber, B.Ch., (Eds.), Sedimentary deposition
894 in rift and foreland basins in France and Spain. Columbia University Press, New
895 York, pp. 420-429.

896 Quirantes, J., 1978. Estudio sedimentológico y estratigráfico del Terciario continental
897 de los Monegros. Institución Fernando el Católico - CSIC, Zaragoza.

898 Regato P., 1988. Contribución al estudio de la flora y la vegetación del “Galacho de La
899 Alfranca” en relación con la evolución del sistema fluvial. Diputación General de
900 Aragón, Zaragoza.

901 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C.,
902 Grootes, P. M., Guilderson, T. P., Hafliðason, H., Hajdas, I., HattĹ, C., Heaton, T. J.,
903 Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., Manning,
904 S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R.
905 A., Turney, C. S. M., van der Plicht, J., 2013. IntCal13 and Marine13 Radiocarbon
906 Age Calibration Curves 0-50,000 Years cal BP. Radiocarbon 55, 1869-1887.

907 Salvany, J.M., 2009. Geología del yacimiento glauberítico de Montes de Torrero.
908 Universidad de Zaragoza, Zaragoza.

909 Salvany J.M., García-Veigas, J., Ortí, F., 2007. Glauberite-halite association of the
910 Zaragoza Gypsum Formation (Lower Miocene, Ebro Basin, NE Spain).
911 Sedimentology 54, 443–467.

912 Sheffer, N.A., Enzel, Y., Benito, G., Grodek, T., Poart, N., Lang, M. Coeur, D.. 2003.
913 Paleofloods and historical floods of the Ardèche River, France. Water Resources
914 Research, 39, 1376.

915 Simón, J.L., Soriano, M.A., Arlegui, L., Caballero, J. 1998. Estudio de riesgos de
916 hundimientos kársticos en el corredor de la carretera de Logroño. Memoria del Plan
917 General de Ordenación Urbana de Zaragoza (P.G.O.U.Z.), Anejo 3.2.
918 <http://www.zaragoza.es/contenidos/urbanismo/pgouz/memoria/anejos/anejo03/anejo>
919 032.pdf

920 Torrescusa, S., Klimowitz, J., 1990. Contribución al conocimiento de las evaporitas
921 Miocenas (Formación Zaragoza) de la Cuenca del Ebro. In: Ortí, F., Salvany, J.M.
922 (Eds.), Formaciones evaporíticas de la Cuenca del Ebro y cadenas periféricas y de la
923 zona de Levante. ENRESA-GPG, Barcelona, pp. 120-122.

924 Turnage, K.M., Lee, S.Y., Foss, J.E., Kim, K.H., Larsen, I.L., 1997. Comparison of soil
925 erosion and deposition rates using radiocesium, RUSLE, and buried soils in dolines
926 in East Tennessee. *Environmental Geology* 29, 1-10.

927 Thorndycraft, V.R., Benito, G., 2006. The Holocene fluvial chronology of Spain:
928 evidence from a newly compiled radiocarbon database. *Quaternary Science Reviews*
929 25, 223-234.

930 Webb, R.H., Blainey, J.B., Hyndman, D.W., 2002. Paleoflood hydrology of the Paria
931 River, Southern Utah and Northern Arizona, USA. In: House, P.K., Webb, R.H.,
932 Baker, V.R., Levish, D.R. (Eds.), *Ancient Floods, Modern Hazards. Principles and*
933 *Applications of Paleoflood Hydrology*. American Geophysical Union. *Water Science*
934 *and Applications* 5, pp. 295-310.

935 Whitmore, T.J., Brenner, M., Curtis, J.H., Dahlin, B.H., Leyden, B.W., 1996. Holocene
936 climatic and human influences on lakes of the Yucatan Peninsula, Mexico: an
937 interdisciplinary, palaeolimnological approach. *The Holocene* 6, 273-287.

938 Williams, G.P., 1984. Paleohydrological equations for rivers. In: Costa, J.E., Fleisher,
939 P.J. (Eds.), *Developments and applications of Geomorphology*. Springer-Verlag.
940 Berlin, pp. 343-367.

941 Zaidner, Y., Frumkin, A., Porat, N., Tsatskin, A., Yeshurun, R., Weissbrod, L., 2014. A
942 series of Mousterian occupations in a new type of site: The Nesher Ramla karst
943 depression, Israel. *Journal of Human Evolution* 66, 1-17.

944 Zarroca, M., Comas, X., Gutiérrez, F., Carbonel, D., Linares, R., Roqué, C., Mozaffari,
945 M., Guerrero, J., Pellicer, X., 2016. The application of GPR and ERI in combination
946 with exposure logging and retrodeformation analysis to characterize sinkholes and
947 reconstruct their impact on fluvial sedimentation. Gállego Valley, NE Spain. Earth
948 Surface Processes and Landforms, in press.
949

Figure captions

Figure 1. Geographical setting of the investigated sinkhole. A: Excerpt of the 1:50,000 scale topographic sheet of Alagón (354) produced by the Dirección General del Instituto Geográfico Catastral in 1930 (1st edition). The map shows the analysed water-filled sinkhole before its anthropogenic infill, as well as numerous sinkholes ponds (locally known as *balsas* and *ojos*) mostly obliterated by man-made deposits and human structures. Inset shows the general geographic location. B: Geomorphological map of a section of the Ebro River valley including the analysed sinkhole (modified from Galve et al., 2009).

Figure 2. Orthoimage from 2002 and topographic profile showing the location of the studied sinkhole, the area covered by the 2003 flood and the estimated limit of the 500-year flood, according to the Ebro River Basin Water Authority.

Figure 3. Examples of sinkhole ponds that intersect the water table in the floodplain of the Ebro Valley. These are the Ojos de Matamala, located downstream of Zaragoza city. The circular sinkhole is 30 m across and is centred at 30N X:688125; Y: 4609114. These actively subsiding depressions with lacustrine deposition may act as traps for detrital sediments during flood events.

Figure 4. Aerial photographs of the investigated sinkhole located in the Ebro River floodplain. This complex sinkhole comprises a large diffuse-edged depression, and a nested collapse sinkhole around 40 m across with a permanent pond. The image from 1970, with higher resolution allows the identification of an additional pond. The

sinkhole was obliterated by artificial filling between 1970 and 1984 and used for the construction of human structures.

Figure 5. Detailed topographic maps of the Municipality of Zaragoza produced in 1969 (1:2,000 scale, 1 m contour interval) and 1971-73 (1:1,000 scale, 2 m contour interval). Both maps depict the sinkhole pond (*balsa*) and a drainage ditch that used to flow into the pond. Dashed contour lines roughly illustrate the topography of the large subsidence depressions.

Figure 6. Map of the investigated complex sinkhole showing the distribution of geomorphic units, subsidence features, location of damage identified on human structures, DInSAR deformation data, trenches, the electrical resistivity section acquired across the buried sinkhole pond and borehole drilling in its central sector.

Figure 7. Trench T1 excavated next to the terrace scarp. It revealed the southern edge of the large depression on the terrace through a deformation band including a down-to-the-north oversteepened fault and a younger fissure fill (shown in inset image). The downthrown block has fine-grained slackwater deposits related to a mid-Holocene Ebro River flood.

Figure 8. Simplified retrodeformation sequence derived from the stratigraphic and structural relationships observed in trench T1. In stage 2 it is assumed that subsidence in the floodplain is counterbalanced by aggradation.

1000 Figure 9. Electrical resistivity section acquired in June 2011 across the buried sinkhole
1001 pond with a Dipole-Dipole array and an interelectrode spacing of 2 m.

1002

1003 Figure 10. Stratigraphic log of the 20 m deep borehole drilled in the central sector of the
1004 buried sinkhole pond.

1005