1 **Dynamic cross-talk between host primary metabolism and**
2
3
4 **viruses during infections in plants**
5
6
7

8
9 César Llave
10
11
12

13
14
15 **Address**
16

17 Department of Environmental Biology, Centro de Investigaciones Biológicas, Consejo
18 Superior de Investigaciones Científicas. Ramiro de Maeztu 9, 2804-Madrid, Spain
19
20
21
22

23
24
25 Corresponding author: Llave, César
26

27 Phone: +34-91-837 31 12 Ext. 4259
28

29 Fax: +34-91-536 04 32
30

31 E-mail: cesarllave@cib.csic.es
32
33
34
35
36
37
38
39

40 Word counting
41

42 Abstract: 112
43
44

45 Text: 2031
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Upon infection plant viruses modulate cellular functions and resources to survive and reproduce. Plant cells in which the virus is replicating are transformed into strong metabolic sinks. This conversion gives rise to a massive reprogramming of plant primary metabolism. Such a metabolic shift involves perturbations in carbohydrates, amino acids and lipids that eventually lead to increase respiration rates, and/or decrease in photosynthetic activity. By doing so, plants provide metabolic acclimation against cellular stress and meet the increased demand for energy needed to sustain virus multiplication and defense responses against viruses. This review will highlight our current knowledge pertaining to the contribution of primary metabolism to the outcome of viral infections in plants.

Introduction

A viral infection is a highly dynamic process in which infected plant cells are transformed into major metabolic consumers (sinks) for products of photosynthesis. During the infection, nitrogen and carbon skeletons are required for the synthesis of new molecules and energy is necessary to fuel biosynthesis. The source-to-sink transition elicits reallocation and increases demands for photosynthetic assimilates, increases respiration rates, or decreases in photosynthetic activity. Such a metabolic shift contributes to sustain viral proliferation but is also necessary to activate defense mechanisms. Due to the uneven nature of viral infections in plants and that different virus-host interactions have been studied, it is complicated to depict an unequivocal generalized picture of the metabolic responses triggered by plant viruses. In addition, many reactions in central metabolism are reversible and complex, hindering the perception of major changes in metabolite levels. Nevertheless, a considerable effort has been made to elucidate alterations in metabolite contents linked to viral infections in plants of which significant commonalities arise (Figure 1). In this review, the current knowledge on the modulation of plant primary metabolism during viral infections and its importance for plant compatibility and/or resistance are discussed.

Metabolomics strategies based on chromatography, mass spectrometry (MS) or nuclear magnetic resonance (NMR) spectroscopy in combination with multivariate data analysis have provided an excellent platform to understand the input of certain metabolites in the plant's response to viral infections [1-7]. Recent studies have gone steps further by using system biology approaches to study primary metabolism in plant-virus interactions. For instance, time-course transcriptomics and GC-MS-based metabolomics supported by functional reverse genetics were used to study the reciprocal influence of primary metabolism and *Tobacco rattle virus* (TRV) infection in *Arabidopsis thaliana* [8]. GC-MS-based metabolomics and gene expression data identified altered and unique metabolic signatures characteristic of two

1 tomato inbred lines that exhibited susceptibility or resistance to *Tomato yellow leaf curl virus*
2 (TYLCV) [9], or in response to mild and aggressive isolates of *Potato virus Y* (PVY) in
3
4 potato leaves at different times of disease development [10].
5
6
7
8

9 **Plant virus, carbohydrate metabolism and photosynthesis**

10 The carbohydrate status has profound implications in mobilization and synthesis of storage
11
12 compounds, symptoms development and defense functions, and its alteration is diagnostic for
13
14 perturbations in photosynthesis and respiration. The accumulation of sugars in the infected
15
16 tissue causes an imbalance in the ratio of nitrogen and carbon, and the sensing of such
17
18 changes results in a feedback transcriptional regulation of photosynthesis genes, and
19
20 occasionally, photosynthetic repression [11-17]. Even though repression of photosynthesis
21
22 and induction of sink metabolism is a general response to viral infection, the effects on sugar
23
24 levels varies considerably between different host-virus interactions. Changes in sugar levels
25
26 involving the accumulation of enlarged starch grains in the chloroplast have been reported for
27
28 some compatible interactions [18-21], whereas starch content decreases for some others
29
30 [22,23]. Interestingly, viral accumulation is unaffected in starch-depleted *Arabidopsis* mutants
31
32 suggesting that starch catabolism is not strictly required for virus multiplication in this species
33
34 [8,20]. Metabolism of sink tissues, where sugar is used, is mainly sustained by sucrose
35
36 synthesized in source leaves and transported through the phloem into sink tissues. Sucrose
37
38 and soluble sugars are abundant in different host species infected with ToMV, *Cauliflower*
39
40 *mosaic virus* (CaMV), and TRV [2,8,24], whereas soluble sugar contents decrease upon
41
42 infection with PVY, *Turnip yellow mosaic virus* (TYMV), *Jatropha mosaic virus* (JMV),
43
44 *Ageratum enation virus* (AEV) or *Squash mosaic virus* (SqMV) [5,6,10,25,26].
45
46
47
48
49
50
51
52
53
54

55 Changes in carbohydrates accumulation respond to different causes that include physical
56
57 disturbance of the transport path (e.g. modification of plasmodesmata by viral movement
58
59
60
61
62
63
64
65

proteins), inhibition of sugar transport proteins, induction of starch hydrolysis or cell wall invertases. *Cucumber mosaic virus* (CMV) increases sucrose concentration in the phloem sap of CMV-infected melon plants likely by altering sucrose localization [27]. Sucrose export routes can be severely affected by callose deposition at the cell-to-cell interfaces observed in multiple plant-virus interactions [28]. Infection of cotyledon of marrow plants by CMV causes a gradual increment of soluble sugars and a detriment of starch likely due to enhanced starch hydrolase activities [29,30]. The elevated hexose levels observed in tobacco plants infected with PVY^N are concomitant with increased invertase activity, which cleaves sucrose into glucose and fructose [31]. Interestingly, cell-wall invertase-overexpressing transgenic tobacco or *Arabidopsis* plants accumulate large amounts of soluble sugars and are resistant against PVY^N and *Tobacco mosaic virus* (TMV), respectively [31]. This observation suggests that high hexoses contribute to make plants less susceptible to viral infection, and highlights the relevance of the regulation of carbohydrate metabolism for defense.

Plant viruses and amino acid metabolism

Viruses are important modulators of the amino acid content in infected cells, and amino acid metabolism is critical in the plant response to infection. For instance, in tobacco, alanine, glutamine and proline levels increase in both locally infected leaves and leaves undergoing systemic acquired resistance to TMV [1]. High concentrations of individual and total amino acids have also been reported in different host species following infection with ZYMV, *Papaya ringspot virus* (PRSV), *Bromo mosaic virus* (BMV), and TRV [2,23,32,33]. Amino acids are significantly abundant at the time points of maximal viral accumulation in TRV-infected *Arabidopsis*, although developmental timing is the dominant source of variance in the amino acid content [8]. In potato, infection with PVY leads to an initial decrease in amino acid concentration followed by a gradual increase over time that is particularly significant in

leaves infected with the aggressive PVY^{NTN} isolate compared to the mild PVY^N isolate [10]. Although the precise role of virus-responsive amino acids during the infection has not been elucidated, it is tempting to propose that they may participate in antiviral responses. For instance, several amino acids and intermediates of the amino acid biosynthetic pathway accumulate to higher levels in TYLCV-resistant tomato lines compared to susceptible lines [9]. In *Arabidopsis*, accumulation of proline is observed during the hypersensitive reaction (HR), a plant response whereby the rapid death of cells at the infection site restricts the growth and spread of pathogens to other parts of the plant. External application of proline produces HR-like cell death symptoms [34,35]. In contrast, the *lht1* (*lysine histidine transporter 1*) *Arabidopsis* mutant, which has reduced contents of alanine, glutamine and proline, shows enhanced resistance to multiple pathogens [36]. Likewise, genetic inactivation of *DIN4* gene, which is critical in branched-chain amino acid (BCAA) metabolism, compromises TRV proliferation in infected plants, suggesting that BCAA metabolism contributes to plant susceptibility [8].

Plant viruses and respiration

Plant defense and stress responses induced upon viral infections implicate the up-regulation of the majority of the energy-associated networks (including the glycolysis, the oxidative pentose phosphate pathway, the TCA cycle, mitochondrial energy transport, and ATP biosynthesis) [37]. A sharp increase in respiration concomitant with reduced net photosynthetic rates has been reported in plant-virus interactions [5,22,38,39]. In this scenario, the elevated levels of carbohydrates and the increasing accumulation of amino acids during viral infections fuel the energy-generating TCA pathway to maintain high respiration rates [40]. Furthermore, several organic acids connected to the TCA cycle exhibit positive responses to plants viruses in different host species [2,5-7,10]. This supports the notion that

primary metabolism is largely reconfigured during pathogen infections to satisfy the energy demand required for plant defense and, likely, virus multiplication [41].

Plant viruses and polyamines

Polyamine metabolism, including the most common putrescine, spermidine and spermine, undergoes dynamic changes with polyamine levels increasing several fold during plant-virus interactions [42,43]. Polyamines play a key regulatory role on virus replication [42,44], and have a stimulatory effect on the *in vitro* translation of viral RNA [45,46]. A growing body of evidence suggests that polyamines also transduce defense responses. Elevated levels of free and conjugated putrescine and spermidine and elevated biosynthetic activities occur in TMV-infected tobacco leaves undergoing the HR response, but not in TMV-infected susceptible plants [47-50]. Yamakawa et al. [51] found that free spermine, that accumulates to high levels during the HR in the intercellular fluids of the necrotic lesion-forming leaves, provides resistance against TMV. Induction in polyamine biosynthetic genes and spermedine-responsive genes has been reported during CMV-elicited HR in *Arabidopsis* [52]. Putrescine levels are also augmented in compatible plant-virus interactions [33]. In agreement with a role in defense, high putrescine in *Arabidopsis* provides protection against excessive TRV proliferation, as both the *adc1* and *adc2* mutants impaired in putrescine biosynthesis are hypersusceptible to TRV accumulation [8]. Polyamines are also more abundant in resistant tomato cultivars to TYLCV infection than in susceptible cultivars [9]. In contrast to the above observations, putrescine concentration is reduced in plants infected with *Citrus exocortis viroid* (CEVd) [53].

Plant viruses and lipid metabolism

The significance of cellular lipids in viral infections has long been appreciated because lipids provide energy for metabolism, participate in multiple defense signaling cascades, and are structural components of intracellular membranes in which replication of positive strand RNA viruses take place [54-58]. Membranes of different organelles also protect the viral components from the innate immune system as well as from RNA silencing-based antiviral defense [59,60]. During infection, viruses interfere with lipid (i.e. sterols and phospholipids) and fatty acid (FA) metabolism and biosynthesis to promote changes in the fluidity and/or plasticity of membranes that are required for the proper formation of viral replication complexes [59,61-65]. Genetic inactivation of *INO2*, a transcription activator involved in phospholipid biosynthesis, reduces *Tomato bushy stunt virus* (TBSV) replication and inhibits the activity of the viral replicase complex in yeast model host [66]. The sterol biosynthesis genes *ERG25* and *ERG4* affect the replication of TBSV in yeast [67], while silencing of *N. benthamiana* *SMO1* and *SMO2* genes, which are orthologs of *ERG25*, also result in a reduction in TBSV accumulation, supporting the roles of sterols in virus replication in plants [67]. In yeast, a mutation in the *OLE1* gene encoding $\Delta 9$ fatty acid desaturase (*Ole1p*) severely inhibits BMV replication [68]. In this mutant, the ER lumenal spherule-associated membranes, in which viral RNA synthesis occurs, are locally depleted in unsaturated FAs [68,69]. TRV stimulates the incorporation of polyunsaturated FAs (linoleic and linolenic acid) in various intermediates in the synthesis of triacylglycerols and viral TRV accumulation is reduced in *fad2 Arabidopsis* mutants that contain reduced levels of unsaturated FA [8]. These observations suggest that viral replication is highly sensitive to reduced unsaturated FA levels. In contrast, TRV accumulates to high levels in *acc1 Arabidopsis* mutants, in which the initial step in the biosynthesis of very-long chain fatty acids is partly inhibited [8]. Elevated levels of linolenic acid have been reported in tobacco leaves infected with TMV [1]. A recent study shows that the marine virus *E. huxleyi virus* (EhV) induces profound transcriptome

remodeling in the alga *Emiliania huxley* targeted toward FA synthesis to support viral assembly [70].

Conclusions and perspectives

Viral infections cause profound perturbations in primary metabolism. Although inferences can be made from transcriptomics and metabolomics studies, little is known about the mechanisms used by viruses to interfere with metabolism in both compatible and incompatible interactions and the manner that the metabolite content contributes to viral infection and pathogenesis. Massive reprogramming of primary metabolism aims to meet the increased demand for energy needed to sustain viral multiplication and defense responses against viruses. The emerging view, however, is that accumulation of protective metabolites serve to alleviate the cellular stress imposed by the virus. As a result, viral infections confer a stage of metabolic acclimation that enables plants to cope with other environmental stresses. For instance, CMV and BMV infection of beet and rice, respectively, improve plant tolerance to freezing and drought stress, which correlates with increased osmoprotectant and antioxidant levels in infected plants [33]. Likewise, the enhanced biosynthesis of putrescine in the compatible TRV-*Arabidopsis* interaction makes infected plants more tolerant to freezing stress than non-infected plants [8]. Therefore, plants benefit from the vast array of infection-associated metabolic responses by improving their tolerance to stress.

Systems biology strategies continue to increase our understanding of the dynamic role of primary metabolism during viral infection. However, there are still many questions to answer that concern the precise roles of photosynthesis, sugar partitioning, source-to-sink regulation, respiration and photorespiration in different plant-virus interactions. Future studies should also be expanded to include the use of genetically engineered plants with altered metabolite

1 levels to determine the precise function of virus-responsive metabolites in host antiviral
2 responses or acclimation to environmental stresses.
3
4
5
6

7 **Acknowledgements**

8
9 I wish to thank the editor for the invitation to write this review, and Virginia Ruiz-Ferrer,
10 Francisco Tenllado and José Manuel Martínez-Rivas for helpful comments. I apologize for
11 work that was not adequately discussed or cited in this review due to space constraints. This
12 work is supported by grant BIO2012-39973 from the Spanish Ministry of Economy and
13 Competitiveness.
14
15
16
17
18
19
20
21
22
23

24 **References and recommended reading**

25 Papers of particular interest, published within the period of review, have been highlighted as:
26

- 27 • of special interest
- 28
- 29 •• of outstanding interest
- 30
31
32
33
34
35
36

- 37 1. Choi YH, Kim HK, Linthorst HJ, Hollander JG, Lefeber AW, Erkelens C, Nuzillard JM,
38 Verpoorte R: **NMR metabolomics to revisit the tobacco mosaic virus infection in**
39 **Nicotiana tabacum leaves.** *J Nat Prod* 2006, **69**:742-748.
40
41
- 42 2. Lopez-Gresa MP, Lison P, Kim HK, Choi YH, Verpoorte R, Rodrigo I, Conejero V, Belles
43 JM: **Metabolic fingerprinting of Tomato Mosaic Virus infected Solanum**
44 **lycopersicum.** *J Plant Physiol* 2012, **169**:1586-1596.
45
46
- 47 3. López-Gresa MP, Maltese F, Bellés JM, Conejero V, Kim HK, Choi YH, Verpoorte R:
48 **Metabolic response of tomato leaves upon different plant-pathogen interactions.**
49 *Phytochem Anal* 2010, **21**:89-94.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
4. Mandala R, Kathiriac P, Psychogiosa N, Bouatraa S, Krishnamurthya R, Wisharta D, Kovalchukc I: **Progeny of tobacco mosaic virus-infected *Nicotiana tabacum* plants exhibit trans-generational changes in metabolic profiles.** *Biocatal Agric Biotechnol* 2012, **1**:115-123.
5. Sidhu OP, Annarao S, Pathre U, Snehi SK, Raj SK, Roy R, Tuli R, Khetrapal CL: **Metabolic and histopathological alterations of *Jatropha* mosaic begomovirus-infected *Jatropha curcas* L. by HR-MAS NMR spectroscopy and magnetic resonance imaging.** *Planta* 2010, **232**:85-93.
6. Srivastava S, Bisht H, Sidhu OP, Srivastava A, Singh PC, Pandey RM, Raj SK, Roy R, Nautiyal CS: **Changes in the metabolome and histopathology of *Amaranthus hypochondriacus* L. in response to *Ageratum enation* virus infection.** *Phytochemistry* 2012, **80**:8-16.
7. Bazzini AA, Manacorda CA, Tohge T, Conti G, Rodriguez MC, Nunes-Nesi A, Villanueva S, Fernie AR, Carrari F, Asurmendi S: **Metabolic and miRNA profiling of TMV infected plants reveals biphasic temporal changes.** *PLoS ONE* 2011, **6**:e28466.
8. Fernandez-Calvino L, Osorio S, Hernandez ML, Hamada IB, Del Toro FJ, Donaire L, Yu A, Bustos R, Fernie AR, Martinez-Rivas JM, et al.: **Virus-Induced Alterations in Primary Metabolism Modulate Susceptibility to Tobacco rattle virus in *Arabidopsis*.** *Plant Physiol* 2014, **166**:1821-1838.
- This work integrates transcriptomics, metabolomics and reverse genetics to study the contribution of TRV-responsive metabolic pathways to the outcome of virus infection and prove smetabolic acclimatation to stress in TRV-infected plants.
9. Sade D, Shriki O, Cuadros-Inostroza A, Tohge T, Semel Y, Haviv Y, Willmitzer L, Fernie AR, Czosnek H, Brotman Y: **Comparative metabolomics and transcriptomics of**

1 **plant response to Tomato yellow leaf curl virus infection in resistant and**
2 **susceptible tomato cultivars.** *Metabolomics* 2015, **11**:81-97.

- 3
- 4
- 5 • This study reveals distinctive systemic metabolic signatures to TYLCV infection between
- 6 resistant and susceptible tomato plants, including alterations in the abundance of amino
- 7 acids, polyamines and defense compounds.
- 8
- 9
- 10
- 11
- 12 10. Kogovsek P, Pompe-Novak M, Petek M, Fragner L, Weckwerth W, Gruden K: **Primary**
- 13 **Metabolism, Phenylpropanoids and Antioxidant Pathways Are Regulated in**
- 14 **Potato as a Response to Potato virus Y Infection.** *PLoS One* 2016, **11**:e0146135.
- 15
- 16
- 17
- 18
- 19 • This work studies changes in the metabolome and transcriptome of potato leaves infected
- 20 with mild and aggressive isolates of PVY.
- 21
- 22
- 23
- 24 11. Reinero A, Beachy RN: **Reduced Photosystem II Activity and Accumulation of Viral**
- 25 **Coat Protein in Chloroplasts of Leaves Infected with Tobacco Mosaic Virus.**
- 26 *Plant Physiol* 1989, **89**:111-116.
- 27
- 28
- 29
- 30
- 31 12. Balachandran S, Osmond CB, Makino A: **Effects of Two Strains of Tobacco Mosaic**
- 32 **Virus on Photosynthetic Characteristics and Nitrogen Partitioning in Leaves of**
- 33 **Nicotiana tabacum cv Xanthi during Photoacclimation under Two Nitrogen**
- 34 **Nutrition Regimes.** *Plant Physiol* 1994, **104**:1043-1050.
- 35
- 36
- 37
- 38
- 39
- 40
- 41 13. Rahoutei J, García-Luque I, Barón M: **Inhibition of photosynthesis by viral infection:**
- 42 **Effect on PSII structure and function.** *Physiol Plantarum* 2000, **110**:286-292.
- 43
- 44
- 45
- 46 14. Stare T, Ramsak Z, Blejec A, Stare K, Turnsek N, Weckwerth W, Wienkoop S, Vodnik
- 47 D, Gruden K: **Bimodal dynamics of primary metabolism-related responses in**
- 48 **tolerant potato-Potato virus Y interaction.** *BMC Genomics* 2015, **16**:716.
- 49
- 50
- 51
- 52
- 53 • This work shows the dynamic changes affecting the photosynthetic apparatus upon PVY
- 54 infection in potato.
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

15. Seo S, Okamoto M, Iwai T, Iwano M, Fukui K, Isogai A, Nakajima N, Ohashi Y:
Reduced levels of chloroplast FtsH protein in tobacco mosaic virus-infected tobacco leaves accelerate the hypersensitive reaction. *Plant Cell* 2000, **12**:917-932.
16. Sampol B, Bota J, Riera D, Medrano H, Flexas J: **Analysis of the virus-induced inhibition of photosynthesis in malmsey grapevines.** *New Phytol* 2003, **160**:403-412.
17. von Schaewen A, Stitt M, Schmidt R, Sonnewald U, Willmitzer L: **Expression of a yeast-derived invertase in the cell wall of tobacco and Arabidopsis plants leads to accumulation of carbohydrate and inhibition of photosynthesis and strongly influences growth and phenotype of transgenic tobacco plants.** *EMBO J* 1990, **9**:3033-3044.
18. Cohen J, Loebenstein G: **An electron microscope study of starch lesions in cucumber cotyledons infected with tobacco mosaic virus.** *Phytopathology* 1975, **65**:32-39.
19. Tomenius K, Oxelfelt P: **Ultrastructure of pea leaf-cells infected with three strains of red clover mottle virus.** *J Gen Virol* 1982, **61**:143-147.
20. Handford MG, Carr JP: **A defect in carbohydrate metabolism ameliorates symptom severity in virus-infected Arabidopsis thaliana.** *J Gen Virol* 2007, **88**:337-341.
21. Otulak K, Chouda M, Bujarski J, Garbaczewska G: **The evidence of Tobacco rattle virus impact on host plant organelles ultrastructure.** *Micron* 2015, **70**:7-20.
22. Shalitin D, Wolf S: **Cucumber mosaic virus infection affects sugar transport in melon plants.** *Plant Physiol* 2000, **123**:597-604.
- This study shows how the altered levels of carbohydrates in infected plants was accompanied by increased respiration and decreased photosynthetic rates.

23. Blua MJ, Perring TM, Madore MA: **Plant virus-induced changes in aphid population development and temporal fluctuations in plant nutrients.** *J Chem Ecol* 1994, **20**:691-707.
24. Love AJ, Martin T, Graham IA, Milner JJ: **Carbohydrate partitioning and sugar signalling in Cauliflower mosaic virus-infected turnip and Arabidopsis.** *Phys Mol Plant Pathol* 2005, **67**:83-91.
25. Bedbrook JR, Matthews RE: **Changes in the flow of early products of photosynthetic carbon fixation associated with the replication of TYMV.** *Virology* 1973, **53**:84-91.
26. Magyarosy AC, Buchanan BB, Schurmann P: **Effect of a systemic virus infection on chloroplast function and structure.** *Virology* 1973, **55**:426-438.
27. Herbers K, Takahata Y, Melzer M, Mock HP, Hajirezaei M, Sonnewald U: **Regulation of carbohydrate partitioning during the interaction of potato virus Y with tobacco.** *Mol Plant Pathol* 2000, **1**:51-59.
28. De Storme N, Geelen D: **Callose homeostasis at plasmodesmata: molecular regulators and developmental relevance.** *Front Plant Sci* 2014, **5**:138.
29. Tecsli LI, Maule AJ, Smith AM, Leegood RC: **Metabolic alterations in cotyledons of Cucurbita pepo infected with Cucumber mosaic virus.** *J Exp Bot* 1994, **45**:1541-1551.
30. Tecsli LI, Smith AM, Maule AJ, Leegood RC: **A Spatial Analysis of Physiological Changes Associated with Infection of Cotyledons of Marrow Plants with Cucumber Mosaic Virus.** *Plant Physiol* 1996, **111**:975-985.
31. Herbers K, Meuwly P, Frommer WB, Metraux JP, Sonnewald U: **Systemic Acquired Resistance Mediated by the Ectopic Expression of Invertase: Possible Hexose Sensing in the Secretory Pathway.** *Plant Cell* 1996, **8**:793-803.

32. Wijendra WAS, Ranaweera SS, Salim N: **The effect of Papaya ringspot virus infection on the nitrogen metabolism of Carica papaya L.: Part II. Composition of free amino acids in the leaves.** *Vidyodaya J of Sci* 1995, **5**:131-138.
33. Xu P, Chen F, Mannas JP, Feldman T, Sumner LW, Roossinck MJ: **Virus infection improves drought tolerance.** *New Phytol* 2008, **180**:911-921.
- This excellent study shows that virus infection improves plant tolerance to abiotic stress by increasing the levels of osmoprotectant and antioxidant compounds in infected plants.
34. Deuschle K, Funck D, Forlani G, Stransky H, Biehl A, Leister D, van der Graaff E, Kunze R, Frommer WB: **The role of [Delta]1-pyrroline-5-carboxylate dehydrogenase in proline degradation.** *Plant Cell* 2004, **16**:3413-3425.
35. Fabro G, Kovacs I, Pavet V, Szabados L, Alvarez ME: **Proline accumulation and AtP5CS2 gene activation are induced by plant-pathogen incompatible interactions in Arabidopsis.** *Mol Plant Microbe Interact* 2004, **17**:343-350.
36. Liu G, Ji Y, Bhuiyan NH, Pilot G, Selvaraj G, Zou J, Wei Y: **Amino acid homeostasis modulates salicylic acid-associated redox status and defense responses in Arabidopsis.** *Plant Cell* 2010, **22**:3845-3863.
37. Less H, Angelovici R, Tzin V, Galili G: **Coordinated gene networks regulating Arabidopsis plant metabolism in response to various stresses and nutritional cues.** *Plant Cell* 2011, **23**:1264-1271.
38. Leal N, Lastra R: **Altered metabolism of tomato plants infected with tomato yellow mosaic virus.** *Physiol Plant Pathol* 1984, **24**:1-7.
39. Jensen SG: **Photosynthesis, respiration and other physiological relationships in barley infected with barley yellow dwarf virus.** *Phytopathology* 1967, **58**:204-208.
40. Berger S, Sinha AK, Roitsch T: **Plant physiology meets phytopathology: plant primary metabolism and plant-pathogen interactions.** *J Exp Bot* 2007, **58**:4019-4026.

- Good review of how plant-pathogen interactions affect photosynthesis, sugar accumulation and sink metabolism
41. Bolton MD: **Primary metabolism and plant defense--fuel for the fire.** *Mol Plant Microbe Interact* 2009, **22**:487-497.
 - This excellent review provides a comprehensive overview on changes in host physiology and primary metabolism during incompatible plant-pathogen interactions and describes how plants recruit energy for the defense response.
 42. Balint R, Cohen SS: **The incorporation of radiolabeled polyamines and methionine into turnip yellow mosaic virus in protoplasts from infected plants.** *Virology* 1985, **144**:181-193.
 43. Torget R, Lapi L, Cohen SS: **Synthesis and accumulation of polyamines and S-adenosylmethionine in Chinese cabbage infected by turnip yellow mosaic virus.** *Biochem Biophys Res Commun* 1979, **87**:1132-1139.
 44. Balint R, Cohen SS: **The effects of dicyclohexylamine on polyamine biosynthesis and incorporation into turnip yellow mosaic virus in Chinese cabbage protoplasts infected in vitro.** *Virology* 1985, **144**:194-203.
 45. Benicourt C, Haenni AL: **In vitro synthesis of turnip yellow mosaic virus coat protein in a wheat germ cell-free system.** *J Virol* 1976, **20**:196-202.
 46. Morch MD, Benicourt C: **Polyamines stimulate suppression of amber termination codons in vitro by normal tRNAs.** *Eur J Biochem* 1980, **105**:445-451.
 47. Negrel J, Vallee JC, Martin C: **Ornithine decarboxylase activity and the hypersensitive reaction of tobacco to tobacco mosaic virus in *Nicotiana tabacum*.** *Phytochemistry* 1984, **23**:2747-2751.

48. Torrigiani P, Rabiti AL, Bortolotti C, Betti L, Marani F, Canova A, Bagni N: **Polyamine synthesis and accumulation in the hypersensitive response to TMV in *Nicotiana tabacum***. *New Phytol* 1997, **135**:467–473.
49. Rabiti AL, Betti L, Bortolotti C, Marini F, Canova A, Bagni N, Torrigiani P: **Short-term polyamine response in TMV-inoculated hypersensitive and susceptible tobacco plants**. *New Phytol* 1998, **139**:549–553.
50. Marini F, Betti L, Scaramagli S, Biondi S, Torrigiani P: **Polyamine metabolism is upregulated in response to tobacco mosaic virus in hypersensitive, but not in susceptible, tobacco**. *New Phytol* 2001, **149**:301-309.
- This work shows that in NN tobacco plants the response to TMV implies an increase in polyamine turnover with elevated biosynthetic enzyme activities and free and conjugated putrescine and sperimidine levels.
51. Yamakawa H, Kamada H, Satoh M, Ohashi Y: **Spermine is a salicylate-independent endogenous inducer for both tobacco acidic pathogenesis-related proteins and resistance against tobacco mosaic virus infection**. *Plant Physiol* 1998, **118**:1213-1222.
52. Mitsuya Y, Takahashi Y, Berberich T, Miyazaki A, Matsumura H, Takahashi H, Terauchi R, Kusano T: **Spermine signaling plays a significant role in the defense response of *Arabidopsis thaliana* to cucumber mosaic virus**. *J Plant Physiol* 2009, **166**:626-643.
53. Belles JM, Perez-Amador MA, Carbonell J, Conejero V: **Correlation between Ornithine Decarboxylase and Putrescine in Tomato Plants Infected by Citrus Exocortis Viroid or Treated with Ethephon**. *Plant Physiol* 1993, **102**:933-937.

54. Van Der Heijden MW, Carette JE, Reinhoud PJ, Haegi A, Bol JF: **Alfalfa mosaic virus replicate proteins P1 and P2 interact and colocalize at the vacuolar membrane.** *J Virol* 2001, **75**:1879-1887.
55. Schwartz M, Chen J, Janda M, Sullivan M, den Boon J, Ahlquist P: **A positive-strand RNA virus replication complex parallels form and function of retrovirus capsids.** *Mol Cell* 2002, **9**:505-514.
56. Weber-Lotfi F, Dietrich A, Russo M, Rubino L: **Mitochondrial targeting and membrane anchoring of a viral replicase in plant and yeast cells.** *J Virol* 2002, **76**:10485-10496.
57. Turner KA, Sit TL, Callaway AS, Allen NS, Lommel SA: **Red clover necrotic mosaic virus replication proteins accumulate at the endoplasmic reticulum.** *Virology* 2004, **320**:276-290.
58. Rubino L, Weber-Lotfi F, Dietrich A, Stussi-Garaud C, Russo M: **The open reading frame 1-encoded ('36K') protein of Carnation Italian ringspot virus localizes to mitochondria.** *J Gen Virol* 2001, **82**:29-34.
59. Heaton NS, Randall G: **Multifaceted roles for lipids in viral infection.** *Trends Microbiol* 2011, **19**:368-375.
60. Ahlquist P, Noueiry AO, Lee WM, Kushner DB, Dye BT: **Host factors in positive-strand RNA virus genome replication.** *J Virol* 2003, **77**:8181-8186.
61. Dufourc EJ: **Sterols and membrane dynamics.** *J Chem Biol* 2008, **1**:63-77.
62. Schneiter R, Kohlwein SD: **Organelle structure, function, and inheritance in yeast: a role for fatty acid synthesis?** *Cell* 1997, **88**:431-434.
63. Jiang Y, Serviène E, Gal J, Panavas T, Nagy PD: **Identification of essential host factors affecting tombusvirus RNA replication based on the yeast Tet promoters Hughes Collection.** *J Virol* 2006, **80**:7394-7404.

- 20
21
22 67. Sharma M, Sasvari Z, Nagy PD: **Inhibition of sterol biosynthesis reduces tombusvirus**
23
24 **replication in yeast and plants.** *J Virol* 2010, **84**:2270-2281.
25
26 68. Lee WM, Ishikawa M, Ahlquist P: **Mutation of host delta9 fatty acid desaturase**
27 **inhibits brome mosaic virus RNA replication between template recognition and**
28 **RNA synthesis.** *J Virol* 2001, **75**:2097-2106.
29
30
31
32
33 69. Lee WM, Ahlquist P: **Membrane synthesis, specific lipid requirements, and localized**
34 **lipid composition changes associated with a positive-strand RNA virus RNA**
35 **replication protein.** *J Virol* 2003, **77**:12819-12828.
36
37
38
39
40
41 70. Rosenwasser S, Mausz MA, Schatz D, Sheyn U, Malitsky S, Aharoni A, Weinstock E,
42
43 Tzfadia O, Ben-Dor S, Feldmesser E, et al.: **Rewiring Host Lipid Metabolism by**
44 **Large Viruses Determines the Fate of *Emiliana huxleyi*, a Bloom-Forming Alga**
45 **in the Ocean.** *Plant Cell* 2014, **26**:2689-2707.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legend:

Figure 1. A simplified schematic representation of plant primary metabolism showing the major compounds that are altered in response to viral infections. Further details are given within the text and literature cited herein. ACP (acyl carrier protein), ER (endoplasmic reticulum), FA (fatty acids), LPA (lysophosphatidic acid), PA (phosphatidic acid). Glycolipids: DAG (diacylglycerol), DGDG (digalactosyl diacylglycerol), MGDG (monogalactosyl diacylglycerol), SQDG (sulfoquinovosyl diacylglycerol). Phospholipids: PC (phosphatidylcholine), PE (phosphatidylethanolamine), PI (phosphatidylinositol), PG (phosphatidylglycerol),

Figure
[Click here to download Figure: Current Opinion Llave 2016 Figure 1.pdf](#)