

Trifunctional squaramide catalyst for efficient enantioselective Henry reaction activation

Juan V. Alegre-Requena,^a Eugenia Marqués-López^a and Raquel P. Herrera^{a*}

^a Laboratorio de Organocatálisis Asimétrica. Departamento de Química Orgánica. Instituto de Síntesis Química y Catálisis Homogénea (ISQCH) CSIC-Universidad de Zaragoza. C/ Pedro Cerbuna 12, 50009 Zaragoza (Spain). E-mail: raquelph@unizar.es

Abstract. A new class of trifunctional squaramide catalyst acting by mean of multiple interactions has been found in the study of the Henry reaction. Enantiomerically enriched nitroaldol products were obtained in good yields and high enantioselectivities under mild conditions using one of the smallest amount of organocatalyst reported so far for this reaction (0.25 mol%).

The catalyst was able to generate hydrogen bonding and anion- π /hydrogen- π interactions with the substrates, responsible of the improvement in the reactivity and the enantioselectivity of this process. Computational calculations support a mechanistic hypothesis based on an anion- π effect, being this the first example reported in asymmetric catalysis.

Keywords: aldehyde; organocatalysis; Henry; nitroalkanes; squaramide; trifunctional.

Introduction

Multifunctional scaffolds have received special attention in the last decade, with the increasing interest for the synthesis of more complex catalytic systems with a high organization grade and a multidentate activation, to offer a cooperative effect resembling the role of enzymes.^[1,2] These structures could facilitate the formation of cooperative non-covalent interactions in a synergic way and thereby significantly improve their catalytic activity. This would overcome the drawback of high catalyst loading that commonly are used in organocatalysis, which is still a great challenge.^[3] These studies have focused on hydrogen bonding, electrostatics effects, π - π , cation- π , hydrophobic and Van der Waals forces. All these interactions are believed to be involved in stabilizing the transition states by lowering the energetic barrier with a remarkable increase in the rate of the reaction. These binding could also be responsible of the improvement in the stereoselectivity of the process by differentially stabilizing the diastereomeric pathways created in the transition state between the catalyst and the reacting components. Additionally, non-covalent interactions operating in a synergic way can afford the conformational restriction required to induce high enantioselectivity.

The reaction between an *in situ* generated nitronate species and a carbonyl compound, known as Henry (nitroaldol) reaction,^[4] is an important carbon-carbon bond-forming method in organic synthesis.^[5] This process represents a powerful and useful tool for the

synthesis of valuable β -nitroalcohols providing, after further transformations of the versatile nitro group, efficient access to interesting and highly functionalized intermediates, like β -aminoalcohols and α -hydroxy carboxylic acids.^[6,7] The Henry reaction may be promoted under many different conditions and using diverse catalytic systems providing from good-to-excellent enantioselectivities.^[8,9] In fact, many efforts have been invested for improving this method, and different kinds of organocatalysts have been explored in order to increase the pioneering results reported by Nájera and co-workers in 1994.^[10]

Therefore, the development of new asymmetric Henry strategies is still important to address the construction of interesting building blocks as a crucial step in total synthesis (Figure 1).




Figure 1. Biologically active compounds bearing a β -aminoalcohol motif.

Herein, we report a new class of trifunctional squaramide catalysts acting through multidentate activation, which efficiently catalyzes the asymmetric Henry reaction with high yields and high enantioselectivities (up to 94% ee). A low catalyst loading is required, routinely 2 mol%, and this can even be decreased to as low as 0.25 mol% without detriment of the enantioselectivity. Interestingly, this is one of the lowest loading used so far in

organocatalysis for this reaction. Moreover, this is the first reported example of a reaction using squaramides where an anion- π interaction and an H- π interaction have been found in the transition state (TS) and could justify the efficiency of the used catalyst. To the best of our knowledge, this kind of activation has not been explored so far in asymmetric catalysis, and therefore this work could represent a pivotal contribution in this field.^[11,12]

In this respect, many reactions are efficiently promoted using multifunctional catalysts, where nucleophile and electrophile are simultaneously coordinated to the different functional groups present in the catalyst structure.^[13] In the case of the Henry reaction, both components could be efficiently approached in the TS following a bifunctional coordination and represents an attractive model to explore this concept. Based on previous developed studies, we envisaged the importance of having in the same structure a hydrogen bond moiety and a basic part in order to obtain a more rigid transition state at the moment of the carbon-carbon bond formation. For this aim, we chose squaramide structures **1a-j** (Figure 2)^[14] (see Supporting information for the rest of tested structures and screening), synthesized following our one-pot developed procedure.^[15]





Figure 2. Representative squaramide-based organocatalysts tested.

Results and Discussion

We started the investigation of the viability of this process examining the efficiency of catalysts **1** in the model reaction between 4-nitrobenzaldehyde (**2a**) and nitromethane (**3a**) (see supporting information for the complete outcomes). After a very exhaustive

screening of the reaction conditions, novel catalyst **1j** (2 mol%) was found to be the catalyst of choice in terms of reactivity and enantioselectivity (>95% yield, 82% ee, Table 1, entry 1); nitromethane with no extra solvent resulted the best reaction media, at -24 °C. The efficiency of the process was further studied for a range of different aldehydes **2a-o** (Table 1).

Table 1. Scope of the squaramide-catalyzed Henry reaction.^[a]


Entry	R	1j (%)	R'	t (h)	product	yield ^[b] (%)	ee ^[c] (%)
1	4-NO ₂ Ph (2a)	2	H (3a)	24	4aa	>95	83
2	3-NO ₂ Ph (2b)	2	H (3a)	24	4ba	>95	94
3	4-ClPh (2c)	2	H (3a)	91	4ca	81	86
4	3-ClPh (2d)	2	H (3a)	86	4da	62	90
5	4-BrPh (2e)	2	H (3a)	91	4ea	75	86
6	4-CNPh (2f)	2	H (3a)	71	4fa	96	82
7	1-Naphthyl (2g)	2	H (3a)	86	4ga	50	85
8	Ph (2h)	2	H (3a)	86	4ha	59	82
9	4-Ph-Ph (2i)	2.5	H (3a)	96	4ia	50	90
10	2-Pyridyl (2j)	2.5	H (3a)	96	4ja	>95	80
11	3-Pyridyl (2k)	2	H (3a)	92	4ka	95	92
12	2-Furyl (2l)	2	H (3a)	88	4la	74	92
13	2-Thiophenyl (2m)	2	H (3a)	92	4ma	55	92
14	4-MePh (2n)	2	H (3a)	144	4na	20	84
15	PhCH ₂ OCH ₂ (2o)	2	H (3a)	92	4oa	66	76
16 ^[d]	4-NO ₂ Ph (2a)	2	Me (3b)	88	4ab	75 ^[f]	72 ^[g]
17 ^[e]	3-NO ₂ Ph (2b)	2	Me (3b)	91	4bb	74 ^[f]	88 ^[g]

^[a] Experimental conditions: to a mixture of catalyst **1j** (0.0044 mmol) in MeNO₂ (1.1 mL), aldehyde **2a-o** (0.22 mmol) was further added in a test tube at -24 °C. ^[b] After isolation by column chromatography. ^[c] Determined by chiral HPLC analysis. ^[d] d.r. 1:1.3 *anti:syn*. ^[e] d.r. 1:1.4 *anti:syn*. ^[f] Yield as mixture of both diastereoisomers. ^[g] Enantiomeric excess (ee) for the major diastereoisomer.

The Henry reaction took place rendering the desired β -nitroalcohols **4** in good to excellent yields (up to >95%) and high enantioselectivities (up to 94%) with very clean reaction crudes. The effectiveness of the developed procedure is well accounted since it was successfully applied to a representative set of aldehydes **2a-o**. The enantioselectivity was not dependent on the electronic effects of the aldehydes. However, the reactivity suggests a close correlation with the electronegativity of the aldehyde since those with an electro-withdrawing group in the aromatic ring exhibited more reactivity. In fact, in the case of aldehyde **2n** the reaction was slower, although keeping the good enantioselectivity of the process (entry 14). At this point, it is important to remark that

we have observed that the presence of traces of acid in the aldehydes could inactivate the catalyst used at this small scale, and many of the aldehydes were previously treated in order to avoid such inactivation. This fact is in agreement with the lack of background for this reaction in the absence of a base. The absolute configuration of final adducts **4** was determined by comparison of their optical rotation values with those previously reported in the literature for the same products and it was found to be *S* (see experimental section for optical rotation values).

We further evaluated the grade of effectiveness of the catalytic system lowering even more the catalyst loading for four representative aldehydes (**2a,b,f,p**) (Table 2).


Table 2. Study of the catalyst loading for the Henry reaction.^[a]

Entry	R	1j (%)	t (h)	yield ^[b] (%)	ee ^[c] (%)
1	4-NO ₂ Ph (2a)	1	24	>95	82
2 ^[d]	4-NO ₂ Ph (2a)	0.5	76	>95	82
3 ^[e]	4-NO ₂ Ph (2a)	0.25	92	86	82
4	3-NO ₂ Ph (2b)	1	24	>95	94
5 ^[d]	3-NO ₂ Ph (2b)	0.5	76	>95	94
6 ^[e]	3-NO ₂ Ph (2b)	0.25	95	92	94
7 ^[f]	3-NO ₂ Ph (2b)	0.25	96	>95	94
8	4-CNPh (2f)	1	95	>95	82
9 ^[d]	4-CNPh (2f)	0.5	100	>95	82
10 ^[e]	4-CNPh (2f)	0.25	92	59	82
11 ^[d]	F ₅ C ₅ (2p)	0.5	45	>95	86
12 ^[e]	F ₅ C ₅ (2p)	0.25	120	>95	86

^[a] Experimental conditions: to a mixture of catalyst **1j** in MeNO₂ (1.1 mL), aldehyde **2a,b,f,p** (0.22 mmol) was further added in a test tube at -24 °C. After the reaction time, adduct **4** was isolated by flash chromatography. ^[b] After isolation by column chromatography. ^[c] Determined by chiral HPLC analysis. ^[d] Conditions for the use of 0.44 mmol of aldehyde. ^[e] Conditions for the use of 0.88 mmol of aldehyde. ^[f] Reaction performed at higher scale to obtain 1 gram of product.

As disclosed in Table 2, interestingly, we were able to decrease the catalyst loading to 0.25 mol%, one of the lowest value used in a Henry protocol and one of the lowest amounts employed so far in organocatalysis.^[3] In all cases, the same enantioselectivity was found, although after longer reaction times. It is worth noting that this system allows scaling up the reaction, since the same excellent results, in terms of enantioselectivity and reactivity, were afforded when it was set up to obtain 1 gram of final product (Table 2, entry 7).



We continued our investigations exploring the effect of some designed changes in the structure of the catalyst, in order to understand the role played by novel catalyst **1j**, which bears different functionalities (Scheme 1). Thus, catalyst **1k**, which has the opposite configuration (*S*) on the binaphthyl scaffold compared to catalyst **1j** (*R*), was prepared in order to study the possible match/mismatch effect. Interestingly, no reversal of the sense in the asymmetric induction was observed, giving rise to the same absolute enantiomer in the final product. However, final product **4ba** was achieved with lower ee value compared to catalyst **1j** (81% vs 94% ee). This suggests that the sense of the enantioinduction of this process seems to be directly depending on the 2-(1-piperidinyl)cyclohexylamine moiety, but the different value of enantioselectivity obtained reveals an influence of the binaphthyl ring in the different reaction pathways.

**Scheme 1.** Additional squaramide-based organocatalysts **1k-m** tested.

The reaction using catalyst **1l** (with a OMe group instead of a free OH) leads to the same good results obtained with catalyst **1j**. Therefore, it seems that the binaphthyl skeleton could be crucial in stabilizing the TS to properly induce better enantioselectivity, even when the participation of the OR (R = H or Me) group in the process is still unclear at this stage. Interestingly, when we performed the reaction with catalyst **1m** (with a naphthyl moiety instead of the binaphthyl fragment present in **1j**), the reactivity and the enantioselectivity of the process drastically dropped (10% yield, 60% ee). Also, catalyst **1j** showed much better results compared to those obtained with other squaramides whose structures

only differ in their aromatic moieties, such as **1a-c** and **1e** (entry 6 in Table S2 and entries 1, 7 and 8 in Table S3, respectively). This fact confirms the necessity of the complete binaphthyl structure in catalyst **1j** for the success of the process.

In order to explain the role of the multifunctional catalyst **1j**, based on the experimental results and computational calculations (see ESI), a reasonable reaction pathway is proposed for the Henry reaction between 4-cyanobenzaldehyde (**2f**) and nitromethane (**3a**) (only the rate-limiting step is shown for simplicity) (Figure 2).



TSc

Figure 2. **TSa:** catalyst-substrates complex before nitronate attack, **TSb:** transition state (TS) of the nitronate attack, and **TSc:** catalyst-intermediate product complex after the nitronate attack. All DFT calculations were carried out with the Gaussian 09 software^[16] at the IEFPCM-SMD(MeNO₂)/wB97XD/6-31G(d) level (see ESI for more information about computational details).

The O atom of the aldehyde, which is developing a negative charge during the advance of the reaction, and the H atom of the aldehyde are interacting with the π -system of the catalyst in the most energetically favorable reaction pathway (see ESI for additional TSs).

In order to support these π interactions, the density maps of the non-covalent interactions for **TSa-c** have been also calculated (Figures 3-5). These maps show hydrogen bonds between the aldehyde and the squaramide moiety through double hydrogen bonding with the NH groups and between the basic nitrogen atom on the piperidine and the nitromethane. These non-covalent interactions are represented in blue color.

Moreover, the π interactions between the aldehyde and the binaphthyl moiety are also disclosed. These interactions are shown in green color, being weaker than those represented in blue color (Figures 3-5).




Figure 3. Electronic density map of system **TSa**: green and blue regions represent attractive VdW interactions.^[17] Grid data for $\text{sign}(\lambda_2)\rho$ and RDG was generated using Multiwfn.^[18] The image was created using VMD.^[19]




Figure 4. Electronic density map of **TSb**: green and blue regions represent attractive VdW interactions.^[17] Grid data for $\text{sign}(\lambda_2)\rho$ and RDG was generated using Multiwfn.^[18] The image was created using VMD.^[19]




Figure 5. Electronic density map of **TSc**: green and blue regions represent attractive VdW interactions.^[17] Grid data for $\text{sign}(\lambda_2)\rho$ and RDG was generated using Multiwfn.^[18] The image was created using VMD.^[19]

Thus, catalyst **1j** could be acting in a trifunctional fashion. First, the aldehyde would be activated by the squaramide moiety through double hydrogen bonding with the NH groups. At the same time, the π -system of the binaphthyl scaffold would stabilize the TS and would activate the aldehyde for the subsequent attack. Additionally, the basic nitrogen atom on the piperidine ring would activate the nitromethane, allowing the attack of the nitronate form over the *Re* face of the aldehyde. This attack would afford the *S* absolute configuration in all final products, which is consistent with the observed results. Moreover, this attack would be in agreement with the control of the sense in the enantioinduction by the 2-(1-piperidinyl)cyclohexylamine moiety.

Beyond the steric influence caused by the binaphthyl moiety that can orientate the disposition of the reagents in the TS, we believe that the non-covalent interactions between this group and the aldehyde influence both, the reactivity and the enantioselectivity of the process. This anion- π interaction would stabilize the energetic barrier of the TS, increasing the reaction rate of the process and, consequently, the origin of the catalysis would also rely in this weak but effective effect. This interaction ensures a more energetically favorable TS where the chiral catalyst remains in close proximity to the electrophile during the enantioselectivity-determining step of the catalytic cycle. The presence of these secondary binding is able to provide a higher degree of organization in the transition state, necessary for a high enantioinduction. Moreover, the lower enantioselectivity observed with catalyst **1m** compared with **1j**, would also be in agreement with the importance of the presence of the binaphthyl scaffold to organize a more stable TS.

Conclusion

In conclusion, a new class of chiral squaramide catalyst acting by mean of multiple interactions has been developed to efficiently catalyze the Henry reaction with very good results. Novel trifunctional catalyst **1j** is able to generate hydrogen bonding and anion- π interactions with the substrates, which are responsible of the improvement in the reactivity and the stereoselectivity of the process. This active catalyst was found to provide excellent values of enantioselectivity and reactivity even at 0.25 mol% catalyst loading, one of the lowest amounts for this reaction using organocatalysts. A unique example of anion- π effect has been described for the first time in asymmetric catalysis, and this feature makes catalyst **1j** a plausible trifunctional catalyst. Further investigations of the efficacy of this organocatalyst in other catalytic asymmetric reactions, as well as

additional computational and NMR studies, are ongoing in our lab in order to support the singular activation mode proposed herein.

Experimental Section

General experimental methods. Purification of reaction products was carried out either by filtration or by flash chromatography using silical-gel (0.063-0.200 mm). Analytical thin layer chromatography was performed on 0.25 mm silical gel 60-F plates. ESI ionization method and mass analyzer type MicroTof-Q were used for the ESI measurements. $^1\text{H-NMR}$ spectra were recorded at 300 and 400 MHz; $^{13}\text{C-APT-NMR}$ spectra were recorded at 75 and 100 MHz; CDCl_3 , CD_3CN and $\text{DMSO-}d_6$ were used as the deuterated solvents. Chemical shifts were reported in the δ scale relative to residual CHCl_3 (7.26 ppm), MeCN (1.94 ppm) and DMSO (2.50 ppm) for $^1\text{H-NMR}$ and to the central line of CDCl_3 (77 ppm), CD_3CN (1.24 ppm) and $\text{DMSO-}d_6$ (39.43 ppm) for $^{13}\text{C-APT-NMR}$.

Materials. Spectral data for **1b**,^[20] **1c**,^[21] **1d**,^[22] **1e**,^[23] **1f**,^[22] **1g**,^[22] **1h**,^[22] **1i**,^[24] **1j**,^[21] **1n**,^[21] **1o**,^[21] **1p**,^[21] and **1q**^[21] are consistent with values previously reported in the literature. For the synthesis and spectra of catalysts **1k-m** see supporting information.

Representative procedure for the squaramide-organocatalyzed Henry reaction of aldehydes

To a mixture of catalyst **1j** (0.0044 mmol unless otherwise stated in Table 1) and aldehyde **2a-o** (0.22 mmol), was added MeNO_2 (1.1 mL) in a test tube at $-24\text{ }^\circ\text{C}$. After the reaction time (see Table 1 and 2), adducts **4** were isolated by flash chromatography (SiO_2 , using Hex:EtOAc 9:1 to Hex:EtOAc 7:3). Yields and enantioselectivities are reported in Table 1. If acid traces were observed in the aldehydes by NMR, these aldehydes were previously purified by column chromatography (very short column, eluted with CH_2Cl_2 or MeCN) or extraction (dissolving these aldehydes in CH_2Cl_2 and washing with a 0.3 M solution of NaOH). Then, the CH_2Cl_2 was evaporated in vacuum and the aldehydes were used within 2-5 minutes to avoid acid formation.

(S)-2-Nitro-1-(4-nitrophenyl)ethanol (4aa)^[25] Following the general procedure, compound **4aa** was obtained after 24 h of reaction at $-24\text{ }^\circ\text{C}$ as a dark green oil in >95% yield. The ee of the product was determined to be 83% by HPLC using a Daicel Chiralpak IA column (*n*-hexane/*i*-PrOH = 80:20, flow rate 1 mL min^{-1} , $\lambda = 230.3\text{ nm}$): $\tau_{\text{major}} = 15.6\text{ min}$; $\tau_{\text{minor}} = 12.1\text{ min}$. $[\alpha]_{\text{D}}^{28} = +23.2$ (*c* 1.30, CHCl_3 , 82% ee) {lit.,^[25] $[\alpha]_{\text{D}}^{24} -30.4$ (*c* 0.53, CHCl_3) for (*R*)-**4aa**, 88% ee}.

(S)-2-Nitro-1-(3-nitrophenyl)ethanol (4ba)^[26] Following the general procedure, compound **4ba** was obtained after 24 h of reaction at $-24\text{ }^\circ\text{C}$ as a dark green solid in >95% yield. The ee of the product was determined to be 94% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 80:20, flow rate 1 mL min^{-1} , $\lambda = 281.7\text{ nm}$): $\tau_{\text{major}} = 10.5\text{ min}$; $\tau_{\text{minor}} = 9.9\text{ min}$. $[\alpha]_{\text{D}}^{23} = +27.2$ (*c* 0.33, CHCl_3 , 94% ee) {lit.,^[26] $[\alpha]_{\text{D}}^{26} -27.4$ (*c* 0.87, CH_2Cl_2) for (*R*)-**4ba**, 96% ee}.

(S)-1-(4-Chlorophenyl)-2-nitroethanol (4ca)^[25] Following the general procedure, and purifying the aldehyde by column chromatography (eluted with CH_2Cl_2), compound **4ca** was obtained after 91 h of reaction at

-24 °C as a dark brown oil in 81% yield. The ee of the product was determined to be 86% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 230.1 nm): τ_{major} = 12.6 min; τ_{minor} = 11.1 min. [α]_D²³ = +27.2 (*c* 0.31, CHCl₃, 86% ee) {lit.,^[25] [α]_D²² -38.8 (*c* 0.55, CHCl₃) for (*R*)-**4ca**, 90% ee}.

(S)-1-(3-Chlorophenyl)-2-nitroethanol (4da)^[27]

Following the general procedure, and purifying the aldehyde **2d** by basic washing, compound **4da** was obtained after 86 h of reaction at -24 °C as a dark brown oil in 62% yield. The ee of the product was determined to be 90% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 226.2 nm): τ_{major} = 11.8 min; τ_{minor} = 10.3 min. [α]_D²⁴ = +24.8 (*c* 1.4, CHCl₃, 90% ee) {lit.,^[27] [α]_D²⁷ +31.17 (*c* 1.0, CHCl₃) for 95% ee}.

(S)-1-(4-Bromophenyl)-2-nitroethanol (4ea)^[28]

Following the general procedure, and purifying the aldehyde **2e** by column chromatography (eluted with CH₂Cl₂), compound **4ea** was obtained after 91 h of reaction at -24 °C as a dark brown oil in 75% yield. The ee of the product was determined to be 86% by HPLC using a Daicel Chiralpak IA column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 237.2 nm): τ_{major} = 14.9 min; τ_{minor} = 12.2 min. [α]_D²³ = +20.1 (*c* 0.27, CHCl₃, 86% ee) {lit.,^[28] [α]_D²³ -68.6 (*c* 1.40, CHCl₃) for (*R*)-**4ea**, 89% ee}.

(S)-4-(1-Hydroxy-2-nitroethyl)benzonitrile (4fa)^[29]

Following the general procedure, compound **4fa** was isolated by flash chromatography after 71 h of reaction at -24 °C as a pale brown solid in >95% yield. The ee of the product was determined to be 82% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 243.5 nm): τ_{major} = 27.6 min; τ_{minor} = 25.3 min. [α]_D²⁴ = +36.3 (*c* 0.74, CHCl₃, 81% ee) {lit.,^[30] [α]_D²⁰ -32.8 (*c* 0.50, CH₂Cl₂) for (*R*)-**4fa**, 90% ee}.

(S)-1-(Naphthalen-1-yl)-2-nitroethanol (4ga)^[25]

Following the general procedure, and purifying the aldehyde **2g** by basic washing, compound **4ga** was obtained after 86 h of reaction at -24 °C as a dark brown oil in 50% yield. The ee of the product was determined to be 85% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 254.0 nm): τ_{major} = 14.5 min; τ_{minor} = 11.6 min. [α]_D²⁷ = +19.1 (*c* 0.85, CHCl₃, 85% ee) {lit.,^[25] [α]_D²¹ +24.5 (*c* 0.53, CHCl₃, for (*S*)-**4ga**, 88% ee}.

(S)-2-Nitro-1-phenylethanol (4ha)^[31] Following the general procedure, and purifying the aldehyde **2h** by basic washing, compound **4ha** was obtained after 86 h of reaction at -24 °C as a dark brown oil in 59% yield. The ee of the product was determined to be 82% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 248.1 nm): τ_{major} = 11.3 min; τ_{minor} = 10.1 min. [α]_D²⁵ = +11.9 (*c* 1.1, CH₂Cl₂, 82% ee) {lit.,^[31] [α]_D²¹ -41.6 (*c* 1.03, CH₂Cl₂) for (*R*)-**4ha**, 94% ee}.

(S)-1-([1,1'-Biphenyl]-4-yl)-2-nitroethanol (4ia)^[31]

Following the general procedure, compound **4ia** was isolated by flash chromatography after 96 h of reaction at -24 °C as a yellow solid in 50% yield. Mp. 127-129 °C.^[31] The ee of the product was determined to be 90% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 231.2 nm): τ_{major} = 16.3 min; τ_{minor} = 13.3 min. [α]_D²³ = +25.8 (*c* 0.47, CHCl₃, 88% ee) {lit.,^[31] [α]_D²³ -36.1 (*c* 1.35, CH₂Cl₂) for (*R*)-**4ia**, 91% ee}.

(S)-2-Nitro-1-(pyridin-2-yl)ethanol (4ja)^[32] Following the general procedure, and purifying the aldehyde **2j** by column chromatography (eluted with MeCN), compound **4ja** was isolated by flash chromatography (SiO₂, using Hex:EtOAc 8:2 to Hex:EtOAc 1:1) after 96 h of reaction at -24 °C as a dark brown oil in >95% yield. The ee of the product was determined to be 80% by HPLC using a Daicel Chiralpak IA column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 218.4 nm): τ_{major} = 12.1 min; τ_{minor} = 15.3 min. [α]_D²⁴ = +49.8 (*c* 0.16, CHCl₃, 80% ee).

(S)-2-Nitro-1-(pyridin-3-yl)ethanol (4ka)^[33] Following the general procedure, and purifying the aldehyde **2k** by column chromatography (eluted with MeCN), compound **4ka** was isolated by flash chromatography (SiO₂, using Hex:EtOAc 7:3 to Hex:EtOAc 2:8) after 92 h of reaction at -24 °C as a dark brown oil in 95% yield. The ee of the product was determined to be 92% by HPLC using a Daicel Chiralpak IA column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 240.2 nm): τ_{major} = 24.3 min; τ_{minor} = 28.3 min. [α]_D²⁹ = +35.1 (*c* 0.23, MeCN, 92% ee).

(R)-1-(Furan-2-yl)-2-nitroethanol (4la)^[34] Following the general procedure, and purifying the aldehyde **2l** by column chromatography (eluted with CH₂Cl₂), compound **4la** was obtained after 88 h of reaction at -24 °C as a dark brown oil in 74% yield. The ee of the product was determined to be 92% by HPLC using a Daicel Chiralpak IA column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 224.8 nm): τ_{major} = 10.9 min; τ_{minor} = 10.2 min. [α]_D²⁹ = +36.3 (*c* 0.16, CHCl₃, 92% ee) {lit.,^[29] [α]_D²³ -36.7 (*c* 2.72, CHCl₃) for (*S*)-**4la**, 85% ee}.

(R)-2-Nitro-1-(thiophen-2-yl)ethanol (4ma)^[34] Following the general procedure, and purifying the aldehyde **2m** by column chromatography (eluted with CH₂Cl₂), compound **4ma** was obtained after 92 h of reaction at -24 °C as a dark brown oil in 55% yield. The ee of the product was determined to be 92% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 246.6 nm): τ_{major} = 12.0 min; τ_{minor} = 11.4 min. [α]_D²⁹ = +35.9 (*c* 0.08, CHCl₃, 90% ee) {lit.,^[28] [α]_D²³ -26.4 (*c* 3.11, CHCl₃) for (*S*)-**4ma**, 86% ee}.

(S)-2-Nitro-1-*p*-tolylethanol (4na)^[35] Following the general procedure, and purifying the aldehyde **2n** by column chromatography (eluted with CH₂Cl₂), compound **4na** was obtained after 6 days of reaction at -24 °C as a dark brown oil in 20% yield. The ee of the product was determined to be 84% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 95:5, flow rate 1 mL min⁻¹, λ = 220 nm): τ_{major} = 19.8 min; τ_{minor} = 17.0 min. [α]_D¹⁸ = +26.8 (*c* 0.25, CHCl₃, 84% ee) {lit.,^[35] [α]_D²⁰ +34.1 (*c* = 1.90, CHCl₃) for (*S*)-**4na**, 84% ee}.

(R)-1-(Benzyloxy)-3-nitropropan-2-ol (4oa)^[36] Following the general procedure, compound **4oa** was isolated by flash chromatography after 92 h of reaction at -24 °C as a dark brown oil in 66% yield. The ee of the product was determined to be 76% by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 95:5, flow rate 1 mL min⁻¹, λ = 227.7 nm): τ_{major} = 22.6 min; τ_{minor} = 25.0 min. [α]_D²⁴ = +7.8 (*c* 0.25, CHCl₃, 76% ee) {lit.,^[37] [α]_D +1.5 (*c* 0.9, CH₂Cl₂) for (*R*)-**4oa**, 80% ee}.

(S)-2-Nitro-1-(perfluorophenyl)ethanol (4pa)^[37]

Following the general procedure, compound **4pa** was isolated by flash chromatography (SiO₂, using Hex:EtOAc 95:5 to Hex:EtOAc 9:1) after 120 h of reaction at -24 °C as a yellow oil in >95% yield. The ee of the product was determined to be 86% by HPLC using a Daicel Chiralpak

IA column (*n*-hexane/*i*-PrOH = 95:5, flow rate 1 mL min⁻¹, λ = 232.8 nm): τ_{major} = 10.9 min; τ_{minor} = 12.4 min. $[\alpha]_{\text{D}}^{28}$ = +2.8 (*c* 1.1, CHCl₃, 86% ee) {lit.,^[37] $[\alpha]_{\text{D}}$ +8.9 (*c* 0.8, CH₂Cl₂) for (*S*)-**4oa**, 90% ee}.

2-Nitro-1-(4-nitrophenyl)propan-1-ol (4ab)^[34,38]

Following the general procedure, compound **4ab** was isolated by flash chromatography (SiO₂, using Hex:EtOAc 95:5 to Hex:EtOAc 9:1) after 88 h of reaction at -24 °C as a dark green solid in 75% yield.^[34,38] The diastereomeric ratio (*anti*/*syn*, 1:1.3) was determined by ¹H NMR. The ee of the products was determined to be 72% (*syn isomer*) by HPLC using a Daicel Chiralpak IA column (*n*-hexane/*i*-PrOH = 90:10, flow rate 1 mL min⁻¹, λ = 242.0 nm): τ_{major} = 29.9 min; τ_{minor} = 25.9 min; for the *syn* diastereoisomer.

2-Nitro-1-(3-nitrophenyl)propan-1-ol (4bb)^[34,38]

Following the general procedure, compound **4bb** was isolated by flash chromatography (SiO₂, using Hex:EtOAc 95:5 to Hex:EtOAc 9:1) after 91 h of reaction at -24 °C as a dark green solid in 74% yield. The diastereomeric ratio (*anti*/*syn*, 1:1.4) was determined by ¹H NMR.^[34,38] The ee of the products was determined to be 87% (*syn isomer*) by HPLC using a Daicel Chiralpak IB column (*n*-hexane/*i*-PrOH = 95 : 5, flow rate 1 mL min⁻¹, λ = 238.4 nm): τ_{major} = 35.6 min; τ_{minor} = 31.5 min; for the *syn* diastereoisomer.

Acknowledgements

We thank the Ministerio de Economía y Competitividad (MINECO, Project CTQ2013-44367-C2-1-P), the University of Zaragoza (JIUZ-2014-CIE-07), the High Council of Scientific Investigation (CSIC) (PIE-2015801010) and Government of Aragon DGA (Research Group E-104) for financial support of our research. J.V.A.-R. thanks the DGA for a predoctoral contract.

References

- [1] P. M. Pihko (Ed.), *Hydrogen Bonding in Organic Synthesis*; Wiley-VCH: Weinheim, **2009**.
- [2] F. Liu, *Chirality* **2013**, *25*, 675-683.
- [3] For a review, see: a) F. Giacalone, M. Gruttadauria, P. Agrigento, R. Noto, *Chem. Soc. Rev.* **2012**, *41*, 2406-2447. For a recent example of low catalyst loading in an organocascade reaction, see also: b) C. C. J. Loh, P. Chauhan, D. Hack, C. Lehmann, D. Enders, *Adv. Synth. Catal.* **2014**, *356*, 3181-3186.
- [4] L. Henry, *C. R. Hebd. Séances Acad. Sci.* **1895**, *120*, 1265-1268.
- [5] F. A. Luzzio, *Tetrahedron* **2001**, *57*, 915-945.
- [6] N. Ono (Ed.), *The Nitro Group in Organic Synthesis*; Wiley-VCH: New York, USA, **2001**.
- [7] S. C. Bergmeier, *Tetrahedron* **2000**, *56*, 2561-2576.
- [8] J. Boruwa, N. Gogoi, P. P. Saikia, N. C. Barua, *Tetrahedron: Asymmetry* **2006**, *17*, 3315-3326.
- [9] a) C. Palomo, M. Oiarbide, A. Mielgo, *Angew. Chem. Int. Ed.* **2004**, *43*, 5442-5444; b) C. Palomo, M. Oiarbide, A. Laso, *Eur. J. Org. Chem.* **2007**, 2561-2576; c) Y. Alvarez-Casao, E. Marqués-López, R. P. Herrera, *Symmetry* **2011**, *3*, 220-245.
- [10] R. Chinchilla, C. Nájera, P. Sánchez-Agulló, *Tetrahedron: Asymmetry* **1994**, *5*, 1393-1402.
- [11] To the best of our knowledge, there are not any previous works regarding oxygen- π and/or hydrogen- π interactions in asymmetric catalysis. For the unique example of anion- π catalysis, although non asymmetric, see: Y. Zhao, C. Beuchat, Y. Domoto, J. Gajewy, A. Wilson, J. Mareda, N. Sakai, S. Matile, *J. Am. Chem. Soc.* **2014**, *136*, 2101-2111.
- [12] a) M. Nishio, Y. Umezawa, K. Honda, S. Tsuboyamad, H. Suezawa, *CrystEngComm* **2009**, *11*, 1757-1788. b) O. Takahashi, Y. Kohno, M. Nishio, *Chem. Rev.* **2010**, *110*, 6049-6076.
- [13] a) H. Miyabe, Y. Takemoto, *Bull. Chem. Soc. Jpn.* **2008**, *81*, 785-795. b) S. J. Connon, *Chem. Commun.* **2008**, 2499-2510. c) Y. Wei, M. Shi, *Acc. Chem. Res.* **2010**, *43*, 1005-1018. d) P. Chauhan, S. S. Chimni, *RSC Adv.* **2012**, *2*, 737-758. e) Y. Xiao, Z. Sun, H. Guo, O. Kwon, *Beilstein J. Org. Chem.* **2014**, *10*, 2089-2121. f) X. Fang, C.-J. Wang, *Chem. Commun.* **2015**, *51*, 1185-1197.
- [14] For pivotal reviews concerning squaramides, see: a) J. Alemán, A. Parra, H. Jiang, K. A. Jørgensen, *Chem. Eur. J.* **2011**, *17*, 6890-6899. b) R. I. Storer, C. Aciro, L. H. Jones, *Chem. Soc. Rev.* **2011**, *40*, 2330-2346. c) F. R. Wurm, H.-A. Klok, *Chem. Soc. Rev.* **2013**, *42*, 8220-8236. d) J. V. Alegre-Requena, *Synlett* **2014**, *25*, 298-299. e) P. Chauhan, S. Mahajan, U. Kaya, D. Hack, D. Enders, *Adv. Synth. Catal.* **2015**, *357*, 253-281.
- [15] J. V. Alegre-Requena, E. Marqués-López, R. P. Herrera, *RSC Adv.* **2015**, *5*, 33450-33462.
- [16] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven Jr., J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, N. J. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, All calculations were carried out with the Gaussian 09 suite of programs: Gaussian 09, Revision A.1, Gaussian, Inc., Wallingford, CT, 2009.
- [17] E. R. Johnson, S. Keinan, P. Mori-Sánchez, J. Contreras-García, A. J. Cohen, W. Yang, *J. Am. Chem. Soc.* **2010**, *132*, 6498-6506.
- [18] T. Lu, F. Chen, *J. Comp. Chem.* **2012**, *33*, 580-592.

- [19] W. Humphrey, A. Dalke, K. Schulten, *J. Mol. Graphics* **1996**, *14*, 33-38.
- [20] H. Konishi, T. Y. Lam, J. P. Malerich, V. H. Rawal, *Org. Lett.* **2010**, *12*, 2028-2031.
- [21] E. Marqués-López, J. V. Alegre-Requena, R. P. Herrera, *European Pat.* EP14382260.9 filed Jul. 07, 2014; b) J. V. Alegre-Requena, E. Marqués-López, R. P. Herrera, *RSC Adv.* **2015**, *5*, 33450-33462.
- [22] W. Yang, D.-M. Du, *Org. Lett.* **2010**, *12*, 5450-5453.
- [23] J. P. Malerich, K. Hagihara, V. H. Rawal, *J. Am. Chem. Soc.* **2008**, *130*, 14416-14417.
- [24] H. Jiang, M. W. Paixão, D. Monge, K. A. Jørgensen, *J. Am. Chem. Soc.* **2010**, *132*, 2775-2783.
- [25] M. Steurer, C. Bolm, *J. Org. Chem.* **2010**, *75*, 3301-3310.
- [26] S. Kitagaki, T. Ueda, C. Mukai, *Chem. Commun.* **2013**, *49*, 4030-4032.
- [27] O. Soltani, M. A. Ariger, H. Vázquez-Villa, E. M. Carreira, *Org. Lett.* **2010**, *12*, 2893-2895.
- [28] A. Bulut, A. Aslan, Ö. Dogan, *J. Org. Chem.* **2008**, *73*, 7373-7375.
- [29] C. V.-L. Bray, F. Jiang, X.-F. Wu, J.-B. Sortais, C. Darcel, *Tetrahedron Lett.* **2010**, *51*, 4555-4557.
- [30] E. C. Constable, G. Zhang, C. E. Housecroft, M. Neuburger, S. Schaffner, W.-D. Woggon, J. A. Zampese, *New J. Chem.* **2009**, *33*, 2166-2173.
- [31] D. A. Evans, D. Seidel, M. Rueping, H. W. Lam, J. T. Shaw, C. W. Downey, *J. Am. Chem. Soc.* **2003**, *125*, 12692-12693.
- [32] K. Kanagaraj, P. Sureh, K. Pitchumani, *Org. Lett.* **2010**, *12*, 4070-4073.
- [33] T. Marcelli, R. N. S. van der Haas, J. H. van Maarseveen, H. Hiemstra, *Angew. Chem. Int. Ed.* **2006**, *45*, 929-931.
- [34] K. Kanagaraj, P. Suresh, K. Pitchumani, *Org. Lett.* **2010**, *12*, 4070-4073.
- [35] B. Zheng, M. Wang, Z. Li, Q. Bian, J. Mao, S. Li, S. Liu, M. Wang, J. Zhong, H. Guo, *Tetrahedron: Asymmetry* **2011**, *22*, 1156-1160.
- [36] J. C. Borah, S. Gogoi, J. Boruwa, N. C. Barua, *Synth. Commun.* **2005**, *35*, 873-878.
- [37] M. Bandini, F. Piccinelli, S. Tommasi, A. Umani-Ronchi, C. Ventrici, *Chem. Commun.* **2007**, 616-618.
- [38] R. Boobalan, G.-H. Lee, C. Chen, *Adv. Synth. Catal.* **2012**, *354*, 2511-2520.