**Preservative effect of a previous high-pressure
treatment on the chemical changes related to quality
loss in frozen hake (*Merluccius merluccius*)**

Manuel Vázquez¹, Liliana G. Fidalgo², Jorge A. Saraiva², and
Santiago P. Aubourg^{3,*}

¹ Department of Analytical Chemistry, Faculty of Veterinary Science,
University of Santiago de Compostela, 27002-Lugo, Spain

² Research Unit of Organic Chemistry, Natural and Agro-food Products
(QOPNA), Chemistry Department, Aveiro University, Campus
Universitário de Santiago, 3810-193 Aveiro, Portugal.

³ Department of Food Technology, Marine Research Institute (CSIC),
36208-Vigo, Spain

* Correspondent: saubourg@iim.csic.es; +34986292762 (fax),
+34986231930 (phone)

ABSTRACT

The frozen storage of most Gadiform species is strongly limited by the trimethylamine oxide breakdown into dimethylamine (DMA) and formaldehyde (FA). This study focussed on the quality loss inhibition of hake (*Merluccius merluccius*) during the frozen storage. In it, the effect of a previous high-pressure (HP) treatment (150-450 MPa for 2 min) on the quality of hake kept frozen at -10°C up to 5 months was studied by means of a statistical experiment design. For this purpose, complementary chemical analyses related to quality loss were carried out concerning lipid oxidation and hydrolysis development, volatile amines and formaldehyde (FA) formation and K value. A multifactor ANOVA test showed an inhibitory effect ($p < 0.015$) on dimethylamine (DMA) and free fatty acids formation as a result of the previous HP treatment, this effect increasing with pressure level applied. Additionally, an inhibition ($p < 0.05$) of FA, trimethylamine, total volatile amines and fluorescent compounds formation was obtained according to a one-way ANOVA test. On the contrary, no effect ($p > 0.05$) on the K value, polyene index and formation of peroxides and TBARS was achieved. The work here presented provides for the first time information concerning the employment of HP technology to inhibit the DMA and FA formation during frozen storage.

Running Title: Previous high pressure and frozen hake quality

Key Words: High pressure; *Merluccius merluccius*; frozen storage; formaldehyde; dimethylamine; free fatty acids

INTRODUCTION

Gadiforms is a large group of fish species such as cod, hake, pollack, haddock, whiting, saithe, etc., which represents an important percentage of the overall fish catching and consumption in most European countries (FAO 2007). Thus, in addition to being commercialised in the fresh condition, most frozen fishery products like fillets, surimis and fish fingers are made from whole or minced muscle of these fish. However, fish species are recognised as developing fast and easy deterioration during the post-mortem period as a result of undergoing different kinds of damage pathways such as microbial attack, autolysis and lipid oxidation (Özogul 2010). Moreover, large quantities of trimethylamine oxide demethylase (TMAOase) are included in different body parts of Gadiform species, such as kidney, spleen and microsomal fraction of muscle. This enzyme has been found responsible for the breakdown of trimethylamine oxide into different kinds of deteriorative molecules such as trimethylamine (TMA), formaldehyde (FA) and dimethylamine (DMA) (Leelapongwattana et al. 2005; Malinowska-Pańczyk and Kołodziejska 2016).

Frozen storage has widely been employed to maintain fish properties before consumption or further use in other technological processes. In the case of frozen Gadiform species, protein denaturation has shown to be a major problem during the commercialisation of such kinds of products, this leading to texture losses and off-odour development (Aubourg et al. 2007). Different pathways have been proposed to explain this degradation process (Sotelo and Rehbein 2000): i) partial dehydration of proteins during freezing due to the formation of ice crystals, ii) interaction of proteins with FA and with oxidised and hydrolysed lipids, and iii) modification of the protein micro-environment caused by the concentration of salts in the liquid phase surrounding

79 proteins. In agreement with this high susceptibility to quality loss, substantial efforts are
80 needed to meet the increasing consumer demand for high quality and safe products.

81 Among advanced food processing technologies, high-hydrostatic pressure
82 technology has proved to inactivate microbial populations and endogenous enzymes
83 while retaining the sensory and nutritional properties of foods (Campus 2010; Tabilo-
84 Munizaga et al. 2016). Concerning high pressure (HP) application to seafood, previous
85 research accounts for its practical and beneficial effect when employed prior to a
86 subsequent processing or storage. This strategy has successfully been checked in the
87 cases of further chilled (Hurtado et al. 2001), cold-smoked (Lakshmanan et al. 2003) or
88 semi-dried (Gou et al. 2012) storage. Related to freezing, frozen storage and thawing
89 processes, pressure-shift technology has also been applied (Chevalier et al. 2000; Tironi
90 et al. 2010); as a result of its employment, different kinds of advantages in protein
91 properties have been demonstrated such as a lower denaturation and toughening extent
92 and an increased water-holding capacity. In the case of a further frozen storage, research
93 has been focussed on fatty fish species. Thus, a previous optimised HP treatment led to
94 a quality enhancement of chemical properties of frozen mackerel (Vázquez et al. 2013),
95 horse mackerel (Torres et al. 2013) and sardine (Méndez et al. 2017).

96 European hake (*Merluccius merluccius*) is an abundant Gadiform species of
97 great market demand because of its excellent organoleptic features. Although frozen
98 storage has shown to partially retain its quality and shelf life time, marked chemical
99 changes have proved to be produced and lead to substantial quality losses by increasing
100 the frozen storage time and temperature (Aubourg and Medina 1999). Previous research
101 concerning HP treatment of European hake is scarce. This accounts for its positive
102 effect on physicochemical properties of isolated sarcoplasmic (Villamonte et al. 2016)
103 and myofibrillar (Cando et al. 2014) proteins, and for the inhibition of *Anisakis simplex*

L3 in hake muscle (Vidacek et al. 2009). The present research focusses on the development of chemical changes related to quality loss during the frozen storage of hake. In it, the effect of a previous treatment at 0-300 MPa for 2 min on the lipid oxidation and hydrolysis development, volatile amines and FA formation and K value was studied.

MATERIALS AND METHODS

Raw fish, processing, storage and sampling

European hake (102 individuals) were caught near the Galician Atlantic coast (North-Western Spain) and transported under ice to the laboratory. Whole hake individuals were placed in flexible polyethylene bags (three individuals per bag) and vacuum sealed at 150 mbar (Albipack-Packaging Solutions, Águeda, Portugal). The length and weight of the specimens ranged 27.5-29.5 cm and 180-205 g, respectively.

HP treatments at 150, 169.27, 300, 430.73 and 450 MPa were performed in a 55-L high pressure unit (Hiperbaric 55, Hiperbaric NL, Burgos, Spain) according to the experimental design described in the statistical analysis section; in all cases, a pressure holding time of 2 min was employed. Water applied as the pressurising medium at 3 MPa s⁻¹ yielded 54, 63, 100, 115 and 121 s as the come up times for the 150, 169.27, 300, 430.73 and 450 MPa treatments, respectively, while decompression time was less than 3s. Cold pressurising water was used to maintain temperature conditions during HP treatment at 20 °C.

After HP treatments, hake individuals were kept at -20 °C for 48 hours and then stored at -10 °C with samples analysed after 0.00, 0.32, 2.50, 4.68 and 5.00 months. The experimental design of pressure level/frozen storage time developed in this study included seventeen kinds of batches which were named from T-0 to T-16, in agreement

with the description provided in Table 1. In it, starting fresh fish (T-0 batch), control fish (frozen fish without previous HP treatment; T-1, T-3, T-6, T-12 and T-15 batches) and HP-treated fish (T-2, T-4, T-5, T-7, T-8 T-9, T-10, T-11, T-13, T-14 and T-16 batches) were taken into account.

Three replicates (n=3) for each batch (T-0 to T-16) were analysed independently. Each analysis was based on the corresponding extract from the fish white muscle pooled from two individual fish. A frozen storage temperature (−10 °C) higher than commercial practice (−18 °C) was chosen as an accelerated test condition to estimate the extent of the pressure level and frozen storage effects on the chemical quality indices of frozen hake. All solvents and chemical reagents used were of reagent grade (Merck, Darmstadt, Germany).

Experimental design and statistical analysis

The statistical experiment design was formulated using the Design Expert® 7.1.1 software (Stat-Ease, Inc., Minneapolis, MN, USA). The models were validated through a multifactor ANOVA test. The set of batches (T-0 to T-16) follows a central composite design (CCD), which has three groups of design points: (a) two-level factorial or fractional factorial design points; (b) axial points (also called “star” points); (c) centre points. The present design has 5 levels of independent variables with desirable statistical properties.

Data (n=3) obtained from the different chemical quality indices were subjected to the multifactor ANOVA test to explore differences as a result of pressure level and frozen storage time. Additionally, data (n=3) obtained from the different chemical quality indices were subjected to one-way ANOVA method ($p<0.05$) to explore differences as a result of the freezing step, frozen storage time and pressure level.

Comparison of means was performed using a least-squares difference (LSD) method (Statsoft, Statistica, version 6.0, Tulsa, OK, USA).

Lipid oxidation assessment

Lipids were extracted by the Bligh and Dyer (1959) method employing a 1:1 chloroform-methanol mixture for the single-phase solubilisation of lipids in the fish muscle. Quantification results were expressed as g lipid kg⁻¹ muscle.

The peroxide value (PV) in the lipid extract was determined by peroxide reduction with ferric thiocyanate, according to the Chapman and McKay (1949) method. Results were expressed as meq active oxygen kg⁻¹ lipids.

The thiobarbituric acid index (TBA-i) was determined as described by Vyncke (1970) and based on the reaction between a 5% trichloroacetic acid extract of the fish muscle and thiobarbituric acid. Content on thiobarbituric acid reactive substances (TBARS) was spectrophotometrically measured at 532 nm and results were expressed as mg malondialdehyde kg⁻¹ muscle.

The formation of fluorescent compounds was determined in the aqueous phase obtained during the lipid extraction by measurements at 393/463 nm and 327/415 nm (Aubourg and Medina 1999). Relative fluorescence (RF) was defined as the F/F_{st} ratio where F is the fluorescence measured at each excitation and emission maximum, and F_{st} is the fluorescence intensity of a 1 µg mL⁻¹ quinine sulphate solution in 0.05 M H₂SO₄ at the corresponding wavelength. A fluorescence ratio (FR) was calculated as the ratio between the two RF values, i.e., $FR = RF_{393/463\text{ nm}} / RF_{327/415\text{ nm}}$.

Lipid extracts were converted into fatty acid methyl esters (FAME) by reaction with acetyl chloride and then analysed using a Perkin-Elmer 8700 gas chromatograph equipped with a fused silica capillary column SP-2330 (0.25 mm i.d. x 30 m, 0.20 µm

film, Supelco Inc., Bellefonte, PA, USA) and using nitrogen at 68.95 kPa as carrier gas (linear flow rate of 1.0 mL min⁻¹), a flame ionisation detector at 250 °C, and C 19:0 fatty acid as internal standard for quantitative analysis (Torres et al. 2013). Peaks corresponding to fatty acids were identified by comparison of the retention times of two standards mixtures (Qualmix Fish, Larodan, Malmo, Sweden; FAME Mix, Supelco Inc., Bellefonte, PA, USA). Peak areas were automatically integrated. The polyene index (PI) was calculated as the following fatty acids ratio: C20:5 ω 3+C22:6 ω 3/C 16:0.

Determination of volatile amines

Total volatile base-nitrogen (TVB-N) values were measured by the Antonacopoulos (1960) method with some modifications. Briefly, 10 g fish muscle were extracted with 60 g L⁻¹ perchloric acid in 30 mL water and brought up to 50 mL. An aliquot of the acid extracts was rendered alkaline to pH 13 with 200 g L⁻¹ aqueous NaOH and then steam-distilled. Finally, the TVB-N content was determined by titration of the distillate with 10 mM HCl. Results were expressed as mg TVB-N kg⁻¹ muscle.

Trimethylamine-nitrogen (TMA-N) values were determined by the picrate method, as previously described by Tozawa et al. (1971). This involves the preparation of a 5% trichloroacetic acid extract of fish muscle (10 g/25 ml). Results were expressed as mg TMA-N kg⁻¹ muscle.

Dimethylamine-nitrogen (DMA-N) content was determined in the 5% TCA extracts using the spectrophotometric method of Dowden (1938). This method is based on the specific complex formation from secondary amines in the presence of a copper salt and carbon sulphur in basic medium. Results were expressed as mg DMA-N kg⁻¹ muscle.

Other chemical assessments

Free fatty acids (FFA) content in the lipid extract was determined by the Lowry and Tinsley (1976) method based on complex formation with cupric acetate-pyridine followed by spectrophotometric assessment at 715 nm (Beckman Coulter DU 640, London, UK). Results were expressed as g FFA kg⁻¹ lipids.

Extraction of free formaldehyde (FA) from fish muscle was carried out according to Rey-Mansilla et al. (2001). FA content was determined by the spectrophotometric (Beckman Coulter DU 640 spectrophotometer) method of Nash (1953) and expressed as mg kg⁻¹ muscle.

Nucleotide extracts were prepared and analysed by HPLC according to Aubourg et al. (2005). Standard curves for adenosine 5'-triphosphate (ATP) and each compound involved in its degradation pathway, adenosine 5'-diphosphate (ADP), adenosine 5'-monophosphate (AMP), inosine 5'-monophosphate (IMP), inosine (INO) and hypoxanthine (HX), were constructed in the 0-1 mM range. Results were expressed as the K value (%) that was calculated according to the following concentrations ratio:

$$K \text{ value (\%)} = 100 \times \frac{\text{ATP} + \text{ADP} + \text{AMP} + \text{IMP} + \text{INO} + \text{HX}}{\text{INO} + \text{HX}}$$

RESULTS AND DISCUSSION

Lipid oxidation assessment

Concerning peroxides assessment (Table 1), a slight formation (p<0.05) could be observed as a result of the freezing process according to the comparison between T-0 and T-1 batches. Meantime, samples without previous HP treatment showed a marked peroxide formation at month 2.50 (11.45±3.97 meq active oxygen kg⁻¹ lipids), followed

by a progressive decrease till the end of the experiment (3.08 ± 1.68 meq active oxygen kg^{-1} lipids). Taking into account samples corresponding to each pressure applied (i.e., 169.27, 300 and 430.73 MPa), a PV increase with frozen storage time could be observed in most cases. When the effect of the previous pressure is evaluated, some significant differences ($p < 0.05$) could be observed at each frozen storage time; however, a general trend could not be concluded. Thus, a multifactor ANOVA analysis did not provide a significant ($p > 0.05$) effect of pressure applied on the peroxide formation. In this sense, it can be argued that peroxides are primary lipid oxidation compounds, susceptible to breakdown and lead to lower-molecular-weight molecules, so that their determination cannot always afford a valuable measurement of lipid oxidation development (Aubourg and Medina 1999; Tironi et al. 2002).

For the secondary oxidation compounds (TBA-i; Table 1), a significant increase was detected as a result of the freezing step in agreement with the comparison between T-0 and T-1 batches. As for the peroxide detection, comparison among samples without previous pressure treatment did not provide a definite trend of TBARS formation throughout the frozen storage time. Thus, a marked increase was obtained till month 0.32 (0.80 ± 0.10 mg malondialdehyde kg^{-1} muscle), followed by a progressive decrease till the end of the study (0.15 ± 0.04 mg malondialdehyde kg^{-1} muscle). A decreasing TBA-i was observed with frozen storage time in HP-treated samples corresponding to 169.27, 300 and 430.73 MPa. Regarding to the effect of the previous pressure applied, a definite trend could not be concluded when comparing the different kinds of samples at each frozen storage time. As for the PV, a multifactor ANOVA analysis did not supply a significant ($p > 0.05$) effect of pressure applied to the TBARS formation. In this sense, it can be argued that such compounds can also be considered as highly reactive and

susceptible to interact with nucleophilic compounds present in the fish muscle (Aubourg and Medina 1999; Tironi et al. 2002).

Tertiary lipid oxidation compounds measured by fluorescence detection showed a significant formation as a result of the freezing step (Table 1). However, comparison among samples without previous pressure treatment did not show ($p>0.05$) any changes throughout the frozen storage time; furthermore, a general trend about the effect of the frozen storage time could not be obtained for samples corresponding to 169.27, 300 and 430.73 MPa treatments. However, at each frozen storage time, FR mean values were higher in control samples (i.e., 0.10-MPa batches), although significant ($p<0.05$) differences were only obtained at months 2.50 and 5.00. Therefore, a partial inhibition of tertiary oxidation compounds formation could be concluded as a result of the previous pressure applied. However, a multifactor ANOVA analysis did not provide a significant ($p>0.05$) effect of pressure applied on the FR value.

Finally, the PI score remained quite stable throughout the study for all kinds of samples, being included in the 2.51-3.00 range (Table 1). Thus, no effect ($p>0.05$) of the freezing step, frozen storage or the previous pressure applied could be concluded on this quality index.

In order to explain and justify lipid changes produced in the present research, two different and opposite mechanisms ought to be taken into account. On one side, the inhibitory effect of pressure applied on the endogenous enzymes activity should produce a decrease of lipid oxidation development during the further frozen storage period (Campus 2010; Tabilo-Munizaga et al. 2016). On the other side, iron-bound protein denaturation during HP treatment has been reported to facilitate the increase of free metal ion content, which would lead to a lipid oxidation increase in fish meat after HP treatment (Gudmundsson and Hafsteinsson 2002; Lakshmanan et al. 2003). Thus, in

the present research, a prooxidant effect of HP processing was not observed in frozen hake. In the case of primary and secondary compounds, a definite trend could not be concluded. However, previous HP treatment showed a partial inhibition of tertiary compounds formation, this result being attributed to the damage effect of previous pressure treatment on pro-oxidant enzymes such as lipoxygenases, peroxidases, etc., and leading to an inhibitory effect of their activity.

Previous related research accounts for contradictory results when the effect of HP treatment on lipid oxidation development in frozen fish is evaluated. Thus, Ohshima et al. (1992) found a TBARS formation increase in frozen cod (*Gadus morhua*) and mackerel (*Scomber scombrus*) muscle that had been subjected to HP treatment at 200-600 MPa for 15-30 min. Later on, the use of pressure shift freezing at 140 MPa was compared with air-blast freezing at -20 °C in turbot (*Scophthalmus maximus*) fillets (Chevalier et al. 2000); analysis of TBARS formation during subsequent storage at -20 °C for 75 d did not provide a definite effect of the previous HP processing. Closely related to the present research, a marked inhibition of tertiary lipid oxidation compounds formation in frozen mackerel (Vázquez et al. 2013) and horse mackerel (Torres et al. 2013) by previous HP treatment at 150-450 MPa for 0.0-5.0 min was obtained; this effect was greater when increasing the pressure level or the pressure holding time.

Concerning PUFA damage, present results agree with previous related research. Thus, Ohshima et al. (1992) did not find differences in saturated, monounsaturated and polyunsaturated fatty acids content in cod (*Gadus morhua*) and mackerel (*Scomber scombrus*) muscle after 6 d of storage at -2 °C when previously treated under 200-600 MPa for 15 min. Recently, no significant changes in polyene content were obtained in

frozen mackerel (Vázquez et al. 2013) and horse mackerel (Torres et al. 2013) that was previously treated under 150-450 MPa for 0.0-5.0 min.

Volatile amines formation

Formation of TVB can be considered as relatively low, being all values included in the 194.3-254.6 mg TVB-N kg⁻¹ muscle range (Table 2). The freezing step led to a higher mean value (comparison of T-0 and T-1 batches), although differences were not found significant (p>0.05). Comparison among samples without previous HP treatment did not provide a general trend with the storage time, being obtained the two highest scores in fish corresponding to 2.50 and 5.00 months; a general trend about the effect of the frozen storage time could not be observed for samples corresponding to 169.27, 300 and 430.73 MPa treatments. Concerning the pressure effect, a higher mean score was obtained at most frozen storage times in hake belonging to the 0.10-MPa batch, showing significant differences (p<0.05) at months 0.00 and 2.50. Accordingly, a partial inhibition of TVB formation could be concluded as a result of the previous pressure applied. However, a multifactor ANOVA analysis did not reflect a significant (p>0.05) effect of pressure applied on the total volatile amines formation.

TMA formation revealed a significant increase (p<0.05) as a result of the freezing step (Table 2). Furthermore, a relevant formation was observed with frozen storage time when comparing the different samples corresponding to batches without previous HP treatment, so that a 9.70±2.82 mg TMA-N kg⁻¹ muscle value was reached at the end of the experiment; a general trend concerning the effect of frozen storage time on TMA formation in each of the pressures applied (i.e., 169.27, 300 and 430.73 MPa) could not be reached. When the previous pressure effect is analysed, lower TMA-N values (p<0.05) were detected in HP-treated fish at months 0.00, 2.50 and 5.00. Thus, a

partial inhibition of the TMA-N value could be concluded as a result of the previous pressure applied. However, as for the TVB-N detection, a multifactor ANOVA analysis did not provide a significant ($p>0.05$) effect of pressure applied on the TMA formation.

A marked DMA formation was detected as a result of the freezing step (Table 2). A progressive formation of DMA could be observed in 0.10-MPa samples with frozen storage time for the 0.00-2.50-month period, being then the highest mean value reached at the end of the experiment. An increasing DMA content could also be observed with frozen storage time for samples corresponding to 169.27, 300 and 430.73 MPa. When considering the previous pressure effect, a higher mean DMA-N score was obtained at all frozen storage times in samples corresponding to fish without previous pressure treatment. This effect was found to increase with the previous pressure applied. Accordingly, an inhibitory effect on the formation of this degradative compound was implied by previous pressure treatment.

In agreement with this marked effect of both variables (pressure level and frozen storage time), the multifactor ANOVA analysis indicated an important effect on DMA formation, this yielding a significant ($p<0.0002$) model with an F-value of 27.99. The evaluation of the F-values for the two independent variables confirmed their individual effects. Thus, DMA formation was highly affected by frozen storage time (F-value = 51.66; p-value probability $> F$ was $p<0.0001$); meantime, an important effect of pressure level was also observed (F-value of 4.33; p-value probability $> F$ was $p<0.0710$). The correlation value of the model was $r^2 = 0.8750$ with adjusted and predicted r^2 values of 0.8437 and 0.7405, respectively, and a signal/noise ratio of 13.78. These statistical parameters confirmed that an empirical coded equation (Eq. 1) could be used to model the effect of previous pressure applied and frozen storage time on the DMA formation.

$$Y = 1.8426 - 0.2657 x_1 + 0.9177 x_2 \quad (\text{Eq. 1})$$

In this equation, Y is the DMA-N value (mg kg^{-1} muscle); x_1 is the coded variable for frozen storage time (months) and x_2 is the coded variable for pressure level (MPa). Fig. 1 shows the graph that represents the prediction of the model for the DMA formation.

Degradation compounds resulting from TMAO breakdown exert a strong impact on the quality loss of frozen Gadiform species during commercialisation (Sotelo and Rehbein 2000; Leelapongwattana et al. 2005). In the present research, TMA and DMA, well-known for their characteristic off-flavours, revealed an important formation as a result of the freezing step, as well as throughout the frozen storage time. Since microbial activity is not likely to occur under the actual experimental conditions, TMA formation can be explained on the basis of protein damage (Méndez et al. 2017). Concerning DMA, its formation during the current study has shown to be the most important event concerning volatile amines formation. According to the general behaviour of most Gadiform species under frozen conditions, its formation can be explained on the basis of the TMAO breakdown catalysed by an endogenous enzyme widely present in such fish species (i.e., TMAOase), and be produced equimolecularly with FA. Differently to TMA results, a marked inhibitory effect of previous pressure treatment on DMA formation was proved in the current study, this effect being increased with the pressure level applied. This result can be attributed to the TMAOase damage resulting from the previous pressure treatment, this leading to an inhibitory effect of its activity. To our knowledge, no previous research accounts for the inhibitory effect of HP treatment on DMA formation during a further frozen storage of hake.

Previous research related to volatile amines formation and TMAOase activity inhibition by HP treatment can be considered scarce. Thus, Malinowska-Pańczyk and Kołodziejska (2016) showed that formation of DMA in frozen minced cod (*Gadus morhua*) meat after 40 d at -5°C was strongly reduced if previously HP-treated at 193 MPa. Additionally, Gou et al. (2010) reduced the TMAOase activity and TMA and DMA formation in refrigerated squid (*Todarodes pacificus*) by previous treatment at 300 MPa for 20 min, while Malinowska-Pańczyk and Kołodziejska (2016) showed that formation of TMA in refrigerated minced cod (*Gadus morhua*) meat was strongly reduced if previously HP-treated at 193 MPa. Finally, Gou et al. (2012) reduced the formation of TMA and DMA in semi-dried squid (*T. pacificus*) by previous treatment at 500 MPa for 0-10 min. However, when a pelagic fatty fish species was concerned (sardine; Méndez et al. 2017), a progressive TMA content was obtained throughout a 9-month frozen storage at -18°C , but no inhibitory effect of a previous treatment at 125 MPa was implied.

Assessment of other chemical indices

Comparison between fish corresponding to T-0 and T-1 batches proved a great formation ($p<0.05$) of FFA as a result of the freezing step (Table 3). FFA values obtained for samples corresponding to 0.10-MPa batches revealed a significant ($p<0.05$) increase with the frozen storage time throughout the whole experiment. An increasing FFA content could also be observed with frozen storage time for samples corresponding to 169.27, 300 and 430.73 MPa. Concerning the pressure effect, at each frozen storage time, a lower FFA value could be obtained by increasing the previous pressure applied, being the differences significant ($p<0.05$) in most cases.

In agreement with this important effect of both process variables on FFA formation, the multifactor ANOVA analysis yielded a significant ($p < 0.0001$) model with an F-value of 36.33. The evaluation of the F-values for the two independent variables confirmed their individual effects. Thus, FFA formation was highly affected by frozen storage time (F-value = 63.10; p-value probability $> F$ was $p < 0.0001$); meantime, an important effect of pressure level could be observed (F-value of 9.55; p-value probability $> F$ was $p < 0.0149$). The correlation value of the model was $r^2 = 0.9008$ with adjusted and predicted r^2 values of 0.8760 and 0.8061, respectively, and a signal/noise ratio of 16.41. These statistical parameters confirmed that an empirical coded equation (Eq. 2) could be used to model the effect of previous pressure applied and frozen storage time on the FFA formation.

$$Y = 10.2267 - 0.0148 x_1 + 2.2885 x_2 \quad (\text{Eq. 2})$$

In this equation, Y is the FFA value (g kg^{-1} lipids); x_1 is the coded variable for frozen storage time (months) and x_2 is the coded variable for pressure level (MPa). Fig. 2 shows the graph that represents the prediction of the model for the FFA formation.

In agreement with previous research, hake fish has shown a great FFA formation throughout the frozen storage as a result of endogenous enzyme activity such as lipases and phospholipases (Aubourg and Medina 1999). Accumulation of FFA has been related to some extent to lack of acceptability, because FFA are known to have detrimental effects on ATPase activity, protein solubility, relative viscosity and texture deterioration (Aubourg et al. 2007). Present results concerning the inhibition of FFA formation in frozen hake can be explained on the basis of the lipases and

phospholipases damage produced during the previous HP treatment, so that a lower activity of such enzymes would be developed during the frozen storage.

Reported research concerning the effect on FFA formation after HP treatment followed by a frozen storage is scarce. Thus, Ohshima et al. (1992) found that enzymatic degradation of phospholipids (i.e., phospholipases action) in cod (*Gadus morhua*) muscle was successfully inhibited during storage at -2°C for 6 days when previously treated at pressures above 400 MPa for 15 and 30 min; however, no effect was observed when applying a pressure of 200 MPa. Later on, freezing by pressure-shift freezing at 140 MPa was compared with air-blast freezing at -20°C in turbot (*Scophthalmus maximus*) fillets (Chevalier et al. 2000); as a result, FFA formation during subsequent storage at -20°C for 75 days did not provide differences. Finally, Atlantic mackerel (*Scomber scombrus*; Vázquez et al. 2013) and horse mackerel (*Trachurus trachurus*; Torres et al. 2013) were subjected to HP treatment at 150-450 MPa for 0.0-5.0 min prior to frozen storage at -10°C for 3 months; the results revealed a marked reduction of FFA content for all HP pre-treated samples, with the reduction effect increasing with the previous pressure level and the pressure holding time employed.

No FA formation could be observed as a result of the freezing step (Table 3). However, evaluation of frozen hake without previous HP treatment led to a progressive FA formation with storage time; thus, a marked FA formation was reached at month 2.50, while the highest mean value was obtained at the end of the experiment. An increasing FA content could also be observed with frozen storage time for samples corresponding to 169.27, 300 and 430.73 MPa treatments. Concerning the effect of the applied pressure, higher mean FA values were obtained in fish corresponding to batches

without previous pressure treatment (0.32-5.00-month period). Thus, a marked inhibition of FA formation was obtained in frozen hake with previous applied pressure.

In agreement with this important effect of both process variables on FA formation, the multifactor ANOVA analysis yielded a significant ($p < 0.0001$) model with an F-value of 41.42. The evaluation of the F-values for the two independent variables showed that FA formation was highly affected by frozen storage time (F-value = 81.77; p-value probability $> F$ was $p < 0.0001$); meantime, a not significant effect of pressure could be observed (F-value 0.62; p-value probability $> F$ was $p < 0.4564$). The correlation value of the model was $r^2 = 0.9221$ with adjusted and predicted r^2 values of 0.8998 and 0.8623, respectively, and a signal/noise ratio of 16.44. These statistical parameters confirmed that an empirical coded equation (Eq. 3) could be used to model the effect of previous pressure applied and frozen storage time on the FA formation.

$$Y = 150.3200 - 6.1268 x_1 + 70.2936 x_2 \quad (\text{Eq. 3})$$

In this equation, Y is the FA content (mg kg^{-1} muscle); x_1 is the coded variable for frozen storage time (months) and x_2 is the coded variable for pressure level (MPa). Fig. 3 shows the graph that represents the prediction of the model for the FA formation.

As previously mentioned, FA is produced equimolecularly with DMA as a result of TMAO breakdown by TMAOase catalysis (Sotelo and Rehbein 2000; Benjakul et al. 2004). FA is well known as a reactive aldehyde, susceptible to produce inter- and intramolecular protein crossing, this leading to different kinds of damages such as denaturation enhancement, aggregates formation and texture losses (Rey-Mansilla et al. 2001). In agreement with its high reactivity, FA formation in the current research did not provide such an accurate assessment of TMAO breakdown as DMA. However, and

in agreement with previous comments on inhibition of DMA formation, the lower FA content found in HP-treated fish can be explained on the basis that previous pressure treatment has led to a TMAOase damage, so that an inhibitory effect on the activity of such enzyme was produced during the frozen storage.

Previous research concerning the effect of HP treatment on FA formation can be considered very scarce. Thus, Malinowska-Pańczyk and Kołodziejaska (2016) showed that formation of FA and inhibition of TMAOase activity in frozen minced cod (*Gadus morhua*) meat were strongly reduced if previously treated at 193 MPa at –20 °C.

Mean K value showed a marked increase as a result of the freezing step (Table 3), although differences were not significant ($p>0.05$). Evaluation of samples without previous HP treatment showed some increasing tendency with frozen storage time, leading to a 29.61 ± 2.50 score at the end of the storage; similarly, an increasing trend with frozen storage time was implied in samples corresponding to 169.27, 300 and 430.73 MPa. When analysing the effect of the previous pressure applied, significant differences were not obtained at any of the frozen storage times considered, so that it could be concluded that the previous HP treatment did not affect the K value in frozen hake. Accordingly, a multifactor ANOVA analysis did not provide a significant ($p>0.05$) model about the effect of the two variables of the process, i.e., pressure level and frozen storage time.

During post-mortem fish storage, muscle nucleotides are known to degrade in a series of stages as a result of endogenous biochemical changes, and the level of adenine nucleotides and their related compounds have extensively been used as an index (K value) of the freshness of fish muscle (Özogul et al. 2010). This index is found very sensitive as a freshness marker during the first stages of a refrigerated storage; however,

when a frozen storage period is concerned, its validity as an accurate quality tool has been found rather limited.

Previous research accounts for a K-value decrease in refrigerated and chilled fish by means of inactivation of enzymes that catalyse ATP degradation by previous HP treatment. Such results concern tilapia (*Oreochromis niloticus*) fillets (Ko and Hsu 2001), yellowfin tuna (*Thunnus albacares*) chunks (Kamalakanth et al. 2011) and white prawn (*Fenneropenaeus indicus*) (Ginson et al. 2013). However, no information is available to our knowledge concerning the HP effect on the K value in frozen fish.

CONCLUSIONS

The effect of a previous HP treatment at 150-450 MPa for 2 min on frozen hake quality was studied by means of a statistical experiment design. A multifactor ANOVA test showed an inhibitory effect ($p < 0.015$) on DMA and FFA formation as a result of the previous HP treatment, this effect increasing with pressure level applied. Additionally, an inhibition ($p < 0.05$) of FA, TMA, TVB-N and fluorescent compounds formation was obtained according to a one-way ANOVA test. This inhibitory effect on the formation of such molecules can be explained on the basis of the damage caused to different kinds of enzymes such as TMAOase, lipases and phospholipases, so that their activity during the subsequent frozen storage would decrease. On the contrary, no effect ($p > 0.05$) on the K value, PI and formation of peroxides and TBARS was observed in the present study.

The frozen storage of most Gadiform species is known to be strongly limited by the TMAO breakdown into DMA and FA, two highly deteriorative compounds. The work here presented provides for the first time with information concerning the employment of the HP technology to inhibit the DMA and FA formation during the

frozen storage, as well as the FFA formation. Further research focussed on commercial frozen conditions (−18 °C) and including sensory and nutritional aspects is foreseen.

ACKNOWLEDGMENTS

The authors thank Mr. Marcos Trigo, Mrs. Montserrat López and Mr. Roberto Iglesias for their excellent technical assistance. The work was supported by the Consejo Superior de Investigaciones Científicas (CSIC) (Spain) through the Research Project 2017-70E032, by Fundação para a Ciência e a Tecnologia (FCT Portugal), European Union, QRN, FEDER, COMPETE through founding of the Organic Chemistry Research Unit (QOPNA) (project PEst-C/QUI/UI0062/2013; FCOMP-01-0124-FEDER-037296), and by Formula Grants no. 2011-31200-06041 and 2012-31200-06041 from the USDA National Institute of Food and Agriculture.

REFERENCES

- Antonacopoulos, N. (1960). Verbesserte apparatus zur quantitativer destillation wasserdampf-flüchtiger stoffe. *Zeitschrift für Lebensmittel Untersuchung und Forschung*, 113, 113-160.
- Aubourg, S. P., & Medina, I. (1999). Influence of storage time on lipid deterioration during cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*) frozen storage. *Journal of the Science of Food and Agriculture*, 79, 1943-1948.
- Aubourg, S. P., Lago, H., Sayar, N., & González, R. (2007). Lipid damage during frozen storage of Gadiform species captured in different seasons. *European Journal of Lipid Science and Technology*, 109, 608-616.
- Aubourg, S. P., Piñeiro, C., Gallardo, J. M., & Barros-Velázquez, J. (2005). Biochemical changes and quality loss during chilled storage of farmed turbot (*Psetta maxima*). *Food Chemistry*, 90, 445-452.
- Benjakul, S., Visessanguan, W., & Tanaka, M. (2004). Induced formation of dimethylamine and formaldehyde by lizardfish (*Saurida micropectoralis*) kidney trimethylamine-N-oxide demethylase. *Food Chemistry*, 84, 297-305.
- Bligh, E., & Dyer, W. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37, 911-917.
- Campus, M. (2010). High pressure processing of meat, meat products and seafood. *Food Engineering Reviews*, 2, 256-273.
- Cando, D., Moreno, H., Tovar, C., Herranz, B., & Borderías, A. J. (2014). Effect of high pressure and/or temperature over gelation of isolated hake myofibrils. *Food and Bioprocess Technology*, 7, 3197-3207.

560 Chapman, R., & McKay, J. (1949). The estimation of peroxides in fats and oils by the
 561 ferric thiocyanate method. *Journal of the American Oil Chemists' Society*, 26,
 562 360-363.

563 Chevalier, D., Sequeira-Muñoz, A., Le Bail, A., Simpson, B., & Ghoul, M. (2000).
 564 Effect of pressure shift freezing, air-blast freezing and storage on some
 565 biochemical and physical properties of turbot (*Scophthalmus maximus*).
 566 *Lebensmittel-Wissenschaft und -Technologie*, 33, 570-577.

567 Dowden, H. (1938). The determinations of small amounts of dimethylamine in
 568 biological fluids. *Biochemical Journal*, 32, 455-459.

569 **FAO (2007). Fishery statistics. Commodities. Yearbook 2005, 101, p. 129-131. Food**
 570 **and Agriculture Organization of the United Nations, Rome, Italy.**

571 Ginson, J., Kamalakanth, C., Bindu, J., Venkateswarlu, R., Das, S., Chauhan, O., &
 572 Srinivasa Gopal, T. (2013). Changes in K value, microbiological and sensory
 573 acceptability of high pressure processed Indian white prawn (*Fenneropenaeus*
 574 *indicus*). *Food and Bioprocess Technology*, 6, 1175-1180.

575 Gou, J., Choi, K.-P., He, X., & Ahn, J. (2012). Dimethylamine, trimethylamine, and
 576 biogenic amine formation in high-pressure processed semidried squid
 577 (*Todarodes pacificus*) during refrigerated storage. *Journal of Food Science*, 75,
 578 M489-M495.

579 Gou, J., Lee, H.-Y., & Ahn, J. (2010). Effect of high pressure processing on the quality
 580 of squid (*Todarodes pacificus*) during refrigerated storage. *Food Chemistry*, 119,
 581 471-476.

582 Gudmundsson, M., & Hafsteinsson, H. (2002). Minimal processing in practice: seafood.
 583 In T. Ohlsson, & N. Bengtsson (Eds.), *Minimal processing technologies in the*
 584 *food industry* (pp. 245-266). Boca Raton, FL, USA: CRC Press.

- Hurtado, J., Montero, P., & Borderías, A. J. (2001). Chilled storage of pressurized octopus (*Octopus vulgaris*) muscle. *Journal of Food Science*, 66, 400-406.
- Kamalakanth, C., Ginson, J., Bindu, J., Venkateswarlu, R., Das, S., Chauhan, O., & Srinivasa Gopal, T. (2011). Effect of high pressure on K-value, microbial and sensory characteristics of yellowfin tuna (*Thunnus albacares*) chunks in EVOH films during chill storage. *Innovative Food Science and Emerging Technologies*, 12, 451-455.
- Ko, W-C., & Hsu, K-C. (2001). Changes in K value and microorganisms of tilapia fillet during storage at high-pressure, normal temperature. *Journal of Food Protection* 64, 94-98.
- Lakshmanan, R., Pigott, J., & Paterson, A. (2003). Potential applications of high pressure for improvement in salmon quality. *Trends in Food Science and Technology*, 14, 354-363.
- Leelapongwattana, K., Benjakul, S., Visessanguan W., & Howell, N. (2005). Physicochemical and biochemical changes during frozen storage of minced flesh of lizardfish (*Saurida micropectoralis*). *Food Chemistry*, 90, 141-150.
- Lowry, R., & Tinsley, I. (1976). Rapid colorimetric determination of free fatty acids. *Journal of the American Oil Chemists' Society*, 53, 470-472.
- Malinowska-Pańczyk, E., & Kołodziejska, I. (2016). The effect of high pressure on formation of volatile amines in minced meat of cod (*Gadus morhua*). *European Food Research and Technology*, 242, 415-420.
- Méndez, L., Fidalgo, L., Pazos, M., Lavilla, M., Torres, J. A., Saraiva, J., Vázquez, M., & Aubourg, S. P. (2017). Lipid and protein changes related to quality loss in frozen sardine (*Sardina pilchardus*) previously processed under high-pressure conditions. *Food and Bioprocess Technology*, 10, 296-306.

- Nash, T. (1953). The colorimetric estimation of formaldehyde by means of the Hantzsch reaction. *Biochemical Journal*, 55, 416-421.
- Ohshima, T., Nakagawa, T., & Koizumi, C. (1992). Effect of high hydrostatic pressure on the enzymatic degradation of phospholipids in fish muscle during storage. In E. Bligh (Ed.), *Seafood Science and Technology*, Chapter 8 (pp. 64-75). Oxford, UK: Fishing News Books.
- Özogul, Y. (2010). Methods for freshness quality and deterioration. In Nollet, L., & Toldrá, F. (Eds.), *Handbook of seafood and seafood products analysis* (pp. 189-214). Boca Raton, FL, USA: CRC Press, Taylor & Francis Group.
- Rey-Mansilla, M., Sotelo, C., Aubourg, S. P., Rehbein, H., Havemeister, W., Jørgensen, B., & Nielsen, M. (2001). Localization of formaldehyde production during frozen storage of European hake (*Merluccius merluccius*). *European Food Research and Technology*, 213, 43-47.
- Sotelo, C., & Rehbein, H. (2000). TMAO-degrading enzymes. In N. Haard, & B. Simpson (Eds.), *Seafood enzymes. Utilization and influence on postharvest seafood quality* (pp. 167-190). New York, USA: Marcel Dekker, Inc.
- Tabilo-Munizaga, G., Aubourg, S. P., & Pérez-Won, M. (2016). Pressure effects on seafoods. In V. M. Balasubramanian, G. Barbosa-Cánovas, & H. Lelieveld (Eds.), *High pressure processing of food: Principles, technology and application* (pp. 625-669). Heidelberg, Germany: Springer, Science and Business, Inc.
- Tironi, V., de Lamballerie, M., & Le Bail, A. (2010). Quality changes during the frozen storage of sea bass (*Dicentrarchus labrax*) muscle after pressure shift freezing and pressure assisted thawing. *Innovative Food Science and Emerging Technologies*, 11, 565-573.

634 Tironi, V., Tomás, M., & Añón, M^aC. (2002). Structural and functional changes in
 635 myofibrillar proteins of sea salmon (*Pseudopercis semifasciata*) by interaction
 636 with malondialdehyde (RI). *Journal of Food Science*, 67, 930-935.

637 Torres, J. A., Vázquez, M., Saraiva, J. A., Gallardo, J., & Aubourg, S. P. (2013). Lipid
 638 damage inhibition by previous high pressure processing in white muscle of
 639 frozen horse mackerel. *European Journal of Lipid Science and Technology*, 115,
 640 1454-1461.

641 Tozawa, H., Erokibara, K., & Amano, K. (1971). Proposed modification of Dyer's
 642 method for trimethylamine determination in codfish. In R. Kreuzer (Ed.), *Fish*
 643 *Inspection and Quality Control* (pp. 187-190). London, UK: Fishing News
 644 Books Ltd.

645 Vázquez, M., Torres, J. A., Gallardo, J., Saraiva, J., & Aubourg, S. P. (2013). Lipid
 646 hydrolysis and oxidation development in frozen mackerel (*Scomber scombrus*):
 647 Effect of a high hydrostatic pressure pre-treatment. *Innovative Food Science and*
 648 *Emerging Technologies*, 18, 24-30.

649 Vidacek, S., De las Heras, C., Solas, M., Rodríguez-Mahillo, A., & Tejada, M. (2009).
 650 Effect of high hydrostatic pressure on mortality and allergenicity of *Anisakis*
 651 *simplex* L3 and on muscle properties of infested hake. *Journal of the Science of*
 652 *Food and Agriculture*, 89, 2228-2235.

653 Villamonte, G., Pottier, L., & de Lamballerie, M. (2016). Influence of high-pressure
 654 processing on the physicochemical and the emulsifying properties of
 655 sarcoplasmic proteins from hake (*Merluccius merluccius*). *European Food*
 656 *Research and Technology*, 242, 667-675.

657 Vyncke, W. (1970). Direct determination of the thiobarbituric acid value in
658 trichloroacetic acid extracts of fish as a measure of oxidative rancidity. *Fette,*
659 *Seifen, Anstrichmittel*, 72, 1084-1087.

660

661

FIGURE LEGENDS

Fig. 1: Prediction model obtained for the effect of pressure level (MPa) and frozen storage time (months) on the dimethylamine-nitrogen (DMA-N) content (mg kg^{-1} muscle) in frozen hake muscle previously processed under high-pressure conditions.

Fig. 2: Prediction model obtained for the effect of pressure level (MPa) and frozen storage time (months) on the free fatty acids (FFA) content (g kg^{-1} lipids) in frozen hake muscle previously processed under high-pressure conditions.

Fig. 3: Prediction model obtained for the effect of pressure level (MPa) and frozen storage time (months) on the formaldehyde (FA) content (mg kg^{-1} muscle) in frozen hake muscle previously processed under high-pressure conditions.

TABLE 1: Lipid oxidation assessment* in frozen hake muscle previously processed under different pressure levels

Batch number**	Pressure (MPa)	Storage time (months)	Quality index			
			PV	TBA-i	FR	PI
T-0	Starting fresh fish		0.73 aA (0.47)	0.32 aAB (0.12)	0.58 aA (0.03)	2.67 B (0.06)
T-1	0.10	0.00	1.93 bB (0.10)	0.63 bC (0.16)	0.98 bC (0.20)	2.83 BCD (0.13)
T-2	300	0.00	0.97 aA (0.28)	0.50 bB (0.02)	0.79 abD (0.18)	2.76 (0.05)
T-3	0.10	0.32	1.62 abAB (0.81)	0.80 C (0.10)	0.80 BC (0.23)	2.88 CD (0.08)
T-4	169.27	0.32	0.59 aA (0.26)	0.87 B (0.15)	0.66 (0.10)	2.74 (0.05)
T-5	430.73	0.32	1.46 bA (0.22)	0.61 (0.23)	0.57 (0.08)	2.86 (0.10)
T-6	0.10	2.50	11.45 dC (3.97)	0.35 abB (0.04)	0.87 dBC (0.16)	2.76 BC (0.06)
T-7	150	2.50	2.50 a (0.14)	0.25 a (0.06)	0.54 bc (0.07)	2.86 (0.15)
T-8	300	2.50	3.68 bcB (0.51)	0.50 bB (0.17)	0.61 cCD (0.04)	2.74 (0.17)
T-9	300	2.50	4.75 bcB (0.84)	0.35 abAB (0.07)	0.38 aA (0.02)	2.66 (0.11)
T-10	300	2.50	3.50 abB (0.95)	0.39 bB (0.03)	0.42 abAB (0.10)	2.79 (0.04)
T-11	450	2.50	4.52 bc (1.32)	0.30 ab (0.06)	0.60 abcd (0.21)	2.69 (0.04)
T-12	0.10	4.68	7.65 abC (2.42)	0.17 abA (0.07)	0.75 abBC (0.24)	2.51 aA (0.05)
T-13	169.27	4.68	5.33 aB (0.81)	0.10 aA (0.02)	0.48 a (0.08)	2.75 b (0.10)
T-14	430.73	4.68	8.34 bB (1.47)	0.37 b (0.15)	0.61 b (0.03)	2.80 b (0.05)
T-15	0.10	5.00	3.08 B (1.68)	0.15 aA (0.04)	0.73 bB (0.01)	3.00 bD (0.07)
T-16	300	5.00	2.61 AB (1.45)	0.28 bA (0.02)	0.53 aBC (0.01)	2.73 a (0.11)

* Lipid oxidation indices: PV (peroxide value; meq active oxygen kg⁻¹ lipids), TBA-i (thiobarbituric acid index; mg malondialdehyde kg⁻¹ muscle), FR (fluorescence ratio) and PI (polyene index). For each quality index and for each frozen storage time, mean values followed by different low-case letters denote significant (p<0.05) differences. For each quality index and for each high-pressure level, mean values followed by different capital letters denote significant (p<0.05) differences. No letters are indicated when significant differences are not found (p>0.05).

** Seventeen batches (from T-0 to T-16) were considered in agreement with the statistical experiment design described in the experimental section. Pressure level (0.1, 150, 169.27, 300, 430.73, 450 and 500 MPa) and frozen storage time (0.00, 0.32, 2.50, 4.68 and 5.00 months) values for each batch are described in second and third columns, respectively.

TABLE 2: Volatile amines formation* in frozen hake muscle previously processed under different pressure levels

Batch number**	Pressure (MPa)	Storage time (months)	Quality index		
			TVB-N	TMA-N	DMA-N
T-0	Starting fresh fish		201.62 abA (3.90)	0.20 aA (0.05)	1.83 aA (0.41)
T-1	0.10	0.00	211.36 bAB (5.80)	1.11 cB (0.33)	5.29 bB (1.54)
T-2	300	0.00	196.66 aA (4.80)	0.43 bA (0.00)	3.83 bA (1.10)
T-3	0.10	0.32	217.64 aB (4.95)	1.06 aB (0.07)	11.20 C (2.35)
T-4	169.27	0.32	229.42 b (5.28)	7.66 bB (1.81)	10.90 A (4.82)
T-5	430.73	0.32	213.78 a (9.78)	1.43 a (0.20)	9.46 A (2.85)
T-6	0.10	2.50	254.60 cC (12.83)	5.92 cC (0.54)	46.74 dDE (12.20)
T-7	150	2.50	214.98 b (6.83)	2.05 b (0.29)	19.59 abc (1.72)
T-8	300	2.50	207.37 abAB (11.06)	1.79 bB (0.44)	19.79 bcB (0.53)
T-9	300	2.50	203.38 abAB (8.09)	3.27 bBC (2.31)	18.61 abB (1.71)
T-10	300	2.50	211.38 bB (2.55)	3.08 bBC (1.06)	22.68 cC (2.02)
T-11	450	2.50	194.38 a (10.28)	0.91 a (0.14)	15.91 a (3.11)
B					
T-12	0.10	4.68	220.94 B (13.97)	4.62 bC (1.33)	35.15 bD (7.51)
T-13	169.27	4.68	206.63 (15.30)	4.00 bA (0.83)	34.31 bB (6.88)
T-14	430.73	4.68	198.71 (4.78)	1.60 a (0.29)	22.34 aB (1.84)
T-15	0.10	5.00	252.23 C (10.86)	9.70 bD (2.82)	54.82 bE (6.14)
T-16	300	5.00	235.51 C (7.74)	3.95 aC (0.76)	25.25 aC (4.44)

* Volatile amine indices: TVB-N (total volatile amines-nitrogen; mg kg⁻¹ muscle), TMA-N (trimethylamine-nitrogen; mg TMA-N kg⁻¹ muscle) and DMA-N (dimethylamine-nitrogen; mg DMA-N kg⁻¹ muscle). For each quality index and for each frozen storage time, mean values followed by different low-case letters denote significant (p<0.05) differences. For each quality index and for each high-pressure level, mean values followed by different capital letters denote significant (p<0.05) differences. No letters are indicated when significant differences are not found (p>0.05).

** Description of batches as expressed in Table 1.

TABLE 3: Assessment* of free fatty acids (FFA; g kg⁻¹ lipids) formation, formaldehyde (FA; mg kg⁻¹ muscle) content and K value (%) in frozen hake muscle previously processed under different pressure levels

Batch number**	Pressure (MPa)	Storage time (months)	Quality index		
			FFA	FA	K value
T-0	Starting fresh fish		25.70 aA (1.31)	0.83 A (0.05)	6.98 aA (3.61)
T-1	0.10	0.00	47.48 cB (5.39)	0.70 A (0.07)	16.77 abBC (6.31)
T-2	300	0.00	35.56 bA (1.70)	0.86 A (0.08)	17.67 b (1.86)
T-3	0.10	0.32	88.35 bC (13.25)	0.91 bA (0.10)	16.44 B (2.25)
T-4	169.27	0.32	74.20 bA (5.62)	0.72 aA (0.06)	14.70 (2.23)
T-5	430.73	0.32	52.92 aA (2.85)	0.77 abA (0.08)	16.07 (6.04)
T-6	0.10	2.50	266.00 eD (23.01)	6.60 dBC (2.80)	26.48 D (5.94)
T-7	150	2.50	155.94 d (13.46)	1.66 c (0.15)	20.00 (4.62)
T-8	300	2.50	121.87 cC (2.76)	1.47 bC (0.05)	20.70 (2.62)
T-9	300	2.50	112.32 bB (3.41)	1.17 aB (0.15)	21.06 (4.66)
T-10	300	2.50	138.70 cdC (21.41)	1.67 cD (0.05)	20.83 (2.85)
T-11	450	2.50	99.83 a (2.19)	1.45 ab (0.25)	22.65 (8.04)
T-12	0.10	4.68	327.98 cE (11.75)	4.01 bB (1.24)	22.12 CD (1.16)
T-13	169.27	4.68	171.97 bB (4.24)	4.06 bB (0.89)	21.05 (4.95)
T-14	430.73	4.68	128.96 aB (8.77)	2.05 aB (0.27)	25.27 (2.13)
T-15	0.10	5.00	338.77 bE (12.92)	7.00 bC (1.20)	29.61 D (2.50)
T-16	300	5.00	172.33 aD (16.49)	2.46 aD (0.82)	21.56 (4.32)

* For each quality index and for each frozen storage time, mean values followed by different low-case letters denote significant ($p < 0.05$) differences. For each quality index and for each high-pressure level, mean values followed by different capital letters denote significant ($p < 0.05$) differences. No letters are indicated when significant differences are not found ($p > 0.05$).

** Description of batches as expressed in Table 1.