This is a post print version of:

DITA M-A., RISPAIL N., PRATS E., RUBIALES D., SINGH K.B., 2006

Biotechnology approaches to overcome biotic and abiotic stress
constraints in grain legumes.

Euphytica 147: 1-24.

DOI 10.1007/s10681-006-6156-9

The printed version can be visited at:

<https://link.springer.com/article/10.1007/s10681-006-6156-9>

1 **Biotechnology approaches to overcome biotic and abiotic stress constraints in legumes**

2 Miguel A. Dita*¹, Nicolas Rispail¹, Elena Prats¹, Diego Rubiales¹, Karam B. Singh²

3 ¹Instituto de Agricultura Sostenible, CSIC, Apdo. 4084, E-14080, Córdoba, Spain.
4 (*author for correspondence; e-mail: mangel@vicoso.ufv.br)

5
6 ²CSIRO Plant Industry, Private Bag 5, Wembley, WA 6913, Australia
7

Keywords: metabolomics; mutagenesis; plant transformation; proteomics; tissue culture; transcriptomics

Abstract

Biotic and abiotic stresses cause significant yield losses in legumes and can significantly affect their productivity. Biotechnology tools such as marker-assisted breeding, tissue culture, *in vitro* mutagenesis and genetic transformation can contribute to solve or reduce some of these constraints. However, only limited success has been achieved so far. The emergence of “omic” technologies and the establishment of model legume plants such as *Medicago truncatula* and *Lotus japonicus* are promising strategies for understanding the molecular genetic basis of stress resistance, which is an important bottleneck for molecular breeding. Understanding the mechanisms that regulate the expression of stress-related genes is a fundamental issue in plant biology and will be necessary for the genetic improvement of legumes. In this review, we describe the current status of biotechnology approaches in relation to biotic and abiotic stresses in legumes and how these useful tools could be used to improve resistance to important constraints affecting legume crops.

1 **Introduction**

2 Legumes are among the most important crops worldwide, having major impacts on
3 agriculture, the environment, animal/human nutrition and health (Graham & Vance 2003).
4 Legumes can interact symbiotically with specific soil-borne bacteria, the rhizobia, which
5 allow the plant to fix atmospheric nitrogen and may help to protect them against some fungal
6 pathogens (Chakraborty et al. 2003). As such, they constitute a significant source of nitrogen
7 and consequently play an essential role in both the structure of ecosystems and sustainable
8 agriculture, worldwide. These symbiotic interactions have strongly driven the investigation
9 and application of biotechnology tools for legumes. Nevertheless, a number of biotic (fungi,
10 bacteria, nematodes, viruses, parasitic plants, insects) and abiotic (drought, freezing, salinity,
11 waterlogging) stresses are severely affecting the yield of these crops.

12 The adaptability and productivity of legumes are limited by major abiotic stresses
13 including drought, heat, frost, chilling, waterlogging, salinity and mineral toxicities. Although
14 the type and the severity depend on the specific crop location, abiotic stresses can result in
15 crop damages as high as those caused by biotic stresses. Furthermore, crops under abiotic
16 stress are usually more susceptible to weeds, insects and diseases, which increase
17 considerably the losses (Reddy et al. 2004). An additional factor relevant to the legumes is the
18 response of the symbiotic nitrogen-fixing bacteria to stresses. Application of biotechnology
19 approaches to these crops can contribute efficiently to solve or reduce these problems.

20 Successful application of biotechnology to biotic/abiotic constraints facing legume
21 crops will require both a good biological knowledge of the target species and the mechanisms
22 underlying resistance/tolerance to these stresses. The large genome size and the polyploidy of
23 some legumes have hampered this goal, but in order to solve some of these problems two
24 species, *Medicago truncatula* and *Lotus japonicus*, have emerged as model plants to
25 investigate the genetic of nodulation and other important processes such as resistance or

1 tolerance to stresses. Their respective small and diploid genomes, autogamous nature, short
2 generation times, and prolific seed production were important characteristics for these choices
3 (Cook 1999; Handberg & Stougaard 1992). Since then, powerful genetic and genomic tools
4 have been developed, such as genome sequencing (Kato et al. 2003), isolation of Expressed
5 Sequence Tags (ESTs; Asamizu et al. 2004; Kulikova et al. 2001) and the establishment of
6 genetic and physical maps for each model species (Pedrosa et al. 2002; Thoquet et al. 2002).
7 The increasing wealth of genetic and genomic data and the high degree of synteny between
8 legume genomes (Kalo et al. 2004; Stracke et al. 2004), make these two species valuable
9 models for the molecular genetic study of the biotic and abiotic constraints that hamper
10 legume crop yield.

11 Much of the research on plant stress responses in this area has been conducted with
12 Arabidopsis as a model system. Substantial similarities between the defence responses of
13 Arabidopsis and legumes exist, however, there are also significant differences (Anderson et
14 al. 2005). Thus, it is necessary to increase our understanding of the specific aspects of the
15 defence/stress responses in legumes in order to solve some of the major constraints facing
16 these crops. In this review, relevant advances in marker-assisted breeding, tissue culture,
17 genetic transformation, and gene expression, including large-scale approaches and functional
18 analyses are presented and discussed as a way to overcome biotic and abiotic stresses in
19 legumes.

21 **Major biotic and abiotic stress targets for improvement in legumes**

22 Legume production is greatly constrained by numerous biotic and abiotic stresses.
23 Many of the diseases, pest and abiotic stresses are common to all legume crops; however,
24 their incidence and importance vary according to the legume crop, management practices and
25 regions.

Biotic stresses

The major biotic stresses affecting legumes are fungal diseases although insects, nematodes, viruses, bacteria and parasitic weeds can drastically decrease legume production. Weeds are also a problem for many legume crops but will not be covered here.

The relative importance of aerial fungal diseases and their effect on yield varies among years and cropping regions. However, some of them affect large areas in all the countries where legumes are cultivated and cause considerable losses in quality and quantity. Foliar diseases caused by biotrophic pathogens, such as rusts, downy mildews and powdery mildews, are major limiting factors in legume production and the most important of these are present in all areas where legumes are cultivated (Sillero et al. 2005, this issue). Several rust species can infect grain and forage legumes, most of them belonging to the genus *Uromyces*, such as *U. appendiculatus* on common bean, *U. ciceris-arietini* on chickpea, *U. pisi* on pea, *U. striatus* alfalfa, *U. viciae-fabae* on faba bean, lentil and common vetch and *U. vignae* on cowpea. Also rust species belonging to other genera can be major problems on legumes such as *Phakopsora pachyrhizi* and *P. meibomiae* on soybean or *Puccinia arachidis* on groundnut (Rubiales et al. 2002). Asian rust (*Phakopsora pachyrhizi*) is a severe disease that causes important yield losses in soybean and is spreading rapidly around the world (Carmona et al. 2005; du Preez et al. 2005; Pivonia & Yang 2004). Lack of natural sources of resistance (Ramteke et al. 2004) makes this disease a good candidate to be solved using biotechnology. Normally, legume rust epidemics begin late in the season, when pod filling has started, so yield components are only slightly little affected by the infection and losses are usually low. However, when the infection starts early in the season severe epidemics can occur (Rashid & Bernier 1991).

1 Powdery mildew is an important fungal disease in several legumes caused by *Erysiphe*
2 *pisi* (Sillero et al., 2005, this issue). Powdery mildew of pea has a worldwide distribution
3 being particularly important in climates with warm, dry days and cool nights, adversely
4 affecting yield and quality. Severe infection may cause 25-50% yield losses (Warkentin et al.
5 1996). Downy mildew, caused by *Peronospora viciae* occurs in most places where the crops
6 are grown, but is most frequent and severe in cool, maritime climates (Sillero et al., 2005, this
7 issue).

8 The major necrotrophic fungal diseases are ascochyta blight on various grain legumes,
9 chocolate spot on faba bean and anthracnose of lupin and lentil (Tivoli et al. 2005, this issue).
10 Ascochyta blight, caused by *Ascochyta rabiei*, is the most important fungi disease of
11 chickpea. It affects above-ground parts of the plants causing 100% yield loss in some
12 situations (Nene & Reddy, 1987). Botrytis gray mould caused by *Botrytis cinerea* is of lesser
13 importance in chickpea but also a widespread foliar disease problem. The common foliar
14 diseases on faba bean are Ascochyta blight and Chocolate spot. Ascochyta blight, caused by
15 the fungus *Ascochyta fabae*, is distributed world-wide (Gaunt, 1983). Yield losses of about
16 40% are common, but losses can be as high as 90% in susceptible cultivars (Hanounik, 1980),
17 particularly under wet and cool weather conditions. Chocolate spot, caused by *Botrytis fabae*,
18 is a destructive leaf disease of faba bean that can reduce yields by more than 60 % (Hanounik,
19 1981), particularly in humid regions. Ascochyta blight is considered the most important
20 necrotrophic foliar disease on pea worldwide (Bretag & Ramsey, 2001). It is caused by three
21 related fungal species, commonly referred to as the Ascochyta complex: *Ascochyta pisi*,
22 *Ascochyta pinodes* (teleomorph *Mycosphaerella pinodes*) and *Phoma medicaginis*. The major
23 foliar necrotrophic pathogens on lupins are anthracnose, caused by *Colletotrichum lupine*,
24 followed by Brown spot, caused by *Pleiochaeta setosa* and Phomopsis, caused by *Diaporthe*
25 *toxica* (Sweetingham et al. 1998). Ascochyta blight of lentils, caused by *Ascochyta lentis* has

1 been reported worldwide in most lentil producing countries (Bayaa & Erskine, 1998).
2 Anthracnose of lentils, caused by *Colletotrichum truncatum* is a common and important
3 pathogen on lentil in Canada (Anderson et al., 2000), although of little importance in several
4 countries in Asia and Africa (Bayaa & Erskine 1998).

5 There are several soil-borne diseases that are common among legume crops (Infantino
6 et al. 2005, this issue). Most of these attack the seedling stage of the crop and are referred to
7 as damping-off diseases. For example, damping-off, generally caused by either *Rhizoctonia*
8 *solani* or *Pythium* spp., can result in up to 80% of plant death (Denman et al. 1995; Wang et
9 al. 2003). Fusarium root-rot (caused by *Fusarium* spp.) can also cause severe seedling losses
10 especially in common bean and lentils (Hamwieh et al. 2005; Schneider et al. 2001). In most
11 growing areas of the world, Fusarium wilt (caused by *F. oxysporum*) is a major constraint in
12 the production of pulse crops, chickpea (Navas-Cortés et al. 2000; Nene & Reddy 1987) and
13 lentil (Bayaa et al. 1997) in particular. The disease affects seedlings and adult plants where it
14 causes leaf chlorosis, wilting and death. Other important soil-borne diseases such as southern
15 stem rot (*Sclerotium rolfsii*) and the white mold (*Sclerotinia sclerotiorum*) can cause both
16 seedling and pod rots in warmer and cool weather respectively (Kolkman & Kelly 2003).

17 A number of parasitic plants have become weeds, posing severe constraints to major
18 crops including grain legumes (Rubiales et al. 2005, this issue). *Orobanche crenata* is an
19 important problem in most cool season legumes in the Mediterranean basin and Middle East.
20 Yield loss can be severe with complete loss of crops in severe cases. *O. aegyptiaca* is of
21 importance in the Middle East and Asia. *O. foetida* is widely distributed in natural habitats in
22 the Western Mediterranean area parasitizing wild herbaceous leguminous plants, but is
23 however considered an important agricultural parasite in the faba bean in Beja region of
24 Tunisia. *O. minor* is of economic importance on clover that is grown for seed and has recently
25 become a problem on red clover in Oregon, USA (Rubiales 2001; Rubiales et al. 2005). *Striga*

1 *gesnerioides* and *Alectra vogelii* cause considerable yield reduction of grain legume crops,
2 particularly cowpea, throughout semi-arid areas of sub-Saharan Africa (Parker & Riches
3 1993).

4 Viruses cause yield losses for most legume crops. For example, Bean Common
5 Mosaic Virus (BCMV) and its close relative, Bean Common Mosaic Necrotic Virus
6 (BCMNV) are the most widespread and frequent viruses of common bean leading to
7 significant losses. In addition, over the past two decades, Bean Golden Mosaic Virus
8 (BGMV) has been considered the most important yield limiting disease for bean production in
9 parts of Central America and the lowlands of the Caribbean, with yield losses between 10 and
10 100% (Coyne et al. 2003).

11 Insects are another important biotic stress faced by many legume crops. They cause
12 important damages both through direct feeding, as vectors or by providing infection sites for
13 pathogens (Edwards & Singh 2005, this issue). Examples of important insect pests in grain
14 legumes include aphids like *Aphis glycine*, pod borers such as *Helicoverpa armigera* and *H.*
15 *punctigera* in cool season legumes (Yoshida et al. 1997) and weevils such as *Apion godmani*
16 and *Zabrodes subfasciatus* in warm season legumes (Garza et al. 1996; Romero-Andreas et al.
17 1986).

18 *Abiotic stresses*

19
20 Abiotic stress is a broad term, which includes multiple stresses such as heat, chilling,
21 excessive light, drought, waterlogging, wounding, ozone exposure, UV-B irradiation, osmotic
22 shock and salinity. It has been estimated that only 10% of arable land can be classified under
23 the non-stress category, which implies that crops grown on the other 90% of arable lands
24 experience one or more environmental stresses. Some of these stresses like drought, extreme
25 temperature, and high salinity dramatically limit crop productivity. The prediction is that
26 water deficits will continue to be the major abiotic factor likely to affect crop yields globally

(Sharma & Lavanya 2002). Moreover in many legumes such as peanut (*Arachis hypogaea*), Brazil nuts (*Bertholletia excelsa*) and faba bean (*Vicia faba*), this stress is particularly important because pre-harvest aflatoxin contamination is a common occurrence (Arrus et al. 2005; Mahmoud & Abdalla 1994) that can be reduced in drought tolerant lines (Holbrook et al. 1994). On the other hand, waterlogging due to a combination of unfavourable weather conditions and suboptimal soil and irrigation techniques can result in severe yield losses (Dennis et al. 2000). Waterlogging limits the oxygen diffusion of the soil and as a consequence nitrification is replaced as the most important N-transforming process, by denitrification and/or nitrate ammonification (Laanbroek 1990). Furthermore, under waterlogging stress, plant potassium, sodium, iron, and manganese uptake are limited and crops become more susceptible to diseases. For instance, waterlogging peas are more susceptible to *M. pinodes* (McDonald & Dean 1996).

Soil salinity affects total nitrogen uptake and soil nitrogen contribution (van Hoorn et al. 2001), leading to yield reduction. It is also expected that with the decrease in the ozone layer, UV exposure will become an important stress for cropping system (Chimphango et al. 2003).

Several of the abiotic stresses associated with legume crops can also directly affect symbiotic interactions and therefore limit legume growth. *Sinorhizobium meliloti* shows pH-sensitivity below pH 6 reducing *Medicago sativa* development, while *Mesorhizobium loti* is tolerant up to pH 4, facilitating the growth of *Lotus glaber* in more acid soils (Correa et al. 2001). Deficiencies and toxicities of micronutrients are also an important constraint of legume crops. Limitation of growth due to boron toxicity or deficiency has been described for instance in pea or faba bean (Dwivedi et al. 1992; Poulain & Almohammad 1995). In some cases such as soybean, the deficiency or toxicity is more critical for root nodulation than for the direct growth of the plant (Rahman et al. 1999).

To face the threat represented by these stresses several genetic improvement strategies are available, from classical breeding to a more direct physiological-genetic approach. However, only with an understanding of the mechanisms underlying a specific stress, will the later strategy be feasible. In general for the stresses mentioned above, low yields in developing countries are primarily due to a lack of effective disease management practices, particularly the availability of disease-resistant cultivars. Moreover excessive and often inappropriate fungicide usage in many situations, such as occurs with the control of bean rust, can contribute to higher input costs, human health problems and contamination of water supplies and the environment. In this context, biotechnology is a powerful tool that has potential to contribute to sustainable agriculture. Biotechnology approaches such as marker-assisted breeding, tissues cultures, *in vitro* mutagenesis, and genetic transformation can contribute to speed up classical breeding and overcome major problems such as lack of natural sources of resistance and sexual incompatibility.

The fact that many of these stresses such as the pathogens, *Colletotrichum trifolii*, (Torregrosa et al. 2004), *Aphanomyces euteiches* (Nyamsuren et al. 2003), *Uromyces striatus* (Rubiales & Moral 2004), nematodes (Koltai et al. 2001), *Phytophthora megasperma* f. sp. *medicaginis*, *Fusarium* spp., *Ascochyta* spp. (Salzer et al. 2000) and the parasitic plant *O. crenata* (Rodríguez-Conde et al. 2004) also affects the legume model *M. truncatula*, will help increase our understanding of the underlying molecular and genetics basis of resistance, and consequently increase the potential for biotechnology to overcome these stresses in the major legume crops.

Biotechnology tools

Molecular marker-assisted breeding

1 The use of genetic and genomic analysis to help identify DNA regions tightly linked to
2 agronomic traits in crops, the so-called molecular markers, can facilitate breeding strategies
3 for crop improvement. The use of molecular markers for the indirect selection of improved
4 crops speeds up the selection process by alleviating time-consuming approaches direct
5 screening under greenhouse and field conditions. Molecular markers are particularly useful
6 when targeting characters controlled by several genes. The potential to map different
7 Quantitative Trait Loci (QTL) contributing to an agronomical trait and to identify linked
8 molecular markers opens up the possibility to transfer simultaneously several QTLs and to
9 pyramid QTLs for several agronomical traits in one improved cultivar.

10 Numerous molecular marker-related techniques have been used in legumes in relation
11 to biotic and abiotic stresses. Random Amplified Polymorphism (RAPD), Restriction
12 Fragment Length Polymorphism (RFLP), Amplified Fragment Length Polymorphism
13 (AFLP), Simple Sequence Repeat (SSR) and derivatives have been reported both for biotic
14 (Ouedraogo et al. 2002; Román et al. 2002) and abiotic (Kassem et al. 2004; Lee et al. 2004)
15 stresses. As a result, genetic maps for many species were established in which potential
16 resistance and/or tolerance loci or QTLs have been located (Table 1 and 2). This improved the
17 knowledge of the genetic control of specific resistance and/or tolerance in many legumes by
18 providing information on the number, chromosomal location and individual or interactive
19 effects of the QTLs involved. More importantly, these technologies have identified specific
20 molecular markers that may be used in breeding programs through Marker-Assisted Selection
21 (MAS) to enhance stress tolerance.

22 However, the application of molecular markers in breeding programs requires
23 preliminary studies to identify and validate potential markers. In this process, the following
24 factors have to be considered: a) level of polymorphism existing between parental lines, b)
25 unclear expression of some markers inherent to the marker class used, c) false-positive

1 markers, d) discrepancy between the presence of the marker and target gene, which requires
2 testing the gene with conventional screening and e) presence of multiple genes scattered over
3 several linkage groups (Yu et al. 2004).

4 Although the use of MAS may be helpful for crop improvement, its practical application in
5 legumes for the genetic improvement of resistance or tolerance to stress have been limited,
6 being mainly hampered by lack of investment and the genetic complexity of most stress-
7 related traits. There are some exceptions where MAS has already facilitated breeding efforts
8 for several legume crops against important biotic stress. For instance MAS was successfully
9 used for the breeding of resistant soybean to cyst nematode (Diers 2004), of resistant pinto
10 bean to common bacterial blight (Mutlu et al. 2005) and of resistant narrow-leafed lupin
11 (*Lupinus angustifolius* L.) to phomopsis stem blight (Yang et al. 2002) and anthracnose (Yang
12 et al. 2004). Moreover, when resistance is conferred by single genes and/or easily overcome
13 by new pathogens races, the gene pyramiding strategy facilitated by MAS can be an efficient
14 method. Breeding for abiotic stress is much more complicated due to the complexity of the
15 traits involved. Nevertheless, Schneider et al. (1997) showed that MAS may be useful to
16 select drought tolerant common bean.

18 *Gene pyramiding assisted by MAS*

19 Breeding durable resistance to biotic and/or abiotic stresses is a major task for plant
20 breeders and pyramiding different resistance or tolerance genes into a genotype is one way of
21 achieving this. There are numerous examples of introgression and pyramiding of favourable
22 alleles and QTLs in legumes. However, only in a few cases has MAS been used to assist in
23 gene pyramiding to overcome stresses. Most relevant work has been carried out in common
24 bean breeding for rust and anthracnose resistance (Faleiro et al. 2004). There are RAPD
25 markers linked to the 11 genes (*Ur-1* to *Ur-11*) conferring rust resistance and these markers

1 are being used to incorporate and pyramid rust resistance into common bean cultivars, and/or
2 to combine rust resistance with resistance to other diseases, such as BCMV, BGMV, common
3 bacterial blight, and/or anthracnose (Singh 2001; Stavely 2000). Similarly, molecular markers
4 linked to the majority of genes conferring anthracnose resistance (*Co-1* to *Co-10*) have been
5 described, thereby providing the opportunity to pyramid them in a resistant cultivar through
6 MAS (Kelly & Vallejo 2004). In the quest for resistant cultivars to multiple stresses,
7 combining several biotechnological approaches such as transgenesis or mutagenesis and
8 MAS, to pyramid multiple resistance genes appears as a powerful strategy. Such an approach
9 was recently achieved in soybean to manage insect resistance, resulting in the enhancement of
10 resistance levels to corn earworm (*Helicoverpa zea*) and soybean looper (*Pseudoplusia*
11 *includens*) in eight soybean lines in which two major insect-resistance QTLs and a synthetic
12 Bt gene (*cryIAc*) were combined (Walker et al. 2004).

13 The general knowledge of abiotic stress QTLs in legume is still at an early stage so
14 that gene pyramiding has not been applied yet. Nevertheless, advances achieved in non-
15 legume crops such as tomato, in which many salt stress tolerance QTL have been identified
16 and validated, open the possibility to transfer all of them to obtain a single improved cultivar
17 (Foolad 2004).

18 Thus, legume cultivars having appropriate combinations of resistance and/or tolerance
19 genes to biotic and abiotic stresses, achieved through gene pyramiding, could provide durable
20 resistance, and MAS can be a valuable tool to guide and identify the pyramiding of these
21 genes. Nevertheless, it is important to validate the results with resistance or tolerance tests,
22 due to the possibility of gene mutations, background effects, recombinants and adverse
23 interactions among resistance genes that can occur during breeding programs and influence
24 the expected phenotype. In addition, combining molecular markers with other technologies
25 may improve the efficiency of MAS. Recently, the combination of MAS with biolistic

transformation was used in rice to achieve multiple resistance against blast and bacterial blight disease (Narayanan et al. 2004). Moreover, the use of the information generated by gene expression experiments may help to improve MAS (Figure 1). Gene expression analysis helps to increase the understanding of the molecular basis of stress resistance in plants. Generation of markers based on genes with altered expression patterns in response to stresses, could result in more effective and targeted MAS. Some of these genes are described in the gene expression section and may be good candidates for futures MAS studies in legumes.

Tissue culture

In grain legumes, tissue culture has been repeatedly described as difficult. Regeneration from both organogenesis and embryogenesis has been recalcitrant in this plant group (Anand et al. 2001; Chandra & Pental 2003). This recalcitrance towards *in vitro* regeneration is a major constraint in transgenic plant production for many legumes, since advances in molecular genetics, e.g. gene over-expression, gene suppression, promoter analysis and T-DNA tagging, require efficient transformation systems (Somers et al. 2003). Efficient tissue culture is therefore a vital step, required for both the validation and exploitation of data generated by these powerful molecular tools. Implementation of robust protocols for regeneration is therefore a necessary condition for both genetic transformation and other tissue-culture derived techniques to generate genetic diversity such as somaclonal variation, *in vitro* mutagenesis, doubled haploids culture, and wide hybridization.

Somaclonal variation and in vitro mutagenesis

A little explored strategy for legume breeding is the capacity of tissue culture to generate genetic variations. Tissue culture generates a wide range of genetic variation in plants, which can be incorporated in plant breeding programmes (Jain 2001). It is well-known

1 that somaclonal variation involving callus cultivation and somatic embryogenesis has the
2 capacity to generate genetic variation (Larkin & Scowcroft 1981). The possibility of
3 producing agronomically-useful somaclones via organogenesis and somatic embryogenesis
4 has already been reported in pea (Griga et al. 1995) and pigeonpea (Chintapalli et al. 1997).
5 These variations are not desirable for some applications such as genetic transformation or
6 massive micropropagation, but can be useful for breeding. These techniques, separately or
7 combined with chemical or physical mutagenesis, generate diversity, which is a major
8 breeding goal.

9 *In vitro* mutagenesis strategies such as treatment with Ethyl-Methane-Sulphonate
10 (EMS), fast neutron radiation and insertional mutagenesis have been applied in plant
11 breeding. These methods induce point mutations, deletions, or insertions, respectively and
12 have been useful in breeding for biotic (Bhagwat & Duncan 1998; Kowalski & Cassells 1999)
13 and abiotic (Fuller & Eed 2003; Khan et al. 2001) stress in non-legume crops. In legumes
14 most effort has occurred with nitrogen fixation (Sagan et al. 1994), and mutants with
15 resistance or tolerance to stresses have not been described. Efforts in this area have been
16 hampered by the recalcitrance of legumes to regeneration and the low efficiency of finding
17 the desired phenotypes. Nevertheless, the improvement of regeneration protocols for many
18 legumes and the performance of induced mutant crop cultivars indicate that *in vitro*
19 mutagenesis can play an important role in legume breeding. Indeed, combining mutagenesis
20 techniques with MAS through TILLING as described below will make mutagenesis more
21 attractive and applicable for legume improvement. The major difficulty with these techniques
22 is the high quantity of individuals required to find the desired trait. Nevertheless, by using *in*
23 *vitro* selection systems this disadvantage can be minimized.

24
25 *In vitro selection*

1 *In vitro* selection has been used for both biotic and abiotic stress. The best studied
2 biotic stresses have been fungal diseases, using toxins or filtrate culture as selective agents
3 (Svabova & Lebeda 2005). *In vitro* selection resulted in the isolation of resistant lines in
4 carnation to *Fusarium oxysporum* f.sp *dianthi* (Thakur et al. 2002), in strawberry to
5 *Alternaria alternata* (Takahashi et al. 1992), and in wheat to *Fusarium graminearum* (Ahmed
6 et al. 1996). Salinity is the main abiotic stress that has been addressed by *in vitro* selection
7 (Flowers 2004; Zair et al. 2003), although applications to other stresses, such as zinc
8 tolerance, have also been reported (Samantaray et al. 1999).

9 Currently, these techniques are considered to be an important complement to classical
10 breeding methods (Svabova & Lebeda 2005), although they have not been sufficiently
11 explored in legumes. *In vitro* selection in legumes has been carried out mainly in alfalfa
12 (*Medicago sativa*) for selection to *C. trifolii* (Cucuzza & Kao 1986), *F. oxysporum* (Cvikrova
13 et al. 1992) and *Verticillium albo-atrum* (Koike & Nanbu 1997). These studies showed the
14 feasibility of *in vitro* selection in legumes, although no resistant lines were reported. This
15 system can also be coupled to other approaches in addition to somaclonal variation. Putative
16 stress-resistant lines derived from both conventional breeding and transgenic approaches
17 could be screened using *in vitro* selection. This is particularly attractive for some abiotic
18 stresses, where appropriate screening methods are unavailable or have low efficiency.
19 Although the advantages of the recent high-throughput technologies, coupled with genetic
20 transformation, are emerging as attractive approaches, somaclonal variation and *in vitro*
21 mutagenesis following by *in vitro* selection offers an alternative way for breeding.

22 *Double haploids and wide hybridization*

23 Doubled haploid (DH) technology refers to the use of the microspore or anther
24 cultures to obtain haploid embryos. This technology offers breeders a tool for the rapid
25 cultures to obtain haploid embryos. This technology offers breeders a tool for the rapid
26

1 production of homozygous lines. These homozygous lines can be multiplied and released as
2 cultivars, or used as recombinant inbred lines for molecular mapping and/or in breeding
3 programs (Martinez et al. 2002). An efficient DH production technology can greatly reduce
4 the time and cost of cultivar development (Liu et al. 2002a) and some stress-improved non-
5 legume varieties have been produced with this technology (Qian et al. 2000). Since DH is a
6 tissue culture dependent-technique, legumes have been generally recalcitrant and as far we
7 know no commercial legume varieties have been produced using this technology. However,
8 some advances have been achieved in alfalfa (Zagorska et al. 1997), *Lupinus* spp. (Bayliss et
9 al. 2004; Ormerod & Caligari 1994) and soybean (Cardoso et al. 2004; Rodrigues et al. 2005).
10 The importance of this approach for plant breeding in Europe, has led to the COST action 851
11 “Gametic Cells and Molecular Breeding for Crop Improvement”
12 (<http://www.scri.sari.ac.uk/assoc/cost851>) led by Brian P. Forster (Scottish Crop Research
13 Institute) which includes some legumes. Successful application of DH technology to legumes
14 in the near future would be a major achievement. However as not all homozygous lines are of
15 interest, the coupling of DH and MAS technology will be more efficient to select individuals
16 carrying desirable traits.

17 Wide hybridization depends on various factors, and according to Sharma (1995) can
18 be as wide as one can make them. Stress-related characters available in wild germplasms
19 could be introgressed into economic target species through improved wide hybridization
20 techniques such as embryo rescue and protoplast fusion. Efficient embryo rescue has allowed
21 the production of interspecific hybrids in legumes such as faba bean (*Vicia faba* x *V.*
22 *narbonensis*; Lazaridou et al. 1993), grass pea (*Lathyrus odoratus* x *L. belinensis*; Hammett et
23 al. 1994), and pigeonpea (*Cajanus cajan* x *C. platycarpus*; Mallikarjuna & Moss 1995). Stress
24 responses have not been assessed in these hybrids. However, the potential of this technique
25 has been demonstrated in non-legumes crops where successful breeding for stress tolerance

has been reported (Bradshaw et al. 1997; Tonguc & Griffiths 2004). Protoplast fusion also has potential applications for crop improvement by overcoming sexual incompatibility or reproductive barriers, and by generating novel combinations of nuclear and/or cytoplasmic genomes (Liu et al. 2005). Intergeneric somatic hybrid plants between sexually incompatible legume species have been reported in alfalfa (*Medicago sativa* x *Onobrychis viciifolia* and *Medicago sativa* x *Lotus corniculatus*; Kaimori et al. 1998; Li et al. 1993) and pea (*Pisum sativum* x *Lathyrus sativus*; Durieu & Ochatt 2000). Additionally, somatic hybrids between legumes and non-legume species has been developed, for instance the hybrids generated from *Vicia faba* x *Helianthus annuus* (Schnabl et al. 1998) and *Lotus corniculatus* x *Oriza sativa* (Nakajo et al. 1994). Interestingly, some regenerated plants from the hybrid calli of the latter fusion were tolerant to low temperatures and low sunlight intensity. Despite the potential of this technique, limited efforts have been applied to overcome stresses in legumes. Nevertheless, the successful transference of resistance or tolerance achieved in non-legumes crops to biotic (Hansen & Earle 1995) and abiotic (Arumugam et al. 2002; Brewer et al. 1999; Yue et al. 2001) stresses together with advances in tissue culture in legumes, should encourage legume breeders to exploit somatic hybrids. For an excellent overview of advances on intergeneric somatic hybridization and its application to crop genetic improvement see Liu et al. (2005).

Genetic transformation

Crop improvement through genetic engineering has become a reality (Dunwell 2000). It is now possible to transform many grain legumes (Chandra & Pental 2003; Somers et al. 2003) although in some cases the rate of recovery of transgenic lines is still low. The use of *Agrobacterium tumefaciens* as a vector for legume transformation was an important breakthrough. Both micro-particle bombardment (Gulati et al. 2002; Li et al. 2004) and A.

1 *tumefaciens* (De Clercq et al. 2002; Li et al. 2004) have been used for DNA delivery into
2 either embryogenic or organogenic cultures. Transformation has been generally based on
3 infection by *A. tumefaciens*, although *A. rhizogenes* is also used for transformation of some
4 species to produce composite plants with hairy roots or hairy root cultures (Boisson-Dernier
5 et al. 2001; Stiller et al. 1997; Wu & VanEtten 2004). The inserted DNA can be either a
6 specific gene with a specific biochemical function, a regulatory gene that controls a network
7 of other genes, or multiple genes to generate long-term durable resistance. In this review we
8 will describe only examples related to biotic or abiotic stress. For a more comprehensive
9 review about transformation and gene technology in legumes, other references should be
10 consulted (Chandra & Pental 2003; Popelka et al. 2004; Somers et al. 2003).

11 A number of legume cultivars have been transformed in order to enhance the
12 resistance to biotic stresses. Resistance to insects using *Bacillus thuringiensis* genes (Walker
13 et al. 2000) and viruses using pathogen-derived resistance (Aragão et al. 2002), along with the
14 introduction of constitutively expressed genes encoding pathogenesis-related (PR) proteins or
15 phytoalexins (He & Dixon 2000; Samac et al. 2004) have been reported in legumes (Table 3).

16 Abiotic stresses generally involve perturbation of various cellular functions and
17 activation of complex metabolic pathways, and are conferred by polygenic traits (Kassem et
18 al. 2004; Lee et al. 2004; Popelka et al. 2004). This complexity together with the lack of good
19 sources of natural tolerance makes this an area that is not readily amenable for conventional
20 breeding strategies. In plants there is a poor understanding of most abiotic stress responses.
21 Thus, the successful use of genetic transformation requires a better physiological and
22 molecular understanding of these stresses. Recent progress achieved in non-legume plants
23 supports the potential use of transgenic approaches to produce tolerant lines (Jiang et al. 2004;
24 Kasuga et al. 1999; Shou et al. 2004; Sivamani et al. 2000; Umezawa et al. 2004). For
25 instance, the use of transgenic, mutagenic and genetic approaches strongly improved the

1 understanding of the genetic and molecular mechanisms of salinity tolerance in plant, and this
2 will help develop crops, including legumes, with improved tolerance (reviewed in Apse &
3 Blumwald 2002; Foolad 2004; Hasegawa et al. 2000). As a result, it was found that over-
4 expression of a single-gene controlling vacuolar or plasma membrane Na⁺/H⁺ antiport
5 protein, in transgenic Arabidopsis, tomato and rape seed provided them with a high level of
6 salt tolerance under greenhouse conditions (Apse et al. 1999; Shi et al. 2003; Zhang &
7 Blumwald 2001; Zhang et al. 2001). Similarly, manipulating expression of pea DNA
8 Helicase45 or the glyoxalate pathways confers high salinity tolerance in tobacco (Sanan-
9 Mishra et al. 2005; Singla-Pareek et al. 2003). Although transgenic plants are yet to be
10 examined for salt-tolerance in the field, the recent genetic advances suggest there are good
11 prospects for developing transgenic legumes with enhanced salt tolerance in the near future
12 (Foolad 2004; Sharma & Lavanya 2002). On the other hand the increase of tolerance to
13 aluminium and cyanamide toxicity in transgenic alfalfa (Morphew et al. 2004) and soybean
14 (Zhang et al. 2005) demonstrates the potential of this approach in legumes. For a more
15 exhaustive review of the application of transgenesis to overcome abiotic stresses in plants, see
16 Sharma & Lavanya (2002).

17 As described above genetic transformation is an attractive approach to overcome
18 stresses in legumes. Nevertheless, as far we know the only transgenic legumes commercially
19 used for biotic stress is the soybean line carrying the insect resistance gene *cryA* from *B.*
20 *thuringiensis* (Babu et al. 2003). This low number has been influenced by technical
21 (regeneration recalcitrance of most legumes), social (public concern issues) and political
22 (lower rate of investment in legume crops compared to other crops such as rice, wheat and
23 maize) reasons. However, the advances in legume transformation protocols and the increasing
24 interest in legumes as high protein content food should see an increase in the production of
25 genetically modified legumes.

Gene expression

As already mentioned, the efficiency of both MAS and transgenic approaches will be improved by using the information from gene expression studies. Understanding the mechanisms employed by plants to defend themselves against stresses and a more complete knowledge about the genes involved, will allow a more precise use of MAS and transgenics. Sequence information, while valuable and a necessary starting point, is insufficient to answer questions concerning gene function, regulatory networks and the biochemical pathways activated in response to stresses. To address these questions, more comprehensive approaches, including quantitative and qualitative analyses of gene expression products, are necessary at the transcriptomic, proteomic, and metabolomic levels.

Transcriptomics

An important step in the control of stress responses in plants is the transcriptional activation or repression of genes (Chen et al. 2002). Thus, identification of differentially expressed genes is particularly important to understand stresses response in plants. To achieve this objective, tools such as microarrays (Schena et al. 1995), Suppression Subtractive Hybridization library (Diatchenko et al. 1996), Serial Analyse of Gene Expression (Velculescu et al. 1995) and quantitative measurement of transcription factor (TF) expression patterns have been developed in addition to older techniques such as Northern blotting.

In legumes, gene expression patterns following biotic stresses have been more extensively studied than those following abiotic stresses. Large-scale analyses of gene expression patterns in response to pathogens have revealed the differential expression of large numbers of genes. Known defence gene families are usually expressed differentially in these studies independent of the specific legume-pathogen interaction being investigated. Among these genes, phytoalexins such as medicarpin (Blount et al. 1992; He & Dixon 2000),

1 PR-proteins including PR-10, chitinases, glucanases (Salles et al. 2002) and lipoxigenases
2 (*Lox* genes) have been frequently detected (Cho & Muehlbauer 2004; Torregrosa et al. 2004).

3 The coupling of these powerful large-scale gene expression profiling methods with
4 recombinant inbred lines or near isogenic lines in legumes that differ in
5 susceptibility/resistance to key pathogen and pests will greatly facilitate a more
6 comprehensive understanding of genes involved in the defence response of legumes to
7 specific biotic stresses. For more details on legume plant-pathogen interactions or legume-
8 pest interactions, see Torregrosa et al. 2005 and Edwards & Singh 2005, respectively, in this
9 issue.

10 In respect to abiotic stress, gene expression analyses have been mainly based on
11 studies with cloned genes (Singh et al. 2004). Other work has shown that transcriptomic tools
12 are also a good option for legume breeding to environmental stresses. Using a modified
13 c-DNA-AFLP technique in soybean, 140 differentially expressed c-DNA fragments were
14 obtained by comparing control and iso-osmotic treated plants. Some of the responsive genes
15 encoded ion transporters, transcription factors (TFs) and redox enzymes (Umezawa et al.
16 2002). Suppression subtractive hybridization screening was carried out in *Retama raetam*, a
17 C-3 drought tolerant legume. This study revealed that dormancy, key to the survival of many
18 species in arid environments, was followed by accumulation of transcripts encoding a PR-10
19 like protein, a low temperature-inducible dehydrin and a WRKY transcription factor (Pnueli
20 et al. 2002). Similarly, 43 drought-responsive mRNA transcripts were reported to be
21 differentially expressed in peanut following water stress (Jain et al. 2001).

22 Much more extensive expression studies have been performed in Arabidopsis, and the
23 resulting knowledge can also be used in legumes through comparative genomics. For
24 example, Ishitani et al. (2004), selected 100-200 genes from the Arabidopsis database, and
25 showed that at least three *DREB*-like genes, thought to be key transcriptional regulators of

drought and/or cold tolerance were present in common bean. Many other expression studies in Arabidopsis have highlighted the involvement of TFs in stress responses and this has encouraged researchers working with other plants to focus on these proteins.

Transcription factors

Transcription factors are proteins that play an important role in controlling the expression of genes in most biochemical pathways including the response to stress (Eulgem 2005; Kasuga et al. 1999). Genomics studies over the last few years have identified numerous TFs (mainly in Arabidopsis) and revealed a high degree of complexity and overlap in the transcriptional regulation of gene expression in response to many stresses (Shinozaki & Yamaguchi-Shinozaki 2000). The understanding of the role of TF may open new avenues for improving resistance or tolerance to stresses (Singh et al. 2002). While large-scale analysis of TFs have been primarily done in Arabidopsis (see review by Eulgem, 2005), a platform for the study of TFs in *M. truncatula* has been developed by Dr. M. Udvardi (Max Planck Institute for Plant Physiology, Golm, Germany).

An interesting family of TFs for stress responses in plants is the ethylene-responsive-element-binding factors (ERF) of which over 60 members have been described in *M. truncatula* (Anderson et al. 2005); and the closely related DREB/CREB proteins (Singh et al. 2002; Yamaguchi-Shinozaki & Shinozaki 2005). Members of ERF family are responsive to cold, drought, pathogen infection, or wounding. The WRKY family, involved in the regulation of plant stress-response genes such as receptor protein kinases (Asai et al. 2002; Robatzek & Somssich 2002), and bZIP family members that regulate PR-1 and Glutathione S-Transferase genes (Chen & Singh 1999; Fan & Dong 2002; Lebel et al. 1998), or cold/dehydration genes (Yamaguchi-Shinozaki & Shinozaki 2005) are other important stress-

1 responsive TF families. A Krüppel-like TF (*Mtzip2-1*) involved in salt tolerance has also been
2 described in *M. truncatula* (Merchan et al. 2003).

3 A given TF can mediate the response to various stresses (Eulgem 2005; Yamaguchi-
4 Shinozaki & Shinozaki 2005). This characteristic makes the TFs especially attractive for
5 genetic transformation, because a single TF gene can result in resistance or tolerance to
6 various stresses. Following this principle, over-expression of a TF that regulates an ABA-
7 responsive gene conferred multiple stress tolerance in rice (Kim et al. 2004). However,
8 different TFs are also known to respond to the same stress with different but overlapping
9 kinetics (Onate-Sanchez & Singh 2002). On the other hand, attempts to knockout specific TFs
10 have often not resulted in any obvious phenotypes, perhaps due to overlapping function. Thus,
11 use of TFs for genetic improvement requires a comprehensive knowledge of their biological
12 functions.

14 *Proteomics*

15 In parallel to the accumulation of a wealth of genomic and transcriptomic data, recent
16 technological developments have allowed the establishment of valuable methods for
17 quantitative and qualitative protein profiling (Cánovas et al. 2004). This approach is very
18 important in evaluating stress-responses, because mRNA levels do not always correlate with
19 protein accumulation (Gygi et al. 1999). Indeed, large differences in protein turnover and
20 post-translational modifications may lead to large variations between transcriptomic and
21 proteomic data. Thus, protein studies are needed to provide information on their levels and
22 activities (Zivy & de Vienne 2000). To this purpose, proteomic-based techniques that allow
23 large-scale protein profiling are powerful tools for the identification of proteins involved in
24 stress-responses in plants (Gygi & Aebersold 2000).

1 Extensive studies have evaluated changes in protein levels in plant tissues in response
2 to stresses (Cánovas et al. 2004; Kim et al. 2003). Unfortunately, these studies have been
3 mainly focused on non-legume species, such as *Arabidopsis* and rice (Cánovas et al. 2004),
4 and only recently have been enlarged to include some legumes (Jorrín et al. 2005, this issue).
5 As a result only a handful of studies have been carried out in legumes, although in the next
6 few years there should be a significant increase in the number of legume species and stresses
7 analysed. So far, pea has been more intensively studied, with the analysis of induced protein
8 expression in roots in response to salt (Kav et al. 2004), to cadmium stress (Repetto et al.
9 2003) and to infection by the parasitic plant *Orobancha crenata* (Castillejo et al. 2004). In
10 addition, proteomic approaches have been applied to *M. truncatula*, lentils, lupin, common
11 bean, cowpea and soybean to identify proteins involved in the response to different stresses
12 (Colditz et al. 2004; Fecht-Christoffers et al. 2003; Kav et al. 2004; Mithofer et al. 2002;
13 Pinheiro et al. 2005; Repetto et al. 2003). Interestingly, many of the induced proteins from
14 these different stresses were common or belonged to overlapping pathways. For example,
15 members of the PR-10, the phytoalexin biosynthesis enzyme and the peroxidase families were
16 identified in different studies. These observations highlight a potential role of these genes for
17 resistance or tolerance to stress in legumes. Although further studies are needed to determine
18 their exact function in stress-response, these genes could be promising candidates for genetic
19 transformation and/or MAS approaches (Figure 1).

21 *Metabolomics*

22 Transcriptomic and proteomic data are important steps in deciphering a complex
23 biological process, but they are still insufficient to understand them fully since most
24 biological processes are ultimately mediated by cell metabolites. Alternative mRNA splicing,
25 protein turn-over rates and post-translational modifications that modulate protein activity

1 imply that changes in the transcriptome or proteome do not always correspond to alterations
2 in the cell metabolome (Sumner et al. 2003). Therefore, the only way to the complete
3 understanding of both gene function and molecular events controlling complex plant
4 processes is to analyse in parallel the transcriptome, the proteome and the metabolome in an
5 integrative manner (Dixon 2001). In legumes, this kind of approach has been taken in *M.*
6 *truncatula* suspension cells to various stimuli (Bell et al. 2001), and with the characterization
7 of metabolic changes during the nitrogen-fixing symbiotic interaction in *L. japonicus*
8 (Colebatch et al. 2004; Desbrosses et al. 2005; Rispaill 2005).

9 Although large-scale, comprehensive metabolomic studies are difficult, a number of
10 targeted analyses have been performed to assess the involvement of subsets of metabolites in
11 various stresses. Most studies on plant stress responses focused on flavonoid and isoflavonoid
12 phytoalexins. Accumulation of medicarpin, pisatin, glyceolin or sativan has been frequently
13 observed in response to pathogen infection and elicitor treatment in alfalfa, pea, soybean and
14 *L. japonicus* respectively (Baldrige et al. 1998; BorejszaWysocki et al. 1997; Lozovaya et al.
15 2004; Saunders & O'Neill 2004; Shimada et al. 2000). Interestingly, phytoalexin
16 accumulation was also observed after copper or mercury stress (Mithofer et al. 2004). The
17 involvement of these compounds as key defence metabolites has been nicely proven by the
18 modification of resistance levels to *Phoma medicaginis* and *Nectria haematococca* in alfalfa
19 and pea cultivars respectively (He & Dixon 2000; Wu & VanEtten 2004).

20 Other metabolite classes have been described as potential defence or signal molecules.
21 For instance, *L. japonicus* leaves have been shown to emit many volatile terpenoids in
22 response to spider mites. The exact roles of these molecules are still unclear, but they are
23 believed to play a role as attractants for natural predators of herbivorous insects and as
24 systemic signal defence inducers for neighbouring plants (Ozawa et al. 2000). The possibility
25 to attract predatory mites by modulation of sesquiterpene synthesis through transgenesis as

1 recently demonstrated in *Arabidopsis* may have important repercussions in alternative insect-
2 damage protection (Kappers et al. 2005). The triterpene saponins also appear to be important
3 in defence reactions as they accumulate in response to insect attacks in alfalfa (Agrell et al.
4 2004) and are well-known for their allelopathic, antimicrobial and anti-insect activities
5 (Dixon & Sumner 2003). These targeted studies highlight the importance of secondary
6 metabolites in stress resistance in legumes.

7 While the preliminary results from combining metabolic approaches with transgenics
8 indicates the potential of increasing intrinsic stress resistance levels in legume crops and
9 strengthens the potential role of biotechnology in crop improvement (He & Dixon 2000; Wu
10 & VanEtten 2004), it must be remembered that most metabolic pathways are interconnected
11 in highly complex networks. Thus, modulating one metabolic pathway may have negative
12 impacts on another, leading to concomitant deleterious traits in the modified crop. Large-scale
13 metabolic analyses are therefore necessary to observe the metabolic networks important for
14 plant growth and development under a range of environmental conditions.

15 Large-scale analysis by using different “omics” technologies are providing extensive
16 data sets that will help identify potential candidate genes for an increase in intrinsic resistance
17 and/or tolerance levels in important legume crops. Identification of these candidate genes,
18 may allow their direct application in crop improvement through MAS, or genetic engineering.
19 However, in most cases, the roles of these candidate genes remain unknown and it will be
20 important to carry out functional studies as a preliminary step towards their use in genetic
21 improvement.

22 23 **Functional analysis**

24 To date the completion of the *Arabidopsis* and rice genomes have been achieved and
25 the genome of some legumes (*M. truncatula*, *L. japonicus*) together with other plant

1 sequencing projects is underway. The traditional pursuit of a gene starting with a phenotype
2 (forward genetics), has given way to the opposite situation where the gene sequences are
3 known but not their functions. The challenge is now to decipher the function of the thousands
4 of genes identified by genome projects, and reverse genetics methodologies are key tools in
5 this endeavour (Gilchrist & Haughn 2005).

6 The ability to knockout genes or suppress their expression are powerful tools to
7 determine the function of a gene. This can be done by anti-sense RNA suppression, targeted
8 gene replacement, insertional mutagenesis, gene silencing and Targeted Induced Local Lesion
9 In Genome (TILLING) approaches. Anti-sense RNA suppression requires considerable effort
10 for any given target gene before even knowing whether it will be successful (McCallum et al.
11 2000) and targeted gene replacement i.e. via homologous recombination has not yet been
12 reproducibly achieved for higher plants.

13 Collections of random T-DNA (over 225,000 independent *Agrobacterium* T-DNA
14 insertions) or transposable element insertion mutants are currently available for the
15 Arabidopsis community (Alonso 2003). While such a collection does not exist yet in legumes,
16 insertional mutagenesis has been successfully used. For example, in *L. japonicus*
17 identification of regulatory components of nodule induction have been achieved by
18 characterizing a transposon-tagged mutant, *nin*, arrested at the stage of bacterial recognition
19 (Schauser et al. 1999). However, although collections of T-DNA mutants may be very useful,
20 they produce a limited range of allele types and do not always produce null alleles (Rispaill
21 2005; Webb et al. 2000). Recently the use of the tobacco retrotransposon Tnt1 has been
22 successfully applied for large-scale insertional mutagenesis in *M. truncatula* and promises to
23 be a useful tool for functional genomic (Tadege et al. 2005).

24 The term RNA silencing has been adopted to describe phenomena such as post-
25 transcriptional gene silencing in plants, quelling in fungi and RNA interference in animals

(Baulcombe 2004). Researchers have developed different RNA silencing strategies as tools for selective knockout of targeted genes. Despite the successes of this technique in several species, RNA silencing has several drawbacks, i.e. phenotypic instability in later generations (Hannon 2002) and the requirement for a reliable plant transformation system. RNA silencing is believed to be a natural plant defence against viruses. Following this principle, another technique, Virus-Induced Gene Silencing (VIGS), has been developed to suppress plant gene expression through infection with virus vectors that harbour a target region of the host gene (Baulcombe 2004; Britt & May 2003). There are vectors with the ability to support VIGS in plants (Dalmay et al. 2000; Liu et al. 2002b). Nevertheless, in legumes, VIGS has not been extensively used. Interestingly, there are recent reports that a VIGS vector based on the Pea Early Browning Virus (PEBV) can also be used successfully in legume species (Constantin et al. 2004; Van den Boogaart et al. 2004). These findings together with the advances in legume genomic should increase the use of VIGS as a functional genomic tool in the near future.

The limitations of RNA silencing or insertational mutagenesis, previously stated, can be overcome by TILLING. This technique combines chemical mutagenesis with a powerful screening method for potential mutations (Gilchrist & Haughn 2005; Henikoff et al. 2004; McCallum et al. 2000). The generation of phenotypic variants without introducing foreign DNA in the plant makes TILLING very suitable not only for functional analysis, but also for agricultural applications. The TILLING facility for a *L. japonicus* collection of mutants is available from the Parniske group at the John Innes Centre (Perry et al. 2003). This facility has been validated by identification of 15 homozygous mutants representing six different alleles of SYMRK, an important symbiotic gene (Perry et al. 2003). A database comprising information on individual mutant plants in their collection is also accessible at <http://www.lotusjaponicus.org/finder.htm>. In *M. truncatula*, TILLING programs have been set up by Professor D. Cook (U.C. Davis, USA) and by Dr. A. Kondorosi (CNRS, Gif-Sur-

Yvette, France). The advantages of TILLING are resulting in private companies, such as Anawah (<http://anawah.com/programs>), extending TILLING facilities to a wider variety of organisms including soybean and peanuts. The diversity of species for which this technique will be available, opens up new possibilities for legume researchers both for the functional analysis of genes previously identified by the “omic” technologies (Figure 1), as well as the generations of new varieties.

Conclusions

Over the years biotechnology has emerged as a promising tool to overcome stresses in plants, but to date, progress has been limited in legumes. The current advances in tissue-derived techniques, genetic transformation and MAS, together with the advances in powerful new ‘omic’ technologies offer great potential to improve this situation. Indeed, it is now possible to target almost all legume crops with a variety of biotechnological approaches for genetic improvement. As such, the more efficient regeneration protocols recently established for many legumes should encourage legume researchers to resume the use of techniques such as DH, wide hybridization and mutagenesis in breeding programmes. On the other hand, crops without appropriate regeneration protocols may also be improved by mutagenesis through TILLING. It is important to provide breeders with the broadest variety of biotechnology tools as possible since each stress-crop case has its own particularities and so would need one or a combination of specific biotechnological approach(es) to tackle them efficiently. Strategies such as resistance gene pyramiding assisted by MAS could be useful to overcome resistance breakdown by new races of *U. appendiculatus* in common bean, while pathogen-derived genes and heterologous expression of PR-genes may be a better approach to enhance resistance to viruses and polyphagous fungi, respectively.

1 Although the advances in biotechnology greatly facilitate legume improvement, a
2 more comprehensive knowledge of resistance or tolerance mechanisms is required to direct
3 breeding. Indeed, only a better understanding of the underlying mechanisms activated in
4 response to stresses will allow an efficient application of biotechnology in sustainable
5 agriculture. The advent of the ‘omic’ technologies together with the functional genomic tools
6 is a promising approach to achieve this. Most advances in these fields have been performed in
7 Arabidopsis which has provided us with a growing understanding of important stress
8 pathways. Nevertheless, legumes offer a number of attractive features in their own right that
9 are drawing researchers interested in abiotic and biotic stress responses. For example, they
10 provide an excellent system to analyse how plants distinguish between friend and foe. They
11 also have advantages over some of the other plant model species for specific stresses. For
12 example, *M. truncatula* has good genetic resistance to aphids (Klingler et al. 2005),
13 something which is lacking in Arabidopsis. Thus it is important to better characterise legume
14 responses to stress. The establishment of the model legumes *M. truncatula* and *L. japonicus* is
15 starting to provide applicable information for legumes. The integration of knowledge
16 generated by the different approaches described here, should lead to more accurate and
17 efficient breeding of key legume crops. In the case of genetic engineering, this would not only
18 allow the targeting of transgene expression to particular conditions (e.g using stress-
19 responsive or tissue-specific promoters), but also monitoring the effect of the transgene (e.g
20 by proteomic and metabolomic approaches). Additionally, researchers dealing with others
21 strategies such as MAS or even classical breeding will be able to take advantage of the results
22 being gathered from “omic” technologies. However, the delivery of “omic” information
23 should be done in a friendlier mode for plant breeders in order to facilitate its efficient
24 application in genetic improvement. Overall, for biotechnology to fulfill its potential for
25 legume breeding there needs to be good and regular communication between classical

1 breeders and biotechnologists to firstly, make sure that the tools of biotechnology are applied
2 to the most pressing and appropriate problems and secondly, to ensure that pathways for
3 delivery/uptake into breeding programs are in place.

4 5 **Acknowledgment**

6 We thank to Fundación Carolina for M. A. Dita's scholarship. E. Prats is funded by an
7 European Marie Curie Reintegration Grant, N. Risipail by (FP5) Eufaba project. Our work in
8 this area is supported by Spanish CICYT project AGL-2002-03248 and European Union
9 project FP6-2002-FOOD-1-506223. K. Singh's work in this area is supported in part by the
10 Grains Research and Development Corporation (GRDC) and the Department of Education,
11 Science and Training (DEST) in Australia. We also apologize for not being able to include all
12 the important references because of length limitations.

References

- Agrell, J., P. Anderson, W. Oleszek, A. Stochmal & C. Agrell, 2004. Combined effects of elevated CO₂ and herbivore damage on alfalfa and cotton. *J Chem Ecol* 30: 2309-2324.
- Ahmed, H.U., C.C. Mundt, M.E. Hoffer & S.M. Coakley, 1996. Selective influence of wheat cultivars on pathogenicity of *Mycosphaerella graminicola* (Anamorph *Septoria tritici*). *Phytopathology* 86: 454-458.
- Alonso, J.M., 2003. Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301: 1849-1849.
- Anand, R.P., A. Ganapathi, G. Vengadesan, N. Selvaraj, V.R. Anbazhagan & S. Kulothungan, 2001. Plant regeneration from immature cotyledon-derived callus of *Vigna unguiculata* (L.) Walp (cowpea). *Curr Sci* 80: 671-674.
- Anderson, J.P., L.F. Thatcher & K.B. Singh, 2005. Plant defence responses: conservation between models and crops. *Funct Plant Biol* 32: 21-34.
- Anderson, K.L., L. Buchwaldt, G. Chongo, B.D. Gossen, R.A.A. Morrall, & P.G. Pearce, 2000a. Diseases of lentil in Saskatchewan 1999. *Can Plant Dis Surv* 80: 96-98.
- Apse, M.P. & E. Blumwald, 2002. Engineering salt tolerance in plants. *Curr Opin Biotechnol* 13: 146-150.
- Apse, M.P., G.S. Aharon, W.A. Snedden & E. Blumwald, 1999. Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiport in *Arabidopsis*. *Science* 285: 1256-1258.
- Aragão, F.J.L., G.R. Vianna, M.M.C. Albino & E.L. Rech, 2002. Transgenic dry bean tolerant to the herbicide glufosinate ammonium. *Crop Sci* 42: 1298-1302.
- Aragão, F.J.L., S.G. Ribeiro, L.M.G. Barros, A.C.M. Brasileiro, D.P. Maxwell, E.L. Rech & J.C. Faria, 1998. Transgenic beans (*Phaseolus vulgaris* L.) engineered to express viral antisense RNAs show delayed and attenuated symptoms to bean golden mosaic geminivirus. *Mol Breed* 4: 491-499.

1 Arrus, K., G. Blank, D. Abramson, R. Clear & R.A. Holley, 2005. Aflatoxin production by
2 *Aspergillus flavus* in Brazil nuts. J Stored Prod Res 41: 513-527.

3 Arumugam, N., A. Mukhopadhyay, V. Gupta, Y.S. Sodhi, J.K. Verma, D. Pental & A.K.
4 Pradhan, 2002. Synthesis of somatic hybrids (RCBB) by fusing heat-tolerant *Raphanus*
5 *sativus* (RR) and *Brassica oleracea* (CC) with *Brassica nigra* (BB). Plant Breed 121: 168-
6 170.

7 Asai, T., G. Tena, J. Plotnikova, M.R. Willmann, W.L. Chiu, L. Gomez-Gomez, T. Boller,
8 F.M. Ausubel & J. Sheen, 2002. MAP kinase signalling cascade in Arabidopsis innate
9 immunity. Nature 415: 977-983.

10 Asamizu, E., Y. Nakamura, S. Sato & S. Tabata, 2004. Characteristics of the *Lotus japonicus*
11 gene repertoire deduced from large-scale expressed sequence tag (EST) analysis. Plant
12 Mol Biol 54: 405-414.

13 Avila, C.M., J.C. Sillero, D. Rubiales, M.T. Moreno & A.M. Torres, 2003. Identification of
14 RAPD markers linked to the *Uvf-1* gene conferring hypersensitive resistance against rust
15 (*Uromyces viciae-fabae*) in *Vicia faba* L. Theor Appl Genet 107: 353-358.

16 Babu, R.M., A. Sajeena, K. Seetharaman & M.S. Reddy, 2003. Advances in genetically
17 engineered (transgenic) plants in pest management - an over view. Crop Prot 22: 1071-
18 1086.

19 Baldrige, G.D., N.R. O'Neill & D.A. Samac, 1998. Alfalfa (*Medicago sativa* L.) resistance to
20 the root-lesion nematode, *Pratylenchus penetrans*: defense-response gene mRNA and
21 isoflavonoid phytoalexin levels in roots. Plant Mol Biol 38: 999-1010.

22 Baulcombe, D., 2004. RNA silencing in plants. Nature 431: 356-363.

23 Bayaa, B. & W. Erskine, 1998. Diseases of lentil. In: D.J. Allen & L.M. Lenné (Eds.), The
24 Pathology of Food and Pasture Legumes, pp 423-472. Wallingford, UK.

1 Bayaa, B., W. Erskine & M. Singh, 1997. Screening lentil for resistance to fusarium wilt:
2 methodology and sources of resistance. *Euphytica* 98: 69-74.

3 Bayliss, K.L., J.M. Wroth & W.A. Cowling, 2004. Pro-embryos of *Lupinus* spp. produced
4 from isolated microspore culture. *Aust J Agric Res* 55: 589-593.

5 Bell, C.J., R.A. Dixon, A.D. Farmer, R. Flores, J. Inman, R.A. Gonzales, M.J. Harrison, N.L.
6 Paiva, A.D. Scott, J.W. Weller & G.D. May, 2001. The Medicago Genome Initiative: a
7 model legume database. *Nucleic Acids Res* 29: 114-117.

8 Bhagwat, B. & E.J. Duncan, 1998. Mutation breeding of banana cv. Highgate (*Musa* spp.,
9 AAA Group) for tolerance to *Fusarium oxysporum* f. sp. *cubense* using chemical
10 mutagens. *Sci Hortic* 73: 11-22.

11 Blount, J.W., R.A. Dixon & N.L. Paiva, 1992. Stress responses in alfalfa (*Medicago sativa*
12 L). Antifungal activity of medicarpin and its biosynthetic precursors - Implications for the
13 genetic manipulation of stress metabolites. *Physiol Mol Plant Pathol* 41: 333-349.

14 Boisson-Dernier, A., M. Chabaud, F. Garcia, G. Becard, C. Rosenberg & D.G. Barker, 2001.
15 *Agrobacterium rhizogenes*-transformed roots of *Medicago truncatula* for the study of
16 nitrogen-fixing and endomycorrhizal symbiotic associations. *Mol Plant-Microbe Interact*
17 14: 695-700.

18 BorejszaWysocki, W., E. BorejszaWysocka & G. Hrazdina, 1997. Pisatin metabolism in pea
19 (*Pisum sativum* L) cell suspension cultures. *Plant Cell Reports* 16: 304-309.

20 Boukar, O., L. Kong, B.B. Singh, L. Murdock & H.W. Ohm, 2004. AFLP and AFLP-derived
21 SCAR markers associated with *Striga gesnerioides* resistance in cowpea. *Crop Sci* 44:
22 1259-1264.

23 Bradshaw, J.E., D.J. Gemmell & R.N. Wilson, 1997. Transfer of resistance to clubroot
24 (*Plasmodiophora brassicae*) to swedes (*Brassica napus* L var *napobrassica* Peterm) from
25 *B. rapa*. *Ann Appl Biol* 130: 337-348.

1 Bretag T.W. & M. Ramsey, 2001. Foliar diseases caused by fungi: *Ascochyta* spp. p. 24-28.
2 In Compendium of pea disease and pests, second edition. Kraft & Pflieger ed. APS Press.
3 Brewer, E.P., J.A. Saunders, J.S. Angle, R.L. Chaney & M.S. McIntosh, 1999. Somatic
4 hybridization between the zinc accumulator *Thlaspi caerulescens* and *Brassica napus*.
5 Theor Appl Genet 99: 761-771.
6 Britt, A.B. & G.D. May, 2003. Re-engineering plant gene targeting. Trends Plant Sci 8: 90-
7 95.
8 Cánovas, F., E. Dumas-Gaudot, G. Recorbet, J. Jorin, H.-P. Mock & M. Rossignol, 2004.
9 Plant proteome analysis. Proteomics 4: 285-298.
10 Cardoso, M.B., E. Kaltchuk-Santos, E.C. de Mundstock & M.H. Bodanese-Zanettini, 2004.
11 Initial segmentation patterns of microspores and pollen viability in soybean cultured
12 anthers: Indication of chromosome doubling. Braz Arch Biol Technol 47: 703-712.
13 Carmona, M.A., M.E. Gally & S.E. Lopez, 2005. Asian soybean rust: Incidence, severity, and
14 morphological characterization of *Phakopsora pachyrhizi* (uredinia and telia) in
15 Argentina. Plant Dis 89: 109-109.
16 Castillejo, M.A., N. Amieur, E. Dumas-Gaudot, D. Rubiales & J.V. Jorin, 2004. A proteomic
17 approach to studying plant response to crenate broomrape (*Orobancha crenata*) in pea
18 (*Pisum sativum*). Phytochemistry 65: 1817-1828.
19 Chakraborty, U., B. Sarkar & B.N. Chakraborty, 2003. Protection of soybean rot by
20 *Bradyrhizobium japonicum* and *Trichoderma harzianum* associated with changes in
21 enzyme activities and phytoalexin production. J Mycol Plant Pathol 33: 21-25.
22 Chandra, A. & D. Pental, 2003. Regeneration and genetic transformation of grain legumes:
23 An overview. Curr Sci 84: 381-387.

1 Chen, W.Q. & K.B. Singh, 1999. The auxin, hydrogen peroxide and salicylic acid induced
2 expression of the Arabidopsis GST6 promoter is mediated in part by an ocs element. Plant
3 J 19: 667-677.

4 Chen, W.Q., N.J. Provart, J. Glazebrook, F. Katagiri, H.S. Chang, T. Eulgem, F. Mauch, S.
5 Luan, G.Z. Zou, S.A. Whitham, P.R. Budworth, Y. Tao, Z.Y. Xie, X. Chen, S. Lam, J.A.
6 Kreps, J.F. Harper, A. Si-Ammour, B. Mauch-Mani, M. Heinlein, K. Kobayashi, T. Hohn,
7 J.L. Dangel, X. Wang & T. Zhu, 2002. Expression profile matrix of Arabidopsis
8 transcription factor genes suggests their putative functions in response to environmental
9 stresses. Plant Cell 14: 559-574.

10 Chimphango, S.B.M., C.F. Musil & F.D. Dakora, 2003. Effects of UV-B radiation on plant
11 growth, symbiotic function and concentration of metabolites in three tropical grain
12 legumes. Funct Plant Biol 30: 309-318.

13 Chintapalli, P.L., J.P. Moss, K.K. Sharma & J.K. Bhalla, 1997. *In vitro* culture provides
14 additional variation for pigeonpea *Cajanus cajan* (L) Millsp crop improvement. In Vitro
15 Cell Dev Biol-Plant 33: 30-37.

16 Cho, S.H. & F.J. Muehlbauer, 2004. Genetic effect of differentially regulated fungal response
17 genes on resistance to necrotrophic fungal pathogens in chickpea (*Cicer arietinum* L.).
18 Physiol Mol Plant Pathol 64: 57-66.

19 Cho, S.H., W.D. Chen & F.J. Muehlbauer, 2004. Pathotype-specific genetic factors in
20 chickpea (*Cicer arietinum* L.) for quantitative resistance to ascochyta blight. Theor Appl
21 Genet 109: 733-739.

22 Cobos, M.J., M. Fernandez, J. Rubio, M. Kharrat, M.T. Moreno, J. Gil & T. Millan, 2005. A
23 linkage map of chickpea (*Cicer arietinum* L.) based on populations from Kabuli x Desi
24 crosses: location of genes for resistance to fusarium wilt race 0. Theor Appl Genet 110:
25 1347-1353.

1 Colditz, F., O. Nyamsuren, K. Niehaus, H. Eubel, H.P. Braun & F. Krajinski, 2004. Proteomic
2 approach: Identification of *Medicago truncatula* proteins induced in roots after infection
3 with the pathogenic oomycete *Aphanomyces euteiches*. Plant Mol Biol 55: 109-120.

4 Colebatch, G., G. Desbrosses, T. Ott, L. Krusell, O. Montanari, S. Kloska, J. Kopka & M.K.
5 Udvardi, 2004. Global changes in transcription orchestrate metabolic differentiation
6 during symbiotic nitrogen fixation in *Lotus japonicus*. Plant J 39: 487-512.

7 Constantin, G.D., B.N. Krath, S.A. MacFarlane, M. Nicolaisen, I.E. Johansen & O.S. Lund,
8 2004. Virus-induced gene silencing as a tool for functional genomics in a legume species.
9 Plant J 40: 622-631.

10 Cook, D.R., 1999. *Medicago truncatula* - a model in the making! Commentary. Curr Opin
11 Plant Biol 2: 301-304.

12 Correa, O.S., A. Aranda & A.J. Barneix, 2001. Effects of pH on growth and nodulation of two
13 forage legumes. J Plant Nutr 24: 1367-1375.

14 Coyne, D.P., J.R. Steadman, G. Godoy-Lutz, R. Gilbertson, E. Arnaud-Santana, J.S. Beaver
15 & J.R. Myers, 2003. Contributions of the Bean/Cowpea CRSP to management of bean
16 diseases. Field Crop Res 82: 155-168.

17 Cucuzza, J.D. & J. Kao, 1986. Invitro assay of excised cotyledons of alfalfa (*Medicago*
18 *sativa*) to screen for resistance to *Colletotrichum trifolii*. Plant Dis 70: 111-115.

19 Cvikrova, M., P. Binarova, J. Eder & J. Nedelnik, 1992. Accumulation of phenolic-acids in
20 filtrate-treated alfalfa cell-cultures derived from genotypes with different susceptibility to
21 *Fusarium oxysporum*. J Plant Physiol 140: 21-27.

22 Dalmay, T., A. Hamilton, E. Mueller & D.C. Baulcombe, 2000. Potato virus X amplicons in
23 *Arabidopsis* mediate genetic and epigenetic gene silencing. Plant Cell 12: 369-379.

1 De Clercq, J., M. Zambre, M. Van Montagu, W. Dillen & G. Angenon, 2002. An optimized
2 *Agrobacterium*-mediated transformation procedure for *Phaseolus acutifolius* A. Gray.
3 Plant Cell Reports 21: 333-340.

4 de Sousa-Majer, M.J., N.C. Turner, D.C. Hardie, R.L. Morton, B. Lamont & T.J.V. Higgins,
5 2004. Response to water deficit and high temperature of transgenic peas (*Pisum sativum*
6 L.) containing a seed-specific alpha- amylase inhibitor and the subsequent effects on pea
7 weevil (*Bruchus pisorum* L.) survival. J Exp Bot 55: 497-505.

8 Denman, S., P.S. Knoxdavies, F.J. Calitz & S.C. Lamprecht, 1995. Pathogenicity of *Pythium*
9 *irregulare*, *Pythium sylvaticum* and *Pythium ultimum* var. *ultimum* to lucerne (*Medicago*
10 *sativa*). Austral Plant Pathol 24: 137-143.

11 Dennis, E.S., R. Dolferus, M. Ellis, M. Rahman, Y. Wu, F.U. Hoeren, A. Grover, K.P.
12 Ismond, A.G. Good & W.J. Peacock, 2000. Molecular strategies for improving
13 waterlogging tolerance in plants. J Exp Bot 51: 89-97.

14 Desbrosses, G.G., J. Kopka & M.K. Udvardi, 2005. *Lotus japonicus* metabolic profiling.
15 Development of gas chromatography-mass spectrometry resources for the study of plant-
16 microbe interactions. Plant Physiol 137: 1302-1318.

17 Diatchenko, L., Y.F.C. Lau, A.P. Campbell, A. Chenchik, F. Moqadam, B. Huang, S.
18 Lukyanov, K. Lukyanov, N. Gurskaya, E.D. Sverdlov & P.D. Siebert, 1996. Suppression
19 subtractive hybridization: A method for generating differentially regulated or tissue-
20 specific cDNA probes and libraries. Proc Natl Acad Sci USA 93: 6025-6030.

21 Diers, B. 2004. Soybean genetic improvement through conventional and molecular based
22 strategies. In 5th European Conference on Grain Legumes, Dijon, France, 7-11 June, 2004.
23 Ed AEP. pp 147-148.

24 Dixon, R.A. & L.W. Sumner, 2003. Legume natural products: understanding and
25 manipulating complex pathways for human and animal health. Plant Physiol 131: 878-885.

1 Dixon, R.A., 2001. Natural products and plant disease resistance. *Nature* 411: 843-847.

2 Donaldson, P.A., T. Anderson, B.G. Lane, A.L. Davidson & D.H. Simmonds, 2001. Soybean
3 plants expressing an active oligomeric oxalate oxidase from the wheat gf-2.8 (germin)
4 gene are resistant to the oxalate-secreting pathogen *Sclerotinia sclerotiorum*. *Physiol Mol*
5 *Plant Pathol* 59: 297-307.

6 du Preez, E.D., N.C. van Rij & K.F. Lawrance, 2005. First report of soybean rust caused by
7 *Phakopsora pachyrhizi* on dry beans in South Africa. *Plant Dis* 89: 206-206.

8 Dunwell, J.M., 2000. Crop genomics: progress and prospects. *J Chem Technol Biotechnol* 75:
9 913-918.

10 Durieu, P. & S.J. Ochatt, 2000. Efficient intergeneric fusion of pea (*Pisum sativum* L.) and
11 grass pea (*Lathyrus sativus* L.) protoplasts. *J Exp Bot* 51: 1237-1242.

12 Dwivedi, B.S., M. Ram, B.P. Singh, M. Das & R.N. Prasad, 1992. Effect of liming on boron
13 nutrition of pea (*Pisum sativum* L) and corn (*Zea mays* L) grown in sequence in an acid
14 alfisol. *Fertil Res* 31: 257-262.

15 Edwards, O & K.B. Singh, 2005. Resistance to insect pests: What do legumes have to offer?.
16 *Euphytica*, this issue.

17 Eizenberg, H., J. Qolquhoun & C.A. Malory-Smith, 2004. The relationship between
18 temperature and small broomrape (*O. minor*) parasitism in red clover (*Trifolium pratense*).
19 *Weed Sci.* 52:735-741.

20 Eujayl, I., W. Erskine, B. Bayaa, M. Baum & E. Pehu, 1998. Fusarium vascular wilt in lentil:
21 Inheritance and identification of DNA markers for resistance. *Plant Breed* 117: 497-499.

22 Eujayl, I., W. Erskine, M. Baum & E. Pehu, 1999. Inheritance and linkage analysis of frost
23 injury in lentil. *Crop Sci* 39: 639-642.

24 Eulgem, T., 2005. Regulation of the Arabidopsis defense transcriptome. *Trends Plant Sci* 10:
25 71-78.

1 Faleiro, F.G., V.A. Ragagnin, M.A. Moreira & E.G. de Barros, 2004. Use of molecular
2 markers to accelerate the breeding of common bean lines resistant to rust and anthracnose.
3 Breeding of common bean lines resistant to rust and anthracnose aided by molecular
4 markers. *Euphytica* 138: 213-218.

5 Fan, W.H. & X.N. Dong, 2002. *In vivo* interaction between NPR1 and transcription factor
6 TGA2 leads to salicylic acid-mediated gene activation in Arabidopsis. *Plant Cell* 14: 1377-
7 1389.

8 Fecht-Christoffers, M.M., H.P. Braun, C. Lemaitre-Guillier, A. VanDorsselaer & W.J. Horst,
9 2003. Effect of Manganese toxicity on the proteome of the leaf apoplast in cowpea. *Plant*
10 *Physiol* 133: 1935-1946.

11 Flowers, T.J., 2004. Improving crop salt tolerance. *J Exp Bot* 55: 307-319.

12 Foolad, M.R., 2004. Recent advances in genetics of salt tolerance in tomato. *Plant Cell Tissue*
13 *Organ Cult* 76: 101-119.

14 Ford, R., E.C.K. Pang & P.W.J. Taylor, 1999. Genetics of resistance to ascochyta blight
15 (*Ascochyta lentis*) of lentil and the identification of closely linked RAPD markers. *Theor*
16 *Appl Genet* 98: 93-98.

17 Fuller, M.P. & M.H.I. Eed 2003. The development of multiple stress-resistant cauliflower
18 using mutagenesis in conjunction with a microshoot tissue culture technique. *In XXVI*
19 *International Horticultural Congress: Environmental Stress and Horticulture Crops*,
20 Toronto, Canada, 2003. Ed I.A. Horticulturae.

21 Gao, Z., S. Eysers, C. Thomas, N. Ellis & A. Maule, 2004. Identification of markers tightly
22 linked to sbm recessive genes for resistance to Pea Seed-borne Mosaic Virus. *Theor Appl*
23 *Genet* 109: 488-494.

24 Garza, R., C. Cardona & S.P. Singh, 1996. Inheritance of resistance to the bean-pod weevil
25 (*Apion godmani* Wagner) in common beans from Mexico. *Theor Appl Genet* 92: 357-362.

1 Gaunt, R.E., 1983. Shoot diseases caused by fungal pathogens. In: P.D. Hebblethwaite (ed.),
2 The Faba Bean, 463-492. Butterworths, London.

3 Gilchrist, E.J. & G.W. Haughn, 2005. TILLING without a plough: a new method with
4 applications for reverse genetics. *Curr Opin Plant Biol* 8: 211-215.

5 Graham, P.H. & C.P. Vance, 2003. Legumes: importance and constraints to greater use. *Plant*
6 *Physiol* 131: 872-877.

7 Griga, M., J. Stejskal & K. Beber, 1995. Analysis of tissue culture-derived variation in pea
8 (*Pisum sativum* L). Preliminary results. *Euphytica* 85: 335-339.

9 Gulati, A., P. Schryer & A. McHughen, 2002. Production of fertile transgenic lentil (*Lens*
10 *culinaris* Medik) plants using particle bombardment. *In Vitro Cell Dev Biol-Plant* 38: 316-
11 324.

12 Gygi, S.P. & R. Aebersold, 2000. Mass spectrometry and proteomics. *Curr Opin Chem Biol*
13 4: 489-494.

14 Gygi, S.P., Y. Rochon, B.R. Franza & R. Aebersold, 1999. Correlation between protein and
15 mRNA abundance in yeast. *Mol Cell Biol* 19: 1720-1730.

16 Hammett, K.R.W., B.G. Murray, K.R. Markham & I.C. Hallett, 1994. Interspecific
17 hybridization between *Lathyrus odoratus* and *L. belinensis*. *Int J Plant Sci* 155: 763-771.

18 Hamwieh, A., S.M. Udupa, W. Choumane, A. Sarker, F. Dreyer, C. Jung & M. Baum, 2005.
19 A genetic linkage map of *Lens* sp based on microsatellite and AFLP markers and the
20 localization of fusarium vascular wilt resistance. *Theor Appl Genet* 110: 669-677.

21 Handberg, K. & J. Stougaard, 1992. *Lotus japonicus*, an autogamous, diploid legume species
22 for classical and molecular genetics. *Plant J* 2: 487-496.

23 Hannon, G., 2002. Mechanisms and application of RNA interference. *Biol Reprod* 66: 70.

24 Hanounik, S., 1980. Effect of chemical treatments and host genotypes on disease
25 severity/yield relationships of Ascochyta blight in faba beans. *FABIS Newsletter* 2: 50

1 Hanounik, S., 1981. Influence of Ronilan on the severity of chocolate spot and yield of faba
2 bean. FABIS Newsletter 3: 50-51

3 Hanselle, T. & W. Barz, 2001. Purification and characterisation of the extracellular PR-2b
4 beta-1,3-glucanase accumulating in different *Ascochyta rabiei*-infected chickpea (*Cicer*
5 *arietinum* L.) cultivars. Plant Sci 161: 773-781.

6 Hansen, L.N. & E.D. Earle, 1995. Transfer of resistance to *Xanthomonas campestris* pv.
7 *campestris* into *Brassica oleracea* L. by protoplast fusion. Theor Appl Genet 91: 1293-
8 1300.

9 Hasegawa, P.M., R.A. Bressan, J.K. Zhu & H.J. Bohnert, 2000. Plant cellular and molecular
10 responses to high salinity. Annu Rev Plant Physiol Plant Molec Biol 51: 463-499.

11 He, X.Z. & R.A. Dixon, 2000. Genetic manipulation of isoflavone 7-O-methyltransferase
12 enhances biosynthesis of 4'-O-methylated isoflavonoid phytoalexins and disease resistance
13 in alfalfa. Plant Cell 12: 1689-1702.

14 Henikoff, S., B.J. Till & L. Comai, 2004. TILLING. Traditional mutagenesis meets functional
15 genomics. Plant Physiol 135: 630-636.

16 Hipkind, J.D. & N.L. Paiva, 2000. Constitutive accumulation of a resveratrol-glucoside in
17 transgenic alfalfa increases resistance to *Phoma medicaginis*. Mol Plant-Microbe Interact
18 13: 551-562.

19 Holbrook, C.C., D.M. Wilson, K.S. Rucker, C.K. Kvien & J.E. Hook, 1994. Possible role of
20 drought tolerance in reducing aflatoxin contamination of peanut. Agronomy Abstract: 223.

21 Infantino, A, M. Kharrat, L. Riccioni¹, C.J. Coyne, K.E. McPhee & N.J. Grünwald, 2005.
22 Screening techniques and sources of resistance to root diseases in cool season food
23 legumes. Euphytica, this issue.

1 Ishitani, M., I. Rao, P. Wenzl, S. Beebe & J. Tohme, 2004. Integration of genomics approach
2 with traditional breeding towards improving abiotic stress adaptation: drought and
3 aluminum toxicity as case studies. *Field Crop Res* 90: 35-45.

4 Jain, A.K., S.M. Basha & C.C. Holbrook, 2001. Identification of drought-responsive
5 transcripts in peanut (*Arachis hypogaea* L.). *EJB* 4: 59-67.

6 Jain, S.M., 2001. Tissue culture-derived variation in crop improvement. *Euphytica* 118: 153-
7 166.

8 Janila, P. & B. Sharma, 2004. RAPD and SCAR markers for powdery mildew resistance gene
9 er in pea. *Plant Breed* 123: 271-274.

10 Jayasena, K.W., M.R. Hajimorad, E.G. Law, A.U. Rehman, K.E. Nolan, T. Zanker, R.J. Rose
11 & J.W. Randles, 2001. Resistance to Alfalfa mosaic virus in transgenic barrel medic lines
12 containing the virus coat protein gene. *Aust J Agric Res* 52: 67-72.

13 Jiang, D.H., X.J. Chen, K.L. Wu & Z.J. Guo, 2004. Expression of Cryptogein in tobacco
14 plants exhibits enhanced disease resistance and tolerance to salt stress. *Chin Sci Bull* 49:
15 803-809.

16 Jones, A.L., I.E. Johansen, S.J. Bean, I. Bach & A.J. Maule, 1998. Specificity of resistance to
17 pea seed-borne mosaic potyvirus in transgenic peas expressing the viral replicase (NIb)
18 gene. *J Gen Virol* 79: 3129-3137.

19 Jorrín, J.V., D. Rubiales, E. Dumas-Gaudot, G. Recorbet, A. Maldonado, M.A. Castillejo &
20 M. Curto, 2005. Proteomics: a promising approach to study biotic interaction in legumes.
21 A review. *Euphytica*, this issue.

22 Kahraman, A., I. Kusmenoglu, N. Aydin, A. Aydogan, W. Erskine & F.J. Muehlbauer, 2004.
23 QTL mapping of winter hardiness genes in lentil. *Crop Sci* 44: 13-22.

24 Kaimori, N., M. Senda, R. Ishikawa, S. Akada, T. Harada & M. Niizeki, 1998. Asymmetric
25 somatic cell hybrids between alfalfa and birdsfoot trefoil. *Breed Sci* 48: 29-34.

1 Kalo, P., A. Seres, S.A. Taylor, J. Jakab, Z. Kevei, A. Kereszt, G. Endre, T.H.N. Ellis & G.B.
2 Kiss, 2004. Comparative mapping between *Medicago sativa* and *Pisum sativum*. Mol
3 Genet Genomics 272: 235-246.

4 Kappers, I.F., A. Aharoni, T. van Herpen, L.L.P. Luckerhoff, M. Dicke & H.J. Bouwmeester,
5 2005. Genetic engineering of terpenoid metabolism attracts, bodyguards to Arabidopsis.
6 Science 309: 2070-2072.

7 Kassem, M.A., K. Meksem, C.H. Kang, V.N. Njiti, V. Kilo, A.J. Wood & D.A. Lightfoot,
8 2004. Loci underlying resistance to manganese toxicity mapped in a soybean recombinant
9 inbred line population of 'Essex' x 'Forrest'. Plant and Soil 260: 197-204.

10 Kasuga, M., Q. Liu, S. Miura, K. Yamaguchi-Shinozaki & K. Shinozaki, 1999. Improving
11 plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible
12 transcription factor. Nat Biotechnol 17: 287-291.

13 Kato, T., S. Sato, Y. Nakamura, T. Kaneko, E. Asamizu & S. Tabata, 2003. Structural
14 analysis of a *Lotus japonicus* genome. V. Sequence features and mapping of sixty-four
15 TAC clones which cover the 6.4 Mb regions of the genome. DNA Res 10: 277-285.

16 Kav, N.N.V., S. Srivastava, L. Goonewardene & S.F. Blade, 2004. Proteome-level changes in
17 the roots of *Pisum sativum* in response to salinity. Ann Appl Biol 145: 217-230.

18 Kelly, J.D. & V.A. Vallejo, 2004. A comprehensive review of the major genes conditioning
19 resistance to anthracnose in common bean. Hortscience 39: 1196-1207.

20 Kelly, J.D., P. Gepts, P.N. Miklas & D.P. Coyne, 2003. Tagging and mapping of genes and
21 QTL and molecular marker-assisted selection for traits of economic importance in bean
22 and cowpea. Field Crop Res 82: 135-154.

23 Khan, A.J., S. Hassan, M. Tariq & T. Khan, 2001. Haploidy breeding and mutagenesis for
24 drought tolerance in wheat. Euphytica 120: 409-414.

1 Kim, J.B., J.Y. Kang & S.Y. Kim, 2004. Over-expression of a transcription factor regulating
2 ABA-responsive gene expression confers multiple stress tolerance. *Plant Biotechnol J* 2:
3 459-466.

4 Kim, S.T., K.S. Cho, S. Yu, S.G. Kim, J.C. Hong, C.-D. Han, D.W. Bae, M.H. Nam & K.Y.
5 Kang, 2003. Proteomic analysis of differentially expressed proteins induced by rice blast
6 fungus and elicitor in suspension-cultured rice cells. *Proteomics* 3: 2368-2378.

7 Klingler, J., R. Creasy, L. Gao, R.M. Nair, C. Alonso-Suazo, H.S. Jacob, O.R. Edwards &
8 K.B. Singh, 2005. Aphid resistance in *Medicago truncatula* involves antixenosis and
9 phloem-specific, inducible antibiosis, and maps to a single locus flanked by NBS-LRR
10 resistance gene analogs. *Plant Physiol* 137: 1445-1455.

11 Koike, M. & K. Nanbu, 1997. Phenylalanine ammonia-lyase activity in alfalfa suspension
12 cultures treated with conidia and elicitors of *Verticillium albo-atrum*. *Biol Plant* 39: 349-
13 353.

14 Kolkman, J.M. & J.D. Kelly, 2003. QTL conferring resistance and avoidance to white mold in
15 common bean. *Crop Sci* 43: 539-548.

16 Koltai, H., M. Dhandaydham, C. Opperman, J. Thomas & D. Bird, 2001. Overlapping plant
17 signal transduction pathways induced by a parasitic nematode and a rhizobial
18 endosymbiont. *Mol Plant-Microbe Interact* 14: 1168-1177.

19 Kowalski, B. & A.C. Cassells, 1999. Mutation breeding for yield and *Phytophthora infestans*
20 (Mont.) de Bary foliar resistance in potato (*Solanum tuberosum* L.-cv. Golden Wonder)
21 using computerized image analysis in selection. *Potato Res* 42: 121-130.

22 Kulikova, O., G. Gualtieri, R. Geurts, D.J. Kim, D. Cook, T. Huguet, J.H. de Jong, P.F.
23 Fransz & T. Bisseling, 2001. Integration of the FISH pachytene and genetic maps of
24 *Medicago truncatula*. *Plant J* 27: 49-58.

1 Laanbroek, H.J., 1990. Bacterial cycling of minerals that affect plant-growth in waterlogged
2 soils: A review. *Aquat Bot* 38: 109-125.

3 Larkin, P.J. & W.R. Scowcroft, 1981. Somaclonal variation - a novel source of variability
4 from cell-cultures for plant Improvement. *Theor Appl Genet* 60: 197-214.

5 Lazaridou, T.B., D.G. Roupakias & A.S. Economou, 1993. Embryo rescue in *Vicia faba* and
6 *Vicia narbonensis*. *Plant Cell Tissue Organ Cult* 33: 297-301.

7 Lebel, E., P. Heifetz, L. Thorne, S. Uknes, J. Ryals & E. Ward, 1998. Functional analysis of
8 regulatory sequences controlling PR-1 gene expression in Arabidopsis. *Plant Journal* 16:
9 223-233.

10 Lee, G.J., H.R. Boerma, M.R. Villagarcia, X. Zhou, T.E. Carter, Z. Li & M.O. Gibbs, 2004. A
11 major QTL conditioning salt tolerance in S-100 soybean and descendent cultivars. *Theor*
12 *Appl Genet* 109: 1610-1619.

13 Li, H.Y., Y.M. Zhu, Q. Chen, R.L. Conner, X.D. Ding, J. Li & B.B. Zhang, 2004. Production
14 of transgenic soybean plants with two anti-fungal protein genes via *Agrobacterium* and
15 particle bombardment. *Biol Plant* 48: 367-374.

16 Li, Y.D., Y.J. Wang, Y.P. Tong, J.G. Gao, J.S. Zhang & S.Y. Chen, 2005. QTL mapping of
17 phosphorus deficiency tolerance in soybean (*Glycine max* L. Merr.). *Euphytica* 142: 137-
18 142.

19 Li, Y.G., G.J. Tanner, A.C. Delves & P.J. Larkin, 1993. Asymmetric somatic hybrid plants
20 between *Medicago sativa* L (Alfalfa, Lucerne) and *Onobrychis viciifolia* Scop (Sainfoin).
21 *Theor Appl Genet* 87: 455-463.

22 Liu, J.H., X.Y. Xu & X.X. Deng, 2005. Intergeneric somatic hybridization and its application
23 to crop genetic improvement. *Plant Cell Tissue Organ Cult* 82: 19-44.

1 Liu, W.G., M.Y. Zheng, E.A. Polle & C.F. Konzak, 2002a. Highly efficient doubled-haploid
2 production in wheat (*Triticum aestivum* L.) via induced microspore embryogenesis. Crop
3 Sci 42: 686-692.

4 Liu, Y.L., M. Schiff & S.P. Dinesh-Kumar, 2002b. Virus-induced gene silencing in tomato.
5 Plant J 31: 777-786.

6 Livingstone, D.M., J.L. Hampton, P.M. Phipps & E.A. Grabau, 2005. Enhancing resistance to
7 *Sclerotinia minor* in peanut by expressing a barley oxalate oxidase gene. Plant Physiol
8 137: 1354-1362.

9 Lozovaya, V.V., A.V. Lygin, S. Li, G.L. Hartman & J.M. Widhohn, 2004. Biochemical
10 response of soybean roots to *Fusarium solani* f. sp *glycines* infection. Crop Sci 44: 819-
11 826.

12 Magbanua, Z.V., H.D. Wilde, J.K. Roberts, K. Chowdhury, J. Abad, J.W. Moyer, H.Y.
13 Wetzstein & W.A. Parrott, 2000. Field resistance to Tomato Spotted Wilt Virus in
14 transgenic peanut (*Arachis hypogaea* L.) expressing an antisense nucleocapsid gene
15 sequence. Mol Breed 6: 227-236.

16 Mahmoud, A.L.E. & M.H. Abdalla, 1994. Natural occurrence of mycotoxins in broad bean
17 (*Vicia faba* L.) seeds and their effect on *Rhizobium*-Legume symbiosis. Soil Biol.
18 Biochem. 26: 1081-1085.

19 Mallikarjuna, N. & J.P. Moss, 1995. Production of hybrids between *Cajanus platycarpus* and
20 *Cajanus cajan*. Euphytica 83: 43-46.

21 Martinez, V.A., W.G. Hill & S. Knott, 2002. On the use of double haploids for detecting QTL
22 in outbred populations. Heredity 88: 423-431.

23 McCallum, C.M., L. Comai, E.A. Greene & S. Henikoff, 2000. Targeting induced local
24 lesions in genomes (TILLING) for plant functional genomics. Plant Physiol 123: 439-442.

1 McDonald, G.K. & G. Dean, 1996. Effect of waterlogging on the severity of disease caused
2 by *Mycosphaerella pinodes* in peas (*Pisum sativum* L). Aus J Exp Agric 36: 219-222.

3 Merchan, F., C. Breda, J.P. Hormaeche, C. Sousa, A. Kondorosi, O.M. Aguilar, M. Megias &
4 M. Crespi, 2003. A Kruppel-like transcription factor gene is involved in salt stress
5 responses in *Medicago* spp. Plant Soil 257: 1-9.

6 Miklas, P.N., D.P. Coyne, K.F. Grafton, N. Mutlu, J. Reiser, D.T. Lindgren & S.P. Singh,
7 2003. A major QTL for common bacterial blight resistance derives from the common bean
8 great northern landrace cultivar Montana No. 5. Euphytica 131: 137-146.

9 Miklas, P.N., W.C. Johnson, R. Delorme & P. Gepts, 2001. QTL conditioning physiological
10 resistance and avoidance to white mold in dry bean. Crop Sci 41: 309-315.

11 Mithofer, A., B. Muller, G. Wanner & L.A. Eichacker, 2002. Identification of defence-related
12 cell wall proteins in *Phytophthora sojae*-infected soybean roots by ESI-MS/MS. Mol Plant
13 Pathol 3: 163-166.

14 Mithofer, A., B. Schulze & W. Boland, 2004. Biotic and heavy metal stress response in
15 plants: evidence for common signals. FEBS Letters 566: 1-5.

16 Morphew, R., R. Arndt, C. Lauter, M. Tesfaye, D. Samac & G. Temple, 2004.
17 Characterization of altered metabolism in transgenic alfalfa overexpressing malate
18 dehydrogenase to confer aluminum tolerance. Abstr Pap Am Chem Soc 227: 148.

19 Murray, J.D., T.E. Michaels, C. Cardona, A.W. Schaafsma & K.P. Pauls, 2004. Quantitative
20 trait loci for leafhopper (*Empoasca fabae* and *Empoasca kraemeri*) resistance and seed
21 weight in the common bean. Plant Breed 123: 474-479.

22 Mutlu, N., P. Miklas, J. Reiser & D. Coyne, 2005. Backcross breeding for improved
23 resistance to common bacterial blight in pinto bean (*Phaseolus vulgaris* L.). Plant Breed
24 124: 282-287.

1 Nakajo, S., M. Niizeki, T. Harada, R. Ishikawa & K. Saito, 1994. Somatic-cell hybridization in
2 rice (*Oryza sativa* L.) and birdsfoot-trefoil (*Lotus-Corniculatus* L.). *Breed Sci* 44: 79-81.

3 Narayanan, N.N., N. Baisakh, N.P. Oliva, C.M. VeraCruz, S.S. Gnanamanickam, K. Datta &
4 S.K. Datta, 2004. Molecular breeding: marker-assisted selection combined with biolistic
5 transformation for blast and bacterial blight resistance in Indica rice (cv. CO39). *Mol*
6 *Breed* 14: 61-71.

7 Narayanan, R.A., R. Atz, R. Denny, N.D. Young & D.A. Somers, 1999. Expression of
8 soybean cyst nematode resistance in transgenic hairy roots of soybean. *Crop Sci* 39: 1680-
9 1686.

10 Navas-Cortés, J.A., B. Hau & R.M. Jimenez-Diaz, 2000. Yield loss in chickpeas in relation to
11 development of *Fusarium* wilt epidemics. *Phytopathology* 90: 1269-1278.

12 Nene, Y.L. & M.V. Reddy, 1987 Chickpea diseases and their control. In: M.C. Saxena &
13 K.B. Singh (Eds). In the chickpea, pp. 233-270. CAB International, Oxon, UK.

14 Nyamsuren, O., F. Colditz, S. Rosendahl, M. Tamasloukht, T. Bekel, F. Meyer, H. Kuester, P.
15 Franken & F. Krajinski, 2003. Transcriptional profiling of *Medicago truncatula* roots after
16 infection with *Aphanomyces euteiches* (oomycota) identifies novel genes upregulated
17 during this pathogenic interaction. *Physiol Mol Plant Pathol* 63: 17-26.

18 Onate-Sanchez, L. & K.B. Singh, 2002. Identification of Arabidopsis ethylene-responsive
19 element binding factors with distinct induction kinetics after pathogen infection. *Plant*
20 *Physiol* 128: 1313-1322.

21 Ormerod, A.J. & P.D.S. Caligari, 1994. Anther and microspore culture of *Lupinus albus* in
22 liquid culture-medium. *Plant Cell Tissue Organ Cult* 36: 227-236.

23 Ornatowski, W., J. Jayaraj, T.C. Todd, W.T. Schapaugh, S. Muthukrishnan & H.N. Trick,
24 2004. Introduction and constitutive expression of a tobacco hornworm (*Manduca sexta*)
25 chitinase gene in soybean. *In Vitro Cell Dev Biol-Plant* 40: 260-265.

1 Ouedraogo, J.T., J.B. Tignegre, M.P. Timko & F.J. Belzile, 2002. AFLP markers linked to
2 resistance against *Striga gesnerioides* race 1 in cowpea (*Vigna unguiculata*). *Genome* 45:
3 787-793.

4 Ozawa, R., T. Shimoda, M. Kawaguchi, G. Arimura, J. Horiuchi, T. Nishioka & J.
5 Takabayashi, 2000. *Lotus japonicus* infested with herbivorous mites emits volatile
6 compounds that attract predatory mites. *J Plant Res* 113: 427-433.

7 Parker, C. & C.R. Riches, 1993. Parasitic weeds of the world: biology and control. CAB
8 International, Wallingford, UK.

9 Pedrosa, A., N. Sandal, J. Stougaard, D. Schweizer & A. Bachmair, 2002. Chromosomal map
10 of the model legume *Lotus japonicus*. *Genetics* 161: 1661-1672.

11 Perry, J.A., T.L. Wang, T.J. Welham, S. Gardner, J.M. Pike, S. Yoshida & M. Parniske, 2003.
12 A TILLING reverse genetics tool and a web-accessible collection of mutants of the
13 legume *Lotus japonicus*. *Plant Physiol* 131: 866-871.

14 Pinheiro, C., J. Kehr & C.P. Ricardo, 2005. Effect of water stress on lupin stem protein
15 analysed by two-dimensional gel electrophoresis. *Planta* 221: 716-728.

16 Pivonia, S. & X.B. Yang, 2004. Assessment of the potential year-round establishment of
17 soybean rust throughout the world. *Plant Dis* 88: 523-529.

18 Pnueli, L., E. Hallak-Herr, M. Rozenberg, M. Cohen, P. Goloubinoff, A. Kaplan & R. Mittler,
19 2002. Molecular and biochemical mechanisms associated with dormancy and drought
20 tolerance in the desert legume *Retama raetam*. *Plant J* 31: 319-330.

21 Popelka, J.C., N. Terryn & T.J.V. Higgins, 2004. Gene technology for grain legumes: can it
22 contribute to the food challenge in developing countries? *Plant Sci* 167: 195-206.

23 Poulain, D. & H. Almohammad, 1995. Effects of boron deficiency and toxicity on faba bean
24 (*Vicia faba* L.). *Eur J Agron* 4: 127-134.

1 Qian, Q., D.L. Zeng, P. He, X.W. Zheng, Y. Chen & L.H. Zhu, 2000. QTL analysis of the rice
2 seedling cold tolerance in a double haploid population derived from anther culture of a
3 hybrid between indica and japonica rice. *Chin Sci Bull* 45: 448-453.

4 Rahman, M.H.H., Y. Arima, K. Watanabe & H. Sekimoto, 1999. Adequate range of boron
5 nutrition is more restricted for root nodule development than for plant growth in young
6 soybean plant. *Soil Sci Plant Nutr* 45: 287-296.

7 Ramteke, R., G.K. Gupta & O.P. Joshi, 2004. Evaluation of rust resistance in soybean
8 (*Glycine max*) under field condition. *Indian J Agric Sci* 74: 623-624.

9 Rashid, K.Y. & C.C. Bernier, 1991. The effect of rust on yield of faba bean cultivars and slow
10 rusting populations. *Can J Plant Sci* 71: 967-972.

11 Reddy, A.R., K.V. Chaitanya & M. Vivekanandan, 2004. Drought-induced responses of
12 photosynthesis and antioxidant metabolism in higher plants. *J Plant Physiol* 161: 1189-
13 1202.

14 Reddy, M.S.S., S.A. Ghabrial, C.T. Redmond, R.D. Dinkins & G.B. Collins, 2001. Resistance
15 to Bean pod mottle virus in transgenic soybean lines expressing the capsid polyprotein.
16 *Phytopathology* 91: 1238-1238.

17 Repetto, O., G. Bestel-Corre, E. Dumas-Gaudot, G. Berta, V. Gianinazzi-Pearson & S.
18 Gianinazzi, 2003. Targeted proteomics to identify cadmium-induced protein modifications
19 in *Glomus mosseae*-inoculated pea roots. *New Phytol* 157: 555-567.

20 Risipail, N. 2005. Molecular and metabolic characterisation of symbiotic interactions in *Lotus*
21 *japonicus* PhD Thesis Institute of Grassland and Environmental Research (IGER),
22 University of Wales, Aberystwyth.

23 Robatzek, S. & I.E. Somssich, 2002. Targets of AtWRKY6 regulation during plant
24 senescence and pathogen defense. *Genes Dev* 16: 1139-1149.

1 Rodrigues, L.R., J.M.S. Oliveira, J.E.A. Mariath & M.H. Bodanese-Zanettini, 2005.
2 Histology of embryogenic responses in soybean anther culture. Plant Cell Tissue Organ
3 Cult 80: 129-137.

4 Rodríguez-Conde, M.F., M.T. Moreno, J.I. Cubero & D. Rubiales, 2004. Characterization of
5 the *Orobanche-Medicago truncatula* association for studying early stages of the parasite-
6 host interaction. Weed Res 44: 218-223.

7 Román, B., A.M. Torres, D. Rubiales, J.I. Cubero & Z. Satovic, 2002. Mapping of
8 quantitative trait loci controlling broomrape (*Orobanche crenata* Forsk.) resistance in faba
9 bean (*Vicia faba* L.). Genome 45: 1057-1063.

10 Román, B., Z. Satovic, C.M. Avila, D. Rubiales, M.T. Moreno & A.M. Torres, 2003.
11 Locating genes associated with *Ascochyta fabae* resistance in *Vicia faba*. Aust J Agric Res
12 54: 85-90.

13 Romero-Andreas, J., B.S. Yandell & F.A.L. Bliss, 1986. Inheritance of a novel seed protein of
14 *Phaseolus vulgaris* L. and its effect on seed composition. Theor Appl Genet 72: 123-128.

15 Rubiales, D, A. Pérez-de-Luque, J.C. Sillero, B. Román, M. Kharrat, S. Khalil, D.M. Joel &
16 C. Riches, 2005. Screening techniques and sources of resistance against parasitic
17 weeds in grain legumes. Euphytica, this issue.

18 Rubiales, D. & A. Moral, 2004. Prehaustorial resistance against alfalfa rust (*Uromyces*
19 *striatus*) in *Medicago truncatula*. Eur J Plant Pathol 110: 239-243.

20 Rubiales, D., 2001. Parasitic plants: an increasing threat. Grain Legumes, 33: 10-11

21 Rubiales, D., A.A. Emeran & J.C. Sillero, 2002. Rusts on legumes in Europe and North
22 Africa. Grain Legumes, 37: 8-9.

23 Sagan, M., T. Huguet & G. Duc, 1994. Phenotypic characterization and classification of
24 nodulation mutants of pea (*Pisum sativum* L.). Plant Sci 100: 59-70.

25 Salles, II, J.W. Blount, R.A. Dixon & K. Schubert, 2002. Phytoalexin induction and beta-1,3-
26 glucanase activities in *Colletotrichum trifolii* infected leaves of alfalfa (*Medicago sativa*
27 L.). Physiol Mol Plant Pathol 61: 89-101.

1 Salzer, P., A. Bonanomi, K. Beyer, R. Vogeli-Lange, R.A. Aeschbacher, J. Lange, A.
2 Wiemken, D. Kim, D.R. Cook & T. Boller, 2000. Differential expression of eight chitinase
3 genes in *Medicago truncatula* roots during mycorrhiza formation, nodulation, and
4 pathogen infection. *Mol Plant-Microbe Interact* 13: 763-777.

5 Samac, D.A. & A.C. Smigocki, 2003. Expression of oryzacystatin I and II in alfalfa increases
6 resistance to the root-lesion nematode. *Phytopathology* 93: 799-804.

7 Samac, D.A., M. Tesfaye, M. Dornbusch, P. Saruul & S.J. Temple, 2004. A comparison of
8 constitutive promoters for expression of transgenes in alfalfa (*Medicago sativa*).
9 *Transgenic Res* 13: 349-361.

10 Samantaray, S., G.R. Rout & P. Das, 1999. *In vitro* selection and regeneration of zinc tolerant
11 calli from *Setaria italica* L. *Plant Sci* 143: 201-209.

12 Sanan-Mishra, N., X.H. Pham, S.K. Sopory & N. Tuteja, 2005. Pea DNA helicase 45
13 overexpression in tobacco confers high salinity tolerance without affecting yield. *Proc Natl*
14 *Acad Sci USA* 102: 509-514.

15 Sanyal, I., A.K. Singh, M. Kaushik & D.V. Amla, 2005. *Agrobacterium*-mediated
16 transformation of chickpea (*Cicer arietinum* L.) with *Bacillus thuringiensis* cry1Ac gene
17 for resistance against pod borer insect *Helicoverpa armigera*. *Plant Sci* 168: 1135-1146.

18 Sarmah, B.K., A. Moore, W. Tate, L. Molvig, R.L. Morton, D.P. Rees, P. Chiaiese, M.J.
19 Chrispeels, L.M. Tabe & T.J.V. Higgins, 2004. Transgenic chickpea seeds expressing high
20 levels of a bean alpha-amylase inhibitor. *Mol Breed* 14: 73-82.

21 Saunders, J. & N. O'Neill, 2004. The characterization of defense responses to fungal infection
22 in alfalfa. *BioControl* 49: 715-728.

23 Schauser, L., A. Roussis, J. Stiller & J. Stougaard, 1999. A plant regulator controlling
24 development of symbiotic root nodules. *Nature* 402: 191-195.

1 Schena, M., D. Shalon, R.W. Davis & P.O. Brown, 1995. Quantitative monitoring of gene-
2 expression patterns with a complementary-DNA microarray. *Science* 270: 467-470.

3 Schnabl, H., T.L. Mahaworasilpa, H.G.L. Coster & A. von Keller, 1998. Production of hybrid
4 cells from single protoplasts of sunflower hypocotyl and broad bean guard cells by
5 electrical fusion. *Plant Cell Tissue Organ Cult* 55: 59-62.

6 Schneider, K.A., K.F. Grafton & J.D. Kelly, 2001. QTL analysis of resistance to fusarium
7 root rot in bean. *Crop Sci* 41: 535-542.

8 Schneider, K.A., M.E. Brothers & J.D. Kelly, 1997. Marker-assisted selection to improve
9 drought resistance in common bean. *Crop Sci* 37: 51-60.

10 Schröder, H.E., S. Gollasch, A. Moore, L.M. Tabe, S. Craig, D.C. Hardie, M.J. Chrispeels,
11 D. Spencer & T.J.V. Higgins, 1995. Bean alpha-amylase inhibitor confers resistance to the
12 pea weevil (*Bruchus pisorum*) in transgenic peas (*Pisum sativum* L.). *Plant Physiol* 107:
13 1233-1239.

14 Sharma, H.C., 1995. How wide can a wide cross be?. *Euphytica* 82: 43-64.

15 Sharma, K.K. & M. Lavanya, 2002. Recent developments in transgenics for abiotic stress in
16 legumes of the semi-arid tropics. *JIRCAS Working Report*: 61-73.

17 Shi, H.Z., B.H. Lee, S.J. Wu & J.K. Zhu, 2003. Overexpression of a plasma membrane
18 Na⁺/H⁺ antiporter gene improves salt tolerance in *Arabidopsis thaliana*. *Nat Biotechnol*
19 21: 81-85.

20 Shimada, N., T. Akashi, T. Aoki & S. Ayabe, 2000. Induction of isoflavonoid pathway in the
21 model legume *Lotus japonicus*: molecular characterization of enzymes involved in
22 phytoalexin biosynthesis. *Plant Sci* 160: 37-47.

23 Shinozaki, K. & K. Yamaguchi-Shinozaki, 2000. Molecular responses to dehydration and low
24 temperature: differences and cross-talk between two stress signaling pathways. *Curr Opin*
25 *Plant Biol* 3: 217-223.

1 Shou, H.X., P. Bordallo, J.B. Fan, J.M. Yeakley, M. Bibikova, J. Sheen & K. Wang, 2004.
2 Expression of an active tobacco mitogen-activated protein kinase kinase enhances
3 freezing tolerance in transgenic maize. *Proc Natl Acad Sci USA* 101: 3298-3303.

4 Sillero J.C., S. Fondevilla, J. Davidson, M.C. Vaz Patto, T.D. Warkentin, J. Thomas & D.
5 Rubiales, 2005. Screening techniques and sources of resistance to rusts and mildews in
6 grain legumes. *Euphytica*, this issue.

7 Singh, B.N., R.N. Mishra, P.K. Agarwal, M. Goswami, S. Nair, S.K. Sopory & M.K. Reddy,
8 2004. A pea chloroplast translation elongation factor that is regulated by abiotic factors.
9 *Biochem Biophys Res Commun* 320: 523-530.

10 Singh, K.B., R.C. Foley & L. Onate-Sanchez, 2002. Transcription factors in plant defense and
11 stress responses. *Curr Opin Plant Biol* 5: 430-436.

12 Singh, S.P., 2001. Broadening the genetic base of common bean cultivars: A review. *Crop Sci*
13 41: 1659-1675.

14 Singla-Pareek, S.L., M.K. Reddy & S.K. Sopory, 2003. Genetic engineering of the glyoxalase
15 pathway in tobacco leads to enhanced salinity tolerance. *Proc Natl Acad Sci USA* 100:
16 14672-14677.

17 Sivamani, E., C.W. Brey, W.E. Dyer, L.E. Talbert & R.D. Qu, 2000. Resistance to wheat
18 streak mosaic virus in transgenic wheat expressing the viral replicase (NIb) gene. *Mol*
19 *Breed* 6: 469-477.

20 Sledge, M.K., J.H. Bouton, M. Dall'Agnoll, W.A. Parrott & G. Kochert, 2002. Identification
21 and confirmation of aluminum tolerance QTL in diploid *Medicago sativa* subsp. *coerulea*.
22 *Crop Sci* 42: 1121-1128.

23 Somers, D.A., D.A. Samac & P.M. Olhoft, 2003. Recent advances in legume transformation.
24 *Plant Physiol* 131: 892-899.

- 1 Stavelly, J.R., 2000. Pyramiding rust and viral resistance genes using traditional and marker
2 techniques in common bean. *Ann Rep Bean Imp Coop* 43: 1-4.
- 3 Stiller, J., L. Martirani, S. Tuppal, R.J. Chian, M. Chiurazzi & P.M. Gresshoff, 1997. High
4 frequency transformation and regeneration of transgenic plants in the model legume *Lotus*
5 *japonicus*. *J Exp Bot* 48: 1357-1365.
- 6 Stoddard, F.L., C. Balko, W. Erskine, H.R. Khan, W. Link & A. Sarker, 2005. Screening
7 techniques and sources of resistance to abiotic stresses in cool-season food legumes.
8 *Euphytica*, this issue
- 9 Stracke, S., S. Sato, N. Sandal, M. Koyama, T. Kaneko, S. Tabata & M. Parniske, 2004.
10 Exploitation of colinear relationships between the genomes of *Lotus japonicus*, *Pisum*
11 *sativum* and *Arabidopsis thaliana*, for positional cloning of a legume symbiosis gene.
12 *Theor Appl Gen* 108: 442-449.
- 13 Sumner, L.W., P. Mendes & R.A. Dixon, 2003. Plant metabolomics: large-scale
14 phytochemistry in the functional genomics era. *Phytochemistry* 62: 817-836.
- 15 Surekha, C., M.R. Beena, A. Arundhati, P.K. Singh, R. Tuli, A. Dutta-Gupta & P.B. Kirti,
16 2005. *Agrobacterium*-mediated genetic transformation of pigeon pea (*Cajanus cajan* (L.)
17 Mill sp.) using embryonal segments and development of transgenic plants for resistance
18 against *Spodoptera*. *Plant Sci* 169: 1074–1080.
- 19 Svabova, L. & A. Lebeda, 2005. *In vitro* selection for improved plant resistance to toxin-
20 producing pathogens. *J Phytopathol* 153: 52-64.
- 21 Sweetingham, M.W., Jones R.A.C. & A.G.P. Brown, 1998. Diseases and Pests. In: “Lupins as
22 Crop Plants: Biology, Production and Utilization” Eds J.S.Gladstones, C.A Pettersen D.S.,
23 C.A. Atkins and J.Hamblin. pp. 263-290. CAB International, UK.
- 24 Tadege, M., P. Ratet & K.S. Mysore, 2005. Insertional mutagenesis: a Swiss army knife for
25 functional genomics of *Medicago truncatula*. *Trends Plant Sci* 10: 229-235.

1 Takahashi, H., T. Takai & T. Matsumoto, 1992. Breeding of strawberry cultivars resistant to
2 *Alternaria* black spot of strawberry. Resistant plants to *Alternaria alternata* strawberry
3 pathotype selected from calliclones of strawberry cultivar Morioka-16. J Jpn Soc Hortic
4 Sci 61: 323-327.

5 Tar'an, B., L. Buchwaldt, A. Tullu, S. Banniza, T.D. Warkentin & A. Vandenberg, 2003a.
6 Using molecular markers to pyramid genes for resistance to ascochyta blight and
7 anthracnose in lentil (*Lens culinaris* Medik). Euphytica 134: 223-230.

8 Tar'an, B., T. Warkentin, D.J. Somers, D. Miranda, A. Vandenburg, S. Blade, S. Woods, D.
9 Bing, A. Xue, D. DeKoeyer & G. Penner, 2003b. Quantitative trait loci for lodging
10 resistance, plant height and partial resistance to mycosphaerella blight in field pea (*Pisum*
11 *sativum* L.). Theor Appl Genet 107: 1482-1491.

12 Tar'an, B., T.E. Michaels & K.P. Pauls, 2001. Mapping genetic factors affecting the reaction
13 to *Xanthomonas axonopodis* pv. *phaseoli* in *Phaseolus vulgaris* L. under field conditions.
14 Genome 44: 1046-1056.

15 Thakur, M., D.R. Sharma & S.K. Sharma, 2002. In vitro selection and regeneration of
16 carnation (*Dianthus caryophyllus* L.) plants resistant to culture filtrate of *Fusarium*
17 *oxysporum* f.sp *dianthi*. Plant Cell Reports 20: 825-828.

18 Thoquet, P., M. Gherardi, E.P. Journet, A. Kereszt, J.M. Ane, J.M. Prosperi & T. Huguet,
19 2002. The molecular genetic linkage map of the model legume *Medicago truncatula*: an
20 essential tool for comparative legume genomics and the isolation of agronomically
21 important genes. BMC Plant Biol 2: 1.

22 Timmerman-Vaughan, G.M., M.D. Pither-Joyce, P.A. Cooper, A.C. Russell, D.S. Goulden,
23 R. Butler & J.E. Grant, 2001. Partial resistance of transgenic peas to alfalfa mosaic virus
24 under greenhouse and field conditions. Crop Sci 41: 846-853.

1 Tivoli, B., A. Baranger, C.M. Avila, S. Banniza, M. Barbetti, W. Chen, J. Davidson, K.
2 Lindeck, M. Kharrat, D. Rubiales, M. Sadiki, J.C. Sillero, M. Sweetingham & F.J.
3 Muehlbauer, 2005. Screening techniques and sources of resistance to foliar diseases
4 caused by major necrotrophic fungi in grain legumes. *Euphytica*, this issue.

5 Tonguc, M. & P.D. Griffiths, 2004. Transfer of powdery mildew resistance from *Brassica*
6 *carinata* to *Brassica oleracea* through embryo rescue. *Plant Breed* 123: 587-589.

7 Torregrosa, C., B. Dumas, F. Krajinski, M.T. Esquerre-Tugaye & C. Jacquet, 2005.
8 Transcriptomic approaches to unravel plant - pathogen interactions in legumes. *Euphytica*,
9 this issue.

10 Torregrosa, C., S. Cluzet, J. Fournier, T. Huguet, P. Gamas, J.M. Prosperi, M.T. Esquerre-
11 Tugaye, B. Dumas & C. Jacquet, 2004. Cytological, genetic, and molecular analysis to
12 characterize compatible and incompatible interactions between *Medicago truncatula* and
13 *Colletotrichum trifolii*. *Mol Plant-Microbe Interact* 17: 909-920.

14 Umezawa, T., K. Mizuno & T. Fujimura, 2002. Discrimination of genes expressed in
15 response to the ionic or osmotic effect of salt stress in soybean with cDNA-AFLP. *Plant*
16 *Cell Environ* 25: 1617-1625.

17 Umezawa, T., R. Yoshida, K. Maruyama, K. Yamaguchi-Shinozaki & K. Shinozaki, 2004.
18 SRK2C, a SNF1-related protein kinase 2, improves drought tolerance by controlling
19 stress-responsive gene expression in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 101:
20 17306-17311.

21 Valderrama, M.R., B. Roman, Z. Satovic, D. Rubiales, J.I. Cubero & A.M. Torres, 2004.
22 Locating quantitative trait loci associated with *Orobanche crenata* resistance in pea. *Weed*
23 *Res* 44: 323-328.

1 Van den Boogaart, T., A.J. Maule, J.W. Davies & G.P. Lomonossoff, 2004. Sources of target
2 specificity associated with the recovery against Pea Seed-borne Mosaic Virus infection
3 mediated by RNA silencing in pea. *Mol Plant Pathol* 5: 37-43.

4 Van Hoorn, J.W., N. Katerji, A. Hamdy & M. Mastrorilli, 2001. Effect of salinity on yield
5 and nitrogen uptake of four grain legumes and on biological nitrogen contribution from the
6 soil. *Agric Water Manage* 51: 87-98.

7 VanToai, T.T., S.K. St Martin, K. Chase, G. Boru, V. Schnipke, A.F. Schmitthenner & K.G.
8 Lark, 2001. Identification of a QTL associated with tolerance of soybean to soil
9 waterlogging. *Crop Sci* 41: 1247-1252.

10 Velculescu, V.E., L. Zhang, B. Vogelstein & K.W. Kinzler, 1995. Serial Analysis of Gene-
11 Expression. *Science* 270: 484-487.

12 Walker, D.R., J.M. Narvel, H.R. Boerma, J.N. All & W.A. Parrott, 2004. A QTL that
13 enhances and broadens Bt insect resistance in soybean. *Theor Appl Genet* 109: 1051-1057.

14 Walker, D.R., J.N. All, R.M. McPherson, H.R. Boerma & W.A. Parrott, 2000. Field
15 evaluation of soybean engineered with a synthetic *cryIAC* transgene for resistance to corn
16 earworm, soybean looper, velvetbean caterpillar (Lepidoptera: Noctuidae), and lesser
17 cornstalk borer (Lepidoptera: Pyralidae). *J Econ Entomol* 93: 613-622.

18 Wang, H., S.F. Hwang, K.F. Chang, G.D. Turnbull & R.J. Howard, 2003. Suppression of
19 important pea diseases by bacterial antagonists. *Biocontrol* 48: 447-460.

20 Wang, X.Y., A.L. Eggenberger, F.W. Nutter & J.H. Hill, 2001. Pathogen-derived transgenic
21 resistance to soybean mosaic virus in soybean. *Mol Breed* 8: 119-127.

22 Webb, K.J., L. Skot, M.N. Nicholson, B. Jorgensen & S. Mizen, 2000. *Mesorhizobium loti*
23 increases root-specific expression of a calcium-binding protein homologue identified by
24 promoter tagging in *Lotus japonicus*. *Mol Plant-Microbe Interact* 13: 606-616.

1 Wu, Q.D. & H.D. VanEtten, 2004. Introduction of plant and fungal genes into pea (*Pisum*
2 *sativum* L.) hairy roots reduces their ability to produce pisatin and affects their response to
3 a fungal pathogen. Mol Plant-Microbe Interact 17: 798-804.

4 Yamaguchi-Shinozaki, K. & K. Shinozaki, 2005. Organization of cis-acting regulatory
5 elements in osmotic- and cold-stress-responsive promoters. Trends Plant Sci 10: 88-94.

6 Yang, H., J.G. Boersma, M.P. You, B.J. Buirchell & M.W. Sweetingham, 2004. Development
7 and implementation of a sequence-specific PCR marker linked to a gene conferring
8 resistance to anthracnose disease in narrow-leaved lupin (*Lupinus angustifolius* L.). Mol
9 Breed 14: 145-151.

10 Yang, H., M. Shankar, B.J. Buirchell, M.W. Sweetingham, C. Caminero & P.M.C. Smith,
11 2002. Development of molecular markers using MFLP linked to a gene conferring
12 resistance to *Diaporthe toxica* in narrow-leaved lupin (*Lupinus angustifolius* L.). Theor
13 Appl Genet 105: 265-270.

14 Ye, G., D.L. McNeil & G.D. Hill, 2002. Breeding for resistance to lentil Ascochyta blight.
15 Plant Breed 121: 185-191.

16 Yoshida, M., S.E. Cowgill & J.A. Wightman, 1997. Roles of oxalic and malic acids in
17 chickpea trichome exudate in host-plant resistant to *Helicoverpa armigera*. J Chem Ecol
18 23: 1195-1210.

19 Yu, K.F., S.J. Park, B.L. Zhang, M. Haffner & V. Poysa, 2004. An SSR marker in the nitrate
20 reductase gene of common bean is tightly linked to a major gene conferring resistance to
21 common bacterial blight. Euphytica 138: 89-95.

22 Yue, W., G.M. Xia, D.Y. Zhi & H.M. Chen, 2001. Transfer of salt tolerance from *Aeluropus*
23 *littoralis* sinensis to wheat (*Triticum aestivum* L.) via asymmetric somatic hybridization.
24 Plant Sci 161: 259-266.

- 1 Zagorska, N., B. Dimitrov, P. Gadeva & P. Robeva, 1997. Regeneration and characterization
2 of plants obtained from anther cultures in *Medicago sativa* L. In Vitro Cell Dev Biol-Plant
3 33: 107-110.
- 4 Zair, I., A. Chlyah, K. Sabounji, M. Tittahsen & H. Chlyah, 2003. Salt tolerance improvement
5 in some wheat cultivars after application of in vitro selection pressure. Plant Cell Tissue
6 Organ Cult 73: 237-244.
- 7 Zambre, M., A. Goossens, C. Cardona, M. Van Montagu, N. Terryn & G. Angenon, 2005. A
8 reproducible genetic transformation system for cultivated *Phaseolus acutifolius* (tepary
9 bean) and its use to assess the role of arcelins in resistance to the Mexican bean weevil.
10 Theor Appl Genet 110: 914-924.
- 11 Zhang, H.X. & E. Blumwald, 2001. Transgenic salt-tolerant tomato plants accumulate salt in
12 foliage but not in fruit. Nat Biotechnol 19: 765-768.
- 13 Zhang, H.X., J.N. Hodson, J.P. Williams & E. Blumwald, 2001. Engineering salt-tolerant
14 Brassica plants: Characterization of yield and seed oil quality in transgenic plants with
15 increased vacuolar sodium accumulation. Proc Natl Acad Sci USA 98: 12832-12836.
- 16 Zhang, X.H., W.Q. Zhong & J.M. Widholm, 2005. Expression of a fungal cyanamide
17 hydratase in transgenic soybean detoxifies cyanamide in tissue culture and in planta to
18 provide cyanamide resistance. J Plant Physiol 162: 1064-1073.
- 19 Zivy, M. & D. de Vienne, 2000. Proteomics: a link between genomics, genetics and
20 physiology. Plant Mol Biol 44: 575-580.

Figure 1. Integrated scheme outlining key steps for plant molecular breeding using biotechnology. VIGS: Virus Induced Gene silencing; RNAi: RNA interference; TILLING: Targeted Induced Lesion IN Genome; TF: Transcription Factors; SAGE: Serial Analyses of Gene expression; SSH: Suppression Subtractive Hybridization; DH: Doubled haploid.

Table 1: List of some QTL identified in important legume crops associated to biotic stresses

Legume	Biotic stresses	Gene(s)/QTL	Associated markers	Marker type	Referente(s)
<i>Cicer arietinum</i>	<i>Ascochyta rabiei</i>	<i>Ar19</i>	GA20 GA24	STMS STMS	Cho et al. (2004)
	<i>Fusarium oxysporum</i> f. sp. <i>cicer</i>		TR59	STMS	Cobos et al. (2005)
<i>Lens culinaris</i>	<i>Ascochyta lentis</i>	<i>ral2</i> <i>AbR1</i>	UBC227 ₁₂₉₀ and OPD10 ₈₇₀ <i>RB18</i> and <i>RV01</i>	RAPD RAPD	Tar'an et al. (2003b)
	<i>Colletotrichum truncatum</i>		OPD6 ₁₂₉₀	RAPD	Tar'an et al. (2003a)
	<i>Fusarium oxysporum</i> f. sp. <i>lentis</i>	<i>FW</i>	<i>OPK15900</i>	<i>RAPD</i>	Eujayl et al. (1998)
<i>Lupinus angustifolius</i>	<i>Diaporthe toxica</i>	<i>Phr1</i>	Ph258M2	SCAR	Yang et al. (2002)
	<i>Colletotrichum gloeosporioides</i>	<i>Lanr1</i>	AntjM1	SCAR	Yang et al. (2004)
<i>Vicia faba</i>	<i>Orobanche crenata</i>	<i>Oc1</i>	OPJ13 ₆₈₆ / OPAC02 ₇₃₀	RAPD	Roman et al. (2002)
		<i>Oc2</i>	OPAC06 ₃₄₂ / OPN07 ₈₄₉	RAPD	
		<i>Oc3</i>	OPAA07 ₈₀₇ / OPAA07 ₈₀₇	RAPD	
	<i>Ascochyta fabae</i>	<i>Af1</i>	OPA11 ₁₀₄₅ / OPAB07 ₁₀₂₆	RAPD	Roman et al. (2003)
		<i>Af2</i>	OPE17 ₁₂₇₂ / OPJ18 ₆₂₆	RAPD	
	<i>Uromyces viciae-fabae</i>	<i>Uvf-1</i>	OPI20 ₉₀₀	RAPD	Avila et al. (2003)
<i>Vigna unguiculata</i>	<i>Striga gesnerioides</i>	<i>Rsg1</i>	SEACTMCAC83/85	SCAR	Boukar et al. (2004)
		<i>Rsg3</i>	E-AGA/M-CTA ₄₆₀	AFLP	Ouedraogo et al. (2002)
		<i>994-Rsg</i>	E-AAG/M-AAC ₄₅₀		

RAPD: Random Amplified Polymorphism DNA; RFLP: Restriction Fragment Length Polymorphism; AFLP: Amplified Fragment Length Polymorphism; SCAR: Sequence Characterized Amplified Region; STMS: Sequence Tagged Microsatellite Site; SSR: Simple Sequence Repeat; STS: Sequence Tagged Site; EST: Expressed Sequence Tag.

Table 1 (continuation). List of some QTLs identified in important legume crops associated to biotic stresses

Legume target	Biotic stresses	Gene(s)/QTL	Associated markers	Marker type	Reference(s)
<i>Phaseolus vulgaris</i>	<i>Xanthomonas axonopodis</i> pv. <i>phaseoli</i>		D1287	SCAR	Kelly et al. (2003)
			BNG71	RFLP	
			BNG21	RFLP	
			G19.1800; <i>PHVPVPK-1</i>	RAPD; SSR	Tar'an et al. (2001)
			PV-tttc001; BNG060	SSR; RFLP	
			BNG191	RFLP	Yu et al. (2004)
	<i>Fusarium solani</i> f. sp. <i>phaseoli</i>	<i>PvPR2</i> <i>PvPR1</i>	SU91 / R7313	SCAR	Miklas et al. (2003)
			AI07.600	RAPD	
			BnG154; BNG25a	RFLP; RFLP	
			P7 ₇₀₀	RAPD	
<i>Pisum sativum</i>	<i>Empoasca fabae</i> <i>Empoasca kraemeri</i>		S8 ₅₀₀	RAPD	Schneider et al. (2001)
			G3 ₂₀₀₀	RAPD	
			PV-atcc003	SSR	Murray et al. (2004)
	<i>Sclerotinia sclerotiorum</i>	<i>fin</i> <i>Phs</i>	U73	RFLP	
			O15.1800	RAPD	Kolkman & Kelly (2003)
			PR-aggcctt85	SSR	
	<i>Orobanche crenata</i>	<i>Ocp1</i> <i>Ocp2</i>	P482	STS	Valderrama et al. (2004)
			OPB11 ₅₄₁	RAPD	
	<i>Erysiphe pisi</i>	<i>er</i>	ScOPD-10 ₆₅₀	SCAR	Janila & Sharma (2004)
	<i>Mycosphaerella pinodes</i>		ccta2	SSR	Tar'an et al. (2003b)
			cccc1	SSR	
			acct1	SSR	
	Pea Seed-Borne Mosaic Virus (PSbMV)	<i>sbm-1</i> <i>sbm-2</i>	e1F4E <i>e1F(iso)4E</i>		Gao et al. (2004)

Table 2: List of major QTLs identified in important legume crops associated to abiotic stress. Avreviation of marker type are as defined for Table 1.

Legume target	Abiotic stress	Gene(s)/ QTL(s)	Associated Markers	Marker type	Reference(s)
<i>Lens culinaris</i>	Cold	<i>Frt</i>	OPS16 ₇₅₀	RAPD	Eujayl et al. (1999)
	Winter hardiness		ubc808_3_ubc807_3	SSR	Kahraman et al. (2004)
			ubc840_3	SSR	
			cs48_1	RAPD	
			ubc808_12	SSR	
			E3M3	AFLP	
<i>Glycine max</i>	Manganese toxicity		BARC_SATT318	SSR	Kassem et al. (2004)
			BARC_Sat092	SSR	
			BARC_SATT305	SSR	
			BARC_SATT239	SSR	
			OE02 ₁₀₀₀	RAPD	
	Salt stress		Sat-091	SSR	Lee et al. (2004)
	Waterlogging		Sat-064	SSR	VanToai et al. (2001)
	Phosphorus deficiency	<i>fsw1</i>	L37_2I_Sat_36a	SSR	Li et al. (2005)
		<i>fsw2 and rp1</i>	K4-2V	RFLP	
		<i>fsw3</i>	B212T	EST	
		<i>rp2</i>	Satt114_Satt334	SSR	
		<i>lp1</i>	Satt252_Satt269	SSR	
		<i>lp2</i>	Satt586-gmrvbp	SSR	
<i>Medicago sativa</i>	Aluminium toxicity		UGAc471_UGAc502	RFLP	Sledge et al. (2002)

Table 3 List of some legumes genetically engineered for biotic stress

Legume target	Biotic stress	Gene(s)	Reference(s)
<i>Arachis hypogaea</i>	Tomato spotted wilt virus (TSWV)	Nucleocapsid from TSWV	Magbanua et al. (2000)
	<i>Sclerotinia minor</i>	Oxalate Oxidase from barley	Livingstone et al. (2005)
<i>Cajanus cajan</i>	<i>Spodoptera litura</i>	<i>cry I E-C</i>	Surekha et al. (2005)
<i>Cicer arietinum</i>	<i>Helicoverpa armigera</i>	<i>cryIAc</i>	Sanyal et al. (2005)
	<i>Callosobruchus maculatus</i>	Alpha-amylase inhibitor from bean	Sarmah et al. (2004)
	<i>Callosobruchus chinensis</i>		
	<i>Sclerotinia sclerotiorum</i>	Germin (gf-2.8) fom wheat	Donaldson et al. (2001)
<i>Glycine max</i>	Bean Pod Mottle Virus (BPMV)	Capsid polyprotein from BPMV	Reddy et al. (2001)
	<i>Heterodera glycines</i>	Chitinase from <i>Manduca sexta</i>	Ornatowski et al. (2004)
	<i>Helicoverpa zea</i>	<i>cryIAc</i>	Walker et al. (2000)
	<i>Anticarsia gemmatalis</i>		
	<i>Pseudoplusia includens</i>		
	<i>Heterodera glycines</i>	Promoter of chalcone synthase from Promoter of Phenylalanine Ammonia Lyase	Narayanan et al. (1999)
	Soybean Mosaic Virus (SMV)	Coat protein gene from SMV	Wang et al. (2001)
<i>Medicago sativa</i>	<i>Phoma medicaginis</i>	Resveratrol synthase from <i>A. hypogaea</i>	Hipskind & Paiva (2000)
	<i>Phoma medicaginis</i>	Isoflavone <i>O</i> -Methyltransferase	He & Dixon (2000)
	<i>Phoma medicaginis</i>	Endochitinase (ech42)	Samac et al. (2004)
	<i>Pratylenchus penetrans</i>	Oryzacystatin I and II	Samac & Smigocki (2003)
<i>Medicago truncatula</i>	Alfalfa Mosaic Virus (AMV)	Virus coat protein from AMV	Jayasena et al. (2001)
<i>Phaseolus acutifolius</i>	<i>Zabrotes subfasciatus</i>	Arcelins-1, Arcelins-5	Zambre et al. (2005)
<i>Phaseolus vulgaris</i>	Bean Golden Mosaic Virus (BGMV)	<i>Rep-TrAP-REn</i> , <i>BCI</i> (viral genes)	Aragão et al. (1998)
<i>Pisum sativum</i>	<i>Bruchus pisorum</i>	Alpha-amylase inhibitor (alpha-AI-1)	Schröder et al. (1995) de Sousa-Majer et al. (2004)
	Pea Seed-borne Mosaic Virus (PSbMV)	Replicase (<i>NIb</i>) from PSbMV	Jones et al. (1998)
	Alfalfa Mosaic Virus (AMV)	Coat protein from AMV	Timmerman-Vaughan et al. (2001)