Woody plant encroachment following grazing abandonment in the subalpine belt. A case study in Northern Spain

Yasmina Sanjuána, José Arnáez^b, Santiago Beguería^c, Noemí Lana-Renault^{b,*}, Teodoro Lasanta^a, Amelia Gómez-Villard^d, Javier Álvarez-Martínez^e, Paz Coba-Pérez^f, José M. García-Ruiz^a

^a Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC)

^b Área de Geografía Física, Universidad de La Rioja

^c Estación Experimental de Aula Dei, CSIC, Campus de Aula Dei

^d Departamento de Geografía y Geología, Facultad de Filosofía y Letras, Universidad de León

^e Departamento de Ingeniería Agrícola y Forestal, Universidad de Valladolid

^f Centro de Investigaciones en Geografía Ambiental, CIGA, Universidad Nacional Autónoma de México

Yasmina Sanjuán ysanjuan@ipe.csic.es

José Arnáez jose.arnaez@unirioja.es

Santiago Beguería santiago.begueria@csic.es

Noemí Lana-Renault noemi-solange.lana-renault@unirioja.es

Teodoro Lasanta fm@ipe.csic.es

Amelia Gómez-Villar agomv@unileon.es

Javier Álvarez-Martínez gestion@iaf.uva.es

Paz Coba-Pérez pcoba@cieco.unam.mx

José M. García-Ruiz humberto@ipe.csic.es

*Corresponding author: N. Lana-Renault. Área de Geografía Física, Departamento de Ciencias Humanas, Universidad de La Rioja. C/ Luis de Ulloa 2, 26004-Logroño (Spain).

E-mail: noemi-solange.lana-renault@unirioja.es. Telf: +34 941299307.

Length of the manuscript: 8603 words (including title, names of authors + affiliations, abstract, main text, figure captions, figures and tables)

Abstract The montane and subalpine belts in European mountains were affected by intense land use/land cover changes during the 20th century. In the case of the subalpine belt, most European mountains were affected by complex deforestation processes from Neolithic times, leading to the expansion of summer grasslands to support grazing by transhumant sheep and goats flocks. This resulted in an altitudinal reduction of the tree line, and the occurrence of severe soil erosion and shallow landsliding. The intense livestock pressure over centuries explains why the landscape remained without major change until the middle of the 20th century. Since then, depopulation, land abandonment, and the disappearance of the transhumant system have resulted in a marked decline in livestock numbers, and subsequent encroachment of shrubs and trees in the subalpine belt. The Urbión Mountains (Iberian Range, northern Spain) provides one of the clearest examples in the Iberian Peninsula, where there was intense deforestation since the Neolithic period to enlarge the area of subalpine summer grasslands. The recent reversal of this situation in the last decades led to (i) a marked trend to dense forest, which resulted in an average 200 m altitudinal advance; and (ii) the spatial contraction of shrublands, which have been replaced by dense forest in the highest areas. Changes will continue in the future, possibly enhanced by global warming.

Keywords: Woody encroachment, Subalpine belt, Deforestation, Land use/Land cover changes, Land abandonment

Introduction

European mountains have been subject to intense land use changes during the last two centuries (MacDonald et al. 2000; Tasser and Tappeiner 2002; Poyatos et al. 2003; Vicente-Serrano et al. 2005; Améztegui et al. 2010; García-Ruiz et al. 2010; Arnáez et al. 2011; García-Ruiz and Lana-Renault 2011; Gouveia et al. 2016) because of population changes, and societal and economic interactions with the lowlands. In most Mediterranean mountains, the highest population density was recorded in the middle of the 19th century, when hillslopes and valley bottoms supported an integrated land use system aimed at ensuring the survival of local populations through cultivation of all

possible areas. Transhumance of large sheep flocks forced the transformation of the upper montane and subalpine belts into grasslands, to enable the enlargement of summer grazing areas (García-Ruiz & Lasanta, 1990). Consequently, in the middle of the 19th century, most European mountains were under extreme human pressure including: (i) deforestation of most slopes above 1500 m a.s.l. to enlarge the area covered by subalpine grasslands; (ii) frequent use of fire to control shrub and forest expansion; (iii) the presence of large (in some cases transhumant) sheep flocks; and (iv) the occurrence of a variety of erosion processes (sheet wash, rilling and gullyng, landslides) (Höllermann 1985; del Barrio and Puigdefábregas 1987; Tasser et al. 2003; García-Ruiz et al. 2010, 2015; Lana-Renault et al. 2011; García-Ruiz 2015); these erosion processes were responsible for fluvial channel aggradation (Gómez-Villar 2014) and alluvial fan enlargement (Gómez-Villar et al. 2006). In the Mediterranean and central European mountains, the subalpine belt had been deforested since Prehistoric times, as confirmed by the presence within the soil or in peat bogs of charcoal from the Neolithic period and the Bronze and Iron ages (e.g., Colombaroli et al. 2008; Guiguet-Covex et al. 2011; Cunill et al. 2012; Lasheras-Álvarez et al. 2013; Pérez-Sanz et al. 2013). This was also the case for the Iberian Range (northern Spain), where the subalpine belt of the Urbión Mountains has been deforested since the Neolithic (García-Ruiz et al. 2016). This process caused a marked lowering of the altitude of the tree line below its natural elevation, which is estimated from other mountains of the Iberian Peninsula to have been an average of 400 m (e.g., Höllermann 1985; García-Ruiz et al. 1990).

Depopulation and land abandonment have been the most significant changes affecting Mediterranean mountains since the mid 20th century, although it was far from uniform. The peak of depopulation in the central and west European mountains occurred at the beginning of the 20th century (Estienne, 1989; Diry 1995). However, in many Mediterranean mountains, particularly on the Iberian Peninsula, depopulation was a slow process from the end of the 19th century, followed by a rapid acceleration that peaked during the 1960s and 1970s, in parallel with a dramatic process of farmland abandonment (Lasanta 1990; Vitte 1992). Since that time, most of the cultivated hillslopes have been abandoned, with the only remaining cultivated areas being the valley bottoms, large alluvial fans, and some perched flats where cereal crops have been replaced by mown meadows (García-Ruiz and Lasanta 1990). The almost total disappearance of transhumant flocks has led to the abandonment of many summer subalpine pastures. Many aspects of mountain Mediterranean landscapes are changing, including runoff

1 generation (Beguería et al. 2003; Viviroli et al. 2003; García-Ruiz et al. 2011), soil
2 erosion and landsliding (Bathurst et al. 2007; García-Ruiz et al. 2010), fluvial dynamics
3 (Liébault et al. 2005), the spatial organization of plant cover, and the ecosystem services
4 that the area provides to local communities and nationally (Egoh et al. 2007; Egarter Vigl
5 et al 2016). Major changes have been detected in the Alps (e.g. Tasser and Tappeiner
6 2002) and in the Pyrenees (e.g. Lasanta-Martínez et al. 2005; Lasanta and Vicente-
7 Serrano 2007; Gartzia et al. 2014). Nevertheless, changes in land cover of the subalpine
8 belt have not been so intensively studied, although they could be extremely important
9 from the point of view of snow accumulation and water resources (López-Moreno 2005).

10 The Urbión Mountains have a long history of landscape transformation. The
11 development of sheep transhumance was extraordinary, and led to enlargement of the
12 subalpine grasslands, generalized deforestation, and the occurrence of intense landsliding
13 at altitudes above 1500 m a.s.l. (García-Ruiz et al. 2016). The recent and rapid decline of
14 the transhumant system, with the consequent decline in sheep flocks, has resulted in rapid
15 encroachment by shrubs and forest. In this study we investigated this process in the
16 Urbión Mountains, and more precisely in the Ormazal Valley. The study areas is an
17 example of what is happening in the subalpine belt of mountains of northern Spain, with
18 the aim of identifying those factors explaining the spatial complexity of woody
19 encroachment during the period 1956-2006.

20 Study area

21 The Urbión Mountains is one of the highest massifs (peaking at 2228 m) of the
22 Iberian Range, in the northern Iberian Peninsula (30N 514118 4657163 m). The massif
23 is mainly composed of Upper Jurassic and Lower Cretaceous small quartz conglomerate,
24 red mudstone, and limestone arranged in a monoclonal relief.

25 The study was carried out in the Ormazal Valley, which has an area of 2513.8 ha
26 and an altitudinal range of 1240–1887 m a.s.l. The landscape is heterogeneous, ranging
27 from the lower montane belt (cultivated and abandoned fields) to the subalpine belt
28 (grasslands and forests). The valley has relatively gentle slopes (in general, <30%) and
29 wide divides.

30 Weather stations are located on the southern face of the massif, at approximately
31 1100–1150 m a.s.l. A temperature increase has been recorded in the last 60 years; thus,
32 the mean annual temperature at Vinuesa was 7.7°C for the period 1957–1965, and 9.6°C
33

for the period 1990–2001, with an average increase in winter of 1.3°C (García de Celis et al. 2008). Mean annual precipitation exceeds 900 mm above 1000 m a.s.l., and is probably 1500–1600 mm in the main divide, based on an estimate for the neighboring Demanda Sierra (Arnáez Vadillo 1987). Precipitation has declined slightly since the mid 20th century, particularly between November and March (García de Celis et al. 2008).

The plant cover is dominated by extensive *Pinus sylvestris* forests on the south-facing slopes, whereas the northern faces are complex landscapes involving *P. sylvestris* and *Fagus sylvatica* forests, grasslands, and shrubs including *Calluna vulgaris*, *Erica cinerea*, *Vaccinium myrtillus*, and *Juniperus communis*. The area has been used in summer for transhumant sheep herds, with two periods of particularly intense activity: the 15th century, and the period between the end of the 17th and the end of the 18th centuries. In 1792 two villages located close to the Ormazal Valley (Viniegra de Abajo and Viniegra de Arriba) accounted for 17,169 (262 per km²) and 19,100 (497 per km²) sheep, respectively (Moreno Fernández 1996). The number of sheep has declined, and now accounts for 840 and 2690, respectively. Population has also declined in both villages: Viniegra de Abajo had 689 inhabitants in 1842, and 97 in 2011; Viniegra de Arriba had 418 inhabitants in 1842, and 49 in 2011. Below 1500 m a large part of the hillslopes was cultivated with cereals, although most of fields were abandoned by the middle of the 20th century, except for some fields located in the valley bottoms close to the villages.

Methods

Aerial photographs of the Ormazal Valley from 1956 (black and white) and ortho-photographs from 2006 (color), all with a scale of 1:18,000, were used to analyze the evolution of land use and land cover (LULC). In order to reduce methodological inconsistencies in the visual interpretation of two qualitatively different sets of images, a first LULC map was produced based on the best set (2006). A second LULC map was then produced for 1956 using the first one as a base layer, so the interpretation was focused in the changes occurred with respect to this base map. Two maps (one for 1956 and one for 2006) were thus generated, including the following LULC classes: (i) cultivated fields; (ii) abandoned fields; (iii) grasslands; (iv) shrublands; (v) clear forest (where voids up to 20 m occur between trees); (vi) dense forest (where the canopy is continuous, without voids between trees); (vii) eroded areas; (viii) rock outcrops and

debris accumulations; and (ix) reforestation areas. Cultivated and abandoned fields were clearly discriminated visually due to the existence of linear features delimiting the fields. Grasslands, shrublands and forests were also clearly identified, as were rock outcrops and debris accumulations. The distinction between clear and dense forests was somehow more subjective, so reference areas were first set and used therefore as discriminating patterns. Delimitation of eroded areas within the grasslands was the most difficult due to their small size and diffuse spatial limits, although reference areas were also set and used for discrimination.

The 1956 and 2006 LULC maps were overlapped to assess changes that occurred between the two years. Measurement of the areas where changes or no changes had occurred enabled computation of a LULC transition matrix.

To determine the influence of environmental factors on the changes observed in LULC categories, a set of topographic variables was derived from a digital elevation model (Regional Government; <http://iderioja.larioja.org/>) having a spatial resolution of 18×18 m and using ArcGIS Desktop. The variables included: (i) altitude; (ii) slope gradient; (iii) slope aspect; (iv) distance to the hydrographic network; (v) slope shape (curvature); vi) solar irradiation; and vii) the topographic index (Beven & Kirkby, 1979).

Discriminant analysis was used to interpret the influence of the variables on the main LULC transitions observed. This involved conversion of the LULC maps to raster format at the same resolution as the topographic data. Discriminant analysis provided information on the topographic variables (independent variables), which enabled differentiation of the groups of pixels classified from the LULC transitions (dependent variables). In total, 16 changes representing 96.3% of all changes recorded were selected for analysis.

Multinomial logistic regression analysis was used to explain the transition of grasslands and shrublands (the two LULC classes most abundant in 1956, and those that experienced the highest rates of change) to other LULC categories. Multinomial logistic regression facilitated calculation of the probabilities of permanence (no LULC change) and change, as a function of covariates (independent variables). The topographic variables selected as covariates were elevation, slope gradient, solar irradiation, and topographic index. The significance of each covariate as an explanatory factor was assessed using the *t*-test at a significance level of $\alpha = 0.05$, and the significance of the overall model was assessed using the likelihood ratio test. The calculations were

performed using the package `mlogit` version 0.2-4 for the R environment (Croissant, 2003). The results were expressed as expected transition probabilities corresponding to average topographic conditions, and as changes in these probabilities as a function of unit changes in the topographic variables. Maps were computed from the multinomial models to visually assess the spatial variability of transition probabilities.

Results

Spatial distribution and changes in LULC between 1956 and 2006

The maps in Figure 1 show the spatial distribution of the LULC categories in 1956 and 2006, and the areas covered by each LULC category in the two years is shown in Table 1. By 1956, some cultivated fields (0.4% of the total study area) remained in the lowest part of the catchment (1300–1500 m a.s.l.), adjacent to fields that had been cultivated some decades previously and then abandoned (58.2 ha; 2.3%). The main LULCs corresponded to grasslands (910.6 ha; 36.2%) and shrublands (853.2 ha; 33.9%). Dense forests occupied only 43.6 ha (1.7%), mainly in the northernmost part of the catchment (small patches of *F. sylvatica* forest), and clear forests (*Quercus pyrenaica* and *P. sylvestris*) occupied 231.1 ha (9.2%), scattered throughout shrublands areas. Eroded areas were generally located over bare limestone thinly covered by very open grasslands, and occupied 405.3 ha (16.1%).

Substantial changes had occurred by 2006. (i) The expansion of dense forests at the expense of the other LULC categories (particularly grasslands and shrublands) was the main change that occurred in the study area between 1956 and 2006, while the eroded area remained relatively constant. (ii) Cultivated fields had been largely abandoned. (iii) Grasslands were still the most extensive land cover (682.0 ha; 27.1% of the total), although there had been marked spatial shrinkage. (iv) Shrublands were the second largest land cover (651.4 ha; 25.9%), although they also had declined in area. (v) The clear forest area had decreased slightly, occupying 174.2 ha (6.9%) in 2006, and the area of dense forests had remarkably increased to 436.4 ha (17.4%). (vi) A small enlargement of the dense *F. sylvatica* forests was also detected in the north of the catchment. (vii) The eroded areas remained relatively constant in 2006, with 438.4 ha (17.4%). (viii) Major changes had occurred in the southernmost part of the study area, where shrublands and clear forests were substituted by a dense forest of *P. sylvestris*. A close spatial relationship was found between the locations of summer grasslands and the eroded areas.

1 The transition matrix (Table 2) shows the changes that occurred in the area
2 occupied by each LULC category. The most outstanding results shown in this table are
3 described below.
4

5 (i) The average rate of change (approximately 50%) was very high, confirming that the
6 area was affected by highly dynamic changes in LULC categories.
7

8 (ii) Dense forest can be considered a stable or absorbing state of the system, as shown by
9 the high rate of permanence of this LULC category (94.5%).
10

11 (iii) For grasslands, 52.5% maintained their status, with the main transitions being to
12 eroded areas (20.1%) and shrublands (17.3%).
13

14 (iv) For eroded areas, 54.8% remained in the same category, and their most common
15 transition was to grasslands. This indicates that a reversible transition occurred between
16 grasslands and eroded areas.
17

18 (v) Shrublands represent a clear transition state. Their permanence rate was relatively low
19 (43.2%), and their transition tended to be to forest (39.4%, including the two forest types).
20

21 (vi) Abandoned fields showed a relatively high rate of permanence (60.4%), with notable
22 transition to grasslands (18.8%) and dense forests (12%).
23

24 (vii) The lowest rate of permanence corresponded to clear forest (12.3%), which
25 transitioned to dense forest (40.9%) and shrublands (32.4%).
26

27 (viii) In 2006 a new category of LULC (reforestation) was identified; this category mainly
28 incorporated former clear forests (12.6%) and shrublands (4.3%).
29

30 **Factors controlling LULC transitions**

31 Figures 2 and 3 show the frequency distribution of four topographic variables
32 (altitude, slope gradient, solar irradiation, and topographic index) for shrublands and
33 forest in 1956 and 2006. It is noteworthy that neither summer grasslands nor eroded areas
34 significantly changed their 'topographic signature' between the two years, other than a
35 small increase in altitude in the case of summer grasslands. Conversely, shrublands (Fig.
36 2) and dense forests (Fig. 3) showed major changes. Thus, while shrublands tended to be
37 relegated to areas of lower altitude, higher slope gradient, and lower levels of irradiation,
38 the dense forests colonized higher altitudes, shallower gradients, and areas affected by
39 higher levels of solar irradiation. This suggests that the substitution of shrublands by
40 dense forests was markedly influenced by topographic factors. The average altitude for
41 dense forest was 1448 m in 1956 and 1638 m in 2006 (an altitudinal advance of almost
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

200 m). In the case of shrublands, the average altitude was 1627 m in 1956 and 1575 m in 2006.

To assess the influence of topographic variables on the main LULC changes we performed a discriminant analysis. The change from shrublands to dense forest was the largest in area (15.4%), followed by the changes from summer grasslands to eroded areas (12.7%), from summer grasslands to shrublands (10.3%), from shrublands to clear forest (8.3%), from eroded areas to summer grasslands (8.3%), and from clear forest to dense forest (7.2%). The discriminant analysis identified three functions that differentiated the selected LULC transitions. An analysis of the contrast of the functions through the Wilks' Lambda and the Chi-square analysis showed a significance level < 0.05 , indicating that the model could distinguish among the LULC transition groups.

Figure 4 shows the results of the discriminant analysis and enables interpretation of the topographical factors explaining most of the major LULC changes in the study area. Function 1 (horizontal axis), strongly correlated with the altitude, clearly discriminated changes in the agricultural areas (cultivated fields to abandoned fields, and abandoned fields to summer grasslands), with negative values for the centroids, and discriminated changes to shrublands, clear forest, and summer grasslands, which became dense forest (positive values). Function 2 (vertical axis) was defined by slope aspect, and to a lesser extent by the slope gradient. This function discriminated changes from clear forest to eroded areas (positive values), opposed to changes recorded in grasslands (summer grasslands to clear forest and dense forest). Based on these results, LULC changes in the Urbión Mountains between 1956 and 2006 were influenced by topographic variables. The discriminant analysis indicated that the previously cultivated areas, located at the lowest altitudes (1300–1400 m), were in 2006 occupied by abandoned fields and summer grasslands. Conversely, at altitudes of 1600–1700 m the shrublands had transitioned to eroded areas, or more commonly to dense forest. Areas of summer grassland and clear forest in 1956 transitioned to dense forest in 2006. Solar irradiation (which depends on slope aspect) and hillslope gradient also discriminated transitions in LULC, particularly the colonization of summer grasslands by dense forest (preferentially on hillslopes having gradients of 14–15° and southwestern exposures), and the degradation of clear forest to eroded areas (on gradients of almost 20° and eastern exposure).

Application of multinomial logistic regression analysis to the areas covered by grasslands in 1956 enabled estimation of the permanence and probability of transition, as

1 a function of topographic covariates. The model was significant overall (likelihood ratio
2 test p-value < 0.05), indicating a degree of topographic control, although only the slope
3 gradient and solar irradiation were found to be significant according to the t-test; the other
4 topographic variables (elevation, topographic index) were not significant and thus
5 removed from the model.
6
7

8
9 The transition probabilities were consistent when compared with the observed
10 transition matrix (Table 3), although the probability of permanence was slightly
11 overestimated (62% vs. 53%), and the probability of transition to eroded areas was
12 underestimated (16% vs. 20%). A slight overestimation of the permanence rate was
13 expected, as it was the reference level with which the other transitions were compared.
14
15
16

17
18 The influence of topographic variables on the transitions from grassland is
19 summarized in Table 4. With respect to the transition to forest, each increase of 10° in
20 slope gradient resulted in an increase of 3.7% in the probability of transition, while an
21 increase of 200 kWh m⁻² yr⁻¹ in solar irradiation reduced the probability by 5.6%. For the
22 transition to eroded areas and shrublands, increasing slope gradient and solar irradiation
23 had stronger positive effects than occurred for forests.
24
25
26
27

28
29 Application of the resulting model to the spatially-explicit topographic variables
30 enabled computation of transition probability maps (Fig. 5). For each grid cell that was
31 occupied by grasslands in 1956, these show the probability of grassland permanence and
32 transition to another LULC category by 2006, as estimated using the regression model.
33 The spatial distribution of transition probabilities varied in accordance with variation in
34 the topographic characteristics.
35
36
37
38
39

40 Application of the same analysis to the areas covered by shrublands in 1956
41 showed an even stronger correlation of permanence and transition probabilities with the
42 observed transition rates (Table 3). Over-prediction of the probability of permanence was
43 reduced (48% vs. the observed rate of 43%), and only the probability of transition to
44 forest was slightly over-predicted (42% vs. 39%). The topographic conditions were only
45 significant for the transition to forest, but revealed a strong influence. For instance, every
46 change of 10° in the slope gradient resulted in an 11.4% increase in the probability of
47 transition to forest, while an increase of 200 kWh m⁻² yr⁻¹ in irradiation decreased the
48 probability by 7.7% (Table 4). The transition probability maps (Fig. 6) reflect this strong
49 topographic control. For instance, the probability of shrubland permanence was highest
50 for the mid-altitude slopes in the northwest of the study area, where deforestation and
51 land abandonment had led to the occurrence of shrubs and clear forest. In contrast, the
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 probability of transition from shrublands to forest was highest for the steep and shady
2 slopes in the southernmost part of the study area, and also on the low altitude and shady
3 slopes close to formerly cultivated areas in the northwest of the study area. In both areas
4 there was also a relatively high probability of transition to grasslands. The transition to
5 eroded areas, besides being very low overall (less than 5%), indicates the instability of
6 shrublands in the northwestern sector of the study area, where the steepest slopes are
7 located, and close to areas affected by human-induced reforestation.
8
9
10
11
12
13
14
15

16 Discussion

17
18 Major changes have occurred in the montane and subalpine belts of the European
19 mountains. In the Urbión Mountains, northern Iberian Peninsula, forest clearing began in
20 Neolithic times and has continued to the present (García-Ruiz et al. 2016). Forest clearing
21 favored the expansion of transhumant sheep flocks and caused: (i) altitudinal lowering of
22 the tree line by at least 400 m, and (ii) the triggering of soil erosion processes and shallow
23 landsliding. The latter is based on the presence of large landslide scars and areas with
24 bare soil, which in other mountain areas including the Pyrenees (García-Ruiz et al. 2010)
25 and the Alps (Tasser et al. 2003; Guiguet-Covex et al. 2011) have been attributed
26 unequivocally to deforestation. In the Pyrenees, a palinological study of Tramacastilla
27 Lake (Gállego Valley) confirmed the occurrence of moderate erosion during small
28 deforestation events approximately 4000 cal yr BP (Montserrat-Martí 1992), and intense
29 soil erosion in the subalpine belt since the Middle Ages, following extensive fires that
30 coincided with the maximum period of transhumance.
31
32
33
34
35
36
37
38
39
40
41

42 Dating of charcoal fragments in the soil in the Urbión Mountains has suggested
43 that forest clearing and grazing in the subalpine belt has been a common practice since
44 the Neolithic and Calcolithic periods, and also during the Bronze, Iron, and Middle ages
45 (García-Ruiz et al. 2016). Subsequently, the deforested landscape and the large areas of
46 summer grasslands were maintained because of the presence of thousands of sheep
47 grazing from the beginning of June to the end of September. By the middle of the 20th
48 century, aerial photographs show that most of the study area had been deforested, and
49 only small patches of dense forest remained in some valley bottoms, indicating that the
50 traditional landscape was the consequence of intense human occupation and management.
51 Apart from a small area dedicated to farmland activity, the remainder of the study area
52 showed the effects of grazing, with clear forests, shrublands, summer grasslands, and
53
54
55
56
57
58
59
60
61
62
63
64
65

eroded areas. These latter were located in close spatial proximity (and similar topographic signatures) to summer grasslands, suggesting that soil erosion was a direct consequence of deforestation.

In 2006, 50 years later, the landscape retained some of the original characteristics, although major changes had occurred. The main reason for the changes was the decline in livestock numbers that had occurred in the preceding few decades, and the subsequent decrease in livestock pressure. This led to a relatively complex process of woody encroachment characterized by: (i) the expansion and average 200 m altitudinal advance of dense forests at the expense of other LULC classes, which is consistent with tree lines being highly sensitivity to environmental changes (Holtmeier and Broll 2005; Wallentin et al. 2008; Pecher et al. 2011; Tasser et al. 2017); (ii) the spatial contraction of shrublands, which were substituted by dense forests at the highest altitudes; (iii) shrublands and clear forest transitioning towards dense forest; and (iv) a change in LULC categories affecting approximately 50% of the study area during the study period, confirming that the landscape configuration was a consequence of human pressure, through livestock grazing, and domestic wood consumption.

Interestingly, it was found that a reversible transition occurred between erosional features (mostly due to shallow landsliding) and pastures, implying that the eroded stage was not a final stable state, at least within the time frame of this study. This result, however, has to be taken with care since the distinction between pastures and eroded pastures in the aerial photographs was not devoid of difficulties, especially in the oldest set (1956).

The increase in altitude of the tree line may also be attributed, at least in part, to the recent increase in temperature, as reported by García de Celis et al. (2008) for the southern slopes of the Urbión Mountains. Nevertheless, the irregular distribution of forest and summer grasslands in Ormazal Valley suggests that forest expansion has been largely a consequence of grassland abandonment. The effect of global warming on forest development can only clearly be detected in the natural upper forest limit, following the altitudinal reduction in the tree line that occurred during the Little Ice Age (Camarero et al. 2015).

Woody encroachment in the subalpine belt has also been reported in many other European mountains. For instance, densification of forest near the tree line has been reported in the eastern Pyrenees (Batllori & Gutiérrez 2008), where the tree line has moved upwards by an average of almost 40 m because of land use changes (Améztegui

et al. 2015). In the central Pyrenees, Gartzia et al. (2014) reported the expansion of shrublands and forest at the expense of summer grasslands. These authors noted the important role of the distance to patches of woody species in explaining the pattern of tree recruitment. In the Ijuez Valley, in the central Pyrenees, Sanjuán et al. (2016) noted the expansion of dense pine forest, a decline in the area of subalpine grasslands because of the advance of pine forest towards the upper part of the basin, and a decline in the area occupied by shrubs and clear pine forest. In the eastern French Pyrenees (Ariège Valley), the population and livestock decline after 1950 was followed by an increase in pine and the presence of *Juniperus* pollen in the peat bog of Orry de Théo (Galop et al. 2011). Summer grasslands have been colonized by heathland (*J. communis*, *C. vulgaris*, and *Cytisus scoparius*).

A remarkable trend towards encroachment has been also reported in the Alps, particularly between 1650 and 2450 m, where 10% of the new forest areas correspond to upward shifts in altitude, and 90% is attributable to densification of clear forests (Gehrig-Fasel et al. 2007). In the Unterlptal valley the shrub cover increased by 18.7%, and the shrub cover by 32.6%, resulting in an increase in resistance to soil erosion. Abandonment is also the main factor explaining the regeneration of forests in the Austrian Alps, particularly in the altitudinal range 1500–2150 m (Fondevilla et al. 2016).

Forest expansion will probably continue in the Urbión Mountains, because not all the potential forest areas in the montane and the subalpine belts have been occupied by forest since land abandonment. This expansion will occur in a haphazard manner, controlled by topographic factors. For example, our analyses demonstrated the importance of slope gradient, altitude, and solar irradiation, the latter depending on the slope gradient and aspect.

The presence of sheep flocks and cattle in summer could delay the advance of dense forest into the subalpine grasslands, whereas the advance will occur over several decades in the shrublands and clear forests. Forest expansion will not be possible into the eroded areas (caused by deforestation since the Neolithic period), so a complex mosaic of bare areas and summer grasslands will remain. These changes will have ecological, hydrological, and geomorphological consequences, including: (i) increasing homogenization of the landscape, with larger patches dominated by dense forest surrounding shrinking summer grasslands and eroded areas; (ii) a decrease in the capacity of the massif for runoff generation, as has been demonstrated worldwide in mountains where dense forest has substituted grasslands and cultivated fields (García-Ruiz et al.

2011; García-Ruiz and Lana-Renault 2011), being this the main reason explaining the declining discharge of the rivers in Mediterranean mountains (Beguiría et al. 2003; López-Moreno et al. 2014; Morán-Tejeda et al. 2014); (iii) a decline in soil erosion and shallow landslides, which in turn will result in reduced sediment supply to the streams as confirmed in Slovenian (Keesstra 2007) French Alps (Liébault and Piégay 2001; Piégay et al. 2004) and Pyrenean rivers (Beguiría et al. 2006; Sanjuán et al. 2016); (iv) a reduction in the variety of functions and ecosystem services provided by summer grasslands, given the spatial contraction of grazing areas (Gartzia et al. 2014); and (v) possible changes in the fauna and floristic composition. The consequences of these landscape changes also include the timing of snowmelt in a landscape dominated by dense forest rather than shrublands and summer grasslands. Many uncertainties about the future of the upper montane and subalpine belts remain, and more research is needed to understand the changes that are occurring and their effects. This is the case of the future ecological, geomorphological and hydrological implications of climate change and, above all, of a declining snow accumulation during the cold season because of global change (López-Moreno 2005; López-Moreno et al. 2009; Navarro-Serrano and López-Moreno 2017; Vicente-Serrano et al. 2017). This is particularly interesting in areas that are critical for water resources management at regional and national scales, and for sediment transport to the lowlands.

Concluding remarks

Deforestation of the subalpine belt in the Urbión Mountains occurred since Neolithic times, resulting in the transformation of forests into summer grasslands to feed the transhumant sheep flocks. This caused severe soil erosion because of shallow landsliding. Recent decline in the number of sheep flocks has resulted in changes in land cover reorganization, including:

- (i) The expansion of dense and clear forests at the expense of shrubs and grasslands.
- (ii) The average 200 m altitudinal advance of dense forests.
- (iii) The progressive spatial contraction of shrublands, mainly substituted by dense forest.
- (iv) Forest expansion will continue in the next future since shrublands and grasslands are potential forest areas. Nevertheless, this trend will be conditioned by the

1 evolution of the number of sheep and cattle flocks, because grazing controls woody
2 encroachment.

3 The land use evolution detected in the subalpine belt of other Mediterranean
4 mountains will result in land cover changes with highly relevant ecological, hydrological
5 and geomorphological consequences.
6
7
8
9

10 **Acknowledgements**

11 Support for this research was provided by the projects ESPAS (CGL2015-65569-R) and
12 INDICA (CGL2011-27753-C02-01 and -02), funded by the Spanish Ministry of
13 Economy and Competitiveness.
14
15
16
17
18
19

20 **References**

- 21 Améztegui A, Brotons L, Coll L (2010) Land use changes as major drivers of mountain
22 pine (*Pinus uncinata* Ram.) expansion in the Pyrenees. *Global Ecology and*
23 *Biogeography* 19: 632–641. <http://dx.doi.org/10.1111/j.1466-8238.2010.00550.x>
24
25 Améztegui A, Coll L, Brotons L, Ninot JM (2015) Land-use legacies rather than climate
26 change are driving the recent upward shift of the mountain tree line in the Pyrenees.
27 *Global Ecology and Biogeography*. <http://dx.doi.org/10.1111/geb.12407>
28
29 Arnáez Vadillo J (1987) Formas y procesos en la evolución de vertientes de la Sierra de
30 la Demanda (Sistema Ibérico). *Cuadernos de Investigación Geográfica* 13: 7–153.
31 <http://dx.doi.org/10.18172/cig.vol13iss0>
32
33 Arnáez J, Lasanta T, Errea MP, Ortigosa L (2011) Land abandonment, landscape
34 evolution, and soil erosion in a Spanish Mediterranean mountain region: The case of
35 Camero Viejo. *Land Degradation & Development* 22: 537–550.
36 <http://dx.doi.org/10.1002/ldr.1032>
37
38 Bathurst JC, Moretti G, El-Hames A, Beguería S, García-Ruiz JM (2007) Modelling the
39 impact of forest loss on shallow landslide sediment yield, Ijuez catchment, Spanish
40 Pyrenees. *Hydrology and Earth System Sciences* 11: 569–583.
41 <http://dx.doi.org/10.5194/hess-11-569-2007>
42
43 Batllori E, Gutiérrez E (2008) Regional tree line dynamics in response to global change
44 in the Pyrenees. *Journal of Ecology* 96: 1275–1288. <http://dx.doi.org/10.1111/j.1365-2745.2008.01429.x>
45
46 Beguería S, López-Moreno JI, Lorente A, Seeger M, García-Ruiz JM (2003) Assessing
47 the effects of climate oscillations and land-use changes on streamflow in the Central
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- Spanish Pyrenees. *Ambio* 32: 283–286. <http://dx.doi.org/10.1579/0044-7447-32.4.283>
- Beguiría S, López-Moreno JI, Gómez-Villar A, Rubio V, Lana-Renault N, García-Ruiz JM (2006) Fluvial adjustments to soil erosion and plant cover changes in the Central Spanish Pyrenees. *Geografiska Annaler* 88A: 177–186. <http://dx.doi.org/10.1111/j.1468-0459.2006.00293.x>
- Beven KJ, Kirkby MJ (1979) A physically based, variable contributing area model of basin hydrology. *Hydrological Science Bulletin* 24: 43–69. <http://dx.doi.org/10.1080/02626667909491834>
- Camarero JJ, García-Ruiz JM, Sangüesa-Barreda G, Galván JD, Alla AQ, Sanjuán Y, Beguiría S, Gutiérrez E (2015) Recent and intense dynamics in a formerly static Pyrenean treeline. *Arctic, Antarctic, and Alpine Research* 47: 773–783. <http://dx.doi.org/10.1657/AAAR0015-001>
- Colombaroli D, Vannière B, Emmanuel C, Magny M, Tinner W (2008) Fire-vegetation interactions during the Mesolithic-Neolithic transition at Lago dell'Accesa, Tuscany, Italy. *The Holocene* 18: 679–692. <http://dx.doi.org/10.1177/0959683608091779>
- Croissant Y (2013). mlogit: multinomial logit model. R package version 0.2-4. <https://CRAN.R-project.org/package=mlogit>
- Cunill R, Soriano JM, Bal MC, Pélachs A, Pérez-Obiol R (2012) Holocene treeline changes on the south slope of the Pyrenees: a pedoanthracological analysis. *Vegetation History and Archaeobotany* 21: 373–384. <http://dx.doi.org/10.1007/s00334-011-0342-y>
- del Barrio G, Puigdefábregas J (1987) Mass wasting features above the timberline in the Central Pyrenees and their topographic controls. *Pirineos* 130: 29–51. <https://dialnet.unirioja.es/servlet/articulo?codigo=135873>
- Diry JP (1995) Moyennes montagnes d'Europe occidentale et dynamiques rurales. *Revue de Géographie Alpine* 83: 15–26. http://www.persee.fr/doc/rga_0035-1121_1995_num_83_3_3814
- Egarter Vigl L, Schipke U, Tasser E, Tappeiner U (2016) Linking long-term landscape dynamics to the multiple interactions among ecosystem services in the European Alps. *Landscape Ecology* 31: 1903–1918. Doi: 10.1007/s10980-016-0389-3
- Egoh B, Rouget M, Reyers B, Knight AT, Cowling RM, van Jaarsveld AS, Welz A (2007) Integrating ecosystem services into conservation assessment: A review. *Ecological Economics* 63: 714–721. <http://dx.doi.org/10.1016/j.ecolecon.2007.04.007>

- 1 Estienne P (1989) Evolution de la population des montagnes françaises au XX^e siècle.
2 Revue de Géographie Alpine 77: 395–405. http://www.persee.fr/doc/rga_0035-1121_1989_num_77_4_2757
3
4
5 Fondevilla C, Colomer MA, Fillat F, Tappeiner U (2016) Using a new PDP modelling
6 approach for land-use and land-cover change predictions: A case study in the Stubai
7 Valley (Central Alps). Ecological Modelling 322: 101–114.
8 <http://dx.doi.org/10.1016/j.ecolmodel.2015.11.016>
9
10 Galop D, Houet T, Mazier F, Leroux G, Rius D (2011) Grazing activities and biodiversity
11 in the Pyrenees: New insights on high altitude ecosystems in the framework of a
12 Human Environment Observatory. PAGES News 19: 53–55. https://halshs.archives-ouvertes.fr/halshs-00750965/file/2011_Galop_et_al_cor.pdf
13
14 García de Celis A, Arroyo Pérez P, Gandía Fernández A (2008) Cambios recientes del
15 límite superior del bosque en Urbión: gestión forestal, ganadería y clima. Zubía
16 Monográfico 20: 97–118. <https://dialnet.unirioja.es/descarga/articulo/2768907.pdf>
17
18 García-Ruiz JM (2010) The effects of land uses on soil erosion in Spain: A review. Catena
19 81: 1–11. <http://dx.doi.org/10.1016/j.catena.2010.01.001>
20
21 García-Ruiz JM (2015) Why Geomorphology is a global science. Cuadernos de
22 Investigación Geográfica 41: 87–105. Doi: 10.18172/cig.2652
23
24 García-Ruiz JM, Lasanta T (1990) Land-use changes in the Spanish Pyrenees. Mountain
25 Research and Development 10: 267–279. <http://dx.doi.org/10.2307/3673606>
26
27 García-Ruiz JM, Lana-Renault N (2011) Hydrological and erosive consequences of
28 farmland abandonment in Europe, with special reference to the Mediterranean region—
29 A review. Agriculture, Ecosystems and Environment 140: 317–338.
30 <http://dx.doi.org/10.1016/j.agee.2011.01.003>
31
32 García-Ruiz JM, Alvera B, del Barrio G, Puigdefábregas J (1990) Geomorphic processes
33 above the timberline in the Spanish Pyrenees. Mountain Research and Development
34 10: 201–214. <http://dx.doi.org/10.2307/3673600>
35
36 García-Ruiz JM, Beguería S, Alatorre LC, Puigdefábregas J (2010) Land cover changes
37 and shallow landsliding in the Flysch Sector of the Spanish Pyrenees. Geomorphology
38 124: 250–259. <http://dx.doi.org/10.1016/j.geomorph.2010.03.036>
39
40 García-Ruiz JM, López-Moreno JI, Vicente-Serrano SM, Lasanta T, Beguería S (2011)
41 Mediterranean water resources in a Global Change scenario. Earth-Science Reviews
42 105: 121–139. <http://dx.doi.org/10.1016/j.earscirev.2011.01.006>
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- García-Ruiz JM, López-Moreno JI, Lasanta T, Vicente-Serrano SM, González-Sampériz P, Valero-Garcés BL, Sanjuán Y, Beguería S, Nadal-Romero E, Lana-Renault N, Gómez-Villar A (2015) Los efectos geocológicos del Cambio Global en el Pirineo Central español: una revisión a distintas escalas espaciales y temporales. *Pirineos* 170, e012. <http://dx.doi.org/10.3989/Pirineos.2015.170005>
- García-Ruiz JM, Sanjuán Y, Gil-Romera G, González-Sampériz P, Beguería S, Arnáez J, Coba-Pérez P, Gómez-Villar A, Álvarez-Martínez J, Lana-Renault N, Pérez-Cardiel E, López de Calle C (2016) Mid and Late Holocene forest fires and deforestation in the Subalpine belt of the Iberian Range, Northern Spain. *Journal of Mountain Science* 13: 1760-1772. Doi: 10.1007/s11629-015-3763-8
- Gartzia M, Alados CL, Pérez-Cabello F (2014) Assessment of the effects of biophysical and anthropogenic factors on woody plant encroachment in dense and sparse mountain grasslands based on remote sensing data. *Progress in Physical Geography* 38: 201–217. <http://dx.doi.org/10.1177/0309133314524429>
- Gehrig-Fasel J, Guisan A, Zimmermann NE (2007) Tree line shifts in the Swiss Alps: Climate change or land abandonment? *Journal of Vegetation Science* 18(4): 571–582. <http://dx.doi.org/10.1111/j.1654-1103.2007.tb02571.x>
- Gómez-Villar A, Álvarez-Martínez J, García-Ruiz JM (2006) Factors influencing the presence or absence of tributary-junction fans in the Iberian Range, Spain. *Geomorphology* 81: 252-264. <http://dx.doi.org/10.1016/j.geomorph.2006.04.011>
- Gómez-Villar A, Sanjuán Y, García-Ruiz JM, Nadal-Romero E, Álvarez-Martínez J, Arnáez J, Serrano Muela MP (2014) Sediment organization and adjustment in a torrential reach of the Upper Ijuez River, Central Spanish Pyrenees. *Cuadernos de Investigación Geográfica* 40: 191-214. <http://dx.doi.org/10.18172/cig.2566>
- Gouveia CM, Páscoa P, Russo A, Trigo RM (2016) Land degradation trend assessment over Iberia during 1982-2012. *Cuadernos de Investigación Geográfica* 42: 89-112. Doi: 10.18172/cig.2945
- Guiguet-Covex C, Arnaud F, Poulenard J., Disnar JR, Delhon C, Francus P, David F, Enters D, Rey PJ, Delannoy JJ (2011) Changes in erosion patterns during the Holocene in a currently treeless subalpine catchment inferred from lake sediment geochemistry (Lake Anterne, 2063 m a.s.l., NW French Alps): The role of climate and human activities. *The Holocene* 21: 651-665. <http://dx.doi.org/10.1177/0959683610391320>
- Höllermann P (1985) The periglacial belt of mid-latitude mountains from a geocological point of view. *Erdkunde* 39: 259-270. DOI: 10.3112/erdkunde.1985.04.02

- Holtmeier FK, Broll G (2005) Sensitivity and response of northern hemisphere altitudinal and polar treelines to environmental change at landscape and local scales. *Global Ecology and Biogeography* 14: 395-410. <http://dx.doi.org/10.1111/j.1466-822x.2005.00168.x>
- Keesstra SD (2007) Impact of natural reforestation on floodplain sedimentation in the Dragonja basin, SW Slovenia. *Earth Surface Processes and Landforms* 32: 49-65. Doi: [10.1002/esp.1360](https://doi.org/10.1002/esp.1360)
- Lana-Renault N, Alvera B, García-Ruiz JM (2011) Runoff and sediment transport during the snowmelt period in a Mediterranean high mountain catchment. *Arctic, Antarctic and Alpine Research* 43: 213-222. <http://dx.doi.org/10.1657/1938-4246-43.2.213>
- Lasanta T (1990) Tendances actuelles de l'organisation spatiale des montagnes espagnoles. *Annales de Géographie* 551: 51-71. http://www.persee.fr/doc/geo_0003-4010_1990_num_99_551_20944
- Lasanta T, Vicente-Serrano SM (2007) Cambios en la cubierta vegetal en el Pirineo aragonés en los últimos 50 años. *Pirineos* 162: 125-154. doi: [10.3989/pirineos.2007.v162.16](http://dx.doi.org/10.3989/pirineos.2007.v162.16) <http://dx.doi.org/10.3989/pirineos.2007.v162.16>
- Lasanta-Martínez T, Vicente-Serrano SM, Cuadrat-Prats JM (2005) Mountain Mediterranean landscape evolution caused by the abandonment of traditional primary activities: a study of the Spanish Central Pyrenees. *Applied Geography* 25: 47-65. <http://dx.doi.org/10.1016/j.apgeog.2004.11.001>
- Lasheras-Álvarez L, Pérez-Sanz A, Gil-Romera G, González-Sampériz P, Sevilla-Callejo M, Valero-Garcés B (2013) Historia del fuego y la vegetación en una secuencia holocena del Pirineo Central: La Basa de la Mora. *Cuadernos de Investigación Geográfica* 39: 77-95. <http://dx.doi.org/10.18172/cig.2000>
- Liébault F, Piégay H, 2001. Assessment of channel changes due to long-term bedload supply decrease, Roubion River, France. *Geomorphology* 36: 167-186. [https://doi.org/10.1016/S0169-555X\(00\)00044-1](https://doi.org/10.1016/S0169-555X(00)00044-1)
- López-Moreno JI (2005) Recent variations of snowpack depth in the Central Spanish Pyrenees. *Arctic, Antarctic, and Alpine Research* 37: 253-260. [http://dx.doi.org/10.1657/1523-0430\(2005\)037\[0253:RVOSDI\]2.0.CO;2](http://dx.doi.org/10.1657/1523-0430(2005)037[0253:RVOSDI]2.0.CO;2)
- López-Moreno JI, Goyett S, Meniston M (2009) Impact of climate change on snowpack in the Pyrenees: Horizontal spatial variability and vertical gradients. *Journal of Hydrology* 374: 384-396. <http://dx.doi.org/10.1016/j.jhydrol.2009.06.049>

- López-Moreno JI, Zabalza J, Vicente-Serrano SM, Revuelto J, Gilaberte M, Azorín-Molina C, Morán-Tejeda E, García-Ruiz JM, Tague C (2014) Impact of climate and land use change on water availability and reservoir management. Scenarios in the Upper Aragón River, Spanish Pyrenees. *Science of the Total Environment* 493: 1222-1231. <http://dx.doi.org/10.1016/scitotenv.2013.09.031>
- MacDonald D, Crabtree JR, Wiesinger G, Dax T, Stamou N, Fleury P, Gutiérrez Lazpita J, Gibon A (2000) Agricultural abandonment in mountain areas of Europe: environmental consequences and policy response. *Journal of Environmental Management* 59: 47-69. <http://dx.doi.org/10.1006/jema.1999.0335>
- Montserrat J (1992) Evolución glacial y postglacial del clima y la vegetación en la vertiente sur del Pirineo: Estudio palinológico. Instituto Pirenaico de Ecología, Zaragoza, 147 pp.
- Morán-Tejeda E, Lorenzo-Lacruz J, López-Moreno JI, Rahman K, Beniston B (2014) Streamflow timing of mountain rivers in Spain: recent changes and future projections. *Journal of Hydrology*. <http://dx.doi.org/10.1016/j.jhydrol.2014.06.053>
- Moreno Fernández JR (1996) La ganadería trashumante en La Rioja 1752-1865. Una revisión bibliográfica y cuantitativa. *Brocar* 20: 277-302. <https://publicaciones.unirioja.es/ojs/index.php/brocar/article/view/1767>
- Navarro-Serrano E, López-Moreno JI (2017) Spatio-temporal analysis of snowfall events in the Spanish Pyrenees and their relationship to atmospheric circulation. *Cuadernos de Investigación Geográfica* 43 (1): 233-254. <http://doi.org/10.18172/cig.3042>
- Pecher C, Tasser E, Tappeiner U (2011) Definition of the potential treeline in the European Alps and its benefit for sustainability monitoring. *Ecological Indicators* 11 (2): 438-447. <https://doi.org/10.1016/j.ecolind.2010.06.015>
- Pérez-Sanz A, González-Sampériz P, Moreno A, Valero-Garcés B, Gil-Romera G, Rieradevall M, Tarrats P, Lasheras-Álvarez L, Morellón M, Belmonte A, Sancho C, Sevilla-Callejo M, Navas A (2013) Holocene climate variability, vegetation dynamics and fire regime in the central Pyrenees: the Basa de la Mora sequence (NE Spain). *Quaternary Science Reviews* 73: 149-169. <http://dx.doi.org/10.1016/j.quascirev.2013.05.010>
- Piégay H, Walling DE, Landon N, He Q, Liébault F, Petiot R (2004) Contemporary changes in sediment yield in an alpine mountain basin due to afforestation (the Upper Drôme in France). *Catena* 55: 183-212. [https://doi.org/10.1016/S0341-8162\(03\)00118-8](https://doi.org/10.1016/S0341-8162(03)00118-8)

- Poyatos R, Latron J, Llorens P (2003) Land use and land cover change after agricultural abandonment. The case of a Mediterranean mountain area (Catalan Pre-Pyrenees). Mountain Research and Development 23: 362-368. [http://dx.doi.org/10.1659/0276-4741\(2003\)023\[0362:LUALCC\]2.0.CO;2](http://dx.doi.org/10.1659/0276-4741(2003)023[0362:LUALCC]2.0.CO;2)
- Sanjuán Y, Gómez-Villar A, Nadal-Romero E, Álvarez-Martínez J, Arnáez J, Serrano-Muela MP, Rubiales JM, González-Sampériz P, García-Ruiz JM (2016) Linking land cover changes in the sub-alpine and montane belts to changes in a torrential river. Land Degradation & Development 27: 179-189. <http://dx.doi.org/10.1002/ldr.2294>
- Tasser E, Tappeiner U (2002) Impact of land use changes on mountain vegetation. Applied Vegetation Science 5: 173-184. <http://dx.doi.org/10.1111/j.1654-109X.2002.tb00547.x>
- Tasser E, Leitinger G, Tappeiner U (2017) Climate change versus land use change – which affects the landscape more? Land Use Policy 60: 60-72. <https://doi.org/10.1016/j.landusepol.2016.10.019>
- Vicente-Serrano SM, Rodríguez-Camino E, Domínguez-Castro F, El Kenawy A, Azorín-Molina C (2017) An updated review of recent trends in observational surface atmospheric variables and their extremes over Spain. Cuadernos de Investigación Geográfica 43 (1): 209-232. <http://doi.org/10.18172/cig.3134>
- Vitte P (1992) La montagne italienne. Annales de Géographie 563: 68-83. http://www.persee.fr/doc/geo_0003-4010_1992_num_101_563_21066
- Viviroli D, Weingartner R, Messerli B (2003) Assessing the hydrological significance of the World's mountains. Mountain Research and Development 23: 32-40. [http://dx.doi.org/10.1659/0276-4741\(2003\)023\[0032:ATHSOT\]2.0.CO;2](http://dx.doi.org/10.1659/0276-4741(2003)023[0032:ATHSOT]2.0.CO;2)
- Wallentin G, Tappeiner U, Strobl J, Tasser E (2008) Understanding alpine tree line dynamics: an individual based model. Ecological Modelling 218: 235-246. <https://doi.org/10.1016/j.ecolmodel.2008.07.005>

FIGURE CAPTIONS

Figure 1. Maps with LULC categories in the Ormazal Valley in 1956 and 2006, and location of the Ormazal valley in Spain.

Figure 2. Density curves for 1956 and 2006, showing the frequency distribution of shrublands as a function of elevation (m), slope gradient (%), solar irradiation ($\text{kWh m}^{-2} \text{yr}^{-1}$) and topographic index (unitless).

Figure 3. Density curves for 1956 and 2006, showing the frequency distribution of forests as a function of elevation (m), slope gradient (%), solar irradiation ($\text{kWh m}^{-2} \text{yr}^{-1}$) and topographic index (unitless).

Figure 4. Centroids in the discriminant analysis. Horizontal axis (Function 1); vertical axis (Function 2). CF to AF: cultivated fields to abandoned fields; AF to G: abandoned fields to summer grasslands; CF to EA: clear forest to eroded areas; S to EA: shrublands to eroded areas; S to DF: shrublands to dense forest; CF to DF: clear forest to dense forest; G to CF: summer grasslands to clear forest; G to DF: summer grasslands to dense forest.

Figure 5. Transition probabilities for the areas covered by grassland in 1956: grassland permanence and transition to forest, eroded areas, and shrubland.

Figure 6. Transition probabilities for the areas covered by shrubland in 1956: shrubland permanence and transition to forest, eroded areas, and grassland.

Dear Prof. Cramer,

Thank you very much for accepting our paper "Woody plant encroachment following grazing abandonment in the subalpine belt. A case study in Northern Spain". I re-uploaded a clean version of the manuscript with the following corrections:

a) as by our instructions, first names of all authors need to be given on the front page. DONE

b) some doi-numbers are missing, although they do exist (Geografiska Annaler, for example).
DONE

This time I did not include the track-changes version.

I hope everything is OK now.

Best regards

Noemí Lana-Renault

On behalf of teh authors

Figure3

Figure4

Figure5

to Grasslands

to Eroded

to Shrublands

to Forest

Figure6

to Shrublands

to Eroded

to Grasslands

to Forest

Table 1. Land use / land covers (LULC) classes in 1956 and 2006.

LULC	1956 (ha)	1956 (%)	2006 (ha)	2006 (%)
Cultivated fields (CF)	10.07	0.4	0	0
Abandoned fields (AF)	58.23	2.3	59.99	2.4
Reforestation (RF)	0	0.0	70.46	2.8
Rock outcrops, debris (R)	1.72	0.1	1.04	<0.1
Eroded areas (EA)	405.33	16.1	438.37	17.4
Grasslands (pastures) (G)	910.55	36.2	681.97	27.1
Shrublands (S)	853.15	33.9	651.39	25.9
Clear forest (CF)	231.12	9.2	174.18	6.9
Dense forest (DF)	43.64	1.7	436.41	17.4
Total	2513.81	100	2513.81	100

Table 2. Transition matrix between LULC classes in 1956 (vertical) and 2006 (horizontal). Values are expressed in percentage over the area covered by each LULC class in 1956. Legend: CF, cultivated fields; AF, abandoned fields; RF, reforestation; R, rock outcrops and debris; EA, eroded areas; G, grasslands; S, shrublands; CF, clear forest; DF, dense forest. The diagonal numbers in bold indicate the percentage of each LULC category that did not change

1956	2006									Total
	CF	AF	RF	R	EA	G	S	CF	DF	
CF	0	99.6	0	0	0	0	0	0	0.4	100
AF	0	60.4	0	0	4.4	18.8	4.3	0	12	100
RF	0	0	0	0	0	0	0	0	0	0
R	0	0	0	0	0	0	0	0	100	100
EA	0	0	0.4	0.1	54.8	26.9	11.6	4.7	1.5	100
G	0	1.3	0.4	0	20.1	52.5	17.3	2.5	5.9	100
S	0	0.3	4.3	0.1	3.2	9.6	43.2	12.2	27.2	100
CF	0	0.1	12.6	0	1.3	0.4	32.4	12.3	40.9	100
DF	0	0	0.2	0	1.1	2.7	1.1	0.5	94.5	100

Table 3. Permanence and transition probabilities of grassland and shrubland areas in 1956, observed and predicted by multinomial logistic regression analysis.

	Probability of permanence	Probability of transition to forest	Probability of transition to eroded	Probability of transition to shrublands
<i>Grasslands</i>				
Predicted	0.62	0.07	0.16	0.15
Observed	0.53	0.08	0.20	0.17
<i>Shrublands</i>				
				to grasslands
Predicted	0.48	0.42	0.04	0.06
Observed	0.43	0.39	0.03	0.10

Table 4. Transition probabilities from grasslands and shrublands to i) forest, ii) eroded and iii) shrubland or grassland, considering average topographic conditions, and variation in the transition probabilities as a function of changes in the topographic variables (only significant effects).

From grasslands...	... to forest	... to eroded	... to shrublands
Average topographic conditions (36%, 1,229 kWh m ⁻² yr ⁻¹)	10,4%	20,3%	20,0%
Slope gradient +10%	+3,7%	+15,8%	+13,8%
Solar irradiation +200 kWh m ⁻² yr ⁻¹	-5,6%	+23,7%	+6,7%
From shrublands ...	... to forest	... to eroded	... to grasslands
Average topographic conditions (36%, 1,223 kWh m ⁻² yr ⁻¹)	46,5%	7,4%	11,3%
Slope gradient +10%	+11,4%		
Solar irradiation +200 kWh m ⁻² yr ⁻¹	-7,7%		