



Short communication

In vivo assay to identify bacteria with β -glucosidase activity



Erwin Strahsburger^{a,*}, Ana Maria Lopez de Lacey^b, Ilaria Marotti^c, Diana DiGioia^c,
Bruno Biavati^d, Giovanni Dinelli^c

^a Facultad de Recursos Naturales Renovables, Universidad Arturo Prat, Avenida Arturo Prat 2120, Iquique, Chile

^b Instituto del Frío-ICTAN (CSIC), José Antonio Novais 10, 28040 Madrid, Spain

^c Department of Agricultural Sciences, University of Bologna, viale Fanin, 42, Bologna, Italy

^d Institute of Earth Systems, Division of Rural Sciences & Food Systems, University of Malta, Msida, Malta

ARTICLE INFO

Article history:

Received 15 July 2016

Accepted 30 August 2017

Available online 25 September 2017

Keywords:

Antimicrobial

Bifidobacterium

Cell lysis

Enzymatic test

Lactobacillus

p-Nitrophenol

p-Nitrophenyl- β -glucopyranoside

Screening

Screening

β -Glucosidase assay

β -Glucosidase producer strain

ABSTRACT

Background: β -Glucosidase assay is performed with purified or semipurified enzymes extracted from cell lysis. However, in screening studies, to find bacteria with β -glucosidase activity among many tested bacteria, a fast method without cell lysis is desirable. In that objective, we report an *in vivo* β -glucosidase assay as a fast method to find a β -glucosidase producer strain.

Results: The method consists in growing the strains for testing in a medium supplemented with the artificial substrate *p*-nitrophenyl- β -glucopyranoside (*p*NPG). The presence of β -glucosidases converts the substrate to *p*-nitrophenol (*p*NP), a molecule that can be easily measured in the supernatant spectrophotometrically at 405 nm. The assay was evaluated using two *Bifidobacterium* strains: *Bifidobacterium longum* B7254 strain that lacks β -glucosidase activity and *Bifidobacterium pseudocatenulatum* B7003 strain that shows β -glucosidase activity. The addition of sodium carbonate during *p*NP measurement increases the sensitivity of *p*NP detection and avoids the masking of absorbance by the culture medium. Furthermore, we show that *p*NP is a stable enzymatic product, not metabolized by bacteria, but with an inhibitory effect on cell growth. The β -glucosidase activity was measured as units of enzyme per gram per minute per dry cell weight. This method also allowed the identification of *Lactobacillus* strains with higher β -glucosidase activity among several *Lactobacillus* species.

Conclusion: This *in vivo* β -glucosidase assay can be used as an enzymatic test on living cells without cell disruption. The method is simple, quantitative, and recommended, especially in studies screening for bacteria not only with β -glucosidase activity but also with high β -glucosidase activity.

© 2017 Pontificia Universidad Católica de Valparaíso. Production and hosting by Elsevier B.V. All rights reserved. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

β -D-glucosidases (EC 3.2.1.21) catalyze the transfer of glycosylic groups between oxygen nucleophiles [1]. The substrates include a wide range of carbohydrate molecules with a β -anomeric conformation binding to aryl-, amino-, and cyanogenic-glucoside or a β -anomeric conformation, which is present in alkyl- β -D-glucoside, oligosaccharide, and disaccharide. According to their substrate specificities, these enzymes could be classified into three groups: aryl β -glucosidases, cellobiohydrolases, and glucosidases with broad substrate specificity. However, this classification is not necessarily in accordance with the relationship between amino acid sequence and catalytic mechanism [2]. An alternative classification that relates structural characteristics with functional mechanisms has been published [2]. Regarding its

biological role in microorganisms, β -glucosidases participate in the degradation of cellulose or other carbohydrates present in the cell wall. However, β -glucosidases have different functions in plants, where they play a role in the biosynthesis of pigments or the cell wall, fruit ripening, and defense mechanism [3]. In mammals, β -glucosidases participate in ceramide hydrolysis, and defects in these genes are associated with the Gaucher's disease in humans [4].

β -Glucosidases also play an important role in the food industry. These enzymes are produced by probiotic bacteria and can remove the glycoside moiety from glycosylated flavonoids present in soybean products [5]. This hydrolysis allows the intestinal absorption of flavonoids, which is necessary to obtain their beneficial health effects [6]. The identification of *Bifidobacterium* and *Lactobacillus* strains with β -glucosidase enzymes [7,8] has contributed to the development of fermented soy bean products enriched in the aglycone form of flavonoids, which improves their uptake and increases their benefits for consumers [5]. Other significant industrial applications of β -glucosidases are related to the cellulose industry and bioethanol production. Many

* Corresponding author.

E-mail address: estrahbs@unap.cl (E. Strahsburger).

Peer review under responsibility of Pontificia Universidad Católica de Valparaíso.

microorganisms have β -glucosidase enzyme activity against diverse carbon sources such as cellulose, rice bran, wheat bran, and lactose [3]. The biomass or cellobiose fermentation process produces bioethanol, a product with an enormous significance in the biofuel fields. Cloning and *in vitro* studies of these β -glucosidases have greatly contributed to our understanding of their enzymatic mechanisms and allowed to develop applications for these enzymes in the food and biofuel industries [1]. The discovery of new bacterial strains that possess a β -glucosidase enzyme with catalytic properties that respond to challenges in these industries is an ongoing pursuit for many researchers around the world.

β -Glucosidase activity can be measured using an artificial substrate, which is converted to a colored product that is easily detected spectrophotometrically. In general, this assay requires cell lysis and protein purification to determine the amount of total protein and calculate the specific enzyme activity. However, in this study, we report an *in vivo* β -glucosidase assay without cell lysis. This method consists of incubating the microorganism with an artificial substrate and measuring the product in the supernatant. In this way, many microorganisms can be screened to find strains with higher β -glucosidase activity. The proposed method was evaluated in *Bifidobacterium* and *Lactobacillus* strains, which produced a stable artificial product that was not metabolized by the bacteria and was easily measured spectrophotometrically. The results demonstrate the effectiveness of this method and suggest it to be an alternative method to identify microorganisms with high β -glucosidase activity.

2. Materials and methods

2.1. Chemicals

The substrate *p*-nitrophenyl- β -glucopyranoside (pNPG) and product *p*-nitrophenol (pNP) were purchased from Sigma-Aldrich (Milan, Italy).

2.2. Bacteria and culture conditions

All strains used are listed in **Table S1** and belong to the cell collection of the Department of Agroenvironmental Science and Technology, University of Bologna, Italy [9]. *Lactobacillus* strains were cultured in “de Man Rogosa Sharpe” (MRS) medium, while *Bifidobacterium* strains were cultured in MRS supplemented with 0.05% L-cysteine hydrochloride or cultured in trypticase-peptone-yeast (TPY) extract [9]. The strains were incubated at 37°C under anaerobic conditions using oxoid gas jars and anaerobic gas packs (BD; Becton, Dickinson and Company) [10].

2.3. β -Glucosidase activity

β -Glucosidase activity was determined by adding various concentrations of pNPG to the medium at the beginning of culture. Briefly, all strains were grown in 7 mL of their respective medium and supplemented with 0, 400, 800, or 1600 $\mu\text{g}\cdot\text{mL}^{-1}$ pNPG. A control condition without bacteria was included. At different times, 150- μL aliquots were collected in triplicate to measure the optical density at 620 nm in a microplate reader (Multiskan, Thermo Electron Oy, Vaanta, Finland). The samples were recovered and centrifuged at 10000 \times g for 5 min, and 100 μL of each supernatant was transferred to a 96-well plate. Then, 50 μL of 0.1-M sodium carbonate was added to each well, and the pNP concentration was determined at 405 nm. A standard curve between 0.001 and 1 mM pNP dissolved in MRS or TPY medium was generated. In general, one unit (U) of enzyme activity is defined as the amount of β -glucosidase that produces 1 μmol of pNP per minute under assay conditions. However, in this assay, the amount of β -glucosidase enzyme was calculated in terms of grams of dry cell weight biomass. Therefore, we define one unit (U) of enzyme activity as 1 g of dry cell weight biomass that produces 1 nmol of pNP per minute under culture conditions. The kinetic studies were performed with

culture data of the B7003 strain obtained at 16, 20, and 24 h of culture in TPY medium. A Lineweaver–Burk plot was generated, and the K_m was calculated. The Lineweaver–Burk plot graphed V^{-1} ($\mu\text{mol}\cdot\text{min}^{-1}$) $^{-1}$ vs. $[S]^{-1}$ (mM^{-1}). The y-intercept (V^{-1} axis) corresponded to the V_{max}^{-1} . The K_m was calculated using the formula $K_m = m \cdot V_{\text{max}}$, where m is the slope of the line [1].

2.4. Statistical analysis

Mean values were compared with Fisher's test using Microsoft Excel software (Microsoft). When treatments produced statistically significant differences according to Fisher's test, the corresponding means were compared with the SNK test for multiple comparisons at the 0.05 level of probability.

3. Results

3.1. *In vivo* determination of pNP produced

The *in vivo* β -glucosidase assay determined the bioconversion of pNPG into pNP in bacterial culture. The enzymatic product was measured directly from the culture supernatant at 405 nm. However, TPY and MRS media also absorb at that wavelength (**Fig. 1a**). To improve the sensitivity of the method, sodium carbonate was added to increase the molar extinction coefficient of pNP (**Fig. 1a**). Therefore, in subsequent experiments, sodium carbonate was always added during pNP measurement.

3.2. pNP stability and inhibitory effect on cell growth

The *in vivo* β -glucosidase activity was assayed in two *Bifidobacterium* strains: *B. longum* strain B7254, which lacks β -glucosidase activity (negative control), and *B. pseudocatenulatum* strain B7003, which has β -glucosidase activity (positive control). The stability of pNP during culture conditions of both strain was tested. In TPY medium, the artificial product was stable and not metabolized by either strain. However, in MRS medium with or without bacteria, a reduction of between 10 and 15% of the total 300–350 $\mu\text{g}\cdot\text{mL}^{-1}$ of pNP was observed, while the concentration of pNP was approximately 700 $\mu\text{g}\cdot\text{mL}^{-1}$ and showed no significant change (**Fig. S1**).

Regarding the pNP effect on cell viability, an inhibitory effect was observed in both media (**Fig. 1b**). In TPY medium, the inhibition was directly proportional to the concentration of pNP, while in MRS medium, the inhibitory effect was the same at both pNP concentrations tested (**Fig. 1b**). Independent of the medium used, the B7254 strain always reached a higher optical density than the B7003 strain (**Fig. 1b**).

3.3. β -Glucosidase activity in bacterial culture

As expected, enzymatic activity was not observed for the B7254 strain, while the pNP produced was directly proportional to pNPG concentrations added to the medium for the B7003 strain (**Fig. 1c**). The enzymatic activity was saturated at 1600 $\mu\text{g}\cdot\text{mL}^{-1}$ of pNPG (**Fig. 1c**). The growth of B7003 was lower in MRS medium than in TPY medium and inversely proportional to the pNPG concentration added to the medium (**Fig. S2**). As pNP concentrations did not show variation between 14 and 24 h of culture, the enzymatic activity was calculated at 14 h (**Table 1**). Although the pNP production was the same in TPY or MRS bacterial culture, the specific enzyme activity was higher in MRS medium because the biomass was lower (**Table 1**). The Michaelis–Menten constants (K_m) were calculated from the analysis of the Lineweaver–Burk plot at 16, 20, and 24 h of culture, which were 12.5, 14.2, and 15.8 mM, respectively (**Table S2**).




Fig. 1. Study of the stability and production of pNP. (a) Spectrophotometric determination of pNP with and without adding sodium carbonate. (b) Effect of pNP on the cell growth of B7003 and B7254 strains cultured in TPY medium or MRS medium. * ($P > 0.5$), ** ($P > 0.05$), *** ($P > 0.005$). (c) β -glucosidase activity of B7003 strain grown in TPY medium supplemented with different concentration of pNPG. The pNP production was measured at different times of culture.

3.4. *In vivo* β -glucosidase assay in *Lactobacillus* strains

The *in vivo* β -glucosidase assay was performed on several *Lactobacillus* strains. These bacteria were cultured in MRS medium without L-cysteine and supplemented with 800 $\mu\text{g mL}^{-1}$ of pNPG. These strains reached similar cell densities after 23 h of culture but showed different pNP production (Fig. S3). The difference between strains of *L. helveticus* was remarkable. *L. helveticus* ATCC strain 15009 showed a lack of β -glucosidase activity, while *L. helveticus* strain M162 showed moderate β -glucosidase activity. Among the other *Lactobacillus* species assayed, all strains showed consistent results according to their classification of null, low, or high β -glucosidase activity (Fig. S3).

4. Discussion

In this study, we measured β -glucosidase activity directly from the bacterial culture of two *Bifidobacteria* strains and several *Lactobacillus* strains by quantifying the bioconversion of pNPG into pNP. The concentration of pNP was measured directly from the supernatant of the culture after adding sodium carbonate to reduce the interference

of MRS and TPY media. In general, sodium carbonate is used to stop the enzymatic reaction and increase the chromatic properties of pNP [7]. However, in this assay, it is not necessary to chemically stop the enzymatic reaction because the enzymatic source (the bacteria) is separated from the substrate by centrifugation. However, owing to the interference of both media at 405 nm, sodium carbonate was incorporated to increase the pNP molar extension coefficient [11].

Both pNPG and pNP are stable in the culture conditions used here. Only in MRS medium at lower concentrations ($<700 \mu\text{g L}^{-1}$), a slight reduction of pNP concentration was observed, likely due to the oxidative effects of L-cysteine [12,13]. In addition, these assays also demonstrate that pNP is not metabolized by bacteria, remaining stable in this culture condition with or without bacteria. Although some soil bacteria have at least one of the two known metabolic pathways to degrade pNP (hydroquinone or nitrocatechol catabolic pathway) [14, 15,16], these pathways are absent in *Lactobacillus* and *Bifidobacterium* species [17].

Regarding cell growth, MRS medium was not optimal for both *Bifidobacterium* strains. Although this medium supplemented with L-cysteine (0.5 g L^{-1}) is usually used to grow *Bifidobacterium* species [18], we suggest TPY medium as the optimal media for this assay. In

Table 1

Bioconversion activity of B7003 strain cultured with pNPG.

Artificial substrate pNPG ($\mu\text{g mL}^{-1}$)	pNP produced ($\mu\text{mol min}^{-1} \text{L}^{-1}$) (\pm SD)		Biomass (g of dry weight L^{-1}) (\pm SD)		Specific activity ($\mu\text{mol min}^{-1} \cdot \text{g of dry weight}^{-1} \text{L}^{-1}$) (\pm SD)	
	TPY	MRS	TPY	MRS	TPY	MRS
400	12.5 (\pm 0.39)	12.4 (\pm 0.23)	0.51 (\pm 0.0013)	0.15 (\pm 0.001)	24.4 (\pm 0.10)	82.8 (\pm 1.6)
800	22.2 (\pm 0.55)	25.5 (\pm 1.86)	0.45 (\pm 0.0047)	0.1 (\pm 0.001)	48.9 (\pm 1.70)	255.3 (\pm 18.6)
1600	33.6 (\pm 0.51)	45.4 (\pm 2.98)	0.39 (\pm 0.0047)	0.06 (\pm 0.0001)	86.9 (\pm 0.93)	756.1 (\pm 49.7)

SD: Standard deviation.

fact, a standard medium to grow *Bifidobacterium* species has not been established [19]. Regarding the effect of *p*NP and *p*NPG on cell viability, we found that *p*NP, but not *p*NPG, has a detrimental effect on both strains. In the B7254 strain (lacking β -glucosidase activity), *p*NPG was innocuous, but in the B7003 strain, the cytotoxic effect could be a result of the enzymatic conversion of *p*NPG into *p*NP. According to the United States Environmental Protection Agency, this molecule affects many organisms, similar to other nitrophenol or aminophenol derivative compounds [20,21,22,23]. This supports the idea that *p*NPG is innocuous and the cytotoxic effect is not associated with the β -glucosidase metabolism because the B7254 strain is also affected by *p*NP.

Regarding the kinetic parameters, the microbial cultures did not show enzymatic variation after 14 h of culture, and enzymatic activity was saturated at $1600 \mu\text{g}\cdot\text{mL}^{-1}$ of *p*NPG. This suggests that the enzymes are synthesized mainly during the exponential phase and remain highly expressed during the stationary phase. In this way, the enzymatic activity remains a function of substrate concentration, but not of biomass, explaining the observation that similar *p*NP production was achieved by the B7003 strain cultured in different media. Previously, β -glucosidase activity was measured in either the exponential or stationary phase, depending on the *Bifidobacterium* strain studied [24]. Although the composition of the culture medium seems to contribute to the β -glucosidase activity [24], this effect was not observed under the conditions used in this study.

Regarding the kinetic analysis, this assay does not correspond exactly with the theory of Michaelis–Menten kinetics. For the substrate to reach the enzyme, it has to pass through the cell membrane. Similarly, before the enzymatic product is measured, it also has to pass through the cell membrane. Therefore, other constants have to be added to the Michaelis–Menten equation. However, these constants do not seem to be limiting for the reaction. The K_m observed in this *in vivo* assay was tenfold higher than the K_m described for an *in vitro* assay with an enzyme purified from the *Bifidobacterium breve* 203 strain using the same substrate (*p*NPG) [25]. However, our K_m value was similar to the K_m values obtained for *B. breve* strain UCC2003 studied with different substrates (trehalose, palatinose, panose, and isomaltotriose) [26]. If the K_m is understood to be an enzymatic affinity constant, we can assume that the cell wall does not constitute a limiting step in the Michaelis–Menten equation, explaining the similar results observed between *in vitro* and *in vivo* β -glucosidase assays.

Finally, the β -glucosidase activities of *L. plantarum* strain M137 and the *L. acidophilus* strain are in correspondence with results reported in the literature [27,28]. However, to our knowledge, this is the first report describing β -glucosidase activity for the other *Lactobacillus* species.

5. Conclusion

In conclusion, the *in vivo* β -glucosidase assay is a consistent method whereby the product *p*NP is not metabolized, allowing the assumption that all *p*NP comes from the enzymatic activity. Although *p*NP affects cell viability, the bacterial culture does not produce high concentrations of *p*NP, thus reducing its detrimental effect if lower concentrations of *p*NPG are used ($<800 \mu\text{g}\cdot\text{mL}^{-1}$). The culture medium recommended for this assay is the optimal medium for each of the tested bacteria. It is recommended that sodium carbonate be added during the *p*NP determination to avoid the masking effect of the culture medium. The incubation time seems to not be a critical point in this assay, and at least 14-h culture can be used if the temporal expression of this enzyme is not known. Thus, this *in vivo* β -glucosidase assay is recommended for routine or high-throughput screening to screen for microorganisms with high β -glucosidase activity.

Conflicts of interest

The authors declare that there is no conflict of interest in this work.

Financial support

The study was funded by the University of Bologna, RFO Program number J61J10000790001. Erwin Strahsburger received a scholarship within the Erasmus Mundus Program. Ana M. Lopez de Lacey received a fellowship within the JAE-CSIC predoctoral Programme. The study was also financially funded by the Universidad Arturo Prat, internal projects number VRIIP0218-15 and VRIIP01-14.

Supplementary data

<https://doi.org/10.1016/j.ejbt.2017.08.010>.

References

- [1] Bhatia Y, Mishra S, Bisaria VS. Microbial β -glucosidases: cloning, properties, and applications. *Crit Rev Biotechnol* 2002;22:375–407. <https://doi.org/10.1080/07388550290789568>.
- [2] Kötzer MP, Hancock SM, Withers SG. Glycosidases: Functions, Families and Folds. eLS. Chichester: John Wiley & Sons, Ltd; 2014. <https://doi.org/10.1002/9780470015902.a0020548.pub2>.
- [3] Singhania RR, Patel AK, Sukumaran RK, Larroche C, Pandey A. Role and significance of beta-glucosidases in the hydrolysis of cellulose for bioethanol production. *Bioresour Technol* 2013;127:500–7. <https://doi.org/10.1016/j.biortech.2012.09.012>.
- [4] Lieberman RL, Wustman BA, Huertas P, Powe AC, Pine CW, Khanna R, et al. Structure of acid β -glucosidase with pharmacological chaperone provides insight into Gaucher disease. *Nat Chem Biol* 2007;3:101–7. <https://doi.org/10.1038/nchembio850>.
- [5] Otieno DO, Shah NP. Endogenous β -glucosidase and β -galactosidase activities from selected probiotic micro-organisms and their role in isoflavone biotransformation in soymilk. *J Appl Microbiol* 2007;103:910–7. <https://doi.org/10.1111/j.1365-2672.2007.03438.x>.
- [6] Izumi T, Piskula MK, Osawa S, Obata A, Tobe K, Saito M, et al. Soy isoflavone aglycones are absorbed faster and in higher amounts than their glucosides in humans. *J Nutr* 2000;130:1695–9.
- [7] Di Gioia D, Strahsburger E, Lopez de Lacey AM, Bregola V, Marotti I. Flavonoid bioconversion in *Bifidobacterium pseudocatenulatum* B7003: a potential probiotic strain for functional food development. *J Funct Foods* 2014;7:671–9. <https://doi.org/10.1016/j.jff.2013.12.018>.
- [8] Gaya P, Peiroten A, Medina M, Landete JM. Isoflavone metabolism by a collection of lactic acid bacteria and *bifidobacteria* with biotechnological interest. *Int J Food Sci Nutr* 2016;67:117–24. <https://doi.org/10.3109/09637486.2016.1144724>.
- [9] Biavati B, Mattarelli P. Genus *Bifidobacterium*. In: Goodfellow M, Kampfer P, Busse HJ, Suzuki K, Ludwig W, Whitman WB, editors. *Bergey's manual of systematic bacteriology*, 5. 2nd edn. New York: Springer; 2012. p. 171–206.
- [10] Payne JF, Morris AEJ, Beers P. Note: Evaluation of selective media for the enumeration of *Bifidobacterium* sp. in milk. *J Appl Microbiol* 1999;86:353–8. <https://doi.org/10.1046/j.1365-2672.1999.00671.x>.
- [11] Biggs AI. A spectrophotometric determination of the dissociation constants of *p*-nitrophenol and papaverine. *Trans Faraday Soc* 1954;50:800–2. <https://doi.org/10.1039/TF9545000800>.
- [12] Yang J, Pan B, Li H, Liao S, Zhang D, Wu M, et al. Degradation of *p*-nitrophenol on biochars: role of persistent free radicals. *Environ Sci Technol* 2015;50:694–700. <https://doi.org/10.1021/acs.est.5b04042>.
- [13] Zhang JJ, Liu H, Xiao Y, Zhang XE, Zhou NY. Identification and characterization of catabolic *para*-nitrophenol 4-monooxygenase and *para*-benzoquinone reductase from *Pseudomonas* sp. strain WBC-3. *J Bacteriol* 2009;191:2703–10. <https://doi.org/10.1128/JB.01566-08>.
- [14] Arora PK, Bae H. Bacterial degradation of chlorophenols and their derivatives. *Microb Cell Fact* 2014;13:31. <https://doi.org/10.1186/1475-2859-13-31>.
- [15] Min J, Zhang JJ, Zhou NY. A two-component *para*-nitrophenol monooxygenase initiates a novel 2-chloro-4-nitrophenol catabolism pathway in *Rhodococcus intechensis* rj300. *Appl Environ Microbiol* 2016;82:714–23. <https://doi.org/10.1128/AEM.03042-15>.
- [16] Chen Q, Tu H, Luo X, Zhang B, Huang F, Li Z, et al. The regulation of *para*-nitrophenol degradation in *Pseudomonas putida* DLL-E4. *PLoS One* 2016;11:e0155485. <https://doi.org/10.1371/journal.pone.0155485>.
- [17] Tirola MA, Wang H, Paulin L, Kulomaa MS. Evidence for natural horizontal transfer of the *pcpB* gene in the evolution of polychlorophenol-degrading *Sphingomonads*. *Appl Environ Microbiol* 2002;68:4495–501. <https://doi.org/10.1128/AEM.68.9.4495-4501.2002>.
- [18] Shu G, Yang H, Tao Q, He C. Effect of ascorbic acid and cysteine hydrochloride on growth of *Bifidobacterium bifidum*. *Adv J Food Sci Technol* 2013;5:678–81.
- [19] Roy D. Media for the isolation and enumeration of bifidobacteria in dairy products. *Int J Food Microbiol* 2001;69:167–82. [https://doi.org/10.1016/S0168-1605\(01\)00496-2](https://doi.org/10.1016/S0168-1605(01)00496-2).
- [20] Zohar S, Kviatkovski I, Masaphy S. Increasing tolerance to and degradation of high *p*-nitrophenol concentrations by inoculum size manipulations of *Arthrobacter* 4H β isolated from agricultural soil. *Int Biodegr Biodegr* 2013;84:80–5. <https://doi.org/10.1016/j.ibiod.2012.05.041>.
- [21] Chirino B, Strahsburger E, Agulló L, González M, Seeger M. Genomic and functional analyses of the 2-aminophenol catabolism pathway and partial conversion of its substrate into picolinic acid in *Burkholderia xenovorans* LB400. *PLoS One* 2013;8:e75746. <https://doi.org/10.1371/journal.pone.0075746>.

- [22] Garrison AT, Abouelhassan Y, Norwood VM, Kallifidas D, Bai F, Nguyen MT, et al. Structure-activity relationships of a diverse class of halogenated phenazines that targets persistent, antibiotic-tolerant bacterial biofilms and *Mycobacterium tuberculosis*. *J Med Chem* 2016;59:3808–25. <https://doi.org/10.1021/acs.jmedchem.5b02004>.
- [23] Al-Adham IS, Ashour H, Al-Kaissi E, Khalil E, Kierans M, Collier PJ. Studies on the kinetics of killing and the proposed mechanism of action of microemulsions against fungi. *Int J Pharm* 2013;454:226–32. <https://doi.org/10.1016/j.ijpharm.2013.06.049>.
- [24] Jeon KS, Ji GE, Hwang IK. Assay of β -glucosidase activity of bifidobacteria and the hydrolysis of isoflavone glycosides by *Bifidobacterium* sp. Int-57 in soymilk fermentation. *J Microbiol Biotechnol* 2002;12:8–13.
- [25] Nunoura N, Ohdan K, Yano T, Yamamoto K, Kumagai H. Purification and characterization of β -D-glucosidase (β -D-fucosidase) from *Bifidobacterium breve* clb acclimated to cellobiose. *Biosci Biotechnol Biochem* 1996;60:188–93. <https://doi.org/10.1271/bbb.60.188>.
- [26] Pokusaeva K, O'Connell-Motherway M, Zomer A, Fitzgerald GF, van Sinderen D. Characterization of two novel α -glucosidases from *Bifidobacterium breve* UCC2003. *Appl Environ Microbiol* 2009;75:1135–43. <https://doi.org/10.1128/AEM.02391-08>.
- [27] Braune A, Blaut M. Bacterial species involved in the conversion of dietary flavonoids in the human gut. *Gut Microbes* 2016;7:216–34. <https://doi.org/10.1080/19490976.2016.1158395>.
- [28] Donkor ON, Shah NP. Production of β -glucosidase and hydrolysis of isoflavone phytoestrogens by *Lactobacillus acidophilus*, *Bifidobacterium lactis*, and *Lactobacillus casei* in soymilk. *J Food Sci* 2008;73:15–20. <https://doi.org/10.1111/j.1750-3841.2007.00547.x>.